

Structural Polymorphism of Alzheimer's β -Amyloid Fibrils as Controlled by an E22 Switch: A Solid-State NMR Study

Matthew R. Elkins,^{†,||} Tuo Wang,^{†,||} Mimi Nick,[‡] Hyunil Jo,[‡] Thomas Lemmin,[‡] Stanley B. Prusiner,[§] William F. DeGrado,[‡] Jan Stöhr,[§] and Mei Hong^{*,†}


[†]Department of Chemistry, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, United States

[‡]Department of Pharmaceutical Chemistry, University of California, San Francisco, San Francisco, California 94158, United States

[§]Institute for Neurodegenerative Diseases, Department of Neurology, University of California, San Francisco, San Francisco, California 94143, United States

Supporting Information

ABSTRACT: The amyloid- β ($A\beta$) peptide of Alzheimer's disease (AD) forms polymorphic fibrils on the micrometer and molecular scales. Various fibril growth conditions have been identified to cause polymorphism, but the intrinsic amino acid sequence basis for this polymorphism has been unclear. Several single-site mutations in the center of the $A\beta$ sequence cause different disease phenotypes and fibrillization properties. The E22G (Arctic) mutant is found in familial AD and forms protofibrils more rapidly than wild-type $A\beta$. Here, we use solid-state NMR spectroscopy to investigate the structure, dynamics, hydration and morphology of Arctic E22G $A\beta$ 40 fibrils. ^{13}C , ^{15}N -labeled synthetic E22G $A\beta$ 40 peptides are studied and compared with wild-type and Osaka E22 Δ $A\beta$ 40 fibrils. Under the same fibrillization conditions, Arctic $A\beta$ 40 exhibits a high degree of polymorphism, showing at least four sets of NMR chemical shifts for various residues, while the Osaka and wild-type $A\beta$ 40 fibrils show a single or a predominant set of chemical shifts. Thus, structural polymorphism is intrinsic to the Arctic E22G $A\beta$ 40 sequence. Chemical shifts and inter-residue contacts obtained from 2D correlation spectra indicate that one of the major Arctic conformers has surprisingly high structural similarity with wild-type $A\beta$ 42. ^{13}C - ^1H dipolar order parameters, ^1H rotating-frame spin-lattice relaxation times and water-to-protein spin diffusion experiments reveal substantial differences in the dynamics and hydration of Arctic, Osaka and wild-type $A\beta$ 40 fibrils. Together, these results strongly suggest that electrostatic interactions in the center of the $A\beta$ peptide sequence play a crucial role in the three-dimensional fold of the fibrils, and by inference, fibril-induced neuronal toxicity and AD pathogenesis.



INTRODUCTION

Insoluble amyloid fibrils formed by a variety of proteins underlie many neurodegenerative diseases, principal among which are Alzheimer's disease (AD), Parkinson's disease, Huntington's disease, and Creutzfeldt-Jakob disease.^{1,2} Elucidating the molecular structures and oligomeric assemblies of amyloid proteins³⁻¹⁰ is the first step in designing therapeutic compounds against these diseases. For AD, two central events in its pathogenesis are the aggregation of two proteins in the brain: the small amyloid- β ($A\beta$) peptide of 37-43 residues that forms extracellular plaques and the hyperphosphorylated tau protein that forms intracellular neurofibrillary tangles. Electron microscopy (EM) and solid-state NMR (SSNMR) spectroscopy have been used extensively to characterize $A\beta$ fibrils,¹¹⁻¹⁸ and show that $A\beta$ structures are highly polymorphic on the micron scale and the molecular scale.^{19,20} On the micron scale, $A\beta$ aggregates can appear as striated ribbons, twisted fibrils, or spherical aggregates, and have variable periodicity of twists and variable widths. On the nanometer scale, SSNMR structures of wild-type (WT) $A\beta$ 40 fibrils show a common strand-loop-

strand motif that forms hydrogen bonds in a parallel in-register fashion²¹ along the fibril axis to form a cross- β structure. However, each repeat motif of the cross- β structure can contain two or three molecules with 2- or 3-fold rotational symmetries along the fibril axis, whose intermolecular interfaces differ significantly.^{12,13} Local torsion angles show moderate variations among different structures, and the registry between the two β -strands of each peptide can shift.²² Further, a brain-derived and seeded $A\beta$ 40 sample exhibits a 3-fold symmetric structure that differs from the in vitro 3-fold structure in the β -strand length and subunit intermolecular packing.¹⁶ These structural variations have so far been largely attributed to fibril growth conditions such as shear forces, pH, temperature and peptide concentration.²³⁻²⁵

While fibril growth conditions can affect $A\beta$ structure, recent experimental data indicate that the amino acid sequence also plays a significant role in $A\beta$ polymorphism, with single- or

Received: April 11, 2016

Published: July 14, 2016

double-residue changes able to dramatically alter the fibril structure. An atomic structural model of A β 42, which is less abundant than A β 40 but has higher toxicity and morphological heterogeneity^{26,27} has recently been reported.¹⁵ With only two extra residues, Ile41 and Ala42, the protein adopts a completely different, S-shaped triple parallel β -sheet structure,^{15,28} stabilized by an intramolecular K28-A42 salt bridge. The chemical shifts of recombinant A β 42 support the distinct β -strand distributions in the A β 42 and A β 40 sequences.²⁹ Another example of the sequence dependence of A β fibril structures comes from a high-resolution structure of the E22 deletion (E22 Δ) mutant, Osaka, of A β 40.¹⁴ A large number of distance and chemical shift constraints showed that this mutant adopts an intricate 2-fold symmetric “cinnamon roll”-like structure that is stabilized by an intermolecular E3-K28 salt bridge.^{14,30} Finally, a D23N mutant of A β 40 was found to adopt both parallel and antiparallel packing, in contrast to the parallel-only structures of wild-type A β 40 fibrils.^{31,32}

The A β structural diversity opens the important questions of whether this polymorphism correlates with and even causes differences in neuronal toxicities, binding stoichiometries of imaging agents, and AD symptoms,^{16,25,33} and what the intrinsic molecular reasons are beyond fibril growth conditions for this polymorphism. Insights into these questions may be gained by comparisons of mutant A β fibrils with each other and with the wild-type fibrils. Many single-site mutations exist within the 43 residues of the A β sequence that are related to familial and early onset AD.³⁴ Interestingly, most of these mutations are clustered in the center of the A β sequence, away from the secretase cleavage sites, in a short segment of A21, E22 and D23. These are Flemish A21G,³⁵ Osaka E22 Δ ,³⁶ Italian E22K,³⁷ Arctic E22G,³⁸ Dutch E22Q,³⁹ and Iowa D23N.⁴⁰ Among these mutants, the Arctic E22G A β shows fast protofibril formation in vitro and in vivo, and causes fibril morphologies distinct from those of wild-type A β 40.^{18,38,41–44} Given the highly charged nature of this segment and the occurrence of multiple charge-altering mutations, we wonder whether electrostatic interactions in this non- β region play an important role in shaping the three-dimensional fold of A β .

In this study, we combine magic-angle-spinning (MAS) SSNMR with EM, fluorescence spectroscopy, and structural stability measurements to characterize the structure, dynamics, and fibrillization properties of Arctic A β 40 fibrils. Synthetic site-specifically labeled E22G A β 40 is examined using 2D correlation SSNMR techniques. To determine whether the observed conformations are intrinsic to the amino acid sequence or dominated by fibrillization conditions, we also measured the spectra of wild-type and Osaka E22 Δ A β 40 fibrils. Under the same fibrillization conditions, we found that Arctic A β 40 exhibits at least four conformations while Osaka and wild-type A β 40 adopt a single or a predominant conformation. Moreover, the chemical shifts of a major Arctic conformer bear close resemblance to reported A β 42 chemical shifts,^{15,28,29} suggesting similarities in the β -strand arrangements of these two fibrils. ¹³C–¹H dipolar order parameters, ¹H relaxation times, and water-to-protein ¹H spin diffusion experiments indicate differences in the mobility and hydration of the three fibrils. These results provide insights into how electrostatic interaction in the center of the A β peptide dictates the three-dimensional fold¹⁴⁵ and consequently the fibrillization properties of this peptide.

MATERIALS AND METHODS

Synthesis and Purification of A β (1–40) Peptides. The peptides were synthesized using a Biotage Initiator+ Alstra automated microwave peptide synthesizer on preloaded HMPB-Chemmatrix resin (0.1 mmol scale). Fmoc deprotection was performed at 70 °C using 0.1 M HOBt in 4-methylpiperidine-DMF (1:4) for 5 min. The resin was then washed with DMF (3 \times). Coupling reactions were performed using *N* α -Fmoc amino acids (5 equiv, 0.5 M), HCTU (4.95 equiv, 0.5 M) and DIEA (10 equiv, 0.5 M) in DMF. All couplings were doubly performed for 5 min at 75 °C except for His (25 min double coupling at RT) and Arg residues (5 min triple coupling at 75 °C). After completion of synthesis, the resin was successively washed with DMF (3 \times) and DCM (3 \times) and dried. For isotope labeled residues, the first coupling was performed using the isotopically labeled *N* α -Fmoc amino acids (1.5 equiv, 0.5 M), HCTU (1.45 equiv, 0.5 M) and DIEA (3 equiv, 0.5 M) in DMF, followed by the second coupling using unlabeled *N* α -Fmoc amino acids (3 equiv, 0.5 M), HCTU (3 equiv, 0.5 M) and DIEA (6 equiv, 0.5 M) in DMF. Peptide cleavage was performed using TFA-DTT-H₂O (95:2.5:2.5) for 3 h at RT. The volatiles were removed by a stream of nitrogen gas and the crude peptide was obtained by precipitation with cold diethyl ether. The crude peptide was dissolved in HFIP and purified by RP-HPLC (Figure S1). A portion of the crude solution was injected into the heated (60 °C) Vydac 300 Å C4 column (214TP1022, 10 μ m, 22 \times 250 mm) running a flow of 10 mL/min with a linear gradient from 15% to 40% of buffer B over 25 min (buffer A: 0.1% TFA in H₂O; buffer B: 0.1% TFA in 99% CH₃CN in H₂O). Peptide mass was analyzed using MALDI (Shimadzu Axima) and electrospray ionization mass spectrometry (Qtrap 3200, ABSCI EX).

Fibrillization. Lyophilized wild-type, E22G, E22 Δ A β 40 peptides were dissolved at 5 mg/mL in HFIP and evaporated into a peptide film. The peptide films were first suspended at 10 mg/mL in DMSO, briefly vortexed and sonicated for 5 min, then diluted to 0.2 mg/mL in 10 mM sodium phosphate buffer, pH 7.2. The solution was shaken at 900 rpm at 37 °C for at least 72 h to form fibrils. This fibrillization method is consistent with our in vivo biological experiments on Arctic A β .^{46,47} To investigate the effect of the solution condition on fibril structure, and to enable comparison with literature results, we also prepared the Osaka and Arctic fibrils using a second method, in which the peptide films were first suspended to 2 mg/mL in 10 mM NaOH and then quickly diluted to 0.2 mg/mL in the dilution buffer, whose final composition is 10 mM H₃PO₄–NaOH, pH 7.4, and 100 mM NaCl.^{14,48} We denote the two Osaka samples prepared in this way as Osaka DMSO and Osaka NaOH. All SSNMR spectra shown here for Arctic and wild-type A β 40 are from the DMSO preparation, but similar data for the NaOH preparation were also acquired.

Electron Microscopy. Wild-type, Arctic, and Osaka A β 40 fibrils (0.2 mg/mL) were adsorbed onto freshly glow-discharged, 200-mesh Formvar/carbon-coated copper grids (Ted Pella) and negative-stained with a 2% (w/v) uranyl acetate solution. Images were taken on a Philips/FEI Tecnai F20 electron microscope at 80 kV.

Thioflavin-T Fluorescence Monitoring of Fibril Formation Kinetics. Thioflavin-T (10 μ M) was added to a freshly prepared solution of wild-type, Arctic, and Osaka A β 40 (0.2 mg/mL) in 10 mM sodium phosphate buffer, pH 7.2. Fluorescence measurements at an excitation wavelength of 485 nm and an emission wavelength of 538 nm were taken every 5 min in a SpectraMax M5 (Molecular Devices) plate reader set to 37 °C with agitation (automix linear shaking) between readings.

Guanidine Hydrochloride Denaturation. One volume of A β fibril sample (0.2 mg/mL) was added to three volumes of concentrated guanidine hydrochloride (GdnHCl) to obtain the desired final concentration (1–6 M GdnHCl) and incubated for 2 h at 22 °C. The fibrils were then centrifuged at 100 000g for 1 h, rinsed with PBS, and ultracentrifuged again. The pellet was resuspended in 4 \times LDS sample buffer, diluted, and then resolved by SDS-PAGE and stained with Coomassie dyes.

Solid-State NMR Spectroscopy. All fibrils were slowly dried in a desiccator to 60–75% water by mass. About 40 mg of these hydrated




Figure 1. Fibrillization of wild-type, Arctic and Osaka $A\beta_{40}$ peptides. (a) Osaka and Arctic $A\beta_{40}$ peptides form fibrils much more rapidly than the wild-type (WT) peptide. Kinetics of fibril formation (0.2 mg/mL in 10 mM sodium phosphate buffer, pH 7.2) was measured by the amyloid-binding dye ThT (10 μ M). $n = 5$, mean \pm SEM. (b) Negative stain TEM images of WT and mutant $A\beta_{40}$ fibrils. Arctic $A\beta_{40}$ fibrils display significant twists with variable periodicities and more pronounced entanglement than wild-type and Osaka fibrils. Scale bar = 100 nm. (c) Mutant fibrils have altered stability to denaturation by GdnHCl. Fibrils were incubated with increasing concentrations (1–6 M) of GdnHCl and then ultracentrifuged at 100 000g to pellet any fibrils that resisted denaturation, which were resolved by SDS-PAGE and Coomassie stain.

fibrils were spun into 3.2 mm MAS rotors. Solid-state NMR experiments were conducted on a Bruker Avance II 800 MHz (18.8 T) spectrometer and an Avance 900 MHz (21.1 T) spectrometer using triple-resonance HCN probes. Typical radiofrequency (rf) field strengths were 63–83 kHz for ^1H , 50–63 kHz for ^{13}C , and 36–45 kHz for ^{15}N . ^{13}C chemical shifts were externally referenced to the methylene signal of adamantane at 40.48 ppm on the DSS-scale. ^{15}N chemical shifts were referenced to the methionine amide resonance (127.88 ppm) of the model peptide *N*-formyl-Met-Leu-Phe-OH (formyl-MLF)^{49,50} on the liquid ammonia scale.

One-dimensional (1D) ^{13}C and ^{15}N cross-polarization (CP) spectra were measured on 800 and 900 MHz NMR spectrometers under 10.5, 12, and 16 kHz MAS. ^1H – ^{13}C and ^1H – ^{15}N contact times were 1 and 2 ms, respectively. The spectra were measured at 273 and 298 K. Two-dimensional (2D) ^{13}C – ^{13}C correlation spectra were measured with 50 ms dipolar-assisted rotational resonance (DARR) mixing⁵¹ for resonance assignment and 1.5 s ^1H -driven spin diffusion (PDS) for detecting long-range correlations. 2D ^{15}N – ^{13}C heteronuclear correlation (HETCOR) spectra were measured in two ways: a rotational echo double resonance (REDOR) transfer with a mixing time of 0.8–1.0 ms allowed the observation of one-bond ^{15}N – $^{13}\text{C}\alpha$ cross peaks,^{49,52} while 2D N(CA)CX⁵³ spectra allowed the detection of ^{15}N to side chain ^{13}C cross peaks. The N(CA)CX experiment used a 2 ms ^1H – ^{15}N CP contact time, a 7 ms ^{15}N – ^{13}C CP contact time, and a 30–50 ms ^{13}C DARR mixing period. The ^{15}N – ^{13}C CP used ^{15}N and ^{13}C spin-lock fields of 27.8 kHz and 38.3 kHz, respectively, and the ^{13}C carrier frequency was set at 55 ppm. ^1H decoupling field strengths were 93 kHz during ^{15}N – ^{13}C CP and 83 kHz during acquisition.

To measure $A\beta_{40}$ hydration, we conducted a water-to-protein ^1H spin diffusion experiment.^{54–58} Following ^1H excitation, a ^1H T_2 filter of 0.6–1.2 ms was applied to suppress the ^1H magnetization of the rigid protein while retaining 80–90% of the ^1H magnetization of mobile water. A ^1H mixing time was then applied to transfer the remaining water ^1H polarization to the protein and then to protein ^{13}C

via CP for detection. The spin diffusion mixing times were varied from 0 to 196 ms to obtain a magnetization transfer buildup curve. The experiment was conducted at 277 K. The 1D experiment was also extended to 2D by adding a ^{13}C evolution period and a 50 ms DARR mixing period to resolve the peptide ^{13}C signals.⁵⁹ For the water-edited 2D experiment, the ^1H mixing time was fixed to 4 ms to detect only well hydrated protein residues.

To investigate peptide dynamics, we measured ^{13}C – ^1H dipolar couplings using the dipolar-chemical shift correlation (DIPSHIFT) experiment.⁶⁰ The samples were spun at 7 kHz at 298 K, and a frequency-switched Lee–Goldburg (FSLG) sequence⁶¹ was applied with a transverse field strength of 80 kHz to remove the ^1H – ^1H dipolar coupling during the evolution period. The FSLG scaling factor was measured on formyl-MLF to be 0.577 for the Osaka and wild-type $A\beta_{40}$ samples and 0.590 for the Arctic $A\beta_{40}$ sample. Apparent ^{13}C – ^1H dipolar couplings were obtained by fitting the t_1 -dependent intensities, then divided by the scaling factor to obtain the true couplings. Order parameters were calculated by dividing the true couplings with the rigid-limit value of 22.7 kHz. DIPSHIFT time curves were simulated using a FORTRAN program,^{62,63} available online at meihonglab.mit.edu/software. An empirical ^1H T_2 relaxation factor was applied to the best-fit curves to reproduce the observed asymmetry in the dipolar oscillation.

^1H rotating-frame spin–lattice ($T_{1\rho}$) relaxation times, which are indicative of microsecond molecular motions,⁶⁴ were measured with ^{13}C detection at 298 and 260 K under 10.5 kHz MAS using a Lee–Goldburg (LG) spin-lock pulse sequence.⁶⁵ The spin lock time varied from 0 to 50 ms, followed by a 0.3 ms Hartman–Hahn CP period. The tilted effective ^1H LG spin-lock field was 62.5 kHz, thus the characteristic time scale probed by this experiment is 16 μ s.

To compare the chemical shifts of the three $A\beta_{40}$ fibrils, we calculated the chemical shift differences for each residue using the following equation:

$$\Delta\delta_{\xi,\kappa} = \sqrt{\frac{1}{n} \sum_{C_i} (\delta_{C_i}^{\xi} - \delta_{C_i}^{\kappa})^2 + \frac{1}{2.5} (\delta_N^{\xi} - \delta_N^{\kappa})^2} \quad (1)$$

where ξ and κ denote different fibrils, n is the number of ^{13}C chemical shifts available for each residue, and the factor of 2.5 scales the ^{15}N chemical shift relative to the ^{13}C chemical shifts due to their relative gyromagnetic ratios. All ^{13}C chemical shifts are reported on the DSS scale (Table S1, S2). Literature values for chemical shifts referenced to TMS were converted to DSS-referenced values by adding 2.0 ppm.

To evaluate the pairwise similarities for the 11 $A\beta$ fibrils, we conducted a correlation analysis by calculating the Pearson product-moment correlation coefficient. The chemical shifts of four atoms, the backbone amide nitrogen, C' , $C\alpha$, $C\beta$, are included. For each strain, the chemical shifts of identical atoms were combined to form a data set. The correlation coefficients between equivalent data sets were then averaged for each pair of strains.

RESULTS

Comparison of $A\beta$ 40 Fibril Formation Kinetics, Fibril Stability and Morphology. WT and mutant $A\beta$ 40 peptides were assembled into fibrils by incubating at 37 °C with constant agitation. Shaking both accelerates fibril formation by fragmenting the fibrils and exposing new ends for the addition of monomers and promotes the formation of equilibrium structures.³² Fibrillization kinetics were monitored using the amyloid-binding dye Thioflavin T (ThT) (Figure 1a). The three peptides show distinct fibril formation rates. The E22G and E22Δ $A\beta$ 40 prepared in DMSO formed at much faster rates than the wild-type peptide. In particular, the Osaka peptide fibrillized on the second to minute time scale and was complete by the first time point of the measurement, which was conducted within 2–3 min of sample preparation. This is manifested by the high initial ThT fluorescence value and the rapid disappearance of the peptide signals in the ^1H solution NMR spectra (Figure S2). In comparison, wild-type $A\beta$ 40 showed a pronounced lag phase and fibrillized in ~20 h with greater kinetic stochasticity. For each sample the final increase in fluorescence intensity was significantly different, indicating different extents of binding and/or the environment of the bound ThT dye molecules. The Arctic E22G mutant fibrillized more slowly than the Osaka peptide, but still more rapidly than WT. These observations are in good agreement with literature reports of the kinetics of Arctic $A\beta$ fibrils^{38,66–68}

We usually first dissolved the peptide in DMSO to prepare fibrils because this method produced fibrils with biological activity in mouse inoculation.⁶⁹ However, we also wanted to prepare the Osaka peptide by dissolving in NaOH to match the published SSNMR structure.¹⁴ The use of NaOH for initial peptide solubilization moderately slowed down fibril formation, but the rates are still much faster than wild-type $A\beta$ 40. There was also a large change in the final intensity between the DMSO and NaOH treated samples; because the peptide sequence is identical in both experiments, we conclude that the two methods give rise to distinct fibril conformations and/or morphologies. Negatively stained TEM images (Figure 1b) show that the wild-type and Osaka fibrils are relatively straight, while the Arctic fibrils display significant twists with a distribution of periodicities and are more entangled than the wild-type and Osaka fibrils.

The mutant $A\beta$ 40 fibrils not only exhibit different fibrillization kinetics and morphologies from the wild-type fibrils, but also have different thermodynamic stabilities. To investigate the fibril stabilities, we incubated the fibrils with increasing concentrations of guanidine hydrochloride and

quantified the amount of insoluble fibrils remaining after a 2 h equilibration time (Figure 1c). We take the midpoint of the solubilization curve as a proxy for relative thermodynamic stability. Arctic E22G fibrils with varying concentrations of GdnHCl showed fibril loss at lower concentrations of GdnHCl than wild-type fibrils. In contrast, the Osaka E22Δ fibrils are more stable against GdnHCl solubilization than the wild-type peptide. The increase in stability also depended on the method of solubilization.

Together, these experiments indicate that the Arctic E22G mutant forms fibrils with a lower activation energy than the WT peptide and adopts a final solid-state structure with lower stability than WT. In comparison, under the same DMSO-solubilized initial conditions, the Osaka mutant forms fibrils with a lower activation energy than WT, but achieves a final conformation that has greater stability than WT. Moreover, both the kinetics and the thermodynamics of the fibril formation process depend on the initial state of the peptide since NaOH-solubilized Osaka peptide formed fibrils more slowly than DMSO-solubilized peptide, and it achieved a final state of greater thermodynamic stability. Finally, the differences in fluorescence intensity of the final states suggest that each sample achieved a different final state. To determine the structural bases for these differences we next turned to SSNMR.

^{13}C and ^{15}N Chemical Shifts of E22G, E22Δ and wild-type $A\beta$ 40 fibrils. 1D ^{13}C and ^{15}N CP-MAS spectra (Figure 2) give qualitative information about the structural homoge-




Figure 2. (a) Amino acid sequences of the three $A\beta$ 40 peptides. Isotopically labeled residues are shown in red. (b,c) 1D ^{13}C (b) and ^{15}N (c) CP-MAS spectra of Arctic, Osaka, and wild-type $A\beta$ 40 fibrils. Spectra were measured on 800 and 900 MHz spectrometers at 273 K under 16 kHz MAS. Assignments obtained from 2D spectra are indicated. All three fibrils show narrow intrinsic line widths, but the number of peaks and the chemical shifts differ significantly.

neity and similarity of the three $A\beta$ 40 fibrils. All samples show sharp ^{13}C and ^{15}N signals, with the Osaka DMSO fibrils having the most resolved spectra. The Arctic spectra have similar intrinsic line widths as the wild-type and Osaka spectra, as seen by resolved side chain signals, but contain more peaks, thus giving more overlapped $C\alpha$ and CO regions. The ^{15}N spectrum of Arctic $A\beta$ 40 shows multiple glycine resonances while the Osaka and wild-type $A\beta$ 40 exhibit only one G38 signal. The spectral overlap is the smallest for Osaka $A\beta$ 40 and the largest




Figure 3. 2D correlation spectra of Arctic $A\beta_{40}$ (E22G) fibrils. (top) 2D ^{13}C – ^{13}C DARR spectrum with 50 ms mixing. For simplicity, assignments are given with lower-case letters for various carbons, and F19 aromatic carbons are designated by “r”. ssb denotes spinning sidebands. (bottom) 2D N(CA)CX ^{15}N – ^{13}C correlation spectrum. Two to four sets of chemical shifts (denoted by subscripts I–IV) are identified for all labeled residues, indicating extensive structural polymorphism. Ellipses guide the eye for chemical shift multiplicity. The DARR spectrum was processed using a QSINE window function with SSB = 2.5, and were plotted with Topspin parameters lev0 = 4, toplev = 90 and nlev = 16. 2D N(CA)CX spectrum was processed using a Gaussian window function with LB = –10 Hz and GB = 0.05, and was plotted using lev0 = 5, toplev = 80 and nlev = 16.

for Arctic $A\beta_{40}$. Spectra measured from 255 to 298 K showed little differences in intensities and line widths, indicating that all fibrils are mostly immobilized at room temperature (data not shown).

2D ^{13}C – ^{13}C DARR and ^{15}N – ^{13}C correlation spectra fully resolved and permitted the assignment of the chemical shifts of the labeled peptides, and revealed extensive polymorphism of the E22G $A\beta_{40}$ fibrils: multiple resonances for each labeled residue are seen in both the DARR and N(CA)CX spectra (Figure 3). F19 exhibits two sets of chemical shifts, A21 and V36 exhibit three, while I32, L34 and G38 have four sets of chemical shifts (Table S1). The ^{13}C chemical shifts vary by up to 3 ppm for each site while ^{15}N chemical shifts vary by as little as 1 ppm for F19 and as much as 20 ppm for L34. The chemical shift differences are much larger than the ^{13}C line widths of 0.6–0.9 ppm and the ^{15}N line widths of less than 2 ppm. The concentration ratio of the four conformers is about 3:3:2:2 and is similar for all residues, as estimated from the integrated peak intensities in the 50 ms DARR spectrum, thus even the two minor conformers are not negligible in concentration. We denote the two sets of chemical shifts with the highest intensities as Arctic I and II. No inter-residue cross peaks were obtained to relate the multiple conformations of one residue to

another, and we assume that the conformer I chemical shifts in each residue belong to the same molecule, and all conformer II chemical shifts belong together to another molecule.

To determine whether the E22G $A\beta_{40}$ polymorphism is intrinsic to this amino acid sequence or due to suboptimal fibrillization conditions, we measured 2D spectra of wild-type and E22 Δ $A\beta_{40}$ fibrils prepared using the same protocol (Figure 4, Figure S3). In dramatic contrast to the E22G mutant, the E22 Δ spectra show a single set of cross peaks, while the wild-type spectra show predominantly one species, with one or two additional conformers observed for residues I32, L34 and V36. The major conformer of wild-type $A\beta_{40}$ represents 50–70% of the whole population. The Osaka and wild-type spectra are therefore much better resolved than the E22G $A\beta_{40}$ spectra. Arctic and wild-type $A\beta_{40}$ prepared in DMSO and NaOH solutions showed no spectral differences (Figure S4), indicating that their conformations are independent of the solvent. Moreover, ^{13}C and ^{15}N spectra (Figure S4a,b) of independently prepared batches of Arctic $A\beta_{40}$ fibrils are indistinguishable, demonstrating that the structural polymorphism is reproducible. For Osaka $A\beta_{40}$, initial peptide solubilization in NaOH before fibrillization in buffer resulted in observable chemical shift differences from the DMSO




Figure 4. Comparison of the chemical shifts of the three $\text{A}\beta 40$ fibrils. (a–c) 2D ^{13}C – ^{13}C DARR spectra with 50 ms mixing for (a) Arctic $\text{A}\beta 40$ (E22G), (b) wild-type $\text{A}\beta 40$, and (c) Osaka $\text{A}\beta 40$ (E22 Δ). (d–f) 2D ^{15}N – ^{13}C correlation spectra for (d) Arctic $\text{A}\beta 40$ (E22G), (e) wild-type $\text{A}\beta 40$, and (f) Osaka $\text{A}\beta 40$ (E22 Δ). A single set of chemical shifts is observed for Osaka $\text{A}\beta 40$, the wild-type fibril shows modest polymorphism from I32 to V36, while the Arctic $\text{A}\beta 40$ fibril has the largest number of polymorphs. All DARR spectra were processed using QSINE window functions with SSB = 2.5 and plotted using lev0 = 4, toplev = 90 and nlev = 16. 2D ^{15}N – ^{13}C correlation spectra were processed using LB = –10 Hz and GB = 0.05, and were plotted using lev0 = 5, toplev = 90 and nlev = 12.

solubilized sample and a modest degree of chemical shift heterogeneity (Figure S3d, Figure S5). Taken together, these results indicate that the pronounced structural polymorphism of Arctic E22G $\text{A}\beta 40$ is an intrinsic property of this amino acid sequence rather than an artifact of fibril growth conditions.

The ^{13}C and ^{15}N chemical shifts vary substantially among the three fibrils studied here and also differ to varying degrees from literature chemical shifts of other $\text{A}\beta$ fibrils. Our Osaka DMSO chemical shifts are in excellent agreement with those reported by Schütz and co-workers,¹⁴ while our wild-type $\text{A}\beta 40$ chemical shifts have moderately good agreement with the 2-fold and 3-fold in vitro wild-type fibrils.¹⁹ In Arctic $\text{A}\beta 40$, I32 and V36 have characteristic β -strand chemical shifts, A21 and L34 show typical random coil chemical shifts, while F19 and G38 exhibit mixed strand and coil chemical shifts. Interestingly, the chemical shifts of the two major Arctic conformers show unexpected agreement with $\text{A}\beta 42$ chemical shifts. For example, the A21 $\text{C}\alpha$ and ^{15}N chemical shift of Arctic I and II are within 1–2 ppm of the chemical shifts of $\text{A}\beta 42$ fibrils^{15,29} (Table S2). The Arctic conformer I L34 $\text{C}\delta 2$ chemical shift of 20.9 ppm matches the reported $\text{A}\beta 42$ values but is more than 2 ppm upfield of the Osaka and wild-type $\text{A}\beta 40$ chemical shifts.²⁹ The Arctic conformer I L34 ^{15}N chemical shift (111.5 ppm) matches that of $\text{A}\beta 42$ but is 12 ppm upfield of the ^{15}N chemical shifts of $\text{A}\beta 40$ fibrils.¹⁵ Arctic G38 chemical shifts show the largest dispersion, 3–5 ppm, among all labeled residues. When we compared the G38 ^{15}N and ^{13}C chemical shifts, which are sensitive to both local conformation and electrostatic effects, to literature values, we find that Arctic conformer II resembles the brain-derived $\text{A}\beta 40$,¹⁶ while no other conformers agree well with any $\text{A}\beta$ peptides.

To obtain quantitative measures of fibril structural similarities, we calculated per-residue average chemical shift differences between the Arctic I conformer and various published $\text{A}\beta$ structures, and between our wild-type $\text{A}\beta 40$ and reported $\text{A}\beta$ structures (Figure 5a,b). The comparison

quantifies the similarity between Arctic $\text{A}\beta 40$ and wild-type $\text{A}\beta 42$, with differences of less than 0.6–1.0 ppm at F19, A21, I32, and L34.²⁹ Near the C terminus, at V36 and G38, the chemical shift differences increase, as expected since the additional two residues in $\text{A}\beta 42$ form part of an extra β -strand that cannot exist in $\text{A}\beta 40$ fibrils.¹⁵ The chemical shifts of our wild-type $\text{A}\beta 40$ fibrils are similar to those of both 2-fold and 3-fold symmetric in vitro WT fibrils, with slightly better agreement with the 3-fold fibrils, principally at I32.^{12,13} Our Osaka DMSO and Osaka NaOH fibrils exhibit non-negligible chemical shift differences, but the Osaka DMSO chemical shifts are identical within experimental uncertainty to the chemical shifts of recombinant Osaka fibrils.¹⁴ To compare all the $\text{A}\beta$ fibrils reported in the literature and the samples studied here, we computed the Pearson product-moment correlation coefficient (Figure 5e, Figure S6). A total of 11 $\text{A}\beta$ fibril samples were analyzed, and four chemical shifts, ^{15}N , C' , $\text{C}\alpha$, and $\text{C}\beta$, were included in the correlation coefficient calculation. The 2D heat maps confirm that Arctic I has the highest correlation with WT $\text{A}\beta 42$, and our Osaka DMSO data has excellent correlation with the literature Osaka structure. The 2D correlation map also revealed that the WT $\text{A}\beta 40$ sample in this study has significant similarity to the Osaka NaOH sample.

Mobility of $\text{A}\beta 40$ Fibrils. To obtain information about the amplitude and rates of motion of the $\text{A}\beta 40$ fibrils, we measured ^{13}C – ^1H dipolar order parameters (S_{CH}) and NMR relaxation times. Representative ^{13}C – ^1H dipolar dephasing data from the 2D DIPSHIFT spectra are shown in Figure 6a. The backbone $\text{C}\alpha$ – $\text{H}\alpha$ dipolar order parameters are at least 0.90, indicating that fast motions are largely absent, while side chain order parameters range from 0.24 to 0.50 (Figure S7). Interestingly, Osaka DMSO and wild-type $\text{A}\beta 40$ exhibit similar order parameters that are smaller than the Arctic $\text{A}\beta 40$ S_{CH} values for both the backbone and side chains. However, the Arctic sample exhibits larger intensity asymmetry in the dipolar




Figure 5. Per-residue average chemical shift differences between different $A\beta$ fibrils, calculated using eq 1. ^{13}C chemical shifts were referenced to DSS for the calculation. (a) Chemical shift differences of Arctic conformer I from published wild-type $A\beta$ 40 and $A\beta$ 42 data. (b) Chemical shift differences of wild-type $A\beta$ 40 studied here from published wild-type $A\beta$ 40 and $A\beta$ 42 data. (c) Chemical shift differences between Osaka DMSO and Osaka NaOH studied here. (d) Chemical shift differences of Osaka DMSO from recombinant Osaka $A\beta$ 40. The very small differences indicate that the synthetic Osaka fibrils have the same structure as fibrils obtained from recombinant peptide.¹⁴ (e) 2D heat map representing the average pairwise Pearson product-moment correlation coefficients between the chemical shifts of $A\beta$ fibrils. Eleven different $A\beta$ fibrils, including five studied here and six literature cases, are compared. The white cross in each row indicates the strain with the highest correlation coefficient. Arctic conformer I has the highest correlation with WT-42 2MXU. PDB codes are given where available to simplify notation. WT-42 corresponds to the study of Colvin et al.,²⁹ Osaka 2MVX corresponds to the study of Schütz et al.,⁷⁰ WT 2M4J is the study of Lu et al.,¹⁶ WT 2LMP/Q is that of Paravastu et al.,¹² and WT 2LMN/O is that of Petkova et al.¹³

oscillation, indicating more pronounced microsecond time scale motion.^{71,72}

^{13}C -resolved ^1H $T_{1\rho}$ relaxation times reveal further differences in the microsecond motions of the three fibrils. At 298 K,




Figure 6. Mobilities of A β 40 fibrils. (a) Representative ^{13}C - ^1H dipolar dephasing curves for Arctic, wild-type and Osaka DMSO A β 40 fibrils. Best-fit couplings and the resulting order parameters (S_{CH}) are given in each panel. All fibrils show mostly immobilized backbones but significant side chain motions. The data were obtained at 298 K under 7 kHz MAS. (b) Representative ^{13}C -detected ^1H $T_{1\rho}$ relaxation data of the three A β 40 fibrils. Top row: Arctic E22G A β 40 shows longer $T_{1\rho}$ than wild-type and Osaka A β 40 at 298 K. Second to fourth rows: temperature dependence of the ^1H $T_{1\rho}$. Arctic A β 40 shows shorter $T_{1\rho}$ at lower temperature, while wild-type and Osaka A β 40 exhibit longer $T_{1\rho}$'s at lower temperature. These indicate that the Arctic A β 40 side chains have faster microsecond motions than the other two fibrils.

Arctic E22G A β 40 shows the longest ^1H $T_{1\rho}$'s while wild-type A β 40 has the shortest relaxation times (Figure 6b, Table S3). To determine whether these relaxation rates reflect motions faster or slower than the ~ 16 μs characteristic time scale of our $T_{1\rho}$ experiment, we measured the temperature dependence of $T_{1\rho}$. Interestingly, decreasing the temperature to 260 K shortened the relaxation times of Arctic A β 40 residues while lengthening those of wild-type A β 40 (Figure S8). Thus, Arctic A β 40 motion occurs on time scales faster than ~ 16 μs at ambient temperature while wild-type A β 40 motion occurs in the slow limit relative to the characteristic time. Osaka A β 40 $T_{1\rho}$ displays a mixed temperature dependence: the F19 side chain exhibits slow motional behavior while I32 and L34 side chains show fast-limit motional characteristics.

Together, these order parameters and relaxation times indicate that, while all three A β samples are relatively rigid, there are non-negligible differences in the peptide motion. Wild-type A β 40 has larger motional amplitudes than the other two peptides but much slower motions, while Arctic E22G A β 40 has the smallest amplitudes for fast motions but has the fastest motion on the microsecond time scale.

Hydration and Inter-residue Contacts of A β 40 Fibrils.

Water contact and long-range inter-residue correlations provide useful constraints for the three-dimensional structures of A β fibrils. Representative 1D and 2D ^{13}C -detected water-to-protein ^1H spin diffusion spectra are shown in Figure 7 and Figure S9.

Compared with the equilibrium spectra measured with 100 ms ^1H mixing, the Osaka fibrils have the highest initial intensities at 4 ms among the three samples, indicating that the Osaka fibrils are the most hydrated. Both the wild-type and Osaka fibrils exhibit similar spectral envelopes at 4 ms as the 100 ms spectra, indicating relatively uniform hydration for these fibrils, but the Arctic sample displays a significantly different intensity pattern at 4 ms compared to 100 ms, with the A21 signals dominating the spectrum, indicating that A21 is preferentially hydrated over other residues. Consistently, when the water-to-protein ^1H spin diffusion is measured as a function of the mixing time, the buildup curves (Figure 7d–f, Figure S5c) show that water polarization transfer to Arctic and wild-type peptides reached equilibrium at ~ 100 ms, while the Osaka A β 40 fibrils intensities have equilibrated at ~ 50 ms, indicating that the Osaka fibrils have a larger water-exposed surface area or more immobilized water molecules. Moreover, wild-type and Osaka fibrils have uniform buildup rates across the different labeled residues, with a narrow intensity distribution at 4 ms, while Arctic A β 40 fibrils exhibit a broader intensity distribution of 10–25% at 4 ms, indicating heterogeneous hydration, consistent with the conformational polymorphism of the Arctic A β fibrils.

Using a 1.5 s mixing ^{13}C - ^{13}C PDSD experiment, we looked for long-range correlations among the labeled residues. In the Arctic spectrum, the F19 side chain shows several cross peaks to the I32 side chain of conformer I but no cross peaks to any




Figure 7. Different hydration behaviors of the three $A\beta_{40}$ fibrils. (a–c) ^{13}C -detected water-to-protein ^1H spin diffusion spectra at 100 and 4 ms for (a) Arctic (E22G) $A\beta_{40}$, (b) wild-type $A\beta_{40}$, and (c) Osaka (E22 Δ) $A\beta_{40}$. The Osaka fibrils show the highest initial intensity. The Arctic initial intensity distribution deviates from the equilibrium intensity more than the wild-type peptide fibrils. (d–f) Water-to-protein ^1H spin diffusion buildup curves for (d) Arctic $A\beta_{40}$, (e) wild-type $A\beta_{40}$, and (f) Osaka $A\beta_{40}$. Vertical dashed line and shaded horizontal bars guide the eye for the magnetization equilibration times and the initial buildup rates. Osaka $A\beta_{40}$ has the shortest equilibration time, indicating the highest hydration, while Arctic $A\beta_{40}$ has the largest distribution of buildup rates.

other residues, while the Osaka spectrum shows F19 cross peaks with L34 and V36 side chains (Figure 8). No F19 cross peaks were observed for the wild-type sample. Figure 9 compares four $A\beta$ fibril structures solved using solid-state NMR, with special emphasis on the hydrophobic contacts




Figure 8. Inter-residue contacts of the $A\beta_{40}$ fibrils from 1.5 s mixing 2D ^{13}C – ^{13}C PDS spectra. In the aromatic region, Arctic $A\beta_{40}$ (E22G) shows several F19–I32 cross peaks for conformer I, while Osaka $A\beta_{40}$ (E22 Δ) shows F19–L34 and F19–V36 cross peaks. The unresolved aromatic carbons of F19 are denoted as F19r.

between F19 and the last ten residues of the peptide. The 2-fold and 3-fold wild-type $A\beta_{40}$ structures exhibit intermolecular F19–L34 contacts: the F19 ring of one peptide packs against the L34 side chain of a different peptide with a short distance of ~ 4 Å.^{12,13} A D23–K28 salt bridge is present in the 2-fold structure and the 3-fold brain-derived structure, but absent in the 3-fold in vitro structure.^{13,16,25} A subtly different 2-fold structural model²² has also been reported using fibrils formed by $A\beta$ (M1–40), which contains an exogenous N-terminal methionine. This model (not shown here) differs in the intramolecular side chain packing and intermolecular contacts between the two cross- β subunits. In the published structure of Osaka $A\beta_{40}$ fibrils derived from recombinant protein, F19 and L34 form intramolecular contacts of ~ 4 Å,^{14,73} which lock the turn from F19 to L34. Compared to these $A\beta_{40}$ structures, $A\beta_{42}$ has a distinct pattern of long F19–L34 distances and short F19–I32 distances of ~ 4 Å.¹⁵ Our observed F19 side chain–L34 $\text{C}\delta 2$ cross peaks in the Osaka PDS spectrum are fully consistent with the reported Osaka $A\beta_{40}$ structure,⁷⁰ while the absence of F19–L34 cross peaks in conjunction with the observed F19–I32 cross peaks in our Arctic PDS spectrum rules out the wild-type $A\beta_{40}$ and Osaka $A\beta_{40}$ structures as possible folds for any of the Arctic conformations. However, the observed F19–I32 cross peaks of Arctic conformer I is consistent with the $A\beta_{42}$ fold. Figure 9 also shows that the A21 methyl group faces inward toward the other strand in the two wild-type $A\beta_{40}$ structures, but points outward to water in the Osaka $A\beta_{40}$ and $A\beta_{42}$ structures. The preferential hydration of A21 observed in our Arctic sample is thus also consistent with the $A\beta_{42}$ fold.

DISCUSSION

These chemical shift data indicate that the E22G mutant of $A\beta_{40}$ is highly polymorphic, and this polymorphism is intrinsic to this particular amino acid sequence. At least four distinct sets




Figure 9. Comparison of four published structures of wild-type and mutant $A\beta$ fibrils. The side chains of residues labeled in this study are shown as red and green sticks. Key charged residues that are known to be important for the three-dimensional fold of the peptides are shown in blue. (a) 2-fold in vitro structure of wild-type $A\beta$ 40 (PDB: 2LMN).¹³ (b) 3-fold in vitro structure of wild-type $A\beta$ 40 (PDB: 2LMQ).¹² (c) Osaka $A\beta$ 40 (E22 Δ) structure.¹⁴ (d) Wild-type $A\beta$ 42 structure (PDB: 2MXU).¹⁵ The shortest F19–I32 and F19–L34 distances are indicated in the structures. Note that the distances in (a) and (b) are intermolecular, while distances in (c) and (d) are intramolecular.

of chemical shifts are resolved, with population ratios of about 3:3:2:2, thus the minor forms are not negligible. This molecular structural polymorphism is reproducible from one sample preparation to another, is unaffected by the initial solvent used in the fibrilization (Figure S4), and occurs despite shaking during fibril growth, which tends to promote self-seeding and conformational homogeneity;^{23,32} thus, the Arctic molecular structure polymorphism is robust. Under the same fibrilization condition, the E22 Δ $A\beta$ 40 fibrils, with initial solubilization in DMSO, exhibited a single set of chemical shifts that are identical within experimental uncertainty to the published Osaka chemical shifts,¹⁴ while wild-type $A\beta$ 40 is dominated by one major conformer. Therefore, the Arctic structural polymorphism is not an artifact of sample preparation but is due to the mutation of the anionic Glu to Gly at residue 22. The chemical shift heterogeneity of the Arctic fibrils is qualitatively consistent with the presence of both straight and twisted fibrils with variable twist periodicities in the TEM images (Figure 1), which in turn agrees well with a detailed TEM and AFM study by Antzutkin and co-workers, who resolved and quantitatively analyzed 4–5 types of Arctic fibril morphologies with different extents of twists and heights. They further measured the mass-length of four of the polymorphs and found them to correspond to 2, 4, and 6 peptide molecules per 0.48 nm repeat of the fibrils,⁶⁶ suggesting that the molecular-level polymorphism of Arctic $A\beta$ may manifest as different numbers of cross- β subunits.

The Arctic $A\beta$ 40 polymorphism is discrete, as shown by the much narrower line widths of each conformer compared to the

chemical shift differences between conformers. This property differs from that of membrane peptides and proteins, which usually exhibit a single broad peak per site, indicating continuous Gaussian distributions of conformations.^{59,74} The discreteness of the Arctic structural polymorphism can be attributed to the quasi-crystalline nature of the fibrils, which preserve and amplify the structure homogeneity within each fibril but leave different fibrils able to adopt different structures that started from a specific monomer or nucleus structure. The narrow line widths of both the “monomorphic” Osaka $A\beta$ 40 and each conformer of the polymorphic Arctic $A\beta$ 40 also indicate that $A\beta$ fibrils obtained from synthetic peptides without seeding can have inherently homogeneous structures that are comparable to the homogeneity of fibrils obtained from recombinant proteins.^{14,17,29}

Since our main motivation in this study is to understand the origins of $A\beta$ polymorphism, we did not try to reduce or eliminate the Arctic structural polymorphism by seeding, and it remains to be seen whether repeated rounds of seeding can render the Arctic fibril structures more homogeneous as it has for other $A\beta$ peptides.^{32,75} Even without a unique structure, chemical shifts and other lines of evidence constrain the structures of the major Arctic conformer I and rule out unlikely structural models. Since different chemical shifts cannot correspond to the same structures, our measured chemical shifts (Figure 5, Table S2) indicate that Arctic conformer I differs from all published $A\beta$ 40 structures so far, but bears significant resemblance to wild-type $A\beta$ 42. This chemical shift similarity is supported by the observed F19–I32 contact, which

is absent in all A β 40 structures but present in the A β 42 structure, and the preferential hydration of A21, which is inward-facing in wild-type A β 40 structures but outward-facing in Osaka A β 40 and wild-type A β 42 structures (Figure 9). Other biophysical and biological data have previously alluded to similar physical properties of Arctic A β 40 and A β 42 fibrils. Both Arctic A β 40 and wild-type A β 42 fibrils have faster and more heterogeneous growth than wild-type A β 40,⁶⁶ and both Arctic A β 40 and wild-type A β 42 fibrils appear in TEM and AFM images as more twisted and entangled than wild-type A β 40 fibrils.^{18,66}

The hydration data provide further insights into the overall fold of Arctic A β 40. The water–protein spin diffusion rates reflect the water-exposed surface area of the fibrils. Ignoring the unstructured N-terminal loop, the external surface area of the fibril core differs for the four A β structures (Figure 9): the approximate cross-sectional areas are 5 × 3 nm for the 2-fold wild-type A β 40, 7 × 7 × 7 nm for the 3-fold wild-type A β 40, 6 × 3 nm for Osaka A β 40, and a compact 4.5 × 3.5 nm for A β 42. The smaller dimension of the Osaka structure is consistent with the observed fast water ¹H spin diffusion to the Osaka A β 40, while the large dimension of the 3-fold A β 40 structure is consistent with the slow water magnetization transfer to the wild-type A β 40. Therefore, the slow water–protein spin diffusion of the Arctic sample suggests a large fibril cross section. If the Arctic conformer I has a similar subunit structure to A β 42, then our data suggest the existence of multiple cross- β subunits in the Arctic I fibril. For the 4.5 × 3.5 nm S-shape triple- β structure of A β 42 reported by Ishii and co-workers, corresponding TEM images showed fibril diameters ranging from 4.5 to 6.0 nm for thin fibrils to 6–13 nm for thicker ones. Thus, while a single subunit was proposed for the structure, a dimer model cannot be excluded.¹⁵ Earlier mass-per-length (MPL) data of A β 42 fibrils showed a broad distribution of 13–50 kDa/nm with a maximum at 19 kDa/nm,⁷⁶ also suggesting that A β 42 fibrils have multiple cross- β subunits, with the dimer being dominant. Interestingly, this A β 42 MPL distribution is much broader than reported wild-type A β 40 fibrils,^{12,25} indicating more pronounced polymorphism, which is also seen for the chemical-shift-similar Arctic A β 40 fibrils. Indeed, during the publication of this study, an atomic-resolution SSNMR structure of A β 42 was reported that revealed a dimer as the repeat motif of the cross- β structure.⁷⁷

In addition to water on the external fibril surface, X-ray fiber diffraction data and MD simulations suggest the presence of water inside the A β fibril core. Comparison of the calculated diffraction patterns for solid versus hollow cylinders with the measured diffraction pattern suggests that the 3-fold in vitro structure of wild-type A β 40 should contain a hollow water-filled core.⁷⁸ MD simulations suggest that this central cavity may have dimensions ranging from 1.6 to 2.3 nm,⁷⁹ and water molecules may diffuse from the fibril ends into the core and may localize in the turn region to solvate the D23–K28 salt bridge.^{80,81} The current water-to-protein ¹H spin diffusion experiment does not explicitly distinguish external versus internal water; but under the condition of our experiments, the external water may be preferentially detected because of its larger quantity. Interior water cavities have also been reported in other amyloid fibrils, including residues 105–115 of transthyretin,⁶ the SH3 domain of PI3 kinase,⁸² and a poly-L-glutamine peptide of huntingtin;⁸³ thus, they may be a general phenomenon of interest. Additional more specialized experi-

ments will be necessary to differentiate external and internal water molecules.

The polymorphism of E22G A β 40 is also consistent with the enhanced microsecond motions of the peptide compared to the wild-type counterpart and the Osaka mutant. Our data indicate that the motional amplitudes in the Arctic fibrils are small for very fast motions but larger for microsecond motions, as seen by the DIPSHIFT intensity asymmetry. Perhaps more importantly, the motional rates are faster when residue 22 is occupied by Gly than if it is occupied by the negatively charged Glu, or if Ala21 is followed directly by Asp23. The replacement of a charged Glu by Gly at residue 22 likely has the simultaneous effect of perturbing a stabilizing electrostatic interaction and conferring greater conformational flexibility to the peptide backbone. Our data here do not indicate the relative importance of these two factors for polymorphism, but comparison with literature suggests that both factors play a role. An intermolecular K16-E22 salt bridge has been observed in WT fibrils prepared quiescently²⁵ while absent in antiparallel fibrils of the D23N mutant.³¹ However, a charged E22 may impact the fibril structure at the nucleation stage, by restricting the initial conformational ensemble, without manifesting a permanent salt bridge in the mature fibril. The fact that the charge-removing D23N mutation at the next residue caused the coexistence of parallel and antiparallel fibrils,^{31,32} which is a dramatic departure from the parallel-only fibrils of the WT peptide, also suggests that removing an electrostatic charge in this region increases the conformational heterogeneity. For the D23N mutant, the parallel fibrils resemble the WT parallel fibril structure, which differ from the E22G mutant, where none of the four polymorphs resembles any existing A β 40 structures based on the chemical shifts. Thus, E22 appears to exert a more significant effect than D23 on the final three-dimensional fold of the fibril. This is generally consistent with the fact that E22 harbors the largest number of single-site mutations in A β peptides.³⁴

CONCLUSIONS

These solid-state NMR data show that mutation of Glu at residue 22 to Gly causes pronounced structural polymorphism and larger-amplitude microsecond motions in A β 40 fibrils. ¹³C and ¹⁵N chemical shifts indicate that at least four distinct molecular conformations exist in this Arctic mutant, none of which resembles the wild-type and other mutant A β 40 structures determined so far. However, chemical shifts and inter-residue contacts of one of the Arctic conformers have significant resemblances to those of A β 42 fibrils, suggesting similar structures. Comparison with the wild-type peptide and with a E22 deletion mutant indicate that the amino acid sequence with the lowest charge density in the region of residues 21–23, which is the Arctic Ala-Gly-Asp, has the highest degree of structural polymorphism and the fastest microsecond motions, while the amino acid sequence with the highest charge density in this region, which is the wild-type Ala-Glu-Asp, displays low polymorphism and the slowest microsecond motion. The intermediate sequence, adopted by the Osaka mutant, Ala-Asp, has a single structure and intermediate motional properties between the wild-type and Arctic A β 40. These results suggest that electrostatic interactions in this non- β region of the A β peptide, together with conformational flexibility, are crucial for determining the number of possible β -strand arrangements and the exact three-dimensional folds of this family of amyloid fibrils.⁷³ Interactions of these structurally

distinct fibrils with other cellular macromolecules can conceivably lead to different AD phenotypes.

■ ASSOCIATED CONTENT

■ Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/jacs.6b03715.

Additional NMR spectra, tables, and sample characterization data include: Peptide purification and early fibrilization data (Figures S1 and S2); Arctic A β chemical shifts and their comparison with other A β fibrils (Tables S1 and S2); Comparative 1D and 2D spectra of A β fibrils (Figures S3, S4, S5); Residue-specific heat maps of chemical shift differences (Figure S6); Additional order parameter and relaxation data (Figures S7, S8, Table S3); 2D water-edited spectra (Figure S9) (PDF)

■ AUTHOR INFORMATION

Corresponding Author

*meihong@mit.edu

Author Contributions

^{||}M.R.E. and T.W. contributed equally to this work.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

The SSNMR part of the work at MIT is supported by National Institutes of Health grants GM088204 and P01 AG002132. 900 MHz NMR spectra were measured at the MIT/Harvard Center for Magnetic Resonance, which is supported by NIH grant EB002026. W.F.D., M.N., S.B.P. and J.S. are supported by grants from the National Institutes of Health (AG02132 and AG10770) and T32 training grant HL07731 (M.N.). J.S. was supported by a fellowship from the Glenn Foundation and the New Investigator Research Grant (NIRG) of the Alzheimer's Association. We thank Dr. Y. Wu for measuring the ¹H solution NMR spectra.

■ REFERENCES

- (1) Tanzi, R. E.; Bertram, L. *Cell* **2005**, *120*, 545–555.
- (2) Hardy, J.; Selkoe, D. J. *Science* **2002**, *297*, 353–356.
- (3) Wasmer, C.; Lange, A.; Van Melckebeke, H.; Siemer, A. B.; Riek, R.; Meier, B. H. *Science* **2008**, *319*, 1523–1526.
- (4) Meier, B. H.; Bockmann, A. *Curr. Opin. Struct. Biol.* **2015**, *30*, 43–49.
- (5) Helmus, J. J.; Surewicz, K.; Surewicz, W. K.; Jaroniec, C. P. *J. Am. Chem. Soc.* **2010**, *132*, 2393–2403.
- (6) Fitzpatrick, A. W.; Debelouchina, G. T.; Bayro, M. J.; Clare, D. K.; Caporini, M. A.; Bajaj, V. S.; Jaroniec, C. P.; Wang, L.; Ladizhansky, V.; Muller, S. A.; MacPhee, C. E.; Waudby, C. A.; Mott, H. R.; De Simone, A.; Knowles, T. P.; Saibil, H. R.; Vendruscolo, M.; Orlova, E. V.; Griffin, R. G.; Dobson, C. M. *Proc. Natl. Acad. Sci. U. S. A.* **2013**, *110*, 5468–5473.
- (7) Debelouchina, G. T.; Bayro, M. J.; Fitzpatrick, A. W.; Ladizhansky, V.; Colvin, M. T.; Caporini, M. A.; Jaroniec, C. P.; Bajaj, V. S.; Rosay, M.; MacPhee, C. E.; Vendruscolo, M.; Maas, W. E.; Dobson, C. M.; Griffin, R. G. *J. Am. Chem. Soc.* **2013**, *135*, 19237–19247.
- (8) Su, Y.; Sarell, C. J.; Eddy, M. T.; Debelouchina, G. T.; Andreas, L. B.; Pashley, C. L.; Radford, S. E.; Griffin, R. G. *J. Am. Chem. Soc.* **2014**, *136*, 6313–6325.
- (9) Tuttle, M. D.; Comellas, G.; Nieuwkoop, A. J.; Covell, D. J.; Berthold, D. A.; Klopper, K. D.; Courtney, J. M.; Kim, J. K.; Barclay, A. M.; Kendall, A.; Wan, W.; Stubbs, G.; Schwieters, C. D.; Lee, V. M.;

George, J. M.; Rienstra, C. M. *Nat. Struct. Mol. Biol.* **2016**, *23*, 409–415.

- (10) Heise, H.; Hoyer, W.; Becker, S.; Andronesi, O. C.; Riedel, D.; Baldus, M. *Proc. Natl. Acad. Sci. U. S. A.* **2005**, *102*, 15871–15876.
- (11) Tycko, R. *Annu. Rev. Phys. Chem.* **2011**, *62*, 279–299.
- (12) Paravastu, A. K.; Leapman, R. D.; Yau, W. M.; Tycko, R. *Proc. Natl. Acad. Sci. U. S. A.* **2008**, *105*, 18349–18354.
- (13) Petkova, A. T.; Yau, W. M.; Tycko, R. *Biochemistry* **2006**, *45*, 498–512.
- (14) Schutz, A. K.; Vagt, T.; Huber, M.; Ovchinnikova, O. Y.; Cadalbert, R.; Wall, J.; Guntert, P.; Bockmann, A.; Glockshuber, R.; Meier, B. H. *Angew. Chem., Int. Ed.* **2015**, *54*, 331–335.
- (15) Xiao, Y. L.; Ma, B. Y.; McElheny, D.; Parthasarathy, S.; Long, F.; Hoshi, M.; Nussinov, R.; Ishii, Y. *Nat. Struct. Mol. Biol.* **2015**, *22*, 499–U497.
- (16) Lu, J. X.; Qiang, W.; Yau, W. M.; Schwieters, C. D.; Meredith, S. C.; Tycko, R. *Cell* **2013**, *154*, 1257–1268.
- (17) Lopez del Amo, J. M.; Schmidt, M.; Fink, U.; Dasari, M.; Fandrich, M.; Reif, B. *Angew. Chem., Int. Ed.* **2012**, *51*, 6136–6139.
- (18) Antzutkin, O. N. *Magn. Reson. Chem.* **2004**, *42*, 231–246.
- (19) Tycko, R. *Protein Sci.* **2014**, *23*, 1528–1539.
- (20) Meinhardt, J.; Sachse, C.; Hortschansky, P.; Grigorieff, N.; Fandrich, M. *J. Mol. Biol.* **2009**, *386*, 869–877.
- (21) Antzutkin, O. N.; Balbach, J. J.; Leapman, R. D.; Rizzo, N. W.; Reed, J.; Tycko, R. *Proc. Natl. Acad. Sci. U. S. A.* **2000**, *97*, 13045–13050.
- (22) Bertini, I.; Gonnelli, L.; Luchinat, C.; Mao, J. F.; Nesi, A. *J. Am. Chem. Soc.* **2011**, *133*, 16013–16022.
- (23) Qiang, W.; Kelley, K.; Tycko, R. *J. Am. Chem. Soc.* **2013**, *135*, 6860–6871.
- (24) Petkova, A. T.; Buntkowsky, G.; Dyda, F.; Leapman, R. D.; Yau, W. M.; Tycko, R. *J. Mol. Biol.* **2004**, *335*, 247–260.
- (25) Petkova, A. T.; Leapman, R. D.; Guo, Z.; Yau, W. M.; Mattson, M. P.; Tycko, R. *Science* **2005**, *307*, 262–265.
- (26) Masuda, Y.; Uemura, S.; Ohashi, R.; Nakanishi, A.; Takegoshi, K.; Shimizu, T.; Shirasawa, T.; Irie, K. *ChemBioChem* **2009**, *10*, 287–295.
- (27) Jarrett, J. T.; Berger, E. P.; Lansbury, P. T. *Biochemistry* **1993**, *32*, 4693–4697.
- (28) Parthasarathy, S.; Inoue, M.; Xiao, Y. L.; Matsumura, Y.; Nabeshima, Y.; Hoshi, M.; Ishii, Y. *J. Am. Chem. Soc.* **2015**, *137*, 6480–6483.
- (29) Colvin, M. T.; Silvers, R.; Frohm, B.; Su, Y.; Linse, S.; Griffin, R. G. *J. Am. Chem. Soc.* **2015**, *137*, 7509–7518.
- (30) Huber, M.; Ovchinnikova, O. Y.; Schutz, A. K.; Glockshuber, R.; Meier, B. H.; Bockmann, A. *Biomol. NMR Assignments* **2015**, *9*, 7–14.
- (31) Qiang, W.; Yau, W. M.; Luo, Y.; Mattson, M. P.; Tycko, R. *Proc. Natl. Acad. Sci. U. S. A.* **2012**, *109*, 4443–4448.
- (32) Qiang, W.; Yau, W. M.; Tycko, R. *J. Am. Chem. Soc.* **2011**, *133*, 4018–4029.
- (33) Mathis, C. A.; Wang, Y. M.; Holt, D. P.; Huang, G. F.; Debnath, M. L.; Klunk, W. E. *J. Med. Chem.* **2003**, *46*, 2740–2754.
- (34) Benilova, I.; Karran, E.; De Strooper, B. *Nat. Neurosci.* **2012**, *15*, 349–357.
- (35) Hendriks, L.; Vanduijn, C. M.; Cras, P.; Cruts, M.; Vanhul, W.; Vanharskamp, F.; Warren, A.; Mcinnis, M. G.; Antonarakis, S. E.; Martin, J. J.; Hofman, A.; Van Broeckhoven, C. *Nat. Genet.* **1992**, *1*, 218–221.
- (36) Tomiyama, T.; Nagata, T.; Shimada, H.; Teraoka, R.; Fukushima, A.; Kanemitsu, H.; Takuma, H.; Kuwano, R.; Imagawa, M.; Ataka, S.; Wada, Y.; Yoshioka, E.; Nishizaki, T.; Watanabe, Y.; Mori, H. *Ann. Neurol.* **2008**, *63*, 377–387.
- (37) Bugiani, O.; Giaccone, G.; Rossi, G.; Mangieri, M.; Capobianco, R.; Morbin, M.; Mazzoleni, G.; Cupidi, G.; Marcon, G.; Giovagnoli, A.; Bizzi, A.; Di Fede, G.; Puoti, G.; Carella, F.; Salmaggi, A.; Romorini, A.; Patruno, G. M.; Magoni, M.; Padovani, A.; Tagliavini, F. *Arch. Neurol.* **2010**, *67*, 987–995.
- (38) Nilsberth, C.; Westlind-Danielsson, A.; Eckman, C. B.; Condron, M. M.; Axelman, K.; Forsell, C.; Stenh, C.; Luthman, J.;

- Teplow, D. B.; Younkin, S. G.; Naslund, J.; Lannfelt, L. *Nat. Neurosci.* **2001**, *4*, 887–893.
- (39) Van Broeckhoven, C.; Haan, J.; Bakker, E.; Hardy, J. A.; Van Hul, W.; Wehnert, A.; Vegter-Van der Vlis, M.; Roos, R. A. *Science* **1990**, *248*, 1120–1122.
- (40) Grabowski, T. J.; Cho, H. S.; Vonsattel, J. P. G.; Rebeck, G. W.; Greenberg, S. M. *Ann. Neurol.* **2001**, *49*, 697–705.
- (41) Englund, H.; Sehlin, D.; Johansson, A. S.; Nilsson, L. N.; Gellerfors, P.; Paulie, S.; Lannfelt, L.; Pettersson, F. E. *J. Neurochem.* **2007**, *103*, 334–345.
- (42) Johansson, A. S.; Berglind-Dehlin, F.; Karlsson, G.; Edwards, K.; Gellerfors, P.; Lannfelt, L. *FEBS J.* **2006**, *273*, 2618–2630.
- (43) Cheng, I. H.; Palop, J. J.; Esposito, L. A.; Bien-Ly, N.; Yan, F. G.; Mucke, L. *Nat. Med.* **2004**, *10*, 1190–1192.
- (44) Lord, A.; Kalimo, H.; Eckman, C.; Zhang, X. Q.; Lannfelt, L.; Nilsson, L. N. *Neurobiol. Aging* **2006**, *27*, 67–77.
- (45) Lam, A. R.; Teplow, D. B.; Stanley, H. E.; Urbanc, B. *J. Am. Chem. Soc.* **2008**, *130*, 17413–17422.
- (46) Stohr, J.; Watts, J. C.; Mensinger, Z. L.; Oehler, A.; Grillo, S. K.; DeArmond, S. J.; Prusiner, S. B.; Giles, K. *Proc. Natl. Acad. Sci. U. S. A.* **2012**, *109*, 11025–11030.
- (47) Watts, J. C.; Condello, C.; Stohr, J.; Oehler, A.; Lee, J.; DeArmond, S. J.; Lannfelt, L.; Ingelsson, M.; Giles, K.; Prusiner, S. B. *Proc. Natl. Acad. Sci. U. S. A.* **2014**, *111*, 10323–10328.
- (48) FINDER, V. H.; Vodopivec, I.; Nitsch, R. M.; Glockshuber, R. *J. Mol. Biol.* **2010**, *396*, 9–18.
- (49) Hong, M.; Griffin, R. G. *J. Am. Chem. Soc.* **1998**, *120*, 7113–7114.
- (50) Rienstra, C. M.; Tucker-Kellogg, L.; Jaroniec, C. P.; Hohwy, M.; Reif, B.; McMahon, M. T.; Tidor, B.; Lozano-Perez, T.; Griffin, R. G. *Proc. Natl. Acad. Sci. U. S. A.* **2002**, *99*, 10260–10265.
- (51) Takegoshi, K.; Nakamura, S.; Terao, T. *Chem. Phys. Lett.* **2001**, *344*, 631–637.
- (52) Hong, M. *J. Biomol. NMR* **1999**, *15*, 1–14.
- (53) Baldus, M.; P, A. T.; Herzfeld, J.; Griffin, R. G. *Mol. Phys.* **1998**, *95*, 1197–1207.
- (54) Kumashiro, K. K.; Schmidt-Rohr, K.; Murphy, O. J. I.; Ouellette, K. L.; Cramer, W. A.; Thompson, L. K. *J. Am. Chem. Soc.* **1998**, *120*, 5043–5051.
- (55) Luo, W.; Hong, M. *J. Am. Chem. Soc.* **2006**, *128*, 7242–7251.
- (56) Ader, C.; Schneider, R.; Seidel, K.; Eitzkorn, M.; Becker, S.; Baldus, M. *J. Am. Chem. Soc.* **2009**, *131*, 170–176.
- (57) White, P. B.; Wang, T.; Park, Y. B.; Cosgrove, D. J.; Hong, M. *J. Am. Chem. Soc.* **2014**, *136*, 10399–10409.
- (58) Williams, J. K.; Hong, M. *J. Magn. Reson.* **2014**, *247*, 118–127.
- (59) Liao, S. Y.; Fritzsche, K. J.; Hong, M. *Protein Sci.* **2013**, *22*, 1623–1638.
- (60) Munowitz, M. G.; Griffin, R. G.; Bodenhausen, G.; Huang, T. H. *J. Am. Chem. Soc.* **1981**, *103*, 2529–2533.
- (61) Bielecki, A.; Kolbert, A. C.; Levitt, M. H. *Chem. Phys. Lett.* **1989**, *155*, 341–346.
- (62) Hong, M.; Gross, J. D.; Rienstra, C. M.; Griffin, R. G.; Kumashiro, K. K.; Schmidt-Rohr, K. *J. Magn. Reson.* **1997**, *129*, 85–92.
- (63) Hong, M. *J. Magn. Reson.* **1999**, *139*, 389–401.
- (64) Rothwell, W. P.; Waugh, J. S. *J. Chem. Phys.* **1981**, *74*, 2721–2732.
- (65) Cady, S. D.; Hong, M. *J. Biomol. NMR* **2009**, *45*, 185–196.
- (66) Norlin, N.; Hellberg, M.; Filippov, A.; Sousa, A. A.; Grobner, G.; Leapman, R. D.; Almqvist, N.; Antzutkin, O. N. *J. Struct. Biol.* **2012**, *180*, 174–189.
- (67) Betts, V.; Leissring, M. A.; Dolios, G.; Wang, R.; Selkoe, D. J.; Walsh, D. M. *Neurobiol. Dis.* **2008**, *31*, 442–450.
- (68) Cloe, A. L.; Orgel, J. P.; Sachleben, J. R.; Tycko, R.; Meredith, S. C. *Biochemistry* **2011**, *50*, 2026–2039.
- (69) Stohr, J.; Condello, C.; Watts, J. C.; Bloch, L.; Oehler, A.; Nick, M.; DeArmond, S. J.; Giles, K.; DeGrado, W. F.; Prusiner, S. B. *Proc. Natl. Acad. Sci. U. S. A.* **2014**, *111*, 10329–10334.
- (70) Schutz, A. K.; Habenstein, B.; Luckgei, N.; Bousset, L.; Sourigues, Y.; Nielsen, A. B.; Melki, R.; Bockmann, A.; Meier, B. H. *Biomol. NMR Assignments* **2014**, *8*, 349–356.
- (71) Cobo, M. F.; Achilles, A.; Reichert, D.; Deazevedo, E. R.; Saalwächter, K. *J. Magn. Reson.* **2012**, *221*, 85–96.
- (72) deAzevedo, E. R.; Saalwachter, K.; Pascui, O.; de Souza, A. A.; Bonagamba, T. J.; Reichert, D. *J. Chem. Phys.* **2008**, *128*, 104505.
- (73) Schledorn, M.; Meier, B. H.; Bockmann, A. *Front. Mol. Biosci.* **2015**, *2*, 14.
- (74) Su, Y.; Hong, M. *J. Phys. Chem. B* **2011**, *115*, 10758–10767.
- (75) Ravotti, F.; Walti, M. A.; Guntert, P.; Riek, R.; Bockmann, A.; Meier, B. H. *Biomol. NMR Assignments* **2016**, DOI: 10.1007/s12104-016-9682-y.
- (76) Antzutkin, O. N.; Leapman, R. D.; Balbach, J. J.; Tycko, R. *Biochemistry* **2002**, *41*, 15436–15450.
- (77) Colvin, M. T.; Silvers, R.; Ni, Q. Z.; Can, T. V.; Sergeyev, I. V.; Rosay, M.; Donovan, K. J.; Michael, B.; Wall, J. S.; Linse, S.; Griffin, R. G. *J. Am. Chem. Soc.* **2016**, DOI: 10.1021/jacs.6b05129.
- (78) McDonald, M.; Box, H.; Bian, W.; Kendall, A.; Tycko, R.; Stubbs, G. *J. Mol. Biol.* **2012**, *423*, 454–461.
- (79) Miller, Y.; Ma, B.; Nussinov, R. *J. Am. Chem. Soc.* **2011**, *133*, 2742–2748.
- (80) Buchete, N. V.; Tycko, R.; Hummer, G. *J. Mol. Biol.* **2005**, *353*, 804–821.
- (81) Prade, E.; Bittner, H. J.; Sarkar, R.; Lopez Del Amo, J. M.; Althoff-Ospelt, G.; Multhaup, G.; Hildebrand, P. W.; Reif, B. *J. Biol. Chem.* **2015**, *290*, 28737–28745.
- (82) Bayro, M. J.; Maly, T.; Birkett, N. R.; MacPhee, C. E.; Dobson, C. M.; Griffin, R. G. *Biochemistry* **2010**, *49*, 7474–7484.
- (83) Perutz, M. F.; Finch, J. T.; Berriman, J.; Lesk, A. *Proc. Natl. Acad. Sci. U. S. A.* **2002**, *99*, 5591–5595.