


Catalytic Synthesis of Saturated Oxygen Heterocycles by Hydrofunctionalization of Unactivated Olefins: Unprotected and Protected Strategies

Hiroki Shigehisa,* Miki Hayashi, Haruna Ohkawa, Tsuyoshi Suzuki, Hiroki Okayasu, Mayumi Mukai, Ayaka Yamazaki, Ryohei Kawai, Harue Kikuchi, Yui Satoh, Akane Fukuyama, and Kou Hiroya

Faculty of Pharmacy, Musashino University, 1-1-20 Shinmachi Nishitokyo-shi, Tokyo 202-8585, Japan

Supporting Information

ABSTRACT: A mild, general, and functional group tolerant intramolecular hydroalkoxylation and hydroacyloxylation of unactivated olefins using a Co(salen) complex, an *N*-fluoropyridinium salt, and a disiloxane reagent is described. This reaction was carried out at room temperature and afforded five- and six-membered oxygen heterocyclic compounds, such as cyclic ethers and lactones. The Co complex was optimized for previously rare medium ring formation by hydrofunctionalization of unactivated olefins. The powerful Co catalyst system also enables the deprotective hydroalkoxylation of *O*-protected alkenyl alcohol and hydroacyloxylation of alkenyl ester to afford cyclic ethers and lactones directly. The substrate scope and mechanistic proof of deprotection were investigated. The experimental evidence supports the concerted transition state of the bond-forming step involving a cationic Co complex.



1. INTRODUCTION

Saturated oxygen heterocycles, such as cyclic ethers and lactones, are found in the structures of many biologically active natural products.¹ Beyond the common five- and six-membered rings, rings of seven or more members, also called medium rings, have also been discovered; examples include polycyclic marine toxins (e.g., ciguatoxin), lauroxanes (e.g., laurencin), heliannuols (e.g., helianane), sodwanone S, zoapatanol, aphanamixoids (e.g., aphanamixoid A), and octalactins (e.g., octalactin A).² From a drug discovery perspective, there is still much room for the development of a chemical space derived from a medium-ring framework.³

In the field of organic synthesis, straightforward methods for accessing the motif of saturated oxygen heterocycles include catalytic regioselective intramolecular hydroalkoxylation and hydroacyloxylation, which refer to the direct coupling of olefins with hydroxyl groups and carboxyl groups, respectively. Consequently, many research groups have reported various examples of attractive Markovnikov selective intramolecular hydroalkoxylation⁴ and hydroacyloxylation^{4b,d,e,5} over the past decade (Scheme 1). For example, some reactions are based on nucleophilic attack by OH groups on the metal-coordinated olefins and protonation of the metal-carbon bond, as presented by Widenhofer,^{4a} He,^{4b} and others.^{4c-f} Nonmetallic systems also enable the hydrofunctionalization of olefins.^{4g,i-k} Catalytic hydrofunctionalizations starting from *O*-nucleophile activation by a metal complex were also reported by Marks,^{4l,m} Duñach,^{4n,o} Hartung,^{4p} and others.^{4q,r} Despite the development


of these elegant approaches, the examples of medium-ring formation by hydroalkoxylation are quite limited, and these scant examples are further limited to the formation of rings of no more than seven members.^{4d,v,aa} Recently, Tiefenbacher reported intramolecular hydroalkoxylation catalyzed by a self-assembled hexameric capsule affording seven-membered products; however, the substrate scope is fundamentally limited by the encapsulation.^{4aa} Nicewicz reported an exclusive anti-Markovnikov selective hydroalkoxylation and an example of oxepane formation.⁶ Meanwhile, to the best of our knowledge, the formation of rings of more than seven members in the hydroacyloxylation of olefins has not been reported. In contrast, haloetherification⁷ and halolactonization^{7b,s} have been thoroughly investigated. Although these approaches have the merit of enabling further manipulation on the halogen atom, removal of the terminal halogen, if unnecessary, entails an additional step.

Medium-ring formation is often fundamentally troublesome because of entropic factors (the approach frequency of two reactive sites), strain of the medium ring, and transannular interactions in the substrate. Therefore, a powerful and selective activation of the olefin in the reaction is highly desired for the development of a general and robust approach based on both hydroalkoxylation and hydroacyloxylation for medium-ring formation with good functional group tolerance.

Received: June 4, 2016


Published: July 27, 2016

Scheme 1. Our Previous Reports and This Work



We recently reported an intermolecular hydroalkoxylation,⁹ an intramolecular hydroamination,¹⁰ and an intramolecular hydroarylation¹¹ using Co(salen) complex **1**, an *N*-fluoro-2,4,6-trimethylpyridinium salt, and a 1,1,3,3-tetramethyldisiloxane [(Me₂SiH)₂O]. In these reactions, it is thought that a cationic intermediate enables bond formation between the olefins and the nucleophile (Scheme 1a–c). The reaction demonstrated a broad substrate scope as a result of its high reactivity and functional group tolerance. In light of these previous observations and recent relevant examples,¹² we envisioned that intramolecular C–O bond formation may occur under the identified or modified reaction conditions to yield both cyclic ethers and lactones with excellent functional group tolerance (Scheme 1d). Previously, we reported a preliminary result in which cyclization-biased alkenyl alcohol **2a** was used to obtain cyclic ether **3a** in excellent yield (Scheme 2).^{9a} Furthermore, TBS (*t*-butyldimethylsilyl)-protected **2aa** was directly cyclized to afford the same product **3a**. In the multistep synthesis of complex molecules, the use of a protective group is often unavoidable because of the functional group reactivity, solubility, and high polarity of the compound. This deprotective cyclization approach has the advantage of being step-economical, allowing the deprotection step to be omitted. Nevertheless, there are few examples of deprotective cyclization for simple substrates under harsh conditions; Niggemann reported only one reaction of benzyl-deprotective hydroalkoxylation (dichloromethane, 80 °C),^{4f} and Duñach reported the hydroacyloxylation of alkenyl ester (dichloroethane or nitromethane, reflux).¹³


Herein, we developed a catalytic, mild, and functional group tolerant intramolecular hydroalkoxylation and hydroacyloxylation of unactivated olefins using a Co(salen) complex **1**, an *N*-

Scheme 2. Preliminary Result of Hydroalkoxylation of Olefins and Deprotective Cyclization^{9a}

fluoro-2,4,6-trimethylpyridinium salt, and (Me₂SiH)₂O to produce various five- and six-membered ring products. Next, the structure–reactivity relationship of the salen ligand was studied, identifying the optimum Co(salen) complex for medium-ring formation (seven- to nine-membered). It was then found that various cyclic ethers or lactones were directly obtained from *O*-protected alkenyl alcohol or alkenyl ester, respectively, by the Co catalyst system at room temperature. The ultimate coproducts of the protective group were also investigated to clarify the mechanism of the deprotection step. Finally, the experimental evidence supported the concerted transition state of the bond-forming step involving a cationic Co complex.


2. RESULTS AND DISCUSSION

Synthesis of Five- and Six-Membered Cyclic Ethers and Lactones. First, the scope of alkenyl alcohol was investigated under the reaction conditions of the intramolecular hydroamination we previously reported, namely, Co complex **1** (3.0 mol %) in toluene and in the presence of *N*-fluoro-2,4,6-trimethylpyridinium trifluoromethanesulfonate (Me₃NFPY·OTf, 2.0 equiv)¹⁴ and (Me₂SiH)₂O (2.0 equiv) (Table 1). We found that various five- to six-membered cyclic ethers were obtained in good-to-excellent yields. Alkenyl alcohol **2a** was smoothly cyclized to afford tetrahydrofuran **3a** in toluene. Substrates **2b** and **2c** (containing a bulky hydroxyl group) and **2d** (1,1-disubstituted olefin) were also subjected to cyclization to afford the desired products **3b**, **3c** (0% yield in ref 4i), and **3d**, respectively, in excellent yield. **3e** and **3f** were obtained from alkenyl alcohol containing the acid-sensitive acetal group **2e** and diol **2f**. A steroidal compound **3g** was also synthesized in good yield. Unfortunately, the phenolic hydroxyl group was found to be unsuitable for the cyclization, resulting in a complex product mixture (**2h**). Concerning the six-membered rings, tetrahydropyran **3i** and isochroman **3j**, **3k** could be synthesized. The investigation of the functional group tolerance of this method revealed that isochromans **3l**–**3n** were obtained from allyl-benzylalcohol-bearing, fluoro-anion-sensitive silyl ether (**2l**); acid-sensitive acetal (**2m**); and benzyl ether (**2n**). Unfortunately, the synthesis of three- or four-ring compounds was found to be ineffective in our conditions (see Supporting Information).

Table 1. Scope of Alkenyl Alcohol for Hydroalkoxylation of Unactivated Olefins^a

^aIsolation yield (0.25 mmol scale, 0.1 M). ^bNMR yield using 1,4-bis(trifluoromethyl)benzene as the internal standard.

Encouraged by this result, we also investigated the hydroacyloxylation of olefins using a series of alkenyl carboxylic acids under the same reaction conditions (Table 2). To our delight, five-membered lactones were obtained from the substrate, including cyclization-biased 4a and 1,1-disubstituted olefins 4b and 4c in excellent yield. Isobenzofuranone 5d was obtained from benzoic acid derivative 4d. Six-membered

Table 2. Scope of Alkenyl Carboxylic Acid for Hydroacyloxylation of Unactivated Olefins^a

^aIsolation yield (0.25 mmol scale, 0.1 M).

products—such as lactones 5e and 5f, 3-isochromanone 5g, and 1-isochromanone 5h—were obtained in good yield.


Synthesis of Medium-Ring Products. We explored the possibility of achieving unprecedented medium-ring formation by hydrofunctionalization under a Co catalyst system. First, we investigated the hydroalkoxylation of alkenyl alcohol 6a containing monosubstituted olefins (Table 3). Because the

Table 3. Optimization of Reaction Condition for Medium-Ring Formation by Hydroalkoxylation of Unactivated Olefins

Reaction scheme for Table 3: Alkenyl alcohol (6a) reacts with a Co catalyst (1, 3 mol %), Me₃NFPY-OTf (2 equiv), and (Me₂SiH)₂O (2 equiv) in CH₃Ph at room temperature for 20 h to form medium-ring product (7a).

entry	Co cat	yield (%) ^a
1	1	30
2	8	20
3	9	40
4	10	43, (41) ^b
5	11	25
6	12	<5
7	13	<5
8	14	22
9	15	8
10	16	57
11	17	60, (50) ^c
12	18	57
13 ^d	17	60

^aNMR yield using 1,4-bis(trifluoromethyl)benzene as internal standard (0.25 mmol scale, 0.1 M). ^bCF₃Ph was used. ^cIsolation yield. ^dConcentration = 0.03 M.




reaction condition for the synthesis of common ring sizes using complex 1 gave oxepane 7a in only 30% yield along with considerable amounts of olefin isomers and recovered 6a (entry 1), we conducted a systematic screening of the salen ligand in the Co complex. We found that decreasing the bulkiness of diamine moiety improved the yield and that the ethylenediamine-containing ligand was effective (entries 2–4). Longer diamines gave worse results (entries 5, 6). Next, we examined the effect of substituents on the aromatic rings. It was found that using Co complexes possessing smaller methyl groups in either the 3 or 5 position (benzaldehyde numbering) dramatically decreased the yield of the desired product 7a (entries 7, 8). Given the importance of both the substituents on the aromatic ring, a kit of Co complexes containing substituents of different sizes was investigated (entries 9–12). Although further screening of Co complexes was investigated, we ultimately identified 17 as being optimal (see Supporting

Information). Dilute conditions gave almost the same result (entry 13).

Having demonstrated the catalytic activity of complex 17, we next examined the scope of other substrates (Table 4). In the

Table 4. Scope of Alkenyl Alcohol and Alkenyl Carboxylic Acid Affording Medium-Ring Products^a




^aIsolation yield (0.25 mmol scale, 0.1 M). ^bCatalyst 1 (3 mol %) was used.

case of synthesizing seven-membered rings by hydroalkoxylation, oxepane 7b and 1,4-dioxepane 7c, 7d were obtained from alkenyl alcohol containing mono- and disubstituted olefin. Nitrogen-containing 1,4-oxazepine 7e was also synthesized by this method. In the case of eight-membered ring formation affording 7f, we reconfirmed the superiority of complex 17 in terms of yield. Furthermore, this method was applicable for nine-membered ring formation affording 7g. Compared to hydroalkoxylation, hydroacyloxylation gave slightly higher yields of oxepanone 7h (vs 7a), benzodioxocanone 7i (vs 7f), and benzodioxonanone 7j (vs 7g). At this stage, some substrates were identified as being unsuitable even when using the optimum catalyst 17 because the isomerization of olefin was unavoidably faster than the cyclization (see Supporting Information). Although the yield of the medium-ring product was inferior to that of the small-ring product, various medium-ring products that had not been achieved previously were obtained by the hydrofunctionalization of olefins.

Deprotective Intramolecular Hydroalkoxylation of Unactivated Olefins. We next investigated the scope of the protective group using cyclization-biased alkenyl alcohol 2aa–2ag by the Co catalyst system (Table 5). In addition to the TBS group previously shown to be effective (Scheme 2), acetal (methoxymethyl acetal (MOM)), methoxyethoxymethyl acetal (MEM), benzyloxymethyl acetal (BOM), and benzyl and methyl groups were found to be applicable. In contrast, the cyclization of 2ag containing an acetyl group did not occur

Table 5. Scope of Protective Group for Hydroalkoxylation of Unactivated Olefins



entry	PG	time (h)	yield (%) ^a
1	TBS (2aa)	0.5	99
2	MOM (2ab)	0.5	99
3	MEM (2ac)	0.5	97
4	BOM (2ad)	0.5	99
5	Bn (2ae)	1.5	93
6	Me (2af)	1.5	87
7	Ac (2ag)	6.5	27

^aIsolation yield (0.25 mmol scale, 0.1 M).

smoothly. Encouraged by this result, we next investigated the substrate scope for three protective groups per substrate (Table 6). Overall, it was found that MOM was much more effective for the deprotective cyclization than either TBS or benzyl groups.¹⁵ For example, in the case of five-membered ring formation, tetrahydrofurans 3b and 3d were obtained in acceptable yield from MOM-protected 2bb¹⁶ and 2db, respectively, whereas the other two protective groups provided the desired products in low yields, if at all. Among the bis-protected diol substrates 2fa–2fe, the yield of the desired product was highest for the MOM-protected 2fb. Although the phenolic hydroxyl group was unsuitable for substrate 2h (Table 1), to our delight, the yield of benzofuran 3h was excellent for any protective group (2ha–2he). A more electron rich substrate 2hb' was applicable. In the case of six-membered ring formation, the yield of tetrahydropyran 3i was also quite high when using the MOM group (2ib) and was exceptionally high when using the benzyl group (2ie). However, isochroman 3j was only obtained in good yield from the MOM group containing substrate 2jb. Considering the result of 3d as well, it was concluded that the MOM group is required when using styrene-type substrates. Indeed, each protective group was effective for the synthesis of other types of isochroman 3k. We next examined the functional group tolerance using the substrate, and various isochromans were obtained from substrate bearing TBS (2lb), MOM (2mb), and benzyl groups (2nb). Unfortunately, the synthesis of medium-ring compounds by deprotective hydroalkoxylation was found to be ineffective, even when using MOM groups. After continued investigation, only the eight-membered cyclic ether 7f was obtained in 26% yield using MOM-protected 6fb.


Deprotective Intramolecular Hydroacyloxylation of Unactivated Olefins. Furthermore, the method of deprotective cyclization was found to be applicable to alkenyl esters. The results of the hydroacyloxylation of alkenyl esters 4aa–4ae are presented (Table 7). This method was applicable to all protective groups examined. In particular, the methyl group, being sterically smallest, gave the highest reactivity. We next investigated the substrate scope for various methyl esters (Table 8). The five-membered lactones 5b and 5d were obtained in good yield. However, 5c was not obtained at all, and instead, the olefin isomer (trisubstituted olefin) was formed. C–O bond cleavage of the oxonium cation to afford a stable carbocation intermediate might be faster than the deprotection step in the case of 1,1-disubstituted styrene-type

Table 6. Substrate Scope of Deprotective Hydroalkoxylation of Unactivated Olefins^a

entry	starting materials	products	yield (%)			entry	starting materials	products	yield (%)		
			TBS	MOM	Bn				TBS	MOM	Bn
1			<10 ^b 23 h	50 ^c 4.5 h	0 ^b 23 h	7			0 ^b 20 h	70 ^c 2.0 h	0 ^b 20 h
	2ba (PG=TMS) 2bb (PG=MOM) 2be (PG=Bn)	3b					2ja (PG=TBS) 2jb (PG=MOM) 2je (PG=Bn)	3j			
2			trace ^b 20 h	59 ^c 1.5 h	0 ^b 20 h	8			82 ^b 20 h	90 ^b 20 h	70 ^b 20 h
	2da (PG=TBS) 2db (PG=MOM) 2de (PG=Bn)	3d					2ka (PG=TBS) 2kb (PG=MOM) 2ke (PG=Bn)	3k			
3			50 ^b 21 h	53 ^b 1.5 h	trace ^b 21 h	9			ND	96 ^b 0.5 h	ND
	2fa (PG=TBS) 2fb (PG=MOM) 2fe (PG=Bn)	3fa (PG=TBS) 3e (PG=MOM) 3fe (PG=Bn)					2lb (OTBS)	3l (OTBS)			
4			84 ^b 19 h	90 ^b 1.5 h	73 ^b 2.5 h	10			ND	86 ^b 0.5 h	ND
	2ha (PG=TBS) 2hb (PG=MOM) 2he (PG=Bn)	3h					2mb (OMOM)	3m (OMOM)			
5			ND	79 ^c 1.5 h	ND	11			ND	89 ^b 2.0 h	ND
	2hb' (OMOM) 2hb'' (OBn)	3h' (OMOM)					2nb (OBn)	3n (OBn)			
6			trace 27 h	84 ^c 22 h	80 ^c 22 h	12 ^d			ND	26 ^c 20 h	ND
	2ia (PG=TBS) 2ib (PG=MOM) 2ie (PG=Bn)	3i					6fb (OMOM)	7f (OMOM)			

^a0.25 mmol scale, 0.1 M. ^bNMR yield using 1,4-bis(trifluoromethyl)benzene as the internal standard. ^cIsolation yield. ^dCatalyst 17 (3 mol %) was used.

Table 7. Scope of Protective Group for Hydroacyloxylation of Unactivated Olefins



entry	PG	time (h)	yield (%) ^a
1	Me (4aa)	0.5	99
2	Et (4ab)	1.0	99
3	Bn (4ac)	19	97
4	PMB (4ad)	3	99
5	^t Bu (4ae)	19	93

^aIsolation yield (0.25 mmol scale, 0.1 M).

substrate. Indeed, replacing the methyl group on olefin moiety of 4ba with the phenyl group did not afford the desired product 5b. Six-membered lactones, such as 5e–5h, were obtained in good to excellent yield. The desired product was obtained from styrene-type substrates 4ga (monosubstituted olefin). The functional group tolerance was also investigated for hydroacyloxylation using alkenyl-ester-bearing TBS (19aa), MOM (19ba), acetyl (19ca), and hydroxyl groups (19da). The eight-membered ring lactone 7i was also obtained from alkenyl ester 6ia.


Reaction Mechanism. The mechanistic implications of this deprotective cyclization merit discussion. First, in the case of the phenolic substrate (2h in Table 1 vs 2ha–2he in Table 6), the yield of desired product 3h was dramatically improved by using a protective group. Second, the investigation of the functional group tolerance using substrates 2lb–2nb (Table 6) and 19aa–19ba (Table 7) shows that the protective groups

Table 8. Scope of Alkenyl Methylester for Hydroacyloxylation of Unactivated Olefins^a

entry	starting materials	products	entry	starting materials	products	entry	starting materials	products
1			5			9		
2			6			10		
3			7			11 ^b		
4			8			12 ^c		

^aIsolation yield (0.25 mmol scale, 0.1 M). ^bConcentration = 0.03 M. ^cCatalyst 17 (3 mol %) was used.

Scheme 3. Plausible Reaction Mechanism



^aNMR yield using dimethylsulfone as the internal standard (0.25 mmol scale). ^bIsolation yield (0.25 mmol scale, 0.1 M).

irrelevant to the cyclization remained intact under the reaction condition. Therefore, cyclization should begin with the formation of an oxonium intermediate, followed by depro-

tection. The experimental results show that the protective group was trapped by 2,4,6-trimethylpyridine (collidine) to generate the alkyldipyridinium salt in the cases of MOM, benzyl,

and methyl groups, although the final product for the TBS group is still unclear (Scheme 3a).

The ligand optimization affected the yield of the desired product, as shown in Table 3, which sheds additional light on the reaction mechanism. Furthermore, the measurable enantioselectivity was observed when using a chiral Co catalyst **24**¹⁷ (Scheme 3b, not fully optimized). We propose that the Co complex could significantly interfere with the transition state of the C–O bond-forming step.

A proposed mechanism that is consistent with the experimental data is provided in Scheme 3c. In this mechanism, hydrogen atom transfer¹⁸ to generate the carbon radical intermediate **A**¹⁹ along with the recovery of the Co complex is plausible according to Shenvi's and Herzon's insightful discussions.^{12f,g} Notably in our mechanism, both single-electron oxidation of the carbon radical by the cationic Co species and intramolecular nucleophilic trapping by the oxygen atom could occur simultaneously to generate oxonium intermediate **B** via a concerted transition state.²⁰ Finally, the protective group (or proton in the case of an unprotected substrate) was transferred from intermediate **B** to 2,4,6-trimethylpyridine, with the formation of the desired product and coproduct.

3. CONCLUSION

We developed Co catalyzed intramolecular hydroalkoxylation and hydroacyloxylation to afford five- and six-membered saturated oxygen heterocycles. The mild reaction condition realized a broad substrate scope and excellent functional group tolerance. The use of the optimum catalyst **17** paved the way for the synthesis of medium rings by the hydrofunctionalization of unactivated olefins. It was found that the high reactivity of this Co catalyst system enabled the deprotective cyclization of protected alkenyl alcohol and alkenyl ester to directly afford oxygen heterocycles. The experimental evidence supported the concerted transition state of the bond-forming step involving a cationic Co complex. On the basis of the results of this study, improving the enantioselectivity is ongoing.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/jacs.6b05720.

Experimental procedures and analytical data (¹H and ¹³C NMR) for all new compounds
(PDF)

¹H and ¹³C NMR data of starting material
(PDF)

¹H, ¹³C NMR, and HPLC data of products
(PDF)

■ AUTHOR INFORMATION

Corresponding Author

*cgehis@musashino-u.ac.jp

Funding

This work was supported by JSPS KAKENHI Grant number 26860017, the Uehara memorial Foundation, Takeda Science Foundation.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We thank Kazuaki Katakawa (Musashino University) for Microwave apparatus. We also thank Nao Shimizu, Natsumi Koseki (Musashino University) for synthesizing some materials.

■ REFERENCES

- (1) Cossy, J. *Synthesis of Saturated Oxygenated Heterocycles I: 5- and 6-Membered Rings*; Springer: Paris, 2014.
- (2) Cossy, J. *Synthesis of Saturated Oxygenated Heterocycles II: 7- to 16-Membered Rings*; Springer: Paris, 2014.
- (3) Bauer, R. A.; Wenderski, T. A.; Tan, D. S. *Nat. Chem. Biol.* **2013**, *9*, 21.
- (4) Olefin activation with a metal catalyst: (a) Qian, H.; Han, X.; Widenhofer, R. A. *J. Am. Chem. Soc.* **2004**, *126*, 9536. (b) Yang, C. G.; Reich, N. W.; Shi, Z.; He, C. *Org. Lett.* **2005**, *7*, 4553. (c) Ito, Y.; Kato, R.; Hamashima, K.; Kataoka, Y.; Oe, Y.; Ohta, T.; Furukawa, I. *J. Organomet. Chem.* **2007**, *692*, 691. (d) Ohta, T.; Kataoka, Y.; Miyoshi, A.; Oe, Y.; Furukawa, I.; Ito, Y. *J. Organomet. Chem.* **2007**, *692*, 671. (e) Adrio, L. A.; Quek, L. S.; Taylor, J. G.; Kuok Hii, K. *Tetrahedron* **2009**, *65*, 10334. (f) Kena Diba, A.; Begouin, J.-M.; Niggemann, M. *Tetrahedron Lett.* **2012**, *53*, 6629. Olefin activation with a nonmetal catalyst: (g) Coulombel, L.; Duñach, E. *Green Chem.* **2004**, *6*, 499. (h) Coulombel, L.; Favier, I.; Duñach, E. *Chem. Commun.* **2005**, 2286. (i) Rosenfeld, D. C.; Shekhar, S.; Takemiya, A.; Utsunomiya, M.; Hartwig, J. F. *Org. Lett.* **2006**, *8*, 4179. (j) Fujita, S.; Abe, M.; Shibuya, M.; Yamamoto, Y. *Org. Lett.* **2015**, *17*, 3822. (k) Sultana, S.; Devi, N. R.; Saikia, A. K. *Asian J. Org. Chem.* **2015**, *4*, 1281. O-nucleophile activation: (l) Dzudza, A.; Marks, T. J. *Chem. - Eur. J.* **2010**, *16*, 3403. (m) Dzudza, A.; Marks, T. J. *Org. Lett.* **2009**, *11*, 1523. (n) Coulombel, L.; Rajzmann, M.; Pons, J.-M.; Olivero, S.; Duñach, E. *Chem. - Eur. J.* **2006**, *12*, 6356. (o) Coulombel, L.; Weíwer, M.; Duñach, E. *Eur. J. Org. Chem.* **2009**, *2009*, 5788. (p) Schuch, D.; Fries, P.; Dönges, M.; Pérez, B. M.; Hartung, J. *J. Am. Chem. Soc.* **2009**, *131*, 12918. (q) Kamiya, I.; Tsunoyama, H.; Tsukuda, T.; Sakurai, H. *Chem. Lett.* **2007**, *36*, 646. (r) Murayama, H.; Nagao, K.; Ohmiya, H.; Sawamura, M. *Org. Lett.* **2015**, *17*, 2039. Only one reaction of hydroalkoxylation: (s) Miller, Y.; Miao, L.; Hosseini, A. S.; Chemler, S. R. *J. Am. Chem. Soc.* **2012**, *134*, 12149. Others: (t) Marotta, E.; Foresti, E.; Marcelli, T.; Peri, F.; Righi, P.; Scardovi, N.; Rosini, G. *Org. Lett.* **2002**, *4*, 4451. (u) Komeyama, K.; Morimoto, T.; Nakayama, Y.; Takaki, K. *Tetrahedron Lett.* **2007**, *48*, 3259. (v) Singh, S. A.; Kabiraj, S.; Khandare, R. P.; Nalawade, S. P.; Upar, K. B.; Bhat, S. V. *Synth. Commun.* **2009**, *40*, 74. (w) Jeong, Y.; Kim, D.-Y.; Choi, Y.; Ryu, J.-S. *Org. Biomol. Chem.* **2011**, *9*, 374. (x) Schlüter, J.; Blazejak, M.; Hintermann, L. *ChemCatChem* **2013**, *5*, 3309. (y) Deka, M. J.; Indukuri, K.; Sultana, S.; Borah, M.; Saikia, A. K. *J. Org. Chem.* **2015**, *80*, 4349. (z) Notar Francesco, I.; Cacciuto, B.; Pucheault, M.; Antoniotti, S. *Green Chem.* **2015**, *17*, 837. (aa) Catti, L.; Tiefenbacher, K. *Chem. Commun.* **2015**, *51*, 892. (ab) Schlüter, J.; Blazejak, M.; Bock, F.; Hintermann, L. *Angew. Chem., Int. Ed.* **2015**, *54*, 4014. (5) (a) Chaminade, X.; Coulombel, L.; Olivero, S.; Duñach, E. *Eur. J. Org. Chem.* **2006**, *2006*, 3554. (b) Komeyama, K.; Mieno, Y.; Yukawa, S.; Morimoto, T.; Takaki, K. *Chem. Lett.* **2007**, *36*, 752. (c) Gooßen, L. J.; Ohlmann, D. M.; Dierker, M. *Green Chem.* **2010**, *12*, 197. (d) Sakuma, M.; Sakakura, A.; Ishihara, K. *Org. Lett.* **2013**, *15*, 2838. (e) Nagamoto, M.; Nishimura, T. *Chem. Commun.* **2015**, *51*, 13466. (6) (a) Hamilton, D. S.; Nicewicz, D. A. *J. Am. Chem. Soc.* **2012**, *134*, 18577. (b) Nicewicz, D.; Hamilton, D. *Synlett* **2014**, *25*, 1191. (7) (a) Brunel, Y.; Rousseau, G. *J. Org. Chem.* **1996**, *61*, 5793. (b) Verma, A.; Jana, S.; Durga Prasad, C.; Yadav, A.; Kumar, S. *Chem. Commun.* **2016**, *52*, 4179. (8) Cheng, Y. A.; Chen, T.; Tan, C. K.; Heng, J. J.; Yeung, Y.-Y. *J. Am. Chem. Soc.* **2012**, *134*, 16492. (9) (a) Shigehisa, H.; Aoki, T.; Yamaguchi, S.; Shimizu, N.; Hiroya, K. *J. Am. Chem. Soc.* **2013**, *135*, 10306. (b) Shigehisa, H.; Kikuchi, H.; Hiroya, K. *Chem. Pharm. Bull.* **2016**, *64*, 371. (10) Shigehisa, H.; Koseki, N.; Shimizu, N.; Fujisawa, M.; Niitsu, M.; Hiroya, K. *J. Am. Chem. Soc.* **2014**, *136*, 13534.

(11) Shigehisa, H.; Ano, T.; Honma, H.; Ebisawa, K.; Hiroya, K. *Org. Lett.* **2016**, DOI: 10.1021/acs.orglett.6b01662.

(12) Hydration: (a) Isayama, S.; Mukaiyama, T. *Chem. Lett.* **1989**, 18, 1071. Mechanism: (b) Tokuyasu, T.; Kunikawa, S.; Masuyama, A.; Nojima, M. *Org. Lett.* **2002**, 4, 3595. Recent examples: (c) Waser, J.; Gaspar, B.; Nambu, H.; Carreira, E. M. *J. Am. Chem. Soc.* **2006**, 128, 11693. (d) Ishikawa, H.; Colby, D. A.; Seto, S.; Va, P.; Tam, A.; Kakei, H.; Rayl, T. J.; Hwang, I.; Boger, D. L. *J. Am. Chem. Soc.* **2009**, 131, 4904. (e) Crossley, S. W. M.; Barabé, F.; Shenvi, R. A. *J. Am. Chem. Soc.* **2014**, 136, 16788. (f) Iwasaki, K.; Wan, K. K.; Oppedisano, A.; Crossley, S. W. M.; Shenvi, R. A. *J. Am. Chem. Soc.* **2014**, 136, 1300. (g) King, S. M.; Ma, X.; Herzon, S. B. *J. Am. Chem. Soc.* **2014**, 136, 6884. (h) Lo, J. C.; Yabe, Y.; Baran, P. S. *J. Am. Chem. Soc.* **2014**, 136, 1304. (i) Obradors, C.; Martinez, R. M.; Shenvi, R. A. *J. Am. Chem. Soc.* **2016**, 138, 4962. (j) Shigehisa, H.; Nishi, E.; Fujisawa, M.; Hiroya, K. *Org. Lett.* **2013**, 15, 5158. (k) Shigehisa, H. *Synlett* **2015**, 26, 2479.

(13) Cacciuttolo, B.; Poulain-Martini, S.; Duñach, E. *Eur. J. Org. Chem.* **2011**, 2011, 3710.

(14) *N*-Fluoro-2,4,6-trimethylpyridinium trifluoromethanesulfonate was used instead of tetrafluoroborate to avoid hydrofluorination. See ref 12j.

(15) The desired bond cleavage is hypothesized to not occur from the postulated oxonium intermediate in the case of TBS and benzyl groups, whereas C–O bond cleavage is assisted by a lone pair of electrons on the oxygen atom of the MOM group.

(16) Use of NFPY·OTf instead of Me₃NFPY·OTf did not improve the yield of the desired product (45% yield with NFPY·OTf).

(17) Nagata, T.; Yorozu, K.; Yamada, T.; Mukaiyama, T. *Angew. Chem., Int. Ed. Engl.* **1995**, 34, 2145.

(18) A deuterium experiment was conducted to determine the source of terminal hydrogen. See ref 9a.

(19) The existence of a carbon radical intermediate was supported by radical clock and O₂ uptake experiments. See ref 9a.

(20) Zhu, R.; Buchwald, S. L. *Angew. Chem., Int. Ed.* **2013**, 52, 12655.