

# Mechanism-Guided Development of a Highly Active Bis-thiourea Catalyst for Anion-Abstraction Catalysis

C. Rose Kennedy,<sup>‡</sup> Dan Lehnher,<sup>‡</sup> Naomi S. Rajapaksa, David D. Ford, Yongho Park, and Eric N. Jacobsen<sup>\*</sup>


Department of Chemistry and Chemical Biology, Harvard University, Cambridge, Massachusetts 02138, United States

**S** Supporting Information

**ABSTRACT:** We describe the rational design of a linked, bis-thiourea catalyst with enhanced activity relative to monomeric analogues in a representative enantioselective anion-abstraction reaction. Mechanistic insights guide development of this linking strategy to favor substrate activation through the intramolecular cooperation of two thiourea subunits while avoiding nonproductive aggregation. The resulting catalyst platform overcomes many of the practical limitations that have plagued hydrogen-bond donor catalysis and enables use of catalyst loadings as low as 0.05 mol %. Computational analyses of possible anion-binding modes provide detailed insight into the precise mechanism of anion-abstraction catalysis with this pseudo-dimeric thiourea.

Chiral dual hydrogen-bond (H-bond) donors such as urea, thiourea, and squaramide derivatives comprise an important class of enantioselective catalysts for reactions proceeding through ion-pair intermediates.<sup>1,2</sup> While these air- and moisture-stable organocatalysts have enabled the development of numerous highly enantioselective transformations, these methods typically suffer from several practical constraints that impede widespread application. These include requirements for high catalyst loadings (5–20 mol %), long reaction times ( $\geq 24$  h), and dilute reaction conditions ( $\leq 0.1$  M in substrate) to achieve maximal levels of enantioinduction.<sup>3</sup>

In an effort to understand the basis for these limitations and thereby develop strategies to overcome them, we recently undertook a detailed mechanistic study of a representative transformation involving anion-abstraction catalysis by a chiral amido-thiourea (**1**, Figure 1A).<sup>2c,4,5</sup> This analysis revealed that thiourea catalyst **1** forms nonproductive dimeric aggregates in the resting state; these aggregates must dissociate before two molecules of catalyst recombine with substrates to promote rate- and enantioselectivity-determining C–C bond formation (Figure 1B).<sup>4a</sup> We hypothesized that appropriate linkage of the monomers should result in stabilization of the dominant transition-state complex and, therefore, enhanced catalytic activity. The linked units would also need to adopt a relative orientation closely resembling that accessed by the untethered monomers in order to promote reactions with similar levels of enantioselectivity. Analogous linking strategies have afforded reactivity enhancements in other systems displaying a bimolecular dependence on chiral catalyst.<sup>6</sup> However, this case poses an additional challenge, because the tether must also



**Figure 1.** Anion-abstraction catalysis involving the cooperative action of two amido-thiourea catalysts. Ar = 4-fluorophenyl, Ar<sup>F</sup> = 3,5-bis(trifluoromethyl)phenyl.


disfavor unproductive ground-state aggregation. Herein, we describe the mechanism-guided development of a highly active bis-thiourea for anion-abstraction catalysis. In addition to providing a platform for future catalyst design, the linking strategy validates computational analyses shedding light on the cooperative mechanism of anion abstraction.

While the cooperative action of thioureas such as **1** in anion-abstraction catalysis has been well established, uncertainty remains regarding the precise mode of electrophile activation by two catalyst units. Two different cooperative arrangements, which we term 2H- and 4H-anion abstraction (Figure 1C), are consistent with experimental data and kinetically indistinguishable.<sup>7</sup> Furthermore, preliminary computational analyses of [1]<sub>2</sub>·Cl<sup>−</sup> complexes using density functional theory (DFT)


Received: September 1, 2016

Published: October 5, 2016


## A. Dimeric Catalyst Complexes in the Solid State



## C. Computed Transition Structures with Monomeric and Linked Catalysts



## B. Design of Linked Bis-Thiourea Catalysts



**Figure 2.** Identification of a linking strategy for bis-thiourea anion-abstraction catalysts. Carbon atoms represented in silver (for **5**), slate blue (for **6**), or light pink (for **7**). Linkage sites highlighted in cyan. Carbon-bound hydrogen atoms omitted for clarity. N = blue, O = red, F = lime, Si = gold, S = yellow, Cl = magenta. Computations performed with B3LYP-D3(BJ)/6-31G(d,p)/PCM(toluene)//B3LYP/6-31G(d). Corrected free energies reported relative to the lowest energy transition state identified with each catalyst. See [Supporting Information](#) for additional discussion.


did not reveal a clear energetic preference for one over the other.<sup>4</sup> We postulated that dimeric catalysts designed to enable  $\text{Cl}^-$  abstraction through only one of these two arrangements could help reveal which of the two activation modes is operative in the enantioselective alkylation of  $\alpha$ -chloroisochroman and in related anion-binding pathways promoted by this class of chiral H-bond donor. Given that these catalysts have been shown to form  $[\text{cat}]_2 \cdot \text{NMe}_4\text{Cl}$  complexes with a 4H- $\text{Cl}^-$ -binding geometry in the solid state (Figure 2A),<sup>4,8</sup> we chose to target linkage strategies that would favor the 4H-activation mode. Introduction of a methyl substituent in the arylpyrrolidine component of **1** has been demonstrated to confer improved reactivity and enantioselectivity by minimizing competing reaction pathways,<sup>8</sup> and accordingly, we used this modified structure (**5a**) as the starting point for further catalyst design.

In order to link two units of **5a** in a manner that might stabilize the reactive conformation of the resulting dimer selectively, we sought to identify linkage points that would allow access to the 4H- $\text{Cl}^-$ -bound complex but disfavor formation of the inactive, self-aggregated complex.<sup>9</sup> The structures of both the nonproductive aggregate (**5a**•**5a**) and a 4H- $\text{Cl}^-$ -bound complex ( $[\text{5a}]_2 \cdot \text{NMe}_4\text{Cl}$ ) have been characterized by X-ray diffraction.<sup>4,8,10</sup> We noted that, in the solid state, the distance between the backbone *tert*-butyl group of one molecule and the aryl thiourea of the other is considerably greater in the nonproductive aggregate than in the  $\text{Cl}^-$ -bound complex (Figure 2A). This observation suggested that linkage

through these sites might fulfill the intended purpose. We deemed that an appropriate linker between these two sites would need to meet the following stringent criteria: (1) it should cause minimal structural and electronic perturbation to the key catalyst features identified in the original optimization efforts;<sup>2c</sup> (2) it must be long enough to allow the same 4H-binding arrangement that is accessible to the unlinked monomers; but, (3) it must be short enough to prevent intramolecular formation of the nonproductive aggregate.

The relative disposition of the **5b** units in the lowest energy rate- and enantiodetermining 4H-alkylation transition structure predicted using DFT was found to be very similar to that in the  $\text{Cl}^-$ -bound complex characterized in the solid state.<sup>8,11–13</sup> We used the computed transition structure to evaluate a variety of hypothetical linked structures according to the criteria outlined above. Through this analysis, pseudo-dimeric thiourea catalyst **6b**, which bears a 5-atom spacer between the amino acid side chain of one thiourea unit and the aryl group of another, was modeled and found to access 4H-transition structures that overlay almost perfectly with the transition structure calculated with **5b** (Figure 2C, left). In contrast, **7b**, which possesses a linker just one methylene unit shorter, was predicted to be incapable of achieving similar conformations (Figure 2C, right). Both catalysts were selected for experimental assessment.

Bis-thioureas **6a** and **7a**, along with their monomeric analogues **5c** and **5d**, were synthesized and compared with **1** and **5a** as catalysts for the alkylation of  $\alpha$ -chloroisochroman with silyl ketene acetal **3b**. While modification of the *tert*-




**Figure 3.** (A) Alkylation of  $\alpha$ -chloroisochroman with **3b** catalyzed by mono- (**1** and **5a,c,d** at 0.5 mol %) and bis-thioureas (**6a** and **7a** at 0.25 mol %) at low loading. Relative rates at 15% conversion. B. Alkylation exhibits a first-order dependence on catalyst **6a** with dramatic rate acceleration over catalyst **1**. See ref 4a. Rates at 30% conversion. C. The enantiomeric excess of product **4a** shows a linear dependence on the ee of bis-thiourea **6a** (0.25 mol %); this stands in contrast to the nonlinear dependence on the ee of catalyst **1** (0.50 mol %).

leucine side chain (as in **5c**) has a negligible effect on enantioselectivity, replacement of the trifluoromethyl group with an ester (as in **5d**) does have a slightly detrimental impact (Figure 3A). Nonetheless, catalyst **6a** displays markedly enhanced reactivity, affording high conversion with a low catalyst loading and short reaction time, while preserving the enantioselectivity observed with **5a**. In contrast, catalyst **7a** exhibits very similar reactivity to the monomeric analogues, consistent with the prediction that the 4-atom linker is too short to enable intramolecular formation of the optimal, cooperative transition structure.

The improved efficiency of catalyst **6a** allows us to address several of the practical limitations presented by the conditions optimized for catalyst **1**. With just 0.1 mol % **6a**, the alkylation of  $\alpha$ -chloroisochroman with silyl ketene acetal **3a** can be run on preparatory scale under relatively concentrated conditions (0.5 M initial substrate concentration) to afford **4a** with excellent yield and enantioselectivity after just 3 h (Scheme 1). This

#### Scheme 1. Improved Volumetric Throughput on Preparatory Scale



stands in stark contrast to the original conditions, in which a 24 h reaction time with 10 mol % **1** at 0.1 M is required to achieve good yield with the same level of enantioinduction.<sup>2c</sup>

The improved efficiency observed with pseudo-dimeric catalyst **6a** provides circumstantial evidence for the validity of the cooperative mechanism for anion-abstraction catalysis outlined in Figure 1B. In order to establish rigorously whether the two linked thiourea subunits act cooperatively within the same molecule, the kinetic dependence of alkylation rate on catalyst concentration was determined by reaction progress kinetic analysis using in situ infrared spectroscopy (Figure 3B).<sup>4a,14</sup> The reaction exhibits a strict first-order dependence on catalyst at low [**6a**], under conditions where a second-order dependence on [**1**] is observed.<sup>4a</sup> The increased activity of **6a** relative to monomeric catalyst **1** is therefore magnified at very low catalyst loadings, and as little as 0.05 mol % **6a** can be

employed while maintaining useful reaction rates. Finally, an absence of a nonlinear effect is observed with catalyst **6a** at low catalyst concentrations (Figure 3C).<sup>15,16</sup> Taken together, these results provide strong evidence that the thiourea subunits in catalyst **6a** cooperate in an intramolecular fashion to promote oxocarbenium alkylation.

Furthermore, the successful development of a highly active pseudo-dimeric catalyst designed around the 4H-activation mode suggests that the 4H-mechanism is most likely operative under the conditions described above. To evaluate whether the 2H-activation mode is also accessible to the dimeric catalysts, the intermediate catalyst-oxocarbenium-Cl<sup>-</sup> complexes and the transition structures for oxocarbenium alkylation promoted by the 2H- and 4H-arrangements of catalyst **6b** were examined computationally. Across a variety of functionals, both the intermediate and transition structures in which the bis-thiourea was forced to access a 2H-geometry were predicted to be significantly higher in energy than the alternative 4H-structures. (See Supporting Information for additional discussion.)

In conclusion, the design of highly active and efficient anion-abstraction catalyst **6a** represents the culmination of a series of mechanistic insights. The mechanism-driven design strategy has afforded a linked, bis-thiourea catalyst system that overcomes many of the practical limitations traditionally associated with H-bond-donor-mediated anion-abstraction catalysis. Key insights enabled formation of a pseudo-dimeric complex for cooperative activation of a chloroether electrophile by a 4H-mechanism while disfavoring nonproductive ground-state aggregation. The high efficiencies enabled by this catalyst design hold promise for application of dual hydrogen-bond-donor-mediated anion-abstraction catalysis to historically challenging transformations. In this vein, the development of highly stereoselective reactions with tertiary alkyl chloride and glycosyl chloride electrophiles is the focus of our ongoing attention.

## ■ ASSOCIATED CONTENT

### Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/jacs.6b09205.

Experimental procedures; spectroscopic data; kinetic data (tabulated and graphical); discussion, geometries, and energies of calculated stationary points; summary of alternative linking strategies (PDF)



X-ray crystallographic structures for synthetic intermediates and select urea and thiourea catalysts (also deposited as CCDC 1478174, 1482963–1482976, and 1501460) (CIF)

## AUTHOR INFORMATION

### Corresponding Author

\*jacobson@chemistry.harvard.edu

### Author Contributions

‡C.R.K. and D.L. contributed equally.

### Notes

The authors declare no competing financial interest.

## ACKNOWLEDGMENTS

This work was supported by the NIH (GM-43214) and through fellowships to C.R.K. (NSF, DGE1144152), D.L. (NSERC PDF), N.S.R. (Boehringer Ingelheim), and D.D.F. (Eli Lilly). The authors thank Dr. Shao-Liang Zheng (Harvard X-ray Laboratory) for collection and refinement of X-ray crystallographic data, Dr. Robert Knowles (Princeton) for early crystallographic analyses, Dr. Alison E. Wendlandt and Andrew J. Bendelsmith (Harvard) for synthetic assistance, and Daniel A. Strassfeld (Harvard) for helpful discussion.

## REFERENCES

- (1) For reviews, see: (a) Zhang, Z.; Schreiner, P. R. *Chem. Soc. Rev.* **2009**, *38*, 1187–1198. (b) Brak, K.; Jacobsen, E. N. *Angew. Chem., Int. Ed.* **2013**, *52*, 534–561. (c) Phipps, R. J.; Hamilton, G. L.; Toste, F. D. *Nat. Chem.* **2012**, *4*, 603–614. (d) Beckendorf, S.; Asmus, S.; Mancheño, O. G. *ChemCatChem* **2012**, *4*, 926–936. (e) Mahlau, M.; List, B. *Angew. Chem., Int. Ed.* **2013**, *52*, 518–533. (f) Seidel, D. *Synlett* **2014**, *25*, 783–794.
- (2) For select examples of enantioselective anion-binding catalysis with dual H-bond donors, see: (a) Taylor, M. S.; Tokunaga, N.; Jacobsen, E. N. *Angew. Chem., Int. Ed.* **2005**, *44*, 6700–6704. (b) Raheem, I. T.; Thiara, P. V.; Peterson, E. A.; Jacobsen, E. N. *J. Am. Chem. Soc.* **2007**, *129*, 13404–13405. (c) Reisman, S. E.; Doyle, A. G.; Jacobsen, E. N. *J. Am. Chem. Soc.* **2008**, *130*, 7198–7199. (d) Klausen, R. S.; Jacobsen, E. N. *Org. Lett.* **2009**, *11*, 887–890. (e) De, C. K.; Klauber, E. G.; Seidel, D. *J. Am. Chem. Soc.* **2009**, *131*, 17060–17061. (f) Xu, H.; Zuend, S. J.; Woll, M. G.; Tao, Y.; Jacobsen, E. N. *Science* **2010**, *327*, 986–990. (g) Knowles, R. R.; Lin, S.; Jacobsen, E. N. *J. Am. Chem. Soc.* **2010**, *132*, 5030–5032. (h) Brown, A. R.; Kuo, W.-H.; Jacobsen, E. N. *J. Am. Chem. Soc.* **2010**, *132*, 9286–9288. (i) Burns, N. Z.; Witten, M. G.; Jacobsen, E. N. *J. Am. Chem. Soc.* **2011**, *133*, 14578–14581. (j) Lin, S.; Jacobsen, E. N. *Nat. Chem.* **2012**, *4*, 817–824. (k) Schafer, A. G.; Wieting, J. M.; Fisher, T. J.; Mattson, A. E. *Angew. Chem., Int. Ed.* **2013**, *52*, 11321–11324. (l) Metz, A. E.; Ramalingam, K.; Kozlowski, M. C. *Tetrahedron Lett.* **2015**, *56*, 5180–5184. (m) Mittal, N.; Lippert, K. M.; De, C. K.; Klauber, E. G.; Emge, T. J.; Schreiner, P. R.; Seidel, D. *J. Am. Chem. Soc.* **2015**, *137*, 5748–5758. (n) Hardman-Baldwin, A. M.; Visco, M. D.; Wieting, J. M.; Stern, C.; Kondo, S.; Mattson, A. E. *Org. Lett.* **2016**, *18*, 3766–3769.
- (3) A few noteworthy exceptions have been reported. For select examples of highly efficient urea or thiourea-catalyzed transformations, see: (a) Kotke, M.; Schreiner, P. R. *Tetrahedron* **2006**, *62*, 434–439. (b) Kotke, M.; Schreiner, P. R. *Synthesis* **2007**, *2007*, 779–790. (c) Zuend, S. J.; Coughlin, M. P.; Lalonde, M. P.; Jacobsen, E. N. *Nature* **2009**, *461*, 968–970. (d) Birrell, J. E.; Desrosiers, J.-N.; Jacobsen, E. N. *J. Am. Chem. Soc.* **2011**, *133*, 13872–13875.
- (4) (a) Ford, D. D.; Lehnher, D.; Kennedy, C. R.; Jacobsen, E. N. *J. Am. Chem. Soc.* **2016**, *138*, 7860–7863. (b) Ford, D. D.; Lehnher, D.; Kennedy, C. R.; Jacobsen, E. N. *ACS Catal.* **2016**, *6*, 4616–4620.
- (5) For representative reports in which novel anion-binding catalysts were validated using the alkylation of  $\alpha$ -chloroisochroman with silyl

ketene acetal nucleophiles as a benchmark reaction, see: (a) Kniep, F.; Jungbauer, S. H.; Zhang, Q.; Walter, S. M.; Schindler, S.; Schnapperelle, I.; Herdtweck, E.; Huber, S. M. *Angew. Chem., Int. Ed.* **2013**, *52*, 7028–7032. (b) Jungbauer, S. H.; Huber, S. M. *J. Am. Chem. Soc.* **2015**, *137*, 12110–12120.

(6) (a) Konsler, R. G.; Karl, J.; Jacobsen, E. N. *J. Am. Chem. Soc.* **1998**, *120*, 10780–10781. (b) Ready, J. M.; Jacobsen, E. N. *J. Am. Chem. Soc.* **2001**, *123*, 2687–2688. (c) Ready, J. M.; Jacobsen, E. N. *Angew. Chem., Int. Ed.* **2002**, *41*, 1374–1377. (d) Denmark, S. E.; Fu, J. *J. Am. Chem. Soc.* **2003**, *125*, 2208–2216. (e) Nakano, K.; Hashimoto, S.; Nozaki, K. *Chem. Sci.* **2010**, *1*, 369–373. (f) Kalow, J.; Doyle, A. G. *J. Am. Chem. Soc.* **2011**, *133*, 16001–16012. (g) Klimczyk, S.; Misale, A.; Huang, X.; Maulide, N. *Angew. Chem., Int. Ed.* **2015**, *54*, 10365–10369.

(7) For examples in which intramolecular H-bonding has been proposed to increase the “2H” anion- or neutral Lewis base-binding ability of a urea or thiourea catalyst, see: (a) Jones, C. R.; Pantoş, G. D.; Morrison, A. J.; Smith, M. D. *Angew. Chem., Int. Ed.* **2009**, *48*, 7391–7394. (b) Probst, N.; Madarász, Á.; Valkonen, A.; Pápai, I.; Rissanen, K.; Neuvonen, A.; Pihko, P. M. *Angew. Chem., Int. Ed.* **2012**, *51*, 8495–8499. (c) ref 2m.

(8) While **1** exists as a slowly interconverting mixture of amide rotamers, **5a** exists exclusively in the more active and enantioselective (*Z*)-rotameric form. This constraint has been demonstrated to lead to improved activity and enantioselectivity in the alkylation of  $\alpha$ -chloroisochroman. See: Lehnher, D.; Ford, D. D.; Bendelsmith, A. J.; Kennedy, C. R.; Jacobsen, E. N. *Org. Lett.* **2016**, *18*, 3214–3217.

(9) The extent of this challenge was further illustrated by the poor efficiencies of symmetrically linked catalysts. See [Supporting Information](#) for details.

(10) The solution-state structures of the nonproductive aggregates **1•1** and **5a•5a** have also been determined via DFT-aided 2D NMR analysis; see refs 4a and 8. These structures are similar to the solid-state structures and position the backbone *tert*-butyl group of one molecule distal from the aryl thiourea of the other.

(11) Catalyst **5b** provides a significantly reduced computational cost relative to catalyst **5a**. For experimental validation, see ref 8.

(12) Frisch, M. J.; et al. *Gaussian 09*, Revision D.02; Gaussian Inc.: Wallingford, CT, 2009; see [Supporting Information](#) for full citation.

(13) (a) Becke, A. D. *J. Chem. Phys.* **1993**, *98*, 5648–5652. (b) Lee, C.; Yang, W.; Parr, R. G. *Phys. Rev. B: Condens. Matter Mater. Phys.* **1988**, *37*, 785–789. (c) Miehlich, B.; Savin, A.; Stoll, H.; Preuss, H. *Chem. Phys. Lett.* **1989**, *157*, 200–206. (d) Frisch, M. J.; Pople, J. A.; Binkley, J. S. *J. Chem. Phys.* **1984**, *80*, 3265–3269. (e) Grimme, S.; Antony, J.; Ehrlich, S.; Krieg, H. *J. Chem. Phys.* **2010**, *132*, 154104.

(14) (a) Blackmond, D. G. *Angew. Chem., Int. Ed.* **2005**, *44*, 4302–4320. (b) Blackmond, D. G. *J. Am. Chem. Soc.* **2015**, *137*, 10852–10866.

(15) (a) Puchot, C.; Samuel, O.; Duñach, E.; Zhao, S.; Agami, C.; Kagan, H. B. *J. Am. Chem. Soc.* **1986**, *108*, 2353–2357. (b) Guillaneux, D.; Zhao, S.; Samuel, O.; Rainford, D.; Kagan, H. B. *J. Am. Chem. Soc.* **1994**, *116*, 9430–9439. (c) Satyanarayana, T.; Abraham, S.; Kagan, H. B. *Angew. Chem., Int. Ed.* **2009**, *48*, 456–494.

(16) A nonlinear effect would be expected if **6a** were undergoing self-aggregation to a measurable extent under the catalytic conditions, or if **6a** achieves cooperative anion abstraction through an intermolecular pathway. In the latter scenario, the magnitude of the nonlinear effect would depend on the degree of stereochemical communication between the thiourea subunits in the ee-determining transition structure.