

Critical Aspects of Heme–Peroxo–Cu Complex Structure and Nature of Proton Source Dictate Metal–O_{peroxo} Breakage versus Reductive O–O Cleavage Chemistry


Suzanne M. Adam,[†] Isaac Garcia-Bosch,^{†,§} Andrew W. Schaefer,[‡] Savita K. Sharma,[†] Maxime A. Siegler,[†] Edward I. Solomon,^{*,‡} and Kenneth D. Karlin^{*,†}

[†]Johns Hopkins University, Baltimore, Maryland 21218, United States

[‡]Department of Chemistry, Stanford University, Stanford, California 94305, United States

Supporting Information

ABSTRACT: The 4H⁺/4e⁻ reduction of O₂ to water, a key fuel-cell reaction also carried out in biology by oxidase enzymes, includes the critical O–O bond reductive cleavage step. Mechanistic investigations on active-site model compounds, which are synthesized by rational design to incorporate systematic variations, can focus on and resolve answers to fundamental questions, including protonation and/or H-bonding aspects, which accompany electron transfer. Here, we describe the nature and comparative reactivity of two low-spin heme–peroxo–Cu complexes, **LS-4DCHIm**, [(DCHIm)F₈Fe^{III}-(O₂²⁻)-Cu^{II}(DCHIm)₄]⁺, and **LS-3DCHIm**, [(DCHIm)F₈Fe^{III}-(O₂²⁻)-Cu^{II}(DCHIm)₃]⁺ (F₈ = tetrakis(2,6-difluorophenyl)-porphyrinate; DCHIm = 1,5-dicyclohexylimidazole), toward different proton (4-nitrophenol and [DMF·H⁺](CF₃SO₃⁻)) (DMF = dimethylformamide) or electron (decamethylferrocene (Fc*)) sources. Spectroscopic reactivity studies show that differences in structure and electronic properties of **LS-3DCHIm** and **LS-4DCHIm** lead to significant differences in behavior. **LS-3DCHIm** is resistant to reduction, is unreactive toward weakly acidic 4-NO₂-phenol, and stronger acids cleave the metal–O bonds, releasing H₂O₂. By contrast, **LS-4DCHIm** forms an adduct with 4-NO₂-phenol, which includes an H-bond to the peroxo O-atom distal to Fe (resonance Raman (rR) spectroscopy and DFT). With addition of Fc* (2 equiv overall required), O–O reductive cleavage occurs, giving water, Fe(III), and Cu(II) products; however, a kinetic study reveals a one-electron rate-determining process, $k_{et} = 1.6 \text{ M}^{-1} \text{ s}^{-1}$ (–90 °C). The intermediacy of a high-valent [(DCHIm)F₈Fe^{IV}=O] species is thus implied, and separate experiments show that one-electron reduction-protonation of [(DCHIm)F₈Fe^{IV}=O] occurs faster ($k_{et2} = 5.0 \text{ M}^{-1} \text{ s}^{-1}$), consistent with the overall postulated mechanism. The importance of the H-bonding interaction as a prerequisite for reductive cleavage is highlighted.



INTRODUCTION

Because of ever-growing global environmental and economic concerns, researchers are pushing to find alternative and, ideally, renewable energy sources to fill the needs of the modern generation.¹ In nature, energy is consumed to make chemical bonds and released when those bonds are broken. Importantly, this concept can be exhibited by the natural processes of photosynthesis and cellular respiration, which make up the life-sustaining oxygen cycle: $\text{O}_2 + 4\text{e}^- + 4\text{H}^+ \rightleftharpoons 2\text{H}_2\text{O}$.^{2,3} Here, there exists a critical interplay between acid–base and redox events⁴ including transformations of high-valent metal–oxo species.^{5,6} These processes have inspired chemists to design and study synthetic constructs that resemble active-site species and/or duplicate function, which can lead to new fundamental insights and the generation of practical systems.^{7–12} Advancements involving Photosystem II-inspired model systems for the purposes of light harvesting and/or

water oxidation to O₂ quickly rise to the forefront of popular science due to their potential applications in solar energy conversion.¹³ However, to complete the biochemical oxygen cycle, the O–O bond is cleaved during cellular respiration to give back water, and the mechanistic similarities of how redox-active transition metal ions make or break the strong O–O bond in these two natural systems³ make Cytochrome *c* Oxidase (CcO) an extremely important enzyme. For this reason, considerable research efforts have been undertaken to understand the mechanistic details of dioxygen reduction, which occurs at the active site.

CcO's heterobinuclear active site's four-electron/four-proton reduction of O₂ to water (Figure 1) is accompanied by membrane proton-translocation leading to ATP synthesis.¹⁴

Received: October 31, 2016

Published: December 28, 2016




Figure 1. Net 4H⁺/4e⁻ O₂ reduction reaction by CcO (top), and the proposed mechanistic involvement of a hydrogen-bonding network in the active site to promote O–O cleavage (bottom).

While there is a consensus about many aspects of the CcO enzymatic mechanism, debate remains concerning the identity of certain intermediates and the details of the process triggering the O–O bond cleavage event, where the release of destructive, partially reduced oxygen species (i.e., H₂O₂, ·OH) is prevented.¹⁵ It has been proposed and widely accepted^{16,17} that a hydrogen-bonding network in the active site provides a channel from the cross-linked tyrosine (Tyr) residue to the putative bridging peroxo moiety (~5 Å away) and facilitates a proton-coupled electron transfer (PCET), which cleaves the O–O bond (Figure 1).^{16,18–22}

(Bio)chemical systems require protons to promote O–O reductive cleavage; however, the specific nature of an acid moiety (e.g., identity, proximity to the O–O moiety, pK_a, H-bonding capability, and number of protons) is crucial.^{23–25} To specifically facilitate O–O heterolytic cleavage in a heme-(hydroperoxo) intermediate, cytochrome P450 monooxygenases employ an acid–alcohol residue pair, located adjacent to the Fe–O–O distal O-atom to provide a key H-bond and proton source. If that H-bonding group is changed (e.g., by mutation), uncoupling occurs, and the outcome is Fe–O cleavage and H₂O₂ production.^{26–28} Nonheme iron and copper proteins are also found to have specific requirements (e.g., proximity, acid strength) to control proton-promoted O–O reductive cleavage chemistries.^{29–32}

A few synthetic heme–Cu complexes can reduce O₂ to water via a (su)peroxo intermediate,^{33–35} yet neither in these cases, nor in CcO, is the nature of H-bonding or protonation reactivity understood with respect to O–O cleavage. To provide the insights and fundamentals needed for this very important reaction, several different approaches can be taken. Studies on a diverse, ever-growing family of model complexes (both synthetic coordination complexes^{11,36–38} and modified protein systems^{12,22,23,39}) have aided researchers in gaining a fundamental understanding about certain aspects of the chemical mechanism of O₂ reduction.^{12,14} Our research group has for many years been systematically studying the O₂ adducts of synthetic model systems in which we can define or control certain aspects and monitor the effects of varying coordination environments/geometries, redox properties, sterics of the Cu and/or iron centers, and nature of a given substrate on the outcomes of different reactions. The bridging heme–peroxo–copper formulation depicted in Figure 1 (center), while not (yet) observed in the enzyme, is considered by some^{11,16,40–42} as a likely intermediate, and several recent computational studies invoke the likelihood of a peroxo-level intermediate in CcO.^{43–45} A heme–peroxo–Cu complex is a convenient starting point from which to study how slight changes in

structure can tune the functional O₂-reduction capabilities of synthetic model complexes.⁴⁶ Alternatively, it is also relevant to consider factors (chemical or structural) that lead to the affiliated nonredox reaction where protonation of O₂-derived atom(s) results in M–O cleavage and release of H₂O₂, although this pathway is importantly avoided in the enzymatic mechanism.

For CcO, or for a given efficient molecular catalyst able to promote four-electron, four-proton O₂-reduction to water ($E^{\circ}_{\text{MeCN}} = 1.21 \text{ V vs Fc}^{+/0}$),⁴⁷ these particular nuances relevant to the promotion of the O–O cleavage process need to be delineated and understood within the framework of reduction/protonation chemistry. The breadth of these PCET studies involves probing different (or perhaps simultaneous) orderings of the events of (i) electron transfer (from Fe, Cu, or the Tyr phenol moiety), and (ii) proton transfer derived from different sources. Yet, it is surely only by investigation of chemical model constructs and/or computational inquiries, where one has the ability to break down such a complicated overall process into individual steps, so that deep fundamental insights can be obtained.^{48,49} The latter are in the future highly relevant to a broad range of fields, that is, (i) the understanding of mechanisms of action of other synthetic and/or metalloenzyme O₂-activating systems, or (ii) the design of practical catalysts for O₂-reduction or substrate oxidation/oxygenation by O₂-derived metal–oxy complexes.

In this report, we investigate the fundamental interactions of model complexes with exogenous substrates as they relate to the enzymatic CcO mechanism, and insights into the structure–function relationship in the O–O versus M–O cleavage pathways are evaluated. Specifically, we detail how different acids and a reductant behave toward two similar model heme-(O₂²⁻)-Cu complexes, LS-3DCHIm and LS-4DCHIm (Figure 2). Such strikingly similar complexes show an interesting difference in their reactivities toward proton and electron sources. In the case of LS-4DCHIm, only when treated with relatively weak acid (4-NO₂-phenol) and reductant (Fc*) is the coveted O–O cleavage reaction accomplished. Spectroscopic and kinetic evidence points toward a phenolic hydrogen-bond-associated intermediate, which triggers the electron transfer from Fc* necessary for O–O scission; a strong acid source instead leads to H₂O₂ release. Interestingly, the related complex, LS-3DCHIm (see Figure 2), shows no reactivity toward the same substrates, and with the support of DFT calculations, we believe this difference in reactivity can be attributed to steric factors and the degree to which the weakly acidic, phenolic substrate can access the bridging peroxo moiety to create a critical, activated H-bonded adduct. We are very interested in the chemistry of metal–oxy interactions with H⁺ and e⁻ sources of varying strengths and the insights these reactions may provide into O–O activation and cleavage chemistry in general.⁵⁰

RESULTS AND DISCUSSION

Reactivity with Fc* and/or Strong Acid. The peroxo complex, LS-4DCHIm, which has been previously characterized,⁵¹ is stable at –90 °C in 2-methyltetrahydrofuran (MeTHF), even in the presence of excess (20 equiv) reductant, Fc* ($E^{\circ}_{\text{MeTHF}} = -270 \text{ mV vs Ag}^{+/0}$).⁵² However, when strong acid, [DMF·H⁺](CF₃SO₃⁻) (pK_a(CH₃CN) = 6.1),⁵³ is added to a solution of LS-4DCHIm, a reaction is immediately observed in the UV–vis spectrum. Rapid disappearance of the low energy absorption features associated with peroxo-to-Fe




Figure 2. In situ generation of low-spin peroxo complexes, LS-3DCHIm and LS-4DCHIm, derived from our “naked” high-spin heme-peroxo-Cu complex, HS-3MeTHF,⁵¹ showing their reactivities toward strong acid ($H^+ = [DMF \cdot H^+](CF_3SO_3^-)$) and electron sources alone, that is, in the absence of proton sources.

CT bands⁵¹ further suggests that the peroxo-bridged structure is no longer intact. In fact, chemical and spectroscopic interrogation of the product mixture reveals that H_2O_2 and mononuclear iron and copper species are produced (Figure 2). Employing the quantitative horseradish peroxidase spectrophotometric assay^{52,54} verifies that 95% of the expected H_2O_2 was released, meaning that both M–O bonds were cleaved via protonation (Figure 2) (see Experimental Section and Table S1). An EPR spectrum of the reaction product solution indicates full formation of a copper(II)-imidazolyl species (see the Supporting Information for further explanation), and a predominantly (>95%) low-spin heme-imidazolyl product (Figure 2, see also Figures S2–4), confirming that the dinuclear complex has come apart.

Thus, the occurrence of metal–oxygen bond cleavage is the result of simple acid–base chemistry. Reductive cleavage of an O–O bond requires protons and electrons, however, Fc^* does not reduce the H_2O_2 present in the product mixture (Figure 2), nor does it reduce the iron(III) or Cu(II) species present; that is, the one-electron oxidized byproduct of a reduction event, the ferrocenium cation Fc^{*+} , is not formed, as would be observed by an increase in absorption at 795 nm.

The analogous complex with three imidazolyl ligands on the copper ion, LS-3DCHIm, shows similar reactivity toward these relatively strong proton or electron sources; that is, there is no observable reaction with Fc^* , and release of H_2O_2 (100%) occurs with addition of $[DMF \cdot H^+](CF_3SO_3^-)$ (Figure 2, Table S1). Because, with both of these heme-peroxo-copper complexes, the addition of strong acid in combination with reductant still leads to only acid–base chemistry, we reasoned that a certain degree of “finesse” was likely necessary with

respect to peroxo- H^+/e^- interactions along the reaction pathway. Therefore, we have probed this idea further by employing milder acidic substrates (phenols) to find conditions that lead to O–O activation and reductive cleavage, but not H_2O_2 release. In support, a recent study involving electrochemical reduction of a synthetic Mn–peroxo species revealed that strong acid favored M–O cleavage (H_2O_2 evolution), whereas employment of a weak acid facilitated O–O reductive cleavage.⁵⁵

Weak (Phenolic) Acid Association. The reactivity of interest here is revealed when the low-spin complex is tested in reactions with phenols. First, we observe no reaction (as monitored by UV–vis) when LS-4DCHIm is tested with phenols having low bond dissociation free energies (BDFE) and low acidities such as 4-MeO-phenol or 4-^tBu-phenol, which are typically used as H-atom sources.^{56,57} With a slightly more acidic phenol, 4-CN-phenol ($pK_a(CH_3CN) = 22.7$),⁵⁸ still, no reaction is observed. However, upon addition of excess (≥ 10 equiv) 4- NO_2 -phenol ($pK_a(CH_3CN) = 20.7$ ⁵⁸ and $pK_a(tetrahydrofuran, THF) = 18$ ^{59,60}) to a solution of LS-4DCHIm (Figure 3, black spectra), a new species is formed, [LS-4DCHIm(ArOH)] (Figure 3, blue spectra) ($\lambda_{max} = 414$ (Soret), 540, and 840(b) nm). The fact that excess phenol is necessary to see full formation of this intermediate leads us to




Figure 3. UV–vis spectra of LS-4DCHIm reacting with 4- NO_2 -phenol to give the [LS-4DCHIm(ArOH)] adduct, and rR difference spectra showing the shifts that occur upon phenolic acid association.

tentatively assign this as an equilibrium process (see kinetic analysis below). The blue-shifted Soret and upshifted ν_4 oxidation state marker band from resonance Raman (rR) spectroscopy of the ArOH-adduct (Figure 3, bottom right, $\lambda_{\text{ex}} = 413$ nm for both species) indicate partial oxidation of the Fe^{III} center (more specifically, decreased donation from Fe (d_{π}) into porphyrin (π^*) orbitals),⁶¹ in agreement with DFT calculations of H-bonding to the peroxo core (vide infra, see Supporting Information). We associate the weak intensity retained in the low-energy bands in the UV-vis spectrum with the binuclear complex remaining intact. The fact that this intermediate is EPR silent (<5% mononuclear Cu^{II} or Fe^{III} ion species), Figure 3, corroborates that conclusion, due to the coupling of the Cu^{II} ($S = 1/2$) and LS-Fe^{III} ($S = 1/2$) through the bridging peroxo moiety.

Evidence that the O–O bond is still intact in [LS-4DCHIm(ArOH)] (Figure 3, top) is provided by experimental quantification of the H₂O₂ produced (85% yield) when this adduct is exposed to a strong acid, here [DMF·H⁺](CF₃SO₃[−]) (horseradish peroxidase assay; see Table S1). Resonance Raman data provide compelling support that the O–O bond is both intact and perturbed by phenol addition, revealing clear differences in the $\nu(\text{Fe–O})$ and $\nu(\text{O–O})$ stretching vibrations between the LS-4DCHIm species ($\lambda_{\text{ex}} = 413$ nm) and the [LS-4DCHIm(ArOH)] adduct ($\lambda_{\text{ex}} = 825$ nm for better selective enhancement) (Figure 3, bottom left). Specifically, the $\nu(\text{Fe–O})$ increases slightly (by ~ 10 cm^{−1}) from ~ 590 to 598 cm^{−1} ($\Delta^{18}\text{O}_2 = 29$ cm^{−1}), while the $\nu(\text{O–O})$ decreases by >40 cm^{−1} from ~ 870 to 827 cm^{−1} ($\Delta^{18}\text{O}_2 = 48$ cm^{−1}) (Figure 3). It is valuable to note that for both LS-4DCHIm and [LS-4DCHIm(ArOH)], the Fe–O and O–O stretches are observed at both high and low energy excitation (413 nm and ~ 800 – 900 nm, Figure S5). This indicates that (1) the chromophore responsible for the low energy absorption bands also has a strong absorption at 413 nm (for the respective samples), (2) the O–O stretches observed for LS-4DCHIm and [LS-4DCHIm(ArOH)] are due to a heme-bound species (considering the significant resonance enhancement at 413 nm with low power (<2 mW)), and (3) the low energy bands around 850 nm can be reasonably assigned as peroxo-to-Fe CT bands (because both the $\nu(\text{Fe–O})$ and the $\nu(\text{O–O})$ are resonantly enhanced).

The phenomenon of H-bonding interactions inducing small (5–10 cm^{−1}) shifts in the Fe–O stretching mode has been reported for dioxygen adducts of cytochrome P450 mutants (not including a copper ion), which contain potential H-bonding residues in various active site positions. Furthermore, it was concluded on the basis of experimental and theoretical studies that a shift to higher energy, as observed here, is indicative of an H-bonding interaction with the distal O-atom with respect to iron.^{62–64} The aforementioned spectral evidence leads us to propose the structure of the intermediate formed to be a 4-NO₂-phenol associated heme-peroxo-copper adduct depicted in Figure 3 wherein the phenolic O–H associates via H-bonding with the Cu-bound O-atom (O_{Cu}).⁷⁸ Considering that it is logically favorable to strengthen the Fe–O bond and weaken the O–O bond to effect O–O reductive cleavage, the weakly acidic phenol is therefore poised to participate in a full PCET reaction, resulting in homolytic O–O bond cleavage.

DFT calculations support the association of 4-NO₂-phenol with LS-4DCHIm to form an adduct (Figure 4, also see the Experimental Section). This H-bonding interaction is indicated




Figure 4. DFT-calculated structures for LS-4DCHIm and [LS-4DCHIm(ArOH)] (with porphyrin Ar^F groups and most H-atoms omitted for clarity). The table shows selected DFT-calculated properties for the H-bonding and H⁺-transfer structures discussed.

in the DFT calculated structure of [LS-4DCHIm(ArOH)] (Figure 4, right) by the dotted green line ($r_{\text{calc}}(\text{H–O}_{\text{Cu}}) = 1.530$ Å; $r_{\text{calc}}(\text{O}_{\text{Ph}}\text{–O}_{\text{Cu}}) = 2.536$ Å). Further calculations comparing possible binding modes (see Supporting Information for details) provide strong support for this assignment, predicting the O_{Cu}–phenol adduct to be ~ 2 kcal/mol lower in energy than the O_{Fe}–phenol adduct. This is expected given that LS-4DCHIm has a greater partial negative charge on O_{Cu} as compared to O_{Fe}, which would favor phenol association on O_{Cu}. Furthermore, phenol binding to O_{Cu} results in a shorter Fe–O bond and greater Fe–O bond order, consistent with the rR observed frequency change, whereas binding to O_{Fe} would give a longer Fe–O bond and lower Fe–O bond order (both cases result in a longer O–O bond and lower O–O bond order, consistent with the lower $\nu(\text{O–O})$ from rR spectroscopy). We note that DFT-calculated frequencies showed substantial mixing of Fe–O, Cu–O, and O–O modes that was sensitive to small changes in core bond angles, therefore complicating interpretation of the individual two-atom centered modes. However, employing Badger's rule to predict a change in the Fe–O stretching frequency on the basis of the change in bond length in Figure 4 yields a $\Delta(\nu(\text{Fe–O}))$ of $+9$ cm^{−1} for the O_{Cu}-bound structure relative to that in the peroxo complex, LS-4DCHIm, providing excellent agreement with the rR data shown in Figure 3 (as compared to $\Delta(\nu(\text{Fe–O})) = -29$ cm^{−1} for the O_{Fe}-bound structure).⁶⁵

Calculations regarding the protonation states along the O–O cleavage pathway have suggested the possible formation of an Fe–OOH species following proton transfer from an active site Tyr residue.⁴⁴ Therefore, we also computationally evaluated the possibility that the H⁺ has transferred from 4-NO₂-PhOH in the adduct, leaving 4-NO₂-PhO[−] strongly bound as an H-bond acceptor (although UV-vis data for 4-NO₂-PhO[−] appearance and kinetic data (vide infra) suggest this is not the case). However, the optimized structure for the H⁺ transferred to the O_{Cu} reveals an O–O bond elongation (relative to LS-

4DCHIm) that is far too large, corresponding to a $\sim 195\text{ cm}^{-1}$ downshift in $\nu(\text{O}-\text{O})$ as predicted by Badger's rule^{65,66} (as compared to an $\sim 92\text{ cm}^{-1}$ downshift when 4-NO₂-PhOH is the H-bond donor). As it is also 2 kcal/mol higher in energy than the H-bonded adduct (Figure 4), a structure in which the H⁺ has transferred to yield a hydroperoxo species is unlikely.

It is interesting to briefly compare the effects of protonating the peroxy moiety (or to a lesser extent, H-bonding to it) in LS-4DCHIm with those observed in Cu₂O₂ and H₂O₂ chemistry, where protonation results in an increase, rather than a decrease, in the O–O stretching frequency (and DFT-predicted bond strength).⁶⁷ This difference in the $\nu(\text{O}-\text{O})$ behavior can be attributed to the fact that Fe(III) has a d_{π} hole, which receives additional (π -symmetry) donation from the O₂²⁻ π^* orbital in the peroxy structure. The interaction with the proton lowers the energy of the O₂²⁻ orbitals, greatly decreasing this donation (thereby increasing π^* occupation and weakening the O–O bond). Note that in the heme–Cu system, protonation on the O_{Cu} atom localizes charge on the O_{Fe} atom (originally in a p_{π} orbital) and strengthens the Fe–O σ bond (thus the net effect is a stronger Fe–O bond).

Reductive O–O Cleavage. The above results show that the H-bonding interaction weakens the O–O bond and “activates” the [LS-4DCHIm(ArOH)] adduct peroxy moiety to be able to accept an electron from a reducing agent (Fc*), leading to a PCET-type, O–O cleavage reaction.

The products of the reaction, obtained in high yield, are shown in the diagram at the top of Figure 5, as deduced from our spectroscopic analysis. The heme-containing product exhibits UV–vis features at $\lambda_{\text{max}} = 414$ (Soret) and 542 nm (Figure 5, green spectrum), characterized as a low-spin, six-coordinate ferric-heme. Two equivalents of decamethylferrocenium ion (Fc*⁺) are formed on the basis of the new absorptions at $\lambda_{\text{max}} = 785, 805\text{ nm}$ (quantified as $\Delta_{\text{Abs}}(785\text{ nm} - 870\text{ nm})$; $\epsilon = 580\text{ M}^{-1}\text{ cm}^{-1}$),⁶⁸ signifying that a two-electron reduction of [LS-4DCHIm(ArOH)] has occurred. Additionally, an increase in absorbance corresponding to the 4-NO₂-PhO⁻ anion appears as a shoulder of the heme Soret band, indicating that proton(s) have also been transferred in significant quantities (see Figure S18).

Further supporting our conclusions about the nature and identity of species in the product mixture is the latter's EPR spectrum showing the emergence of nearly 100% of the Cu^{II} ion expected, and mostly low-spin ($\sim 95\%$) Fe^{III} species, which match an EPR spectrum of an authentic mixture of these products (see Figure 5, bottom). Quantification of the iron(III) and copper(II) products was carried out by comparing the reaction product spectrum with intensity calibration curves obtained by independently generating these species (Figure S11).

A negligible amount of H₂O₂ (<5%, Table S1) is released in this step, and, furthermore, rR data collected at 413 nm for the reaction products show no isotope-sensitive stretching vibrations, and indicate the final heme product is the low-spin bis-imidazole F₈Fe(III) species (Figure S12). While these observations do not definitively prove that water has formed, the consumption of two (stoichiometrically required) reductive equivalents, confirmation of oxidized metal centers in the final product mixture, and lack of H₂O₂ released do provide compelling evidence that the O–O bond has been broken. Recall that both protons and electrons are required for O–O cleavage, as addition of up to 20 equiv of Fc* alone to a solution of LS-4DCHIm causes no spectral changes (Figure 2).




Figure 5. Scheme, UV–vis, and EPR spectra showing the 2-electron reduction of the H-bonded adduct, [LS-4DCHIm(ArOH)], by Fc*. This step includes O–O cleavage and therefore decoupling of the Fe(III) and Cu(II) metal centers.

Kinetic Measurements and Mechanistic Insights. For the processes occurring in Figures 3 and 5, pseudo-first-order kinetic experiments were carried out to determine reaction orders for LS-4DCHIm, 4-NO₂-phenol, and Fc* (see Supporting Information for details). While the reaction rate was found to be independent of [LS-4DCHIm] and linear with respect to [Fc*], saturation behavior was observed with increasing concentrations of 4-NO₂-phenol (Figure 6A,B). This observation is consistent with a reversible equilibrium or formation of an intermediate prior to the rate-determining step.^{52,69,70} In this case, it can be attributed to the association of the 4-NO₂-phenol with the peroxy core. Notably, the UV–vis changes that occur upon addition of 4-NO₂-phenol to LS-4DCHIm are not reversed following addition of strong base; the entire phenolic moiety inserts into the structure of LS-4DCHIm (vide supra). After measuring the rate as a function of Fc* appearance ($\Delta_{\text{Abs}}(785-870\text{ nm})$) over time (also see Figures S14–16) and applying the kinetic model in Figure 6C, we calculate an association constant, $K_{\text{H}^+} = 604 \pm 21\text{ M}^{-1}$, as




Figure 6. Kinetic data for the weak acid/electron reactions with LS-4DCHIm. Dependence of k_{obs} on $[\text{Fc}^*]$ (a) and on $[4\text{-NO}_2\text{-phenol}]$ (b). Kinetic model used for fitting the data (c).

well as a rate constant for the second-order electron transfer step, $k_{\text{et1}} = 1.6 \pm 0.1 \text{ M}^{-1} \text{ s}^{-1}$.

All kinetic experiments were carried out by initial addition of the reductant to the LS-4DCHIm complex, which caused no spectral changes, followed by addition of 4-NO₂-phenol. In all cases, immediate formation of the intermediate species, [LS-4DCHIm(ArOH)], was observed, leading to the formation of the reaction products. This experimental observation is consistent with the proposed mechanism, in which construction of an H-bonding network around the Fe-peroxo-Cu complex precedes and facilitates the electron transfer. The linear dependence of k_{obs} on the concentration of Fc^* indicates that during the rds, only one molecule of reductant is involved, which we believe results in O–O cleavage and transient formation of a $\text{Fe}^{\text{IV}}=\text{O}$, and the second electron transfer occurs immediately following that formation. No H_2O_2 is released when $[\text{DMF}\cdot\text{H}^+](\text{CF}_3\text{SO}_3^-)$ is added to the product mixture (Table S1), indicating that all peroxide has been reductively cleaved and protonated to give water (Figure 5). This result, along with the Cmpd II experimental results (vide infra, Figure 7) and nature of the final products formed ($\text{Fe}(\text{III})$, $\text{Cu}(\text{II})$, and 2 equiv of $\text{Fc}^{*\text{+}}$), support this claim that a transient ferryl must have formed in the initial reduction (1-electron) and protonation.

Deuterated 4-NO₂-phenol(OD) was employed for kinetic isotope effect analysis (see Supporting Information). The resultant KIE values of 1.6 for the phenol-association step ($K_{\text{ArOD}} = 374 \pm 28 \text{ M}^{-1}$) and 1.9 for the rate-determining step ($k_{\text{et(D+)}} = 0.89 \pm 0.1 \text{ M}^{-1} \text{ s}^{-1}$) suggest that the phenolic O–H/D bond is broken, and the proton is transferred in the rate-determining step. This conclusion is consistent with the DFT interrogation of [LS-4DCHIm(ArOH)] (vide supra) and the overall proposed mechanism. While smaller than expected for a phenolic HAT reaction, these KIE values are in the range for PCET reactions involving phenol O–H(D) bond activation by metal–oxygen species.^{71,72}

Mechanistic Insights from Independent $\text{Fe}^{\text{IV}}=\text{O}$ Studies. In CcO, it is proposed that delivery of a H^+ and e^- induces O–O cleavage of an Fe^{III} -peroxo- Cu^{II} moiety, resulting in an $\text{Fe}^{\text{IV}}=\text{O}$ (Cmpd II-type) species, which is subsequently reduced and protonated to give a low-spin Fe^{III} -OH product.⁷³ To probe the possibility that such a ferryl species forms during reduction of [LS-4DCHIm(ArOH)], as implied by the kinetic results just described, initial rates were measured for the protonation/reduction of an authentic




Figure 7. Scheme (top) and UV-vis spectra showing (A) formation and 1e^- reduction reactivity of $\text{Fe}^{\text{IV}}=\text{O}$ in the presence of 4-NO₂-phenol. Initial rates (B) and UV-vis spectra (C) for formation of 1 equiv of $\text{Fc}^{*\text{+}}$ in the reduction of Cmpd II (blue) and 2 equiv of $\text{Fc}^{*\text{+}}$ for [LS-4DCHIm(ArOH)] (green).

solution of $\text{F}_8\text{Fe}^{\text{IV}}=\text{O}^{74}$ by addition of 4-NO₂-phenol and Fc^* .

Indeed, the same low-spin heme product was observed (Figure 7), and both the proton and electron sources were necessary to observe a reaction. However, the rate of reduction (1e^-) of the Cmpd II-type complex is >3 times faster than the reduction of [LS-4DCHIm(ArOH)] (2e^- overall, vide supra); therefore, this explains the finding that we could not spectroscopically (by UV-vis) observe the $\text{Fe}^{\text{IV}}=\text{O}$ intermediate during the LS-4DCHIm reaction because excess reductant is present (note that in CcO and other model complexes,⁷⁵ the stoichiometric supply of electrons can be controlled, allowing observation of a metastable $\text{Fe}^{\text{IV}}=\text{O}$ species). This result demonstrates the mechanistic similarity between O–O cleavage in LS-4DCHIm and CcO, highlighting the relevance of this work among enzyme model complexes.

Impact of Structure on Reactivity. In addition to LS-4DCHIm, our lab has recently characterized a similar heme-peroxo-copper complex, LS-3DCHIm, in which the copper ion is ligated by only three monodentate DCHIm donors.⁵¹ LS-3DCHIm is only stable below $-115 \text{ }^\circ\text{C}$ in MeTHF and is unchanged following addition of 4-NO₂-phenol and/or Fc^* (monitored by UV-vis and rR), illustrating the different chemical properties of the two low-spin imidazolyl complexes. It is important to note that LS-4DCHIm can be cooled below $-115 \text{ }^\circ\text{C}$ and maintain its reactivity (at slower rates). However, no reaction is observed when an excess of a sterically hindered, albeit slightly more acidic, phenol, 2,6-di-*tert*-butyl-4-nitrophenol ($\text{pK}_a(\text{CH}_3\text{CN}) = 19.1$),⁵⁷ is added to a solution of LS-4DCHIm, even in the presence of Fc^* . In contrast, recall that reaction with strong acid, $[\text{DMF}\cdot\text{H}^+](\text{CF}_3\text{SO}_3^-)$, gives identical products for both LS-3DCHIm and LS-4DCHIm (UV-vis, EPR, H_2O_2 quantification, vide supra). To gain insight into this difference in reactivity, we compared the DFT-optimized structures for these two complexes (Figure 8). DFT predicts the lowest energy structure for LS-3DCHIm to have a distorted square planar geometry ($\tau = 0.4$, where 0 = square planar, 1 =




Figure 8. DFT-optimized structures of **LS-4DCHIm** and **LS-3DCHIm** (porphyrin ArF groups and H-atoms omitted for clarity) depicting the relationship between structure and reactivity toward different acids.

tetrahedral) with an Fe–Cu distance of 4.10 Å, and for **LS-4DCHIm** to have a square pyramidal geometry ($\tau = 0.1$, where 0 = square pyramidal, 1 = trigonal bipyramidal)⁷⁶ in which the additional DCHIm ligand occupies the axial position at the copper center, and the Fe–Cu distance is 4.47 Å. While these calculated structures do not give insight into the markedly high O–O stretch observed in rR experiments, it is clear that an additional monodentate ligand induces a structural rearrangement, which influences the accessibility of the peroxo core. In particular, the core geometries of **LS-3DCHIm** and **LS-4DCHIm**, depicted in **Figure 8** (also **Figure S9** for more selected bond lengths and angles), indicate a more accessible peroxo moiety in **LS-4DCHIm**. These results together suggest that steric factors inhibiting H-bond association greatly contribute to the difference in reactivity between the two complexes.

CONCLUSIONS

The structural nature of the CcO-inspired heme–peroxo–copper complexes, and their reactivity in protonation–reduction chemistry, which has been described here, further emphasizes the realization concerning how acutely nature has constructed its systems to achieve the chemical transformations necessary for sustaining aerobic life. Rather than rigorous mimicking of proteins, we have taken the approach of designing synthetically simplified complexes in which we can control or manipulate certain aspects in our pursuit of a fundamental understanding of the factors involved in O₂-reduction processes. This has required the use of cryogenic conditions (e.g. –90 °C) for generation of heme–Cu O₂-adducts, those which are highly reactive in biological systems and currently inaccessible for spectroscopic study.⁷⁷ In doing so, we have shown that both the


structure of the bridging peroxo-complex and the nature of the proton source play a role in determining whether the protonation/reduction mechanism proceeds via acid/base chemistry (metal–O cleavage, release of H₂O₂) or substrate-mediated redox chemistry (O–O bond reductive cleavage). The use of monodentate DCHIm ligands allows for the formation of heme–(O₂²⁻)–Cu complexes possessing comparable yet unique structures, **LS-3DCHIm** and **LS-4DCHIm**, with intriguingly different reactivities.

The two complexes discussed differ in that a phenolic H-bond associated adduct can form with **LS-4DCHIm**, where we have shown that the weak acid, 4-NO₂-phenol, associates with the Cu-bound O_{peroxo}-atom. These observations contrast to the behavior of **LS-3DCHIm**, where a potentially more compressed peroxo core does not allow for formation of such an adduct. If a strong acid is employed, [DMF·H⁺](CF₃SO₃⁻), steric hindrance is no longer a contributing factor, as a (solvated) proton can easily reach either peroxo core to induce metal–O cleavage via acid–base chemistry, and release hydrogen peroxide. Therefore, we have observed that in a fashion similar to the generally proposed CcO enzymatic mechanism, reductive O–O cleavage is effected after first establishing a hydrogen-bonding network close to the peroxo, which then facilitates the proton-coupled electron transfer (in this case, from 4-NO₂-phenol and Fc*, and in CcO, from the cross-linked Tyr residue) (**Scheme 1**). In the enzyme, this efficient manner of transferring the H⁺/e⁻ to the peroxo prevents the production of partially reduced, detrimental, reactive oxygen species.^{15,79,80} Only a few previous CcO model systems have achieved O–O cleavage; however, they have employed rigid structural frameworks or “picket-fence porphyrins”. Now, we have shown that a new, potentially more flexible adduct, when paired with the appropriate reagents, can closer mimic the enzyme’s proposed stepwise transfer of H⁺/e⁻ to an Fe–Cu bridging peroxo moiety. In this way, we continue to uncover the seemingly small, yet critical, chemical and/or structural aspects of transition metal–O₂ assemblies, which greatly impact O–O cleavage chemistry.

EXPERIMENTAL SECTION

General. All reagents and solvents used were of commercially available quality and used without further purification except as noted. Inhibitor-free 2-MeTHF was distilled over Na/benzophenone under Ar and deoxygenated with Ar before use. [Cu^I(CH₃CN)₄(BAR^F)] and F₈Fe^{II} were synthesized as previously described.^{81,82} All UV–vis measurements were carried out using a Hewlett-Packard 8453 diode

Scheme 1. Overall Proposed Mechanism for the Reaction of **LS-4DCHIm** with 4-NO₂-Phenol and Fc*



array spectrophotometer with a quartz Schlenk cuvette cell to monitor changes in Q-band and with a 0.2 cm path length cuvette to monitor changes in Soret band. The spectrometer was equipped with HP Chemstation software and a Unisoku thermostated cell holder for low-temperature experiments. Resonance RAMAN samples were excited at a variety of wavelengths, using either a Coherent I90C-K Kr⁺ ion laser, a Coherent 25/7 Sabre Ar⁺ ion laser, or a Lighthouse Photonics Sprout-G pumped SolsTiS-PX Ti:Saph laser, while the sample was immersed in a liquid nitrogen cooled (77 K) EPR finger dewar (Wilmad). Power was ~2 mW at the sample for 413 nm excitation and 150–200 mW for lower energy excitation. Data were recorded while rotating the sample to minimize photodecomposition. The spectra were recorded using a Spex 1877 CP triple monochromator with either a 600, 1200, or 2400 grooves/mm holographic spectrograph grating, and detected by an Andor Newton CCD cooled to –80 °C (for high-energy excitation) or an Andor IDus CCD cooled to –80 °C (for the low-energy excitation). Spectra were calibrated on the energy axis to toluene. Excitation profiles were intensity calibrated to the solvent (MeTHF) by peak fitting in the program Origin.

UV–Vis. The complexes LS-3DCHIm and LS-4DCHIm were generated as previously described at 0.1 mM concentrations at –125 or –90 °C in a 1 cm path length, rubber septum-capped, quartz Schlenk cuvette.⁵¹ 50 μL of a solution containing the desired number of equivalents of acid (10 for 4-NO₂-phenol, 2 for [DMF·H⁺](CF₃SO₃⁻) and/or reductant (Fc*, 10 equiv) was added via gastight syringe and mixed by bubbling Ar. To obtain conditions optimal for observing Soret peaks, the same concentration (~0.1 mM) was used in a 2 mm path length Schlenk cuvette. For kinetic measurements, spectra were recorded every 1.0 s to obtain “initial rate” data. Reactions were monitored until stable by UV–vis (i.e., no more spectral changes are observed); however, we do note that in reactions with excess Fc*, slow reduction of the metal centers may eventually occur.

EPR. 0.6 mL solutions of the complexes LS-3DCHIm and LS-4DCHIm were generated anaerobically as previously described⁵¹ at 2.0 mM concentrations at –125 °C (pentane/liquid N₂ bath) and –90 °C (acetone/liquid N₂ bath), respectively, in a 5 mm, rubber septum-capped, EPR tube. The desired substrates (acids and/or reductant) were added in the desired amounts, mixed by bubbling Ar, and frozen prior to recording spectra. Spectra were taken with an ER 073 magnet equipped with a Bruker ER041 X-Band microwave bridge and a Bruker EMX 081 power supply: microwave frequency = 9.41 GHz, microwave power = 0.201 mW, attenuation = 30 db, modulation amplitude = 10 G, modulation frequency = 100 kHz, temperature = 10 K.

Raman. The complexes LS-3DCHIm and LS-4DCHIm were generated in the same fashion as EPR samples, above, at 1.0 mM concentrations at –125 °C (pentane/liquid N₂ bath) and –90 °C (acetone/liquid N₂ bath), respectively, in a 9-in., 5 mm, rubber septum-capped, NMR tube using either ¹⁶O₂ or ¹⁸O₂.⁵¹ The desired substrates (acids and/or reductant) were added in the desired amounts and mixed by bubbling Ar. The tubes were then frozen and flame-sealed. Spectra were obtained with spinning tubes at 77 K with various excitation wavelengths noted for each case.

H₂O₂ Quantification by Horseradish Peroxidase (HRP) Test. The spectrophotometric quantification of hydrogen peroxide was achieved by recording the intensity of the diammonium 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonate (AzBTS-(NH₄)₂) peaks (different wavelengths, monitored at 418 nm to minimize error, Figure S1) oxidized by horseradish peroxidase (HRP), which was adapted from published procedures.⁵² In a typical experiment, 3 mL of the desired heme-peroxo-copper complex [0.135 mM] was generated at –90/–125 °C. If required, two sample solutions of 100 μL containing Fc* [2 mM] and 100 μL containing 4-NO₂-phenol [2 mM] were added, and the reaction was allowed to proceed until completion (no further spectral changes were observed). The reaction crude was subjected to the H₂O₂ analysis as described below, both before and after quenching by addition of a 100 μL solution containing 2.5 equiv of [DMF·H⁺](CF₃SO₃⁻) [0.338 mM]. Hydrogen peroxide (H₂O₂) was detected using the procedure described using the following stock solutions: 300 mM sodium phosphate buffer, pH 7.0 (solution A), 1 mg/mL AzBTS-

(NH₄)₂ (solution B), 4 mg of HRP (type II salt free (Sigma)), and 6.5 mg of sodium azide in 50 mL of water (solution C). Quantification of hydrogen peroxide was achieved by adding 100 μL of the cold MeTHF sample solution to a cuvette containing 1.3 mL of water, 500 μL of solution A, 100 μL of solution B, and 50 μL of solution C (all chilled). After being mixed for 15 s, the samples were allowed to incubate for ~2 min until full formation of the 418 nm band was achieved (Table S1).

DFT Calculations. Density functional theory (DFT) calculations were performed with the Gaussian 09, version D.01 software package. All calculations were done using the BP86 functional within the spin-unrestricted formalism and broken-symmetry singlet surface, employing a split basis set as follows: 6-311g* for Fe, Cu, and peroxy O-atoms; 6-31g* for all metal-bound N atoms; and 6-31g for all remaining atoms. Tight SCF convergence and an ultrafine integration grid were used. All cyclohexyl substituents on DCHIm ligands were truncated as isopropyl groups to lower the computational cost, yet still capture some of the inherent steric effects. (A structure for LS-4DCHIm was optimized using the complete DCHIm ligands, which yielded very similar core bond distances and vibrational frequencies, indicating that the truncated model adequately represents the full ligand system for correlating to our experimental data.) Free energies were calculated at 183 K. For each binding orientation (phenol H-bonding to O_{Fe} or O_{Cu}), several positions of the phenol were attempted in which the O–H was directed toward the particular peroxy O-atom. Unconstrained optimization from these starting orientations yielded one minimum for each of the H-bonded structures (O_{Fe}-bound and O_{Cu}-bound). Structures for which the H⁺ was transferred to the respective peroxy O-atoms were obtained by constraining the O_{ph}–H distance at the same distance as the O_{Cu}–H in the H-bonding structure (thus the distance from the H⁺ to the H-bond acceptor remains the same) and optimizing the remaining atoms.

■ ASSOCIATED CONTENT

§ Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/jacs.6b11322.

H₂O₂ quantification details, kinetic and KIE data, strong acid reactivity product analysis, control experimental details, and supplementary DFT calculations (PDF)
X-ray crystallographic data for [(DCHIm)₂Cu^I](BAR^F) (CIF)

X-ray crystallographic data for [(DCHIm)₄Cu^{II}](ClO₄)₂·Et₂O (CIF)

■ AUTHOR INFORMATION

Corresponding Authors

*edward.solomon@stanford.edu

*karlin@jhu.edu

ORCID

Kenneth D. Karlin: 0000-0002-5675-7040

Present Address

§I.G.-B.: Department of Chemistry, Southern Methodist University, Dallas, Texas 75275, United States.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

This research was supported by the U.S. NIH (GM60353 to K.D.K. and DK31450 to E.I.S.). I.G.-B. thanks the European Commission for a Marie Curie IOF Fellowship.

■ REFERENCES

(1) Cook, T. R.; Dogutan, D. K.; Reece, S. Y.; Surendranath, Y.; Teets, T. S.; Nocera, D. G. *Chem. Rev.* **2010**, *110*, 6474–6502.

- (2) Semenza, G. L. *Science* **2007**, *318*, 62–64.
- (3) Hoganson, C. W.; Pressler, M. A.; Proshlyakov, D. A.; Babcock, G. T. *Biochim. Biophys. Acta, Bioenerg.* **1998**, *1365*, 170–174.
- (4) Warren, J. J.; Mayer, J. M. *Biochemistry* **2015**, *54*, 1863–1878.
- (5) Ray, K.; Felix, F.; Wang, B.; Nam, W. *J. Am. Chem. Soc.* **2014**, *136*, 13942–13958.
- (6) Engelmann, X.; Monte-Pérez, I.; Ray, K. *Angew. Chem., Int. Ed.* **2016**, *55*, 7632–7649.
- (7) Collman, J. P.; Boulatov, R.; Sunderland, C. J.; Fu, L. *Chem. Rev.* **2004**, *104*, 561–588.
- (8) Kim, E.; Chufán, E. E.; Kamaraj, K.; Karlin, K. D. *Chem. Rev.* **2004**, *104*, 1077–1133.
- (9) Cracknell, J. A.; Vincent, K. A.; Armstrong, F. A. *Chem. Rev.* **2008**, *108*, 2439–2461.
- (10) Chufán, E. E.; Mondal, B.; Gandhi, T.; Kim, E.; Rubie, N. D.; Moënnelocoz, P.; Karlin, K. D. *Inorg. Chem.* **2007**, *46*, 6382–6394.
- (11) Kieber-Emmons, M. T.; Li, Y.; Halime, Z.; Karlin, K. D.; Solomon, E. I. *Inorg. Chem.* **2011**, *50*, 11777–11786.
- (12) Natri, F.; Chino, M.; Maglio, O.; Bhagi-Damodaran, A.; Lu, Y.; Lombardi, A. *Chem. Soc. Rev.* **2016**, *45*, 5020–5054.
- (13) Cady, C. W.; Crabtree, R. H.; Brudvig, G. W. *Coord. Chem. Rev.* **2008**, *252*, 444–455.
- (14) Yoshikawa, S.; Shimada, A. *Chem. Rev.* **2015**, *115*, 1936–1989.
- (15) Muramoto, K.; Ohta, K.; Shinzawa-Itoh, K.; Kanda, K.; Taniguchi, M.; Nabekura, H.; Yamashita, E.; Tsukihara, T.; Yoshikawa, S. *Proc. Natl. Acad. Sci. U. S. A.* **2010**, *107*, 7740–7745.
- (16) Blomberg, M. R. A.; Siegbahn, P. E. M.; Wikstrom, M. *Inorg. Chem.* **2003**, *42*, 5231–5243.
- (17) Yoshikawa, S.; Muramoto, K.; Shinzawa-Itoh, K.; Mochizuki, M. *Biochim. Biophys. Acta, Bioenerg.* **2012**, *1817*, 579–589.
- (18) Gorbikova, E. A.; Belevich, I.; Wikstrom, M.; Verkhovsky, M. I. *Proc. Natl. Acad. Sci. U. S. A.* **2008**, *105*, 10733–10737.
- (19) Yoshikawa, S.; Muramoto, K.; Shinzawa-Itoh, K. *Biochim. Biophys. Acta, Bioenerg.* **2011**, *1807*, 1279–1286.
- (20) Kaila, V. R. I.; Verkhovsky, M. I.; Wikstrom, M. *Chem. Rev.* **2010**, *110*, 7062–7081.
- (21) Siegbahn, P. E. M.; Blomberg, M. R. A. *Chem. Rev.* **2010**, *110*, 7040–7061.
- (22) Petrik, I. D.; Davydov, R.; Ross, M.; Zhao, X.; Hoffman, B. M.; Lu, Y. *J. Am. Chem. Soc.* **2016**, *138*, 1134–1137.
- (23) Yu, Y.; Lv, X.; Li, J.; Zhou, Q.; Cui, C.; Hosseinzadeh, P.; Mukherjee, A.; Nilges, M. J.; Wang, J.; Lu, Y. *J. Am. Chem. Soc.* **2015**, *137*, 4594–4597.
- (24) Soper, J. D.; Kryatov, S. V.; Rybak-Akimova, E. V.; Nocera, D. G. *J. Am. Chem. Soc.* **2007**, *129*, 5069–5075.
- (25) Barry, B. A. *J. Photochem. Photobiol., B* **2011**, *104*, 60–71.
- (26) Martinis, S. A.; Atkins, W. M.; Stayton, P. S.; Sligar, S. G. *J. Am. Chem. Soc.* **1989**, *111*, 9252–9253.
- (27) Denisov, I. G.; Makris, T. M.; Sligar, S. G.; Schlichting, I. *Chem. Rev.* **2005**, *105*, 2253–2278.
- (28) Shook, R. L.; Borovik, A. S. *Inorg. Chem.* **2010**, *49*, 3646–3660.
- (29) Yamaguchi, S.; Nagatomo, S.; Kitagawa, T.; Funahashi, Y.; Ozawa, T.; Jitsukawa, K.; Masuda, H. *Inorg. Chem.* **2003**, *42*, 6968–6970.
- (30) Augustine, A. J.; Quintanar, L.; Stoj, C. S.; Kosman, D. J.; Solomon, E. I. *J. Am. Chem. Soc.* **2007**, *129*, 13118–13126.
- (31) Tinberg, C. E.; Lippard, S. J. *Acc. Chem. Res.* **2011**, *44*, 280–288.
- (32) Solomon, E. I.; Augustine, A. J.; Yoon, J. *Dalt. Trans.* **2008**, 9226, 3921.
- (33) Collman, J. P.; Devaraj, N. K.; Decréau, R. A.; Yang, Y.; Yan, Y.-L.; Ebina, W.; Eberspacher, T. A.; Chidsey, C. E. D. *Science* **2007**, *315*, 1565–1568.
- (34) Halime, Z.; Kotani, H.; Li, Y.; Fukuzumi, S.; Karlin, K. D. *Proc. Natl. Acad. Sci. U. S. A.* **2011**, *108*, 13990–13994.
- (35) Ghosh, S.; Xie, X.; Dey, A.; Sun, Y.; Scholes, C. P.; Solomon, E. I. *Proc. Natl. Acad. Sci. U. S. A.* **2009**, *106*, 4969–4974.
- (36) Collman, J. P.; Sunderland, C. J.; Berg, K. E.; Vance, M. A.; Solomon, E. I. *J. Am. Chem. Soc.* **2003**, *125*, 6648–6649.
- (37) Kim, E.; Kamaraj, K.; Galliker, B.; Rubie, N. D.; Moënnelocoz, P.; Kaderli, S.; Zuberbühler, A. D.; Karlin, K. D. *Inorg. Chem.* **2005**, *44*, 1238–1247.
- (38) Chufán, E. E.; Pui, S. C.; Karlin, K. D. *Acc. Chem. Res.* **2007**, *40*, 563–572.
- (39) Liu, X.; Yu, Y.; Hu, C.; Zhang, W.; Lu, Y.; Wang, J. *Angew. Chem., Int. Ed.* **2012**, *51*, 4312–4316.
- (40) Blomberg, M. R. A.; Siegbahn, P. E. M.; Babcock, G. T.; Wikström, M. *J. Am. Chem. Soc.* **2000**, *122*, 12848–12858.
- (41) Solomon, E. I.; Heppner, D. E.; Johnston, E. M.; Ginsbach, J. W.; Cirera, J.; Qayyum, M.; Kieber-Emmons, M. T.; Kjaergaard, C. H.; Hadt, R. G.; Tian, L. *Chem. Rev.* **2014**, *114*, 3659–3853.
- (42) Sharma, V.; Wikström, M.; Kaila, V. R. I. *Biochim. Biophys. Acta, Bioenerg.* **2011**, *1807*, 813–818.
- (43) Du, W.-G. H.; Noodleman, L. *Inorg. Chem.* **2013**, *52*, 14072–14088.
- (44) Du, W.-G. H.; Götz, A. W.; Yang, L.; Walker, R. C.; Noodleman, L. *Phys. Chem. Chem. Phys.* **2016**, *18*, 21162–21171.
- (45) Blomberg, M. R. A. *Biochemistry* **2016**, *55*, 489–500.
- (46) Chatterjee, S.; Sengupta, K.; Hematian, S.; Karlin, K. D.; Dey, A. *J. Am. Chem. Soc.* **2015**, *137*, 12897–12905.
- (47) Pegis, M. L.; Roberts, J. A. S.; Wasylenko, D. J.; Mader, E. A.; Appel, A. M.; Mayer, J. M. *Inorg. Chem.* **2015**, *54*, 11883–11888.
- (48) Bhagi-Damodaran, A.; Petrik, I.; Lu, Y. *Isr. J. Chem.* **2016**, *61801*, 773–790.
- (49) Hematian, S.; Garcia-Bosch, I.; Karlin, K. D. *Acc. Chem. Res.* **2015**, *48*, 2462–2474.
- (50) We report pK_a and E⁰ values in MeCN at room temperature (although our chemistry is carried out in MeTHF under cryogenic conditions), as a general guide, because the RT and/or nonaqueous solvent values are documented in the literature, internally consistent, and serve the purpose of providing a relative scale of these parameters in organic solvent.
- (51) Garcia-Bosch, I.; Adam, S. M.; Schaefer, A. W.; Sharma, S. K.; Peterson, R. L.; Solomon, E. I.; Karlin, K. D. *J. Am. Chem. Soc.* **2015**, *137*, 1032–1035.
- (52) Peterson, R. L.; Ginsbach, J. W.; Cowley, R. E.; Qayyum, M. F.; Himes, R. A.; Siegler, M. A.; Moore, C. D.; Hedman, B.; Hodgson, K. O.; Fukuzumi, S.; Solomon, E. I.; Karlin, K. D. *J. Am. Chem. Soc.* **2013**, *135*, 16454–16467.
- (53) Carver, C. T.; Matson, B. D.; Mayer, J. M. *J. Am. Chem. Soc.* **2012**, *134*, 5444–5447.
- (54) Tosha, T.; Kagawa, N.; Arase, M.; Waterman, M. R.; Kitagawa, T. *J. Biol. Chem.* **2008**, *283*, 3708–3717.
- (55) Ching, H. Y. V.; Anxolabéhère-Mallart, E.; Colmer, H. E.; Costentin, C.; Dorlet, P.; Jackson, T. A.; Pícar, C.; Robert, M. *Chem. Sci.* **2014**, *5*, 2304.
- (56) Warren, J. J.; Tronic, T. A.; Mayer, J. M. *Chem. Rev.* **2010**, *110*, 6961–7001.
- (57) Nicoletti, C. R.; Marini, V. G.; Zimmermann, L. M.; Machado, V. G. *J. Braz. Chem. Soc.* **2012**, *23*, 1488–1500.
- (58) Klamt, A.; Eckert, F.; Leito, I.; Kaljurand, I.; Ku, A.; Diedenhofen, M. *J. Comput. Chem.* **2009**, *30*, 799–810.
- (59) Nam, E.; Alokolaro, P. E.; Swartz, R. D.; Gleaves, M. C.; Pikul, J.; Kovacs, J. A. *Inorg. Chem.* **2011**, *50*, 1592–1602.
- (60) For reference and comparison: unsubstituted PhOH, pK_a(H₂O) = 9.95, pK_a(MeCN) = 22.7, pK_a(THF) = 21, and for p-NO₂-ArOH, pK_a(H₂O) = 7.18, pK_a(MeCN) = 20.7, pK_a(THF) = 18 (from refs 58, 59). Also, see ref 12: Vazdar, K.; Kunetskiy, R.; Saame, J.; Kaupmees, K.; Leito, I.; Jahn, U. *Angew. Chem., Int. Ed.* **2014**, *53*, 1435–1438.
- (61) Kitagawa, T.; Abe, M.; Kyogoku, Y. *Chem. Phys. Lett.* **1977**, *48*, 55–58.
- (62) Gregory, M.; Mak, P. J.; Sligar, S. G.; Kincaid, J. R. *Angew. Chem., Int. Ed.* **2013**, *52*, 5342–5345.
- (63) Spiro, T. G.; Soldatova, A. V.; Balakrishnan, G. *Coord. Chem. Rev.* **2013**, *257*, 511–527.
- (64) Denisov, I. G.; Mak, P. J.; Makris, T. M.; Sligar, S. G.; Kincaid, J. R. *J. Phys. Chem. A* **2008**, *112*, 13172–13179.
- (65) Green, M. T. *J. Am. Chem. Soc.* **2006**, *128*, 1902–1906.

(66) Cramer, C. J.; Tolman, W. B. *Acc. Chem. Res.* **2007**, *40*, 601–608.

(67) Root, D. E.; Mahroof-Tahir, M.; Karlin, K. D.; Solomon, E. I. *Inorg. Chem.* **1998**, *37*, 4838–4848.

(68) Kakuda, S.; Peterson, R. L.; Ohkubo, K.; Karlin, K. D.; Fukuzumi, S. *J. Am. Chem. Soc.* **2013**, *135*, 6513–6522.

(69) Wijeratne, G. B.; Corzine, B.; Day, V. W.; Jackson, T. A. *Inorg. Chem.* **2014**, *53*, 7622–7634.

(70) Hatcher, L. Q.; Karlin, K. D. *JBIC, J. Biol. Inorg. Chem.* **2004**, *9*, 669–683.

(71) Osako, T.; Ohkubo, K.; Taki, M.; Tachi, Y.; Fukuzumi, S.; Itoh, S. *J. Am. Chem. Soc.* **2003**, *125*, 11027–11033.

(72) Kundu, S.; Miceli, E.; Farquhar, E. R.; Ray, K. *Dalt. Trans.* **2013**, *43*, 2–5.

(73) Han, S.; Ching, Y.; Rousseau, D. L. *Nature* **1990**, *348*, 89–90.

(74) Garcia-Bosch, I.; Sharma, S. K.; Karlin, K. D. *J. Am. Chem. Soc.* **2013**, *135*, 16248–16251.

(75) Collman, J. P.; Decréau, R. A. *Chem. Commun.* **2008**, 5065.

(76) Marlin, D. S.; Olmstead, M. M.; Mascharak, P. K. *Inorg. Chem.* **2001**, *40*, 7003–7008.

(77) (a) The low temperatures and conditions employed here, somewhat extreme, are however not at all a detriment to the goal of obtaining fundamental insights via modeling. With the constructs employed, and in fact via the experimental approach of cryogenic conditions, we are able to limit the number of factors that vary, and directly interrogate a system where the addition of phenol or acid, or Cu-ligation, are under our control and understood; these limited variations result in dramatic influences upon reactivity (i.e., resulting in O–O reductive cleavage vs breaking metal–O bonds and releasing H₂O₂). The simplification of a complex system, to address limited numbers of key factors pertinent to reactivity (in this case), is basic to modeling in all of science: (i) “...the synthetic modeling approach, whereby inorganic complexes that replicate aspects of the metalloenzyme active site are characterized and their reactivity is examined...”;^{77b} (ii) “...synthetic analogues that approach or achieve one or more significant properties of a protein active site...”;^{77c} and (iii) “...artificial model systems composed of only a few components are being used to further our understanding of...”.^{77d} (b) Tolman, W. B. *Inorg. Chem.* **2013**, *52*, 7307–7310. (c) Holm, R. H.; Solomon, E. I. *Chem. Rev.* **2004**, *104*, 347–348. (d) Marsden, H. R.; Tomatsu, I.; Kros, A. *Chem. Soc. Rev.* **2011**, *40*, 1572–1585 (note: on studies of model systems for membrane fusion).

(78) The previously noted observation that 4-CN-phenol does not interact with **LS-4DCHIm** requires further investigation; it is not sterically inhibited and the small difference in acidity compared to 4-NO₂-phenol is not expected to translate in a major way to H-bonding ability. See Abraham, M. H.; Grellier, P. L.; Prior, D. V.; Duce, P. P.; Morris, J. J.; Taylor, P. J. *J. Chem. Soc., Perkin Trans. 2* **1989**, 699–711.

(79) Proshlyakov, D. A.; Pressler, M. A.; DeMaso, C.; Leykam, J. F.; DeWitt, D. L.; Babcock, G. T. *Science* **2000**, *290*, 1588–1591.

(80) Konstantinov, A. A. *FEBS Lett.* **2012**, *586*, 630–639.

(81) Liang, H.-C.; Kim, E.; Incarvito, C. D.; Rheingold, A. L.; Karlin, K. D. *Inorg. Chem.* **2002**, *41*, 2209–2212.

(82) Ghiladi, R. A.; Kretzer, R. M.; Guzei, I.; Rheingold, A. L.; Neuhold, Y.-M.; Hatwell, K. R.; Zuberbuhler, A. D.; Karlin, K. D. *Inorg. Chem.* **2001**, *40*, 5754–5767.