


Pd-Catalyzed Conjunctive Cross-Coupling between Grignard-Derived Boron "Ate" Complexes and C(sp²) Halides or Triflates: NaOTf as a Grignard Activator and Halide Scavenger

Gabriel J. Lovinger, Mark D. Aparece, and James P. Morken*[✉]

Department of Chemistry, Merkert Chemistry Center, Boston College, Chestnut Hill, Massachusetts 02467, United States

S Supporting Information


ABSTRACT: Catalytic enantioselective conjunctive cross-couplings that employ Grignard reagents are shown to furnish an array of nonracemic chiral organoboronic esters in an efficient and highly selective fashion. The utility of sodium triflate in facilitating this reaction is two-fold: it enables "ate" complex formation and overcomes catalytic inhibition by halide ions.



INTRODUCTION

Chiral organoboron compounds are important synthetic intermediates that can be converted to a broad array of chiral materials by mild stereospecific reactions.¹ For this reason, intense efforts have been directed toward the construction of these compounds in an enantioselective fashion. While ground breaking work by Brown established the utility of chiral organoboranes,² organoboronic esters have become the preferred class of boron reagents in contemporary asymmetric synthesis because of their stability and reactivity properties. With these reagents, powerful stoichiometric asymmetric homologation reactions have enabled a broad array of C–C and C–heteroatom bond constructions directly from the organoboronic ester starting materials.³ Similarly, many catalytic processes have been developed that provide access to chiral organoboronic esters from simple building blocks.⁴ In this vein, our laboratory recently described a catalytic conjunctive cross-coupling reaction (Scheme 1) that operates on alkenyl boronic ester-derived "ate" complexes and produces enantiomerically enriched alkyl boronic esters in an efficient fashion.⁵ Since the alkenyl boron "ate" complex is generated in situ from the reaction of appropriate organoboronic esters and organolithium compounds, the conjunctive cross-coupling represents a three-component reaction that merges simple starting materials to generate chiral organoboronic ester products. Because the conjunctive process depicted in Scheme 1 may offer streamlined and convergent routes for the synthesis of chiral compounds, we sought to enhance the operational utility of the reaction by reengineering it to employ readily available and functional-group-tolerant Grignard reagents in place of organolithium compounds. We also aimed to extend the substrate scope from C(sp²) triflate electrophiles to common and inexpensive C(sp²) halide electrophiles.

Scheme 1. Pd-Catalyzed Conjunctive Cross-Coupling Reaction




In this Article, we describe experiments that reveal the mechanistic challenges associated with extending the conjunctive cross-coupling reaction in these directions, and we present an effective solution that enables broadly useful reactions with aryl halides and that also allows use of Grignard-derived "ate" complexes.

As an operative mechanistic hypothesis to guide development of transition-metal catalyzed conjunctive cross-coupling, it was considered that a π -acidic cationic Pd(II) catalyst is generated by oxidative addition of an L_nPd(0) complex to an aryl triflate (Scheme 2). Subsequent association of the Pd(II) adduct with the alkenylboron "ate" complex is proposed to induce a metalate rearrangement; isotope labeling experiments indicate that this elementary reaction occurs with an

Received: December 13, 2016

Published: February 5, 2017

Scheme 2. Proposed Mechanism for the Pd-Catalyzed Conjointive Cross-Coupling Reaction



antiperiplanar relationship between the migrating group (R_M) and the Pd center (C). Rearranged complex D then releases the conjointive coupling product E upon reductive elimination. While the metal-induced metallate rearrangement step (C \rightarrow D) finds some precedent in the stoichiometric rearrangement reactions of Pt acetylides described by Wrackmeyer⁶ and in Pd-catalyzed reactions of alkenyl alanes studied by Fillion,⁷ this elementary transformation has not been used in the catalytic construction of boronic esters. Indeed, in terms of precedent, the catalytic cycle in Scheme 2 is most closely aligned with a reaction studied by Murakami⁸ involving alkynylborane “ate” complexes, although Murakami proposes alkene carbopalladation followed by invertive reductive displacement of Pd(II) upon 1,2-metalate rearrangement.⁹ Aside from these important precedents, it should be noted that Deng¹⁰ and Ishikura¹¹ have documented relevant catalytic reactions involving outer-sphere addition of alkyne- and indole-derived boron “ate” complexes to palladium allyl and allenyl complexes by a process involving concomitant metallate shift. Similarly, a stoichiometric addition of an alkynyl boronate to an iron(pentadienyl) complex was noted to occur with metallate rearrangement.¹²

RESULTS AND DISCUSSION

Challenges in Conjointive Cross-Coupling: Boron “Ate” Complexes Derived from Grignard Reagents. As mentioned above, while the conjointive cross-coupling is efficient and highly selective, the initial implementation required halide-free organolithium-derived boron “ate” complexes. Because trace amounts of lithium halide salts often arise during the course of lithium-halogen exchange reactions (via elimination), obtaining halide-free organolithium reagents is technically demanding. With a goal of extending the conjointive cross-coupling reaction to readily accessible starting materials that may have greater functional group compatibility than organolithium reagents, we investigated reactions of “ate” complexes derived from commercially available Grignard reagents (Aldrich). As depicted in Figure 1A, when halide-free vinyl lithium was replaced with commercial vinylmagnesium bromide in THF for construction of the “ate” complex from PhB(pin), the conjointive cross-coupling reaction failed to deliver any product. To learn about underlying reasons for the difference in reactivity between the lithium- and magnesium-based reagents, the reaction of the vinyl metal reagents and





Figure 1. (A) Comparison of conjointive cross-coupling with vinyl lithium (eq 1) versus vinylmagnesium bromide (eq 2). (B) ¹¹B NMR analysis of the reaction between vinyl metal reagents and PhB(pin).

PhB(pin) was analyzed by ¹¹B NMR spectroscopy. As depicted in Figure 1B, the reaction between halide-free vinyl lithium and PhB(pin) leads to rapid disappearance of the resonance for PhB(pin) ($\delta = 30.9$ ppm) and appearance of a new resonance at $\delta = 5.8$ ppm consistent with efficient formation of a four-coordinate boron species.¹³ In contrast, the analogous reaction with vinylmagnesium bromide provides ca. 20% conversion to the putative “ate” complex. While formation of “ate” complexes from Grignard reagents has not been studied in significant detail previously, extant reports describing the reaction between Grignard reagents and α -haloboronates¹⁴ imply that access to “ate” complexes as transient reactive intermediates is feasible; however, a report by Blakemore¹⁵ suggests that formation of Grignard-derived “ate” complexes may be more difficult than the analogous reactions of lithium derivatives, as is observed here. Thus, one significant challenge to employing Grignard reagents in conjointive couplings may arise from their diminished nucleophilicity, and hence diminished ability to generate the requisite “ate” complexes, as compared to organolithium reagents.

Challenges in Conjointive Cross-Coupling: Inhibition by Halide Salts. While inefficient “ate” complex formation with vinylmagnesium bromide accounts for diminished efficiency of conjointive cross-coupling with the magnesium-based process, the lack of any conjointive coupling product at all, even when ca. 20% “ate” complex was generated from the

Grignard reagent, suggests other effects might also be operative. In particular, the above-mentioned observation that conjunctive couplings are ineffective with lithium halide-containing “ate” complexes, and that conjunctive couplings are much less effective with aryl and alkenyl halides, suggested that endogenous halide ions might inhibit the catalytic process. To probe the inhibitory effect of halides on conjunctive coupling, the experiments in Table 1 were conducted. Relative

Table 1. Halide Inhibition in Conjunctive Coupling^a




entry	alteration	yield ^b	er ^c
1	none	77	98:2
2	1% LiI	13	98:2
3	1% LiBr	41	98:2
4	1% LiCl	40	98:2
5	1% (<i>n</i> -Bu) ₄ NCl	31	98:2
6	1% LiI, 5% catalyst	69	98:2
7	PhCl instead of PhOTf	<5	nd
8	PhBr instead of PhOTf	9	96:4
9	PhI instead of PhOTf	9	96:4
10	100% LiBr	23	78:22
11	100% <i>n</i> -Bu ₄ NBr	19	92:8

^aThe “ate” complex was prepared by addition of *n*-butyllithium to vinylB(neo) and the conjunctive coupling was conducted at 0.17 M. ^bYield represents isolated yield of purified material. ^cEnantiomer ratio (er) determined by chiral SFC analysis.

to the standard reaction conditions with 1 mol % catalyst loading and LiI free “ate” complex (entry 1, vinyl lithium obtained from tetravinylstannane by Li–Sn exchange¹⁶ or from vinyl iodide and *n*-BuLi, followed by recrystallization from ether¹⁷), addition of 1 mol % LiI leads to a significant erosion of conjunctive cross-coupling efficiency (cf. entries 1 and 2), but an otherwise high level of enantioselectivity. The observation that the inhibitory effect of 1 mol % LiI is ameliorated by conducting the reaction with 5 mol % Pd(OAc)₂/Mandyphos (L1¹⁸) (entry 6) suggests that the effect may be due to interaction of LiI and a catalytically active complex. Considering the inhibitory effect of LiI it is unsurprising that conjunctive cross-coupling of aryl halide electrophiles are inefficient even when lithium-halide-free “ate” complexes are employed: conjunctive cross-coupling itself releases lithium halide as a direct product of the reaction.

The halide inhibition observed in conjunctive coupling reactions is consonant with the working hypothesis for the reaction mechanism wherein π -bonding between a cationic transition metal complex and the reacting “ate” complex is a necessary prerequisite. In this scenario, it is plausible that halide ions outcompete the olefin for binding to palladium (eq 3),



thereby leading to reaction inhibition. In related stoichiometric processes, the presence of halide ions has been shown to inhibit carbopalladation of alkenes by sequestering cationic palladium complexes.¹⁹ Similarly, while catalytic Heck,²⁰ Stille,²¹ and other²² reactions often exhibit acceleration due to the presence of halide ions, this effect is generally traced to an acceleration of oxidative addition; when oxidative addition is not the slow step of catalysis, then halide ions can act as inhibitors.²³ Indeed, halide inhibition of catalysis has been documented in the case of catalytic Heck reactions²⁴ and catalytic activation/cross-coupling of cyclopropanes.²⁵

Effect of Additives on Boron “Ate” Complex Formation from Grignard Reagents and Subsequent Conjunctive Coupling. The above-described studies suggest that development of a strategy for effective “ate” complex formation and concomitant removal of halide ions from the reaction medium might enable conjunctive cross-coupling reactions with Grignard reagents and also allow the use of organic halide electrophiles. It was reasoned that efficient construction of boron “ate” complexes from Grignard reagents might be facilitated by the addition of appropriate additives that enhance the reactivity of magnesium-based reagents. Along these lines we first examined the capacity for LiCl to facilitate “ate” complex formation. Important studies by Knochel have shown that lithium chloride increases the reactivity of Grignard reagents in Mg-halogen exchange reactions²⁶ and increases the reactivity of Hauser bases²⁷ (in the form of so-called “turbo Grignard reagents” and “turbo Hauser bases”). While we considered that addition of LiCl would likely compound the problem of halide inhibition in catalytic conjunctive coupling reactions, it would nonetheless reveal the capacity for additives to enhance “ate” complex formation. As depicted in Figure 2, when vinylmagnesium chloride was added to PhBpin in THF in the presence of 1 equiv of LiCl, complete conversion to the “ate” complex was observed by ¹¹B NMR analysis. Unsurprisingly, the resulting complex was unreactive in conjunctive coupling. In an effort to uncover additives that are less likely to inhibit conjunctive coupling reactions, we studied the capacity




Figure 2. ¹¹B NMR of the reaction between vinylMgCl and PhB(pin) in the presence of additives.

for other weakly basic metal salts to promote “ate” complex formation (see Supporting Information for complete list). It was found that both LiOTf and NaOTf can effect “ate” complex formation, although the later ion pair is less effective and requires 2 equiv to achieve >95% conversion to the “ate” complex. In terms of mechanistic features, it is worth noting the effectiveness of Bu₄NOTf in facilitating conversion to the “ate” complex. While NaOTf has been proposed to activate Grignard reagents by halide abstraction to generate RMgOTf,²⁸ the lack of Lewis acidity of Bu₄NOTf suggests that the ability of additives to facilitate “ate” complex formation stems predominantly from Lewis basicity of the triflate as opposed to either halide abstraction or Lewis acid activation of the boronate through pinacolate O → LA donation.²⁹ Although, it should be noted that NaBPh₄, a nonbasic Lewis acid, also promotes “ate” complex formation, albeit less effectively than Bu₄NOTf; this observation suggests that Lewis acid association may play a beneficial but less significant role in promoting association between the boronic ester and Grignard reagent.


To minimize halide inhibition of catalysis it was considered that cations that are able to ion pair with halide might serve as scavenging agents and facilitate catalysis. While AgOTf was considered to be a reasonable candidate for such a strategy, addition of AgOTf was found to rapidly decompose the halide-free lithium-derived “ate” complex as determined by ¹¹B NMR analysis. We speculated that nonredox active cationic metals whose halide salts are either insoluble or tightly ion-paired in THF solvent might be introduced as metal triflates and, upon anion exchange, act as effective scavengers. In this connection, while the issue of LiCl solubility in THF has received attention,³⁰ the solubility of other metal halides has not been reported. To aid in the interpretation of reaction outcomes, the solubility of a series of metal salts in THF solvent was measured. In these experiments, 1–3 g of anhydrous salt was stirred in 21 mL anhydrous tetrahydrofuran for 24 h at room temperature. The solution was allowed to stand undisturbed overnight, the supernatant then filtered to remove remaining nondissolved salt, and a 10 mL portion of the solution evaporated to constant weight. Using this procedure, the data in Table 2 was collected.

Table 2. Solubility of Metal Salts in Anhydrous THF Solvent^a

salt	solubility (mg/mL)	solubility (M)
LiCl	49.5	1.12
LiBr	388	4.47
LiOTf	473	3.03
NaCl	0.20	0.0036
NaCl (1:1 THF:DMSO) ^b	0.32	0.0055
NaBr	0.15	0.0015
NaOTf	220	1.30
NaOTf (1:1 THF:DMSO) ^b	289	1.68
KCl	0.30	0.0039
KBr	0.30	0.0022
KOTf	4.0	0.0213
MgCl ₂	40.6	0.427
Mg(OTf) ₂	4.4	0.014

^aSee text for procedural details. ^bFor these entries, saturated concentration determined by slowly adding salt to solvent until the solution remained turbid.

Considering the remarkable difference in solubility between NaOTf and NaCl, as well as the ability of NaOTf to promote “ate” complex formation from Grignard reagents, this additive was examined in conjunctive coupling reactions that employ vinylmagnesium chloride. Of note, NaOTf³¹ is a commercially available and inexpensive salt. After optimization of solvent (vide infra for discussion of solvent effects), catalyst loading, stoichiometry, reaction time and temperature (see SI, Table S2), the optimal set of conditions were found to be as depicted in eq 4. As shown, a commercially available solution of



vinylmagnesium chloride in THF (Aldrich) was added to a mixture of PhB(pin) and two equivalents of NaOTf in THF/DMSO, followed by Pd(OAc)₂, ligand L1, and PhOTf. After reaction at 40 °C for 24 h, the conjunctive coupling product was isolated in 81% yield and 96:4 er. This level of selectivity and reaction efficiency is comparable to that obtained with “ate” complexes prepared from halide-free vinylolithium reagents (eq 1, Figure 1). It is worth noting that commercial vinylmagnesium bromide could also be utilized with similar levels of yield and selectivity when KOTf (2 equiv) was employed as an additive. In addition to its ability to facilitate the reaction of Grignard-derived “ate” complexes with aryl or alkenyl triflates, the apparent halide-scavenging ability of NaOTf can also enable the reaction of aryl bromide electrophiles: with an additional equivalent of NaOTf added, these electrophiles also engage in efficient and selective conjunctive coupling with Grignard-derived “ate” complexes (eq 5).³² Aryl iodides behave similarly with PhI reacting in 81% yield and 97:3 er (data not shown).

Formation and Stability of Boron Ate Complexes: Solvent Effects. While NaOTf clearly facilitates formation of boron “ate” complexes from Grignard reagents and can enable conjunctive coupling of Mg-based reagents, two features remained challenging. First, even with NaOTf, formation of “ate” complexes from Grignard reagents and alkyl boronic esters is not efficient (ca. 50% conversion to “ate” complex). Second, whereas the Li-based boron “ate” complexes have good long-term stability, the Mg-based reagents lack stability over a time course comparable to a typical catalytic reaction. These aspects were probed by ¹¹B NMR and the data is depicted in Figure 3. When 3-butenylB(pin) was reacted with vinylolithium, the derived “ate” complex forms immediately (data not shown) and is stable even after 24 h at room temperature. In contrast, even with NaOTf added, vinyl magnesium chloride only converts ca. 50% of the alkylB(pin) substrate to the derived “ate” complex at 1 h and this complex is not stable: after standing for 24 h, the complex is largely converted back to three-coordinate boron species (mixture of (3-butenyl)Bpin and vinylBpin).

Considering that the polarity and/or coordinating ability of the reaction medium might enhance the stability of “ate” complexes, we examined complexation reactions in different





Figure 3. ^{11}B NMR of reactions between vinyl metal reagents and 3-butenylB(pin).

solvents. Most effective was the inclusion of DMSO as a cosolvent: as depicted in Figure 3, the reaction of vinylmagnesium chloride with the alkylB(pin) substrate and NaOTf in THF/DMSO (1:1) solvent mixture proceeds in high conversion and furnishes a boron “ate” complex that persists with little change even after 24 h at room temperature (data not shown) or 40 °C (Figure 3). This observation is expected to aid in the development of practical conjunctive coupling reactions, and may also prove useful in the design of other boron-based processes.

That the increased stability of Grignard-derived boron “ate” complexes in the presence of DMSO cosolvent translates to increased reaction yield, can be ascertained by examining the data in Table 3. Across a small selection of boronic esters and electrophiles, comparison of conjunctive couplings in either

Table 3. Effect of Solvent on Catalytic Conjunctive Coupling Reactions^a



entry	R _M	Ar	solvent	yield ^b	er ^c
1	3-butenyl	Ph	THF	<5%	n/a
2	3-butenyl	Ph	THF/DMSO	73	91:9
3	Ph	Ph	THF	72	94:6
4	Ph	Ph	THF/DMSO	81	96:4
5	Ph	p-MeO-Ph	THF	78	92:8
6	Ph	p-MeO-Ph	THF/DMSO	89	98:2
7	Ph	p-CF ₃ -Ph	THF	67	75:25
8	Ph	p-CF ₃ -Ph	THF/DMSO	85	95:5




^aReactions conducted as described in the text (see SI for additional details) and the conjunctive coupling was conducted at 0.17 M. ^bYield represents isolated yield of purified material. ^cEnantiomer ratio (er) determined by chiral SFC analysis of boronic ester.

THF solvent and THF/DMSO (1:1) revealed that, while reactions are slower in the solvent mixture (24 h vs 12 h reaction time), the selectivity and yield are enhanced in the presence of DMSO. Most strikingly, conjunctive couplings with the alkylB(pin) derivative are ineffective in THF solvent, whereas in THF/DMSO a reasonable yield and selectivity are obtained (entries 1 and 2). Notably, even the reaction of arylB(pin) reagents appear to benefit by the inclusion of DMSO cosolvent (cf. entry 3 and 4), although the beneficial effect is less substantial in these cases. Overall, the data presented in the above two sections argues for the use of NaOTf additive and THF/DMSO solvent mixture for effective and general catalytic conjunctive couplings of Grignard-derived complexes. These conditions were surveyed across a range of substrates (vide infra).

Scope of Catalytic Conjunctive Cross-Coupling with Grignard-Derived “Ate” Complexes. With effective conditions established to employ Grignard-derived “ate” complexes in conjunctive couplings, the scope of the catalytic asymmetric transformation was surveyed. As depicted in Table 4, it was found that the Mg-based system allows conjunctive couplings with a range of aromatic carbocycles, heterocycles and olefinic organotriflate electrophiles. Of note, an aldehyde group attached to the electrophile (product 6) survives the reaction intact, an observation that points to the buffering effect the boron atom imposes on the precursor nucleophilic Grignard reagent. Importantly, labile functional groups such as nitriles, amides, esters, ketones, aldehydes, halides, and unprotected alcohols (employing two equivalents of nucleophile) all survive “ate” complex formation and conjunctive coupling, and are incorporated into nonracemic products selectively and with useful levels of reaction efficiency. It is worth noting that electron-deficient migrating groups such as a *para*-trifluoromethylphenyl group can be employed successfully (product 25) with the current conditions whereas with the lithium-derive “ate” complex in THF solvent the reaction furnished <5% product. Lastly, it should be pointed out that conjunctive couplings involving vinylB(pin) and alkyl or aryl Grignard reagents (method A) also appear to be effective.



A survey of the substrate scope involving organobromide electrophiles is depicted in Table 5. Of note, the yield and selectivity with this substrate class parallels that observed when using organotriflate electrophiles so long as an added equivalent of NaOTf is included in the reaction mixture. In these reactions, it was possible to demonstrate compatibility with furan, thiophene, quinoline, pyridine, pyrimidine, indole, benzothiazole and other functionalized organic bromides suggesting that a large collection of targets may ultimately be available from conjunctive couplings, even when the corresponding organotriflate electrophile is not readily available. It should be noted that under the current conditions, alkenyl bromides are significantly less effective than alkenyl triflate electrophiles (i.e., substrate 8 of Table 4 is prepared in 34% yield, 94:6 er from the alkenyl bromide versus 76% yield, 96:4 er from the triflate).

Conjunctive Cross-Coupling with C(sp₂) Triflates and Li-Based Boron “Ate” Complexes Derived from Li-Halogen Exchange. In addition to enabling reactions of Grignard reagents, it was considered that the halide-scavenging ability of NaOTf might enable the direct use of organolithium reagents generated by lithium-halogen exchange (i.e., without taking efforts to remove lithium halide byproducts). Thus, vinyl lithium, prepared by treatment of vinyl bromide with 2

Table 4. Conjunctive Coupling between Grignard-Derived Boron “Ate” Complexes and Organotriflates^a

^aConjunctive coupling was conducted at 0.17 M. Yields represent isolated yields of purified material. Both the yield and the enantiomer ratio (er) represent the average value for two experiments. ^bProduct isolated as the derived alcohol. ^cReaction conducted at 60 °C. ^dSolvent = THF.


equiv *tert*-butyllithium was directly added to PhB(pin) in the presence of NaOTf. The derived “ate” complex was then engaged in conjunctive coupling with PhOTf. In this experiment (data not shown) the cross-coupling furnished **1** in 70% yield and 95:5 er. While addition of NaOTf was clearly effective, it was found that KOTf performs somewhat better as a LiBr scavenger for aryl triflate electrophiles whereas NaOTf provided higher selectivity when alkenyl triflate electrophiles were employed (Table 5).

Table 5. Conjunctive Coupling between Grignard-Derived Boron “Ate” Complexes and Organic Bromides^a

^aConjunctive coupling was conducted at 0.17 M. Yields represent isolated yields of purified material. Both the yield and the enantiomer ratio (er) represent the average value for two experiments. ^bReaction conducted at 55 °C.



Functional Group Compatibility in Conjunctive Cross-Coupling: Boron-Buffered Nucleophilicity via Kinetic Trapping.

Aspects of the functional group tolerance exhibited during the course of catalytic conjunctive coupling reactions are informative and merit comment. The examples in Tables 4–6 above reveal a range of functional groups—either attached to the electrophile or the “ate” complex—that are compatible with catalytic conjunctive coupling reactions. While compatibility of functional groups attached to the electrophile (i.e., product **6**, Table 5) can be anticipated because of the modest basicity and nucleophilicity of “ate” complexes,³³ the ability to assemble functionalized “ate” complexes by reacting functionalized boronic esters with organometallic compounds (Grignard and organolithium reagents) is less anticipated.³⁴ For example, effective production of compounds **22–24** in Table 6 implies that amides, esters, and alkyl halides survive treatment with organolithium reagents. It was considered that this functional group compatibility likely arises from the boronic ester’s ability to act as both a kinetic and thermodynamic trap that protects against direct reaction of strong nucleophiles with labile functional groups. It was expected that this feature of “ate” complex formation might ultimately allow conjunctive couplings to be operated as a three-component reaction without the need to pregenerate the boron “ate” complex in situ before introduction of the electrophile and the catalyst. To further

Table 6. Conjunctive Coupling with Halide-Containing Organolithium-Derived Boron “Ate” Complexes^a

^aConjunctive coupling was conducted at 0.17 M. Yields represent isolated yields of purified material. Both the yield and the enantiomer ratio (er) represent the average value for two experiments. ^bProduct isolated as the derived alcohol. ^cNaOTf was employed in place of KOTf and the solvent was THF. ^dReaction conducted at $60\text{ }^\circ\text{C}$.

probe the rapidity of “ate” complex formation relative to reaction of organometallic reagents with other functional groups, more challenging competition experiments were conducted. In the first, a mixture of vinylB(pin) and bromobenzene was treated with *tert*-butyllithium at $-98\text{ }^\circ\text{C}$ in ether (eq 6); the solvent was then removed and the ¹¹B



NMR in THF obtained. In this experiment, the ¹¹B NMR resonance corresponding to vinylB(pin) (δ 29.6 ppm) was replaced with a resonance at 7.8 ppm corresponding to “ate” complex **43**; a resonance at 5.8 ppm corresponding to PhLi (generated by Li/Br exchange) addition to vinylB(pin) was not observed. Thus, addition of *tert*-butyllithium to vinylB(pin) appears to outcompete Li–Br exchange. In a similar experiment (eq 7), it was found that addition of *tert*-butyllithium to vinylB(pin) also outcompetes addition of the alkyl lithium to benzaldehyde.

Glovebox-Free and Preparative Scale Procedure. To probe the capacity for conjunctive couplings to be conducted without the aid of a glovebox, as a three-component assembly without preformation of “ate” complexes, and on preparative scale, we examined the reactions shown in Figure 4. In these




Figure 4. Conjunctive couplings conducted without the aid of a glovebox.

experiments, the solid reagents were weighed in the open atmosphere, combined and transferred to a dried flask, and then the headspace of the reaction vessel purged with dry nitrogen gas. After addition of liquid reagents and solvent, the flask was cooled to $0\text{ }^\circ\text{C}$, the Grignard reagent added, and the reaction then allowed to proceed at the indicated temperature overnight. With this straightforward procedure, the derived conjunctive coupling products can be obtained in good yield and outstanding enantioselectivity. As depicted in eq 8, the reaction can be operated in this manner even on preparative scale and provides functionalized products such as **42** in a practical fashion. Lastly, this procedure applies regardless of whether the substrates are both solids (eq 8), both liquids (eq 10) or one of each (eq 9).

CONCLUSIONS

We have identified that key challenges associated with the use of Grignard-derived boronic ester “ate” complexes arise from a combination of ineffective “ate” complex formation and inhibition of conjunctive coupling reactions by the presence of halide ions. The latter problem likely contributes to the diminished reactivity of organic halides in conjunctive couplings as well. The addition of NaOTf or KOTf largely counteracts these problems and provides a convenient and broad-scoped catalytic conjunctive coupling process. We anticipate that these reactions may find use in organic synthesis and that the utility of alkali metal triflates may find use in development of other catalytic processes involving boron reagents.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/jacs.6b12663.

Procedures, characterization and spectral data (PDF)

Crystal data (CIF)

AUTHOR INFORMATION

Corresponding Author

*morken@bc.edu

ORCID

James P. Morken: 0000-0002-9123-9791

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

This work was supported by the NIH (GM-118641). We thank Solvias for supporting this program by providing MandyPhos ligand.

REFERENCES

- (1) Reviews, see: (a) Brown, H. S.; Singaram, B. *Acc. Chem. Res.* **1988**, *21*, 287. (b) *Organoboranes for Synthesis*; Ramachandran, P. V., Brown, H. C., Eds.; American Chemical Society: Washington, D.C., 2001; ACS Symposium Series 783. (c) *Stereodirected Synthesis with Organoboranes*; Matteson, D. S., Ed.; Springer: New York, 1995. (d) Leonori, D.; Aggarwal, V. K. *Acc. Chem. Res.* **2014**, *47*, 3174.
- (2) (a) Brown, H. C.; Ramachandran, P. V. *Pure Appl. Chem.* **1991**, *63*, 307. (b) Brown, H. C.; Ramachandran, P. V. *Acc. Chem. Res.* **1992**, *25*, 16.
- (3) (a) Negishi, E.; Idacavage, M. J. *Org. React.* **1985**, *33*, 1. (b) Aggarwal, V. K.; Fang, G. Y.; Ginesta, X.; Howells, D. M.; Zaja, M. *Pure Appl. Chem.* **2006**, *78*, 215. (c) Leonori, D.; Aggarwal, V. K. *Top. Organomet. Chem.* **2015**, *49*, 271.
- (4) Reviews: (a) Hall, D. G.; Lee, J. C. H.; Ding, J. *Pure Appl. Chem.* **2012**, *84*, 2263. (b) Neeve, E. C.; Geier, S. J.; Mkhaliid, I. A. I.; Westcott, S. A.; Marder, T. B. *Chem. Rev.* **2016**, *116*, 9091. (c) Ferris, G. E.; Mlynarski, S. N.; Morken, J. P. Diborane(4) Compounds (Update 2012). In *Science of Synthesis*; Bode, J. W., Ed.; Thieme, 2012; Vol. 3, p 227.
- (5) Zhang, L.; Lovinger, G. J.; Edelstein, E. K.; Szymaniak, A. A.; Chierchia, M. P.; Morken, J. P. *Science* **2016**, *351*, 70.
- (6) Sebald, A.; Wrackmeyer, B. *J. Chem. Soc., Chem. Commun.* **1983**, 309.
- (7) (a) Fillion, E.; Carson, R. J.; Trépanier, V. E.; Goll, J. M.; Remorova, A. A. *J. Am. Chem. Soc.* **2004**, *126*, 15354. (b) Fillion, E.; Trépanier, V. E.; Heikkinen, J. J.; Remorova, A. A.; Carson, R. J.; Goll, J. M.; Seed, A. *Organometallics* **2009**, *28*, 3518.
- (8) (a) Ishida, N.; Shimamoto, Y.; Murakami, M. *Org. Lett.* **2009**, *11*, 5434. (b) Ishida, N.; Shinmoto, T.; Sawano, S.; Miura, T.; Murakami, M. *Bull. Chem. Soc. Jpn.* **2010**, *83*, 1380. (c) Ishida, N.; Narumi, M.; Murakami, M. *Org. Lett.* **2008**, *10*, 1279. (d) Ishida, N.; Shimamoto, Y.; Murakami, M. *Org. Lett.* **2010**, *12*, 3179. (e) Ishida, N.; Ikemoto, W.; Narumi, M.; Murakami, M. *Org. Lett.* **2011**, *13*, 3008.
- (9) Preliminary DFT studies are supportive of the reaction proceeding by the cycle in Scheme 2 as opposed to the carbopalladation-based pathway.
- (10) Chen, Y.; Li, N.-S.; Deng, M.-Z. *Tetrahedron Lett.* **1990**, *31*, 2405.
- (11) (a) Ishikura, M.; Terashima, M.; Okamura, K.; Date, T. *J. Chem. Soc., Chem. Commun.* **1991**, 1219. (b) Ishikura, M.; Agata, I. *Heterocycles* **1996**, *43*, 1591. (c) Ishikura, M.; Matsuzaki, Y.; Agata, I.; Katagiri, N. *Tetrahedron* **1998**, *54*, 13929. (d) Ishikura, M.; Agata, I.; Katagiri, N. *J. Heterocycl. Chem.* **1999**, *36*, 873. (e) Ishikura, M.; Hino, A.; Yaginuma, T.; Agata, I.; Katagiri, N. *Tetrahedron* **2000**, *56*, 193. (f) Ishikura, M.; Kato, H. *Tetrahedron* **2002**, *58*, 9827. (g) Ishiura, M.; Ida, W.; Yanada, K. *Tetrahedron* **2006**, *62*, 1015.
- (12) Pelter, A.; Gould, K. J. *J. Chem. Soc., Chem. Commun.* **1974**, 1029.
- (13) For ¹¹B NMR of four-coordinate pinacol boron ate complexes, see: Wrackmeyer, B. *Annu. Rep. NMR Spectrosc.* **1988**, *20*, 61.
- (14) Brown, H. C.; De Lue, N. R.; Yamamoto, Y.; Maruyama, K.; Kasahara, T.; Murahashi, S.; Sonoda, A. *J. Org. Chem.* **1977**, *42*, 4088.
- (15) Blakemore, P. R.; Marsden, S. P.; Vater, H. D. *Org. Lett.* **2006**, *8*, 773.
- (16) Seyferth, D.; Weiner, M. A. *J. Am. Chem. Soc.* **1961**, *83*, 3583.
- (17) Shinokubo, H.; Miki, H.; Yokoo, T.; Oshima, K.; Utimoto, K. *Tetrahedron* **1995**, *51*, 11681.
- (18) Almena Perea, J. J.; Lotz, M.; Knochel, P. *Tetrahedron: Asymmetry* **1999**, *10*, 375.
- (19) (a) Kawataka, F.; Shimizu, I.; Yamamoto, A. *Bull. Chem. Soc. Jpn.* **1995**, *68*, 654. (b) Kayaki, Y.; Shimizu, I.; Yamamoto, A. *Chem. Lett.* **1995**, *24*, 1089. (c) Amatore, C.; Carré, E.; Jutand, A.; M'Barki, M. A.; Meyer, G. *Organometallics* **1995**, *14*, 5605. (d) Amatore, C.; Godin, B.; Jutand, A.; Lemaitre, F. *Chem. - Eur. J.* **2007**, *13*, 2002. (e) Amatore, C.; Godin, B.; Jutand, A.; Lemaitre, F. *Organometallics* **2007**, *26*, 1757.
- (20) (a) Jeffrey, T. *J. Chem. Soc., Chem. Commun.* **1984**, 1287. (b) Jeffrey, T. *Tetrahedron Lett.* **1985**, *26*, 2667. (c) Grigg, R.; Stevenson, P.; Worakun, T. *Tetrahedron* **1988**, *44*, 2033. (d) Andersson, C.-M.; Hallberg, A. *J. Org. Chem.* **1988**, *53*, 2112.
- (21) (a) Scott, W. J.; Crisp, G. T.; Stille, J. K. *J. Am. Chem. Soc.* **1984**, *106*, 4630. (b) Scott, W. J.; Stille, J. K. *J. Am. Chem. Soc.* **1986**, *108*, 3033. (c) Scott, W. J.; McMurry, J. E. *Acc. Chem. Res.* **1988**, *21*, 47.
- (22) Beletskaya, I. P. *J. Organomet. Chem.* **1983**, *250*, 511.
- (23) Jutand, A. *Appl. Organomet. Chem.* **2004**, *18*, 574.
- (24) (a) Ozawa, F.; Kubo, A.; Hayashi, T. *J. Am. Chem. Soc.* **1991**, *113*, 1417.
- (25) Aoki, S.; Fujimura, T.; Nakamura, E.; Kuwajima, I. *J. Am. Chem. Soc.* **1988**, *110*, 3296.
- (26) (a) Krasovskiy, A.; Knochel, P. *Angew. Chem., Int. Ed.* **2004**, *43*, 3333. (b) Barl, N. M.; Werner, V.; Sämann, C.; Knochel, P. *Heterocycles* **2014**, *88*, 827. (c) Klatt, T.; Markiewicz, J. T.; Sämann, C.; Knochel, P. *J. Org. Chem.* **2014**, *79*, 4253. (d) Tilly, D.; Chevallier, F.; Mongin, F.; Gros, P. C. *Chem. Rev.* **2013**, *114*, 1207. (e) Bao, R. L.-Y.; Zhao, R.; Shi, L. *Chem. Commun.* **2015**, *51*, 6884.
- (27) (a) Krasovskiy, A.; Krasovskaya, V.; Knochel, P. *Angew. Chem., Int. Ed.* **2006**, *45*, 2958. (b) Lin, W.; Baron, O.; Knochel, P. *Org. Lett.* **2006**, *8*, 5673. (c) Neufed, R.; Teuteberg, T. L.; Herbst-Imer, R.; Mata, R. A.; Stalke, D. *J. Am. Chem. Soc.* **2016**, *138*, 4796.
- (28) (a) Kendall, A. J.; Salazar, C. A.; Martino, P. F.; Tyler, D. R. *Organometallics* **2014**, *33*, 6171. See also (b) Reetz, M. T.; Harmat, N.; Mahrwald, R. *Angew. Chem., Int. Ed. Engl.* **1992**, *31*, 342.
- (29) For examples of boronate activation by Lewis acid-oxygen binding, see: (a) Corey, E. J.; Barnes-Seeman, D.; Lee, T. W. *Tetrahedron: Asymmetry* **1997**, *8*, 3711. (b) Midland, M. *J. Org. Chem.* **1998**, *63*, 914. (c) Rauniyar, V.; Hall, D. G. *J. Am. Chem. Soc.* **2004**, *126*, 4518.
- (30) Krasovskiy, A.; Knochel, P. *Synthesis* **2006**, 2006, 890.
- (31) Surya Prakash, G. K.; Mathew, T. *e-EROS Encyclopedia of Reagents for Organic Synthesis* **2010**, DOI: 10.1002/047084289X.rm01137.
- (32) For examples of halide abstraction by NaOTf, see: From L_nRuCl (a) Quebatte, L.; Scopelliti, R.; Severin, K. *Eur. J. Inorg. Chem.* **2005**, *2005*, 3353. From L_nI(III)Cl (b) Brantley, J. N.; Samant, A. V.; Toste, F. D. *ACS Cent. Sci.* **2016**, *2*, 341. L_nAuCl (c) Serra, D.; Moret, M.-E.; Chen, P. *J. Am. Chem. Soc.* **2011**, *133*, 8914. L_nPdCl (d) Delis, J. G. P.; Groen, J. H.; Vrieze, K.; van Leeuwen, P. W. N. M.; Veldman, N.; Spek, A. L. *Organometallics* **1997**, *16*, 551.
- (33) (a) Feeney, K.; Berionni, G.; Mayr, H.; Aggarwal, V. K. *Org. Lett.* **2016**, *17*, 2614. (b) Berionni, G.; Leonov, A. I.; Mayer, P.; Ofial, A. R.; Mayr, H. *Angew. Chem., Int. Ed.* **2015**, *54*, 2780.
- (34) For addition of alkyllithium reagents to B(pin) derivatives bearing esters and amides, see: (a) Odachowski, M.; Bonet, A.; Essafi, S.; Conti-Ramsden, P.; Harvey, J. N.; Leonori, D.; Aggarwal, V. K. *J. Am. Chem. Soc.* **2016**, *138*, 9521. (b) Armstrong, R. J.; García-Ruiz, C.; Myers, E. L.; Aggarwal, V. K. *Angew. Chem., Int. Ed.* **2017**, *56*, 786. (c) Wang, Y.; Noble, A.; Myers, E. L.; Aggarwal, V. K. *Angew. Chem., Int. Ed.* **2016**, *55*, 4270. Review: (d) Matteson, D. J. *J. Organomet. Chem.* **1999**, *581*, 51.