


Nickel-Catalyzed Cross Couplings of Benzylic Ammonium Salts and Boronic Acids: Stereospecific Formation of Diarylethanes via C–N Bond Activation

Prantik Maity,[†] Danielle M. Shacklady-McAtee,[†] Glenn P. A. Yap, Eric R. Sirianni, and Mary P. Watson*

Department of Chemistry and Biochemistry, University of Delaware, Newark, Delaware 19716, United States

S Supporting Information

ABSTRACT: We have developed a nickel-catalyzed cross coupling of benzylic ammonium triflates with aryl boronic acids to afford diarylmethanes and diarylethanes. This reaction proceeds under mild reaction conditions and with exceptional functional group tolerance. Further, it transforms branched benzylic ammonium salts to diarylethanes with excellent chirality transfer, offering a new strategy for the synthesis of highly enantioenriched diarylethanes from readily available chiral benzylic amines.



Diarylmethanes and diarylethanes are important molecules in organic synthesis and pharmaceutical development.¹ These valuable targets can be prepared via cross couplings of classic electrophiles, such as benzylic halides, with Grignard, organozinc, and organoboron reagents.² In fact, Carretero has shown that enantioenriched diarylethanes can be formed via stereospecific cross couplings of benzylic bromides with aryl Grignard reagents (Scheme 1A-1).^{2c} However, these classic electrophiles are highly reactive, can decompose upon prolonged storage, typically mandate that the cross coupling occur at the beginning of a synthetic sequence, and are difficult to prepare in high enantiopurity. Cross couplings of benzylic carbonates, acetates, and phosphates have also been developed to enable use of more stable starting materials.³ Recent reports have also demonstrated that the use of benzylic ethers and alcohols as coupling partners overcomes some of these challenges, offering greater substrate stability and orthogonality.⁴ Further, the Jarvo group has shown that nickel-catalyzed cross couplings of benzylic ethers with Grignard reagents occurs in a stereospecific fashion (Scheme 1A-2).^{5,6} In contrast, Tian has reported that copper-catalyzed couplings of benzylic sulfonimides proceed with only low levels of chirality transfer (Scheme 1A-3).⁷ Notably, almost all cross couplings of benzylic electrophiles to deliver enantioenriched diarylethanes require Grignard or organozinc coupling partners.^{2b–d,5} No examples of enantioselective couplings of benzylic electrophiles with boronic acids are yet known, and only a single stereospecific coupling of a benzylic α -cyanohydrin mesylate with a boronic acid partner has been reported to our knowledge.⁸

In an alternative bond-disconnection approach to the synthesis of enantioenriched diarylethanes, Crudden has shown that enantioenriched benzylic boronic esters undergo stereospecific cross couplings (Scheme 1A-4).⁹ Stereospecific protodeborylation of tertiary boronic esters also delivers enantioenriched diarylethanes.^{6c}

Despite these impressive advances in stereospecific cross couplings of benzylic reagents, several challenges remain, including identification of a class of benzylic reagents generally available in exceptional enantiopurity and the ability to employ commercially available and functional group tolerant coupling partners. In considering these challenges, we have been drawn to the use of benzylic ammonium salts as substrates.¹⁰ Benzylic ammonium salts are readily prepared from amine precursors.¹¹ Further, these substrates offer a functional group handle orthogonal to both halides and ethers, and both amines and ammonium salts are stable to long-term storage. Importantly, highly enantioenriched benzylic amines are readily available via classical resolution or a variety of catalytic asymmetric methods.¹² Csáký et al. have elegantly demonstrated the rhodium-catalyzed cross couplings of boronic acids with ammonium iodides derived from gramines (3-aminomethylindoles) to form achiral 3-benzyl- and 3-allylindoles.^{10d} However, to our knowledge, a general method for the cross coupling of benzylic ammonium salts to yield diarylmethanes and enantioenriched diarylethanes has not yet been realized.

We anticipated that nickel catalysts would enable the cross coupling of a wide variety of benzylic ammonium salts with mild and highly functional group compatible coupling partners, such as commercially available boronic acids. Because ammonium salts can be readily prepared with a wide variety of counterions, we envisioned facile tuning of the reactivity of these electrophiles. In particular, we predicted that weakly coordinating counterions, such as triflate, would be highly effective. In contrast to halide or alkoxide counterions formed in cross couplings with benzylic halides and ethers, the byproducts of oxidative addition into an ammonium triflate are neutral trimethylamine and the triflate counterion. We envisioned that the resulting oxidative addition intermediate


Received: September 8, 2012

Published: December 26, 2012

Scheme 1. Stereospecific Cross Couplings To Form Diarylethanes

A. Prior Art


(1) Benzylic Bromides (Carretero)




(2) Benzylic Ethers (Jarvo)




(3) Benzylic Sulfonimides (Tian)



(4) Benzylic Boronates (Crudden)



B. This Work: Benzylic Ammonium Triflates



may thus more readily undergo transmetalation with a boronic acid. Such an effect has been observed in cross couplings of aryl ammonium triflates.¹³

Herein we report the first example of a general method for the cross coupling of benzylic ammonium salts with boronic acids (Scheme 1B). To our knowledge, this is the first cross coupling of an aryl boronic acid with a benzylic amine derivative other than gramine and the first stereospecific coupling of an acyclic benzylic amine derivative with excellent chirality transfer.^{7,14,15} This method enables the preparation of diarylmethanes with broad substrate scope and diarylethanes with excellent enantioenrichment.

We began by investigating the cross coupling of *p*-tolyl boronic acid and benzyl ammonium salts **1**, which are easily prepared in quantitative yield via methylation of dimethyl benzyl amine.¹¹ Using conditions similar to those reported for Suzuki reactions of aryl ammonium salts,^{13a} low yields of diaryl methane **2** were observed when iodide **1a** was employed (Table 1, entry 1). In contrast, use of monodentate phosphine ligands, particularly mixed aryl/alkyl phosphines, provided increased yields (entries 2–4). As we had anticipated, the use of the less coordinating triflate counterion resulted in greater reactivity. Quantitative yield was observed when triflate **1b** was employed as the substrate (entry 5). Lowering the reaction temperature to 40 °C, PPh₂Cy proved to be the best ligand (entries 6–8). K₃PO₄ was as effective as CsF, resulting in nearly quantitative yield (entry 10). Notably, no product is observed in the absence of Ni catalyst (entries 12 and 13).¹⁶ Re-

Table 1. Optimization of Reaction Conditions^a

10 mol % Ni(cod)₂
ligand

base (1.3 equiv)
dioxane, temp, 24 h

1a, X = I
1b, X = OTf

2

entry	X	ligand (mol %)	base	temp (°C)	yield (%) ^b
1	I	IMes·HCl (11)	CsF	60	21
2	I	PCy ₃ (22)	CsF	60	30
3	I	PPhCy ₂ (22)	CsF	60	59
4	I	PPh ₂ Cy (22)	CsF	60	54
5	OTf	PPh ₂ Cy (22)	CsF	60	100
6	OTf	PPh ₂ Cy (22)	CsF	40	96
7	OTf	PPhCy ₂ (22)	CsF	40	16
8	OTf	PPh ₃ (22)	CsF	40	82
9	OTf	PPh ₂ Cy (22)	CsF	r.t.	68
10	OTf	PPh ₂ Cy (22)	K ₃ PO ₄	40	99
11	OTf	PPh ₂ Cy (22)	K ₃ PO ₄	r.t.	96
12 ^c	OTf	none	CsF	40	0
13 ^c	OTf	none	K ₃ PO ₄	40	0
14	OTf	IMes·HCl (11)	CsF	40	28


^aConditions: ammonium triflate **1** (0.10 mmol, 1.0 equiv), boronic acid (1.2 equiv), Ni(cod)₂ (10 mol %), ligand, CsF or K₃PO₄ (1.3 equiv), dioxane (0.33 M), 24 h, unless noted otherwise. ^bDetermined by ¹H NMR analysis using 1,3,5-trimethoxybenzene as internal standard. ^cNo Ni(cod)₂ used.

examination of the *N*-heterocyclic carbene IMes·HCl under the optimized conditions again showed phosphines to be superior ligands.


Using the optimized conditions (Table 1, entries 6 and 10), we observed broad substrate scope for both the boronic acid and ammonium triflate in the cross couplings of unbranched benzylic ammonium triflates. We generally observed similar results with CsF and K₃PO₄. However, for products with base-sensitive functional groups, such as **6** (Scheme 2), CsF proved advantageous. For the boronic acid, substitution was well tolerated at the *ortho*, *meta*, and *para* positions of the aromatic ring. High yields were observed for arylboronic acids with both electron-donating (**4**) and electron-withdrawing (**5–9**) substituents. Further, these mild reaction conditions tolerated a wide variety of functional groups, including ethers (**4**, **10**, **14**, **17**, **18**, **21**), fluorides (**5**, **12**, **13**, **15**), esters (**6**), secondary amides (**7**), sulfones (**8**), trifluoromethyls (**9**), free alcohols (**11**), and acetals (**19**). Naphthyl, pyridyl, and quinolinyl boronic acids were also effective partners (**20–23**).¹⁷

For the ammonium triflate partner, a variety of functional groups were tolerated at the *ortho*, *meta*, and *para* positions of the phenyl group (Scheme 3). Products containing ether (**24**, **26–28**), trifluoromethyl (**25**), ketone (**29**), and fluoride (**30**) substituents were formed in high yields. Notably, these reaction conditions selectively activate the benzylic C–N bond in the presence of either aryl or benzylic C–O bonds (**24**, **26–28**), highlighting the complementarity of the benzylic ammonium salts to ethers. This method also provides orthogonal reactivity to aryl halides. Although chloride-substituted benzyl ammonium triflates undergo cross coupling at both the C–Cl and C–NMe₃ bonds, aryl bromides can be effectively cross coupled in the presence of benzylic dimethylamino groups.¹⁸ Finally, naphthyl and indolyl^{10d} substrates are also effective in this reaction (**31**, **32**).

Encouraged by this broad substrate scope, we turned to the cross coupling of branched benzylic ammonium triflate **33** (Np

Scheme 2. Scope in Boronic Acid^a

^aConditions: ammonium triflate **1b** (0.30 mmol, 1.0 equiv), boronic acid (1.2 equiv), Ni(cod)₂ (10 mol %), PPh₂Cy (22 mol %), CsF or K₃PO₄ (1.3 equiv), dioxane (0.33 M), 40 °C, 24 h. Average isolated yield of duplicate experiments reported (±0–12%), unless otherwise noted. ^bResult of a single experiment.

Scheme 3. Scope in Ammonium Salt^a

^aConditions: ammonium triflate (0.20 mmol, 1.0 equiv), boronic acid (1.2 equiv), Ni(cod)₂ (10 mol %), PPh₂Cy (22 mol %), CsF or K₃PO₄ (1.3 equiv), dioxane (0.33 M), 40 °C, 24 h. Average isolated yield of duplicate experiments reported (±0–9%), unless otherwise noted. ^bResult of a single experiment.

= 2-naphthyl), which was readily prepared from (1S)-1-(2-naphthyl)ethanamine, purchased in 99.6% ee. Although application of the previously optimized conditions gave no desired product at 40 °C, high yields of diarylethane **34** were realized at a higher reaction temperature (Table 2, entry 1).

Table 2. Optimization of Branched Ammonium Salt^a

Reaction scheme for Table 2: $\text{Np-CH}_2\text{-NMe}_3^+\text{OTf}^- + (\text{HO})_2\text{B-p-Tol} \xrightarrow[\text{K}_3\text{PO}_4 (1.3 \text{ equiv}), \text{dioxane}, \text{temp}, 4 \text{ h}]{10 \text{ mol } \% \text{ Ni(cod)}_2, \text{ligand}}$

Product: **34**, 99.6% ee

entry	ligand (mol %)	temp (°C)	yield (%) ^b	ee (%) ^c
1	PPh ₂ Cy (22)	100	84	81
2	PPhCy ₂ (22)	100	97	79
3	PPh ₃ (22)	100	83	81
4	<i>t</i> -Bu-XantPhos (12)	100	(91)	98
5	XantPhos (12)	100	15	40
6	P(<i>o</i> -Tol) ₃ (22)	100	71	98
7	P(<i>o</i> -Tol) ₃ (22)	70	94	98
8 ^d	P(<i>o</i> -Tol) ₃ (22)	70	95	98
9 ^e	none	70	0	n.d. ^f
10 ^e	none	100	0	n.d. ^f

^aConditions: ammonium triflate **33** (0.10 mmol, 1.0 equiv), boronic acid (1.2 equiv), Ni(cod)₂ (10 mol %), ligand, K₃PO₄ (1.3 equiv), dioxane (0.4 M), 4 h, unless noted otherwise. ^bDetermined by ¹H NMR analysis using 1,3,5-trimethoxybenzene as internal standard. ^cDetermined by chiral HPLC. ^dPhMe replaced dioxane as solvent. ^eNo Ni(cod)₂ used. ^fn.d. = not determined.

However, only moderate chirality transfer was observed. A screen of ligands revealed that ligand affects the chirality transfer with 9,9-dimethyl-4,5-bis(di-*tert*-butylphosphino)-xanthene (*t*-Bu-XantPhos) and tri(*o*-tolyl)phosphine proving best (entries 4 and 6). Due to ligand cost, we selected P(*o*-Tol)₃ for further optimization and found that the reaction proceeds in high yield and enantiospecificity at 70 °C in either dioxane (entry 7) or PhMe (entry 8). Under these conditions, the reaction proceeds with excellent chirality transfer and yield, providing a powerful route to enantioenriched diarylethanes. Notably, no product is observed in the absence of nickel catalyst at these elevated temperatures (entries 9 and 10).¹⁶

With these conditions in hand, we examined the scope of this transformation. Upon increasing the reaction scale, we found that the catalyst loading could be lowered to 3 mol % without detrimental effect on yield or enantioselectivity of diarylethane **34** (Table 3, entry 1). In some cases, lower catalyst loading resulted in higher enantiospecificities, suggesting a possible Ni-mediated epimerization pathway (entries 1 vs 2, 12 vs 13). Electron-rich and electron-neutral aryl boronic acids perform well in this reaction (entries 1–11). Further, a wide variety of functional groups are tolerated, including ethers, olefins, fluorides, esters, nitriles, and sulfones. In terms of electron-poor boronic acids, we were surprised that the cross coupling of 4-fluorophenyl boronic acid and ammonium triflate **33** resulted in low yield when K₃PO₄ was used as base (entry 6). However, exceptional yield and chirality transfer were observed when CsF was employed (entry 7). Notably, the successful formation of ester **39** (entry 8), nitrile **40** (entry 9), and sulfone **41** (entry 10) particularly highlights the advantage of using a boronic acid as the coupling partner. For some of these base-sensitive functional groups, we replaced K₃PO₄ with CsF (entry 8). Finally, we were pleased to observe that the reaction of vinyl

Table 3. Formation of Enantioenriched Diarylethanes^a

entry	product	mol % Ni	ligand (mol %)	yield (%) ^b	prod ee (%) ^c
1		3	P(<i>o</i> -Tol) ₃ (7)	60	99
2 ^{d,e}		10	P(<i>o</i> -Tol) ₃ (22)	72	97
3		3	P(<i>o</i> -Tol) ₃ (7)	82	99
4		3	P(<i>o</i> -Tol) ₃ (7)	51	95
5		3	P(<i>o</i> -Tol) ₃ (7)	68	98
6		10	P(<i>o</i> -Tol) ₃ (22)	(15)	n.d. ^f
7 ^{d,g}		10	P(<i>o</i> -Tol) ₃ (22)	94	98
8 ^g		10	P(<i>o</i> -Tol) ₃ (22)	71	98
9 ^{d,h}		1	P(<i>o</i> -Tol) ₃ (3)	76	95
10		3	P(<i>o</i> -Tol) ₃ (7)	94	97
11 ^{d,i}		10	P(<i>o</i> -Tol) ₃ (22)	53	52
12 ^j		2	P(<i>o</i> -Tol) ₃ (5)	96	99
13 ^{d,i}		10	P(<i>o</i> -Tol) ₃ (22)	(91)	77
14 ^{d,e,k}		10	<i>t</i> -Bu-XantPhos (12)	46	98
15 ^e		10	<i>t</i> -Bu-XantPhos (12)	37	95
16 ^e		15	P(<i>o</i> -Tol) ₃ (32)	56	98
17 ^e		15	P(<i>o</i> -Tol) ₃ (32)	54	91

^aConditions: ammonium triflate (0.26 mmol, 1.0 equiv), boronic acid (1.2 equiv), Ni(cod)₂, P(*o*-Tol)₃, K₃PO₄ (1.3 equiv), dioxane (0.4 M), 70 °C, 6 h unless otherwise noted. Starting materials were ≥99% ee, unless otherwise noted. ^bAverage isolated yield of duplicate experiments (±5%). ^cYields in parentheses determined by ¹H NMR analysis using 1,3,5-trimethoxybenzene as internal standard. ^dAverage ee from duplicate experiments as determined by chiral HPLC analysis (±1%). ^eResult of a single experiment. ^fPerformed on 0.2 mmol scale. ^gn.d. = not determined. ^hCsF used instead of K₃PO₄. ⁱPerformed on 0.5 mmol scale. ^jPerformed on 0.1 mmol scale. ^kPerformed on 1.36 mmol scale. ^lStarting material was 98% ee.

boronic acids also proceeds in high yield and enantiospecificity (entry 12). Such enantioenriched allyl arenes are known

precursors to α -arylpropanoic acid analgesics, such as naproxen.¹⁹

With respect to the scope of the ammonium salt, electron-poor aryl groups are well tolerated (entries 14–17).²⁰ In every case, the benzylic ammonium triflates are available in greater than 98% ee, highlighting one of the advantages of using benzylic amine derivatives in these stereospecific cross couplings. However, with these substrates, higher catalyst loadings are required. In some cases, we also observed higher yields and greater chirality transfer when *t*-Bu-XantPhos was employed as ligand (entries 14 and 15). Although we do not yet fully understand the basis for the effect of ligand on the chirality transfer, the benefit of higher catalyst loadings for these substituted benzene derivatives suggests that a nickel-catalyzed epimerization pathway is not significant for these substrates.

We established the absolute configuration of diarylethane **35**, whose racemate displays anti-cancer activity,^{1d,e} by comparison of its optical rotation to the reported value.^{5b} The absolute configuration of (*R*)-**38** was determined by X-ray crystallography (Figure 1).^{11,21} Collectively, these assignments demonstrate that these reactions proceed with overall inversion of configuration.






Figure 1. Crystal structure of (*R*)-**38**. Molecular diagram of (*R*)-**38** with ellipsoids at 30% probability. Tertiary H-atom depicted with arbitrary radius. All other H-atoms and a second symmetry-unique compound molecule are omitted for clarity.

To begin to understand the mechanism of this transformation, we studied the stoichiometric reaction of ammonium triflate **47** with Ni(cod)₂ and PPh₂Cy. This reaction produced alkylnickel(II) triflate **48** in 51% isolated yield (Scheme 4). The structure of this complex was confirmed by X-ray crystallography, as well as ¹H, ¹³C, ¹⁹F, and ³¹P NMR analysis.^{11,21} It is consistent with oxidative addition of nickel into the benzylic C–N bond. The chelating carbonyl likely stabilizes this complex in the η^1 form. Upon treatment with *p*-tolylboronic acid, complex **48** is converted to diarylmethane **29** in 95% yield (¹H NMR, eq 1). Further, complex **48** is




catalytically competent; product **29** was formed in quantitative yield (¹H NMR) when **48** was used as catalyst (eq 2). These results suggest that complex **48** is a viable intermediate in the catalytic reaction. We thus propose that these reactions proceed

Scheme 4. Synthesis and Crystal Structure of Oxidative Addition Complex 48^a

^aMolecular diagram of 48 with ellipsoids at 30% probability. H-atoms omitted for clarity.

via oxidative addition of the electron-rich Ni(0) complex into the C–N bond to generate either an η^1 - or η^3 -bound benzylic nickel(II) intermediate.²² Transmetalation with the activated boronic acid and subsequent reductive elimination then delivers the diaryl product.


We have attempted to use a similar substrate to determine whether the oxidative addition of nickel into the C–N bond occurs with retention or inversion of configuration. However, the cross coupling of branched ammonium triflate 49 occurs with poor chirality transfer, delivering diarylethane in only 33% ee (eq 3). This result suggests that the carbonyl group



promotes an epimerization pathway. Unfortunately, attempts to isolate the oxidative addition adducts of substrates without such chelating groups have been unsuccessful to date. Nonetheless, retention of configuration during transmetalation and reductive elimination is well precedented for alkyl metal species.²³ Thus, we propose that the oxidative addition of the nickel catalyst into the benzylic C–N bond likely occurs via an S_N2 mechanism, resulting in inversion of configuration of the benzylic stereocenter (Scheme 5). This mechanistic proposal is consistent with the overall inversion of configuration observed in this cross coupling reaction.

In summary, we have developed mild conditions for a high-yielding cross coupling of benzyl ammonium triflates with aryl boronic acids to give diarylmethanes. This nickel-catalyzed reaction displays exceptional functional group tolerance and substrate scope in both the ammonium salt and boronic acid. Further, the nickel-catalyzed cross coupling of chiral ammonium salts was developed to yield diarylethanes in high enantiospecificities. To our knowledge, this is the first Suzuki

Scheme 5. Proposed Reaction Mechanism



coupling of a benzylic electrophile that provides highly enantioenriched products. Given the ready availability of highly enantioenriched chiral amines, this method offers a powerful approach to the synthesis of enantioenriched diarylethanes. Studies to determine further mechanistic details of this cross coupling as well as to expand the scope of this transformation are underway in our laboratories.

■ ASSOCIATED CONTENT

Supporting Information

Experimental procedures, characterization data and spectra of new compounds, and X-ray crystal structures of (*R*)-38 and 48. This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

mpwatson@udel.edu

Author Contributions

[†]P.M. and D.M.S.-M. contributed equally.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

Acknowledgement is made to NIH COBRE (P20 RR017716) and the Donors of the American Chemical Society Petroleum Research Fund for generous support of this research. We also thank AstraZeneca for donated boronic acids. NMR and other data were acquired at UD on instruments obtained with the assistance of NSF and NIH funding (NSF MIR 0421224, NSF CRIF MU CHE0840401, NIH P20 RR017716).

■ REFERENCES

- (a) Zhang, J.; Bellomo, A.; Creamer, A. D.; Dreher, S. D.; Walsh, P. J. *J. Am. Chem. Soc.* **2012**, *134*, 13765. (b) Cheltsov, A. V.; Aoyagi, M.; Aleshin, A.; Yu, E. C.-W.; Gilliland, T.; Zhai, D.; Bokov, A. A.; Reed, J. C.; Liddington, R. C.; Abagyan, R. *J. Med. Chem.* **2010**, *53*, 3899. (c) Huang, Z.; Ducharme, Y.; Macdonald, D.; Robichaud, A. *Curr. Opin. Chem. Biol.* **2001**, *5*, 432. (d) Messaoudi, S.; Hamze, A.; Provot, O.; Tréguier, B.; Rodrigo De Losada, J.; Bignon, J.; Liu, J.-M.; Wdziczak-Bakala, J.; Thoret, S.; Dubois, J.; Brion, J.-D.; Alami, M. *ChemMedChem* **2011**, *6*, 488. (e) Alami, M.; Messaoudi, S.; Hamze, A.; Provot, O.; Brion, J.-D.; Liu, J.-M.; Bignon, J.; Bakala, J. Dihydro-Iso-Ca-4 and Analogues: Potent Cytotoxics, Inhibitors of Tubulin Polymerization. WO2009/147217, 2009. (f) Moree, W. J.; Li, B.-F.; Jovic, F.; Coon, T.; Yu, J.; Gross, R. S.; Tucci, F.; Marinkovic, D.; Zamani-Kord, S.; Malany, S.; Bradbury, M. J.; Hernandez, L. M.; O'Brien, Z.; Wen, J.; Wang, H.; Hoare, S. R. J.; Petroski, R. E.; Saccaan, A.; Madan, A.; Crowe, P. D.; Beaton, G. *J. Med. Chem.* **2009**, *52*, 5307. (g) Beaton, G.; Moree, W. J.; Jovic, F.; Coon, T.; Yu, J. Sleep Inducing Compound and Methods Relating Thereto. US2006/14797, 2006.
- (2) For examples, see: (a) Molander, G. A.; Elia, M. D. *J. Org. Chem.* **2006**, *71*, 9198. (b) Arp, F. O.; Fu, G. C. *J. Am. Chem. Soc.* **2005**, *127*,

10482. (c) López-Pérez, A.; Adrio, J.; Carretero, J. C. *Org. Lett.* **2009**, *11*, 5514. (d) Binder, J. T.; Cordier, C. J.; Fu, G. C. *J. Am. Chem. Soc.* **2012**, *134*, 17003.

(3) (a) Kuwano, R. *J. Synth. Org. Chem., Jpn.* **2011**, *69*, 1263. (b) Kuwano, R.; Yokogi, M. *Chem. Commun.* **2005**, 5899. (c) Kuwano, R.; Yokogi, M. *Org. Lett.* **2005**, *7*, 945. (d) Kuwano, R.; Yu, J.-Y. *Heterocycles* **2007**, *74*, 233. (e) Ohsumi, M.; Kuwano, R. *Chem. Lett.* **2008**, *37*, 796. (f) Yu, J.-Y. J.; Kuwano, R. *Org. Lett.* **2008**, *10*, 973. (g) Kofink, C. C.; Knochel, P. *Org. Lett.* **2006**, *8*, 4121. (h) McLaughlin, M. *Org. Lett.* **2005**, *7*, 4875.

(4) (a) Guan, B.-T.; Xiang, S.-K.; Wang, B.-Q.; Sun, Z.-P.; Wang, Y.; Zhao, K.-Q.; Shi, Z.-J. *J. Am. Chem. Soc.* **2008**, *130*, 3268. (b) Yu, D.-G.; Wang, X.; Zhu, R.-Y.; Luo, S.; Zhang, X.-B.; Wang, B.-Q.; Wang, L.; Shi, Z.-J. *J. Am. Chem. Soc.* **2012**, *134*, 14639.

(5) (a) Taylor, B. L. H.; Harris, M. R.; Jarvo, E. R. *Angew. Chem., Int. Ed.* **2012**, *51*, 7790. (b) Taylor, B. L. H.; Swift, E. C.; Waetzig, J. D.; Jarvo, E. R. *J. Am. Chem. Soc.* **2011**, *133*, 389. (c) Greene, M. A. M.; Yonova, I. M. I.; Williams, F. J. F.; Jarvo, E. R. *Org. Lett.* **2012**, *14*, 4293.

(6) For other methods to form enantioenriched diarylethane derivatives, see: (a) Bolshan, Y.; Chen, C.-y.; Chilenski, J. R.; Gosselin, F.; Mathre, D. J.; O'shea, P. D.; Roy, A.; Tillyer, R. D. *Org. Lett.* **2004**, *6*, 111. (b) Fessard, T. C.; Andrews, S. P.; Motoyoshi, H.; Carreira, E. M. *Angew. Chem., Int. Ed.* **2007**, *46*, 9331. (c) Lewis, C. A.; Chiu, A.; Kubryk, M.; Balsells, J.; Pollard, D.; Esser, C. K.; Murry, J.; Reamer, R. A.; Hansen, K. B.; Miller, S. J. *J. Am. Chem. Soc.* **2006**, *128*, 16454. (d) Pathak, T. P.; Gligorich, K. M.; Welm, B. E.; Sigman, M. S. *J. Am. Chem. Soc.* **2010**, *132*, 7870. (e) Nave, S.; Sonawane, R. P.; Elford, T. G.; Aggarwal, V. K. *J. Am. Chem. Soc.* **2010**, *132*, 17096. (f) Roesner, S.; Casatejada, J. M.; Elford, T. G.; Sonawane, R. P.; Aggarwal, V. K. *Org. Lett.* **2011**, *13*, 5740.

(7) Li, M.-B.; Tang, X.-L.; Tian, S.-K. *Adv. Synth. Catal.* **2011**, *353*, 1980.

(8) He, A.; Falck, J. R. *J. Am. Chem. Soc.* **2010**, *132*, 2524.

(9) (a) Imao, D.; Glasspoole, B. W.; Laberge, V. S.; Crudden, C. M. *J. Am. Chem. Soc.* **2009**, *131*, 5024. (b) Dreher, S. D.; Dormer, P. G.; Sandrock, D. L.; Molander, G. A. *J. Am. Chem. Soc.* **2008**, *130*, 9257. (c) Sandrock, D. L.; Jean-Gérard, L.; Chen, C.-y.; Dreher, S. D.; Molander, G. A. *J. Am. Chem. Soc.* **2010**, *132*, 17108.

(10) For examples of allylic and benzylic ammonium salts in other reactions, see: (a) Langlois, Y.; Van Bac, N.; Fall, Y. *Tetrahedron Lett.* **1985**, *26*, 1009. (b) Dressaire, G.; Langlois, Y. *Tetrahedron Lett.* **1980**, *21*, 67. (c) Gupton, J. T.; Layman, W. J. *J. Org. Chem.* **1987**, *52*, 3683. (d) de la Herrán, G.; Segura, A.; Csáký, A. G. *Org. Lett.* **2007**, *9*, 961. (e) Hosomi, A.; Hoashi, K.; Tominaga, Y.; Otaka, K.; Sakurai, H. *J. Org. Chem.* **1987**, *52*, 2947.

(11) See Supporting Information for details.

(12) (a) Nugent, T. C. *Chiral Amine Synthesis*; Wiley-VCH Verlag GmbH & Co. KGaA: Weinheim, 2010. (b) Mereyala, H. B.; Pola, P. *Tetrahedron: Asymmetry* **2003**, *14*, 2683. (c) Savile, C. K.; Janey, J. M.; Mundorff, E. C.; Moore, J. C.; Tam, S.; Jarvis, W. R.; Colbeck, J. C.; Krebber, A.; Fleitz, F. J.; Brands, J.; Devine, P. N.; Huisman, G. W.; Hughes, G. J. *Science* **2010**, *329*, 305. (d) Vries, T.; Wynberg, H.; van Echten, E.; Koek, J.; ten Hoeve, W.; Kellogg, R. M.; Broxterman, Q. B.; Minnaard, A.; Kaptein, B.; van der Sluis, S.; Hulshof, L.; Kooistra, J. *Angew. Chem., Int. Ed.* **1998**, *37*, 2349.

(13) (a) Blakey, S.; MacMillan, D. *J. Am. Chem. Soc.* **2003**, *125*, 6046. (b) Wenkert, E.; Han, A.-L.; Jenny, C.-J. *Chem. Commun.* **1988**, 975. (c) Reeves, J. T.; Fandrick, D. R.; Tan, Z.; Song, J. J.; Lee, H.; Yee, N. K.; Senanayake, C. H. *Org. Lett.* **2010**, *12*, 4388. (d) Buszek, K. R.; Brown, N. *Org. Lett.* **2007**, *9*, 707. (e) Xie, L.-G.; Wang, Z.-X. *Angew. Chem., Int. Ed.* **2011**, *50*, 4901. (f) Zhang, X.-Q.; Wang, Z.-X. *J. Org. Chem.* **2012**, *77*, 3658. (g) Blessley, G.; Holden, P.; Walker, M.; Brown, J. M.; Gouverneur, V. *Org. Lett.* **2012**, *14*, 2754.

(14) For an example of an enantioselective reaction of a benzylic amine derivative, see: Weng, Z.-T.; Li, Y.; Tian, S.-K. *J. Org. Chem.* **2011**, *76*, 8095.

(15) High stereospecificity has been observed in Negishi cross couplings of indene-derived aziridines. See: Huang, C.-Y.; Doyle, A. G. *J. Am. Chem. Soc.* **2012**, *134*, 9541.

(16) These control reactions were performed with new stir bars.

(17) Lower yields were observed for pyridine boronic acids without substituents ortho to the nitrogen, such as 3-pyridinylboronic acid. Ortho substituents likely hinder pyridine from binding the Ni catalyst.

(18) For an example, see the preparation of 1-(biphenyl-4-yl)-*N,N*-dimethylethanamine in the Supporting Information.

(19) Li, C. C.; Xing, J. J.; Zhao, J. J.; Huynh, P. P.; Zhang, W. W.; Jiang, P. P.; Zhang, Y. J. *Org. Lett.* **2012**, *14*, 390.

(20) The cross coupling of the 1-naphthyl-substituted ammonium salt proceeded in only moderate enantiospecificity under the optimized reaction conditions (67% ee). See Supporting Information for more detail.

(21) CCDC-899360 ((*R*)-**38**) and CCDC-899359 (**48**) contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via http://www.ccdc.cam.ac.uk/data_request/cif.

(22) Both the η^1 and η^3 forms of benzyl nickel(II) intermediates have been observed. See: Matsubara, R.; Gutierrez, A. C.; Jamison, T. F. *J. Am. Chem. Soc.* **2011**, *133*, 19020.

(23) (a) Stille, J. K. *Oxidative Addition and Reductive Elimination. In The Chemistry of the Metal–Carbon Bond*, Vol. 2; Hartley, F. R., Patai, S., Eds.; John Wiley & Sons, Ltd.: New York, 1985; p 625. (b) Netherton, M. R.; Fu, G. C. *Angew. Chem., Int. Ed.* **2002**, *41*, 3910.