


Substrate-Triggered Addition of Dioxygen to the Diferrous Cofactor of Aldehyde-Deformylating Oxygenase to Form a Diferric-Peroxide Intermediate

Maria E. Pandelia,^{†,⊥} Ning Li,^{‡,§,⊥} Hanne Nørgaard,^{†,||} Douglas M. Warui,[‡] Lauren J. Rajakovich,[‡] Wei-chen Chang,[†] Squire J. Booker,^{*,†,‡} Carsten Krebs,^{*,†,‡} and J. Martin Bollinger, Jr.^{*,†,‡}

[†]Departments of Chemistry and [‡]Biochemistry and Molecular Biology, The Pennsylvania State University, University Park, Pennsylvania 16802, United States

S Supporting Information

ABSTRACT: Cyanobacterial aldehyde-deformylating oxygenases (ADOs) belong to the ferritin-like diiron-carboxylate superfamily of dioxygen-activating proteins. They catalyze conversion of saturated or monounsaturated C_n fatty aldehydes to formate and the corresponding C_{n-1} alkanes or alkenes, respectively. This unusual, apparently redox-neutral transformation actually requires four electrons per turnover to reduce the O_2 cosubstrate to the oxidation state of water and incorporates one O-atom from O_2 into the formate coproduct. We show here that the complex of the diiron(II/II) form of ADO from *Nostoc punctiforme* (*Np*) with an aldehyde substrate reacts with O_2 to form a colored intermediate with spectroscopic properties suggestive of a $Fe_2^{III/III}$ complex with a bound peroxide. Its Mössbauer spectra reveal that the intermediate possesses an antiferromagnetically (AF) coupled $Fe_2^{III/III}$ center with resolved subsites. The intermediate is long-lived in the absence of a reducing system, decaying slowly ($t_{1/2} \sim 400$ s at 5 °C) to produce a very modest yield of formate (<0.15 enzyme equivalents), but reacts rapidly with the fully reduced form of 1-methoxy-5-methylphenazinium methylsulfate (^{MeO}PMS) to yield product, albeit at only ~50% of the maximum theoretical yield (owing to competition from one or more unproductive pathway). The results represent the most definitive evidence to date that ADO can use a *diiron* cofactor (rather than a homo- or heterodinuclear cluster involving another transition metal) and provide support for a mechanism involving attack on the carbonyl of the bound substrate by the reduced O_2 moiety to form a $Fe_2^{III/III}$ -peroxyhemiacetal complex, which undergoes reductive O–O-bond cleavage, leading to C1–C2 radical fragmentation and formation of the alk(a/e)ne and formate products.



INTRODUCTION

Cyanobacterial aldehyde-deformylating oxygenases (ADOs), originally designated by Schirmer et al. as “aldehyde decarboxylases,”¹ catalyze conversion of C_n fatty aldehydes to the corresponding C_{n-1} alk(a/e)nes and formate in a reaction requiring one molecule of O_2 and four electrons per turnover.^{2–4} The ADO substrate is provided in vivo by the preceding enzyme in the pathway, acyl–acyl carrier protein (ACP) reductase, which reductively cleaves its fatty acid thioester substrate using a reduced nicotinamide.¹ Because ADOs catalyze the final step in a pathway that converts physiologically abundant fatty acids into combustible hydrocarbons, they have received much recent attention for their potential biotechnological applications.^{1,5–10} Insight into their structure and mechanism might facilitate their deployment in, for example, bioprocesses to produce renewable hydrocarbon fuels in cyanobacteria or other heterologous bacterial hosts.^{5,11,12} Particularly for the latter approach, an understanding of the details of the reaction, including the identities and modes of participation of accessory factors, might be

necessary to make the bioprocess efficient enough to be economically viable.

ADOs belong to the ferritin-like protein structural superfamily (see PDB accessions 2OC5 and 4KVQ for structural details), which comprises a number of well-characterized oxidases and oxygenases known to employ their carboxylate-bridged nonheme-diiron cofactors to activate O_2 for oxidation (in many cases, oxygenation)^{13–19} of their substrates. The ADO reaction is, however, unusual for a member of this enzyme superfamily, in that it results in no net oxidation of its primary substrate. Indeed, its similarity to the reactions of the other members is revealed only by the obligatory requirement for a reducing system^{1–4} and incorporation of a single atom of oxygen from O_2 into the formate coproduct.^{3,4} Despite these idiosyncrasies, it has been presumed that ADO also employs a *diiron* cofactor in catalysis. The fact that no ADO has, to our knowledge, been isolated from its native source, and the

Received: May 20, 2013

Published: August 29, 2013

growing recognition that members of the structural superfamily can employ metal ions other than iron,^{20–23} have left the nature of the cofactor still somewhat in doubt. Obviously, knowledge of the identity of the cofactor could impact efforts to make the enzyme function at high efficiency in an engineered bioprocess.

In this study, the *Nostoc punctiforme* (*Np*) ADO has been overexpressed in *Escherichia coli* (*Ec*) under culture conditions designed to favor incorporation of exclusively iron (minimal medium supplemented with Fe). A combination of metal analyses and Mössbauer spectroscopy has been used to show that these preparations do indeed harbor predominately (>95%) diiron clusters. Combined with the demonstration that the ADO prepared in this manner reacts, under the proper conditions, almost homogeneously with O₂ to form an intermediate that is capable of yielding products, these results provide the best evidence to date for the previously presumed catalytic competence of the diiron form of ADO.


Iron-dependent oxidases and oxygenases often exhibit the apparently adaptive characteristic of becoming activated toward reaction with O₂ upon binding of either the substrate [as in the Fe^{II}/ α -ketoglutarate-dependent hydroxylases and halogenases^{24–26} and stearyl-ACP Δ^9 desaturase (Δ^9 D)²⁷] or an effector protein [as in soluble methane monooxygenase (sMMO) and toluene/*o*-xylene monooxygenase (ToMOH)^{28–32}]. We have adopted the term “substrate triggering” to refer to the cases in which the substrate acts as the relevant activator.²⁶ This effect may have evolved to limit self-inflicted damage to the enzyme resulting from formation of potentially oxidizing intermediates in the absence of the oxidation target. The activating binding event is typically associated with a change in cofactor coordination. In many cases, dissociation of a water ligand creates an open coordination site to which O₂ can add to initiate the reaction.^{33–35}

In this study, we show that the *Np* ADO is triggered by short-chain (C₆–C₁₀) *n*-aldehyde substrates⁴ for addition of O₂ to its Fe₂^{II/III} cofactor. The reaction yields an intermediate with an absorption feature at ~450 nm, a diamagnetic ground state (*S* = 0), and Mössbauer parameters indicative of two distinct, AF coupled, high-spin Fe^{III} subsites. Its spectroscopic properties and reactivity suggest that the intermediate is a Fe₂^{III/III} complex with a bound peroxide. This intermediate is long-lived (*t*_{1/2} ~ 400 s at 5 °C) in the absence of a reductant, but effects rapid (*t*_{1/2} ~ 0.14 s) oxidation of the reduced form of 1-methoxy-5-methylphenazinium methylsulfate (MeO⁺PMS). Its decay in the absence of reductant produces little formate (<0.15 equiv), but its reductant-promoted decay generates formate with a stoichiometry of ~0.5^{MeO}PMS (approximately half the theoretical maximum yield). These results provide strong support for two key features of our previously proposed mechanism for ADO (Scheme 1):^{3,4} (1) formation of a Fe₂^{III/III} intermediate with a bound peroxide moiety by addition of O₂ to the reduced Fe₂^{II/II} cofactor of the ADO•aldehyde complex; and (2) reductive cleavage of the peroxide to initiate a free-radical C1–C2 β -scission reaction. The results emphasize the importance of controlled delivery of electrons to the ADO cofactor during turnover and thereby identify another aspect of its function that might be tunable for increased efficiency in associated bioprocesses.

EXPERIMENTAL SECTION

Materials. Tris-(2-carboxyethyl)phosphine (TCEP), technical grade (>85% purity) sodium hydrosulfite (dithionite), reduced β -nicotinamide adenine nucleotide ($\geq 97\%$) (NADH), 2-nitrophenylhy-

Scheme 1. Proposed Catalytic Mechanism for the Conversion of Aldehyde to Alk(a/e)ne and Formate by ADO^{3,4a}



^aThe new hydrogen atom incorporated into the alk(a/e)ne product ultimately comes from solvent, but the proximal donor could be either a solvent ligand (green arrows) or an amino acid residue (depicted as H–X; purple arrows).

drazine (97%), and short-chain aldehyde substrates (*n*-hexanal, *n*-octanal, and *n*-decanal) were purchased from Sigma-Aldrich (St. Louis, MO). 1-(3-Dimethylaminopropyl)-3-ethylcarbodiimide (98%) was purchased from Alfa Aesar (Ward Hill, MA). 1-Methoxy-5-methylphenazinium methylsulfate (MeO⁺PMS) was obtained from Acros Organics (Fair Lawn, NJ). 1-[¹³C]-Octanal was synthesized according to a previously described procedure.³ All other chemicals used for cell growth and protein purification were purchased from Sigma-Aldrich (St. Louis, MO), unless stated otherwise.

Preparation of *Np* ADO. The plasmid used to direct overexpression of the ADO protein from *Nostoc punctiforme* PCC 73102 (*Npun*_R1711; accession number YP_001865325) has previously been described.³ The protein was produced in *Escherichia coli* BL21(DE3) (Invitrogen; Carlsbad, CA) by a procedure modified from the original, published procedure, as noted below. The cells were grown at 37 °C in M9 minimal medium containing 50 mg/L kanamycin, 2 mM MgSO₄, 0.2% (v/v) glucose, 100 μ M CaCl₂, and 250 μ M (NH₄)₂Fe(SO₄)₂·6H₂O until they reached an OD₆₀₀ of 0.6–0.8. Expression of the ADO was induced by addition of 250 μ M isopropyl- β -D-1-thiogalactopyranoside (IPTG), and the cultures were incubated at 18 °C for 20–24 h. All buffers used in the anaerobic purification of *Np* ADO contained 1 mM tris-(2-carboxyethyl)-phosphine (TCEP). The anaerobically isolated ADO (shown below to contain primarily the Fe₂^{II/II} form of the cofactor) was obtained by carrying out cell lysis and immobilized metal-ion affinity chromatography in a Coy anaerobic chamber (Grass Lake, MI). Cell lysis by sonication of ~40 g of cells suspended in ~200 mL of lysis buffer (50 mM sodium phosphate buffer pH 8.0, 10 mM imidazole, 300 mM NaCl) was followed by ultracentrifugation in sealed, gastight tubes at 50,000 \times g for 40 min at 4 °C. The supernatant was applied to a Ni^{II}-nitrilo-tris-acetate (NTA) column (30 mL resin per 200–220 mL of crude lysate). The application was followed by washing with 5 column volumes of 50 mM sodium phosphate buffer (pH 8.0) containing 40 mM imidazole and 300 mM NaCl. The protein was eluted with 50 mM sodium phosphate buffer (pH 8.0) containing 250 mM imidazole and 100 mM NaCl. The protein was concentrated, buffer-exchanged by using a prepacked PD-10 desalting column (G-25 Sephadex medium, GE Healthcare) in 50 mM sodium 2-[4-(2-hydroxyethyl)-piperazin-1-yl]ethanesulfonate (HEPES) buffer (pH 7.5, 10% glycerol), transferred into gastight Eppendorf tubes, and flash frozen in liquid N₂, where it was stored until it was used in experiments. Aerobically isolated enzyme (shown below to contain primarily the Fe₂^{III/III} form of the cofactor) was obtained by carrying out the same

procedure in air. Iron and other trace metals in the samples were quantified by inductively coupled plasma atomic emission spectroscopy (ICP-AES) by Mr. Henry Gong at the Penn State Materials Research Institute. Purity of the ADO was estimated on the basis of SDS-PAGE and Coomassie staining to be >90%, and the concentration was calculated by measuring the absorbance at 280 nm and assuming a molar absorption coefficient of 22,920 mM⁻¹ cm⁻¹, which was calculated from the amino acid sequence using the ProtParam tool (<http://web.expasy.org/protparam/>).

Stopped-Flow Absorption (SF-Abs) Experiments and Data Analysis. SF-Abs measurements were carried out with a SX20 stopped-flow spectrophotometer from Applied Photophysics Ltd. (Leatherhead, UK) that was housed in an anoxic chamber (Labmaster, MBraun, Stratham, NH). All experiments were carried out at 5 °C. In single-mix (two-syringe) experiments, the ADO reactant solution was diluted by 2-fold, and in the sequential-mix (four-syringe) experiments, the protein was diluted by 4-fold. In the single-mix experiments, an O₂-free solution of Fe₂^{II/III}-ADO (0.20–2.5 mM), with or without aldehyde substrate, was mixed with O₂-saturated (on ice) 50 mM sodium HEPES buffer (pH 7.5) containing 10% glycerol, giving an estimated O₂ concentration after mixing of 0.9 mM.³⁶ The optical path length used was 10 mm or 2 mm, depending on the protein concentration. For initial characterization of the reaction, the photodiode array (PDA) detector was used to acquire time-resolved absorption spectra. However, the Fe₂^{III/III}-peroxide intermediate was found to be photolytically labile under the intense white light required for use of the multiwavelength detector. Therefore, the true kinetics of the intermediate were determined by using monochromatic 450 nm light and the photomultiplier tube (PMT) detector.

For a system of two consecutive, irreversible reactions converting reactant, R, to product, P, via intermediate, I, Beer's law gives the absorbance of the solution at 450 nm as a function of time as

$$A_{450}(t) = \epsilon_R[R]_t + \epsilon_I[I]_t + \epsilon_P[P]_t \quad (1)$$

The time dependencies of the reaction components obtained by integrating the relevant differential rate equations are obtained as^{37,38}

$$[R]_t = [R]_0 e^{-k_1 t} \quad (2)$$

$$[I]_t = [R]_0 \frac{k_1}{k_2 - k_1} (e^{-k_1 t} - e^{-k_2 t}) \quad (3)$$

$$[P]_t = [R]_0 + \frac{[R]_0}{k_2 - k_1} (k_1 e^{-k_2 t} - k_2 e^{-k_1 t}) \quad (4)$$

from which the expression for $\Delta A_{450 \text{ nm}}$ as a function of time (eq 5) can be obtained as

$$\Delta A_{450}(t) = [R]_0 \left(\frac{(\epsilon_R - \epsilon_P)k_2 + (\epsilon_I - \epsilon_R)k_1}{k_2 - k_1} \right) e^{-k_1 t} + [R]_0 \frac{k_1}{k_2 - k_1} (\epsilon_P - \epsilon_I) e^{-k_2 t} \quad (5)$$

Because the kinetic traces suggest that the reactant exists in two forms, R and R', which can both convert to intermediate I, eq 5 can be expanded to include an additional term for the step R' → I with its unique rate constant, k'. Alternatively, neglecting the minor absorption at 450 nm from the product complex ($\epsilon_P \sim 0$) yields the simpler expression:

$$\Delta A_{450}(t) = \frac{[R]_0 k_1}{k_2 - k_1} (e^{-k_1 t} - e^{-k_2 t}) + \frac{[R']_0 k_1'}{k_2 - k_1'} (e^{-k_1' t} - e^{-k_2' t}) \quad (6)$$

which, when used in the regression analyses, was found to return essentially the same values of the rates constants as the more rigorous, complex expression that takes into account the minor absorbance of the product complex.

ΔA_{388} -versus-time traces reporting oxidation of the Me^oPMS reductant in sequential-mixing SF-Abs experiments were analyzed

according to eq 7, which corresponds to two parallel first-order reactions with different amplitudes (ΔA_1 and ΔA_2) and rate constants (k_1 and k_2).

$$\Delta A_{388}(t) = \Delta A_1(1 - e^{-k_1 t}) + \Delta A_2(1 - e^{-k_2 t}) \quad (7)$$

Preparation of Freeze-Quench (FQ) Samples. Freeze-quench experiments were carried out at 5 °C as previously described.³⁹

Assays for Formate Production. Formate produced in the ADO reaction was quantified as its 2-nitrophenylhydrazide derivative by liquid chromatography and mass spectrometry (LC-MS), as previously described.^{3,40} The reactions were carried out in the anoxic chamber by a procedure designed to mimic the sequential-mixing SF-Abs experiments. A 50 μ L aliquot containing 1.49 mM anaerobically isolated Fe₂^{II/II}-ADO and 16.6 mM 1-[¹³C]-octanal was mixed in a sealed Eppendorf tube via a gastight syringe with an equal volume of O₂-saturated buffer. After a 32 s delay for accumulation of the intermediate, a 100 μ L aliquot of Me^oPMS that had previously been reduced by equimolar sodium dithionite was rapidly injected into this solution. Immediately (1–2 s later), 10 μ L of this solution was quenched into 90 μ L of a solution containing 0.032 mM 1-ethyl-3-[3-dimethylaminopropyl]carbodiimide, 0.013 mM 2-nitrophenylhydrazine (2NPH), and 28 mM HCl. A 10 μ L aliquot of 1 mM propionic acid was then added as an internal standard, giving a final concentration of 0.1 mM. The reaction was incubated for 15 min at 70 °C. The precipitated protein was pelleted by centrifugation at 17,000× g, and the supernatant was then filtered through a 0.2 μ M PVDF filter (National Scientific, Rockwood, TN) by centrifugation. The solution was analyzed on a 6410 LC/MS Agilent QQQ spectrometer by injecting 2 μ L onto a Hamilton PRP-1 analytical column. An isocratic mobile phase (2:1 methanol/H₂O containing 0.05% acetic acid) was used. Single-ion monitoring (SIM) was carried out at *m/z* values of 180 (¹²CHO-2NPH), 181 (¹³CHO-2NPH), and 208 (propionyl-2NPH).³

Mössbauer Spectroscopy. Mössbauer spectra were recorded on spectrometers from WEB Research (Edina, MN) that have been described previously.⁴¹ The spectrometer used to acquire the weak-field spectra is equipped with a Janis SVT-400 variable-temperature cryostat, whereas the spectrometer used to acquire the strong-field spectra is equipped with a Janis 8TMOSS-OM-12SVT variable-temperature cryostat. The external magnetic field was applied parallel to the γ -beam. All isomer shifts quoted are relative to the centroid of the spectrum of α -iron metal at room temperature. Simulation of the Mössbauer spectra was carried out by using the WMOSS spectral analysis software (www.wmoss.org, WEB Research, Edina, MN).

RESULTS

Expression and Purification of *Np* ADO to Favor Iron Incorporation. Since the initial report on the cyanobacterial alk(a/e)ne-production pathway ending with the ADO (formerly aldehyde decarbonylase) reaction,¹ the identities of the functional metal ions in the enzyme's dinuclear cofactor have remained somewhat in doubt. Initially, a *diiron* cofactor was presumed, because it is the most common among the ferritin-like proteins, but the growing recognition that *manganese* is functional in some members of the structural superfamily^{20–23} left the question open. However, our recent characterization of the reaction as a novel oxygenation process made a *diiron* cofactor the most likely possibility^{3,4} and led us to express and purify the enzyme under conditions designed both to favor incorporation of only iron and to yield a maximal fraction of the O₂-reactive state.

Following its overexpression in minimal medium with ⁵⁷Fe supplementation, the as-isolated *Np* ADO (with *N*-terminal His₆ affinity tag comprising 20 additional amino acids) contains 1.15 ± 0.05 Fe ions per monomer, as determined both by inductively coupled plasma atomic emission spectroscopy (ICP-AES) and a spectrophotometric assay based on the Fe^{II}

chelator, ferrozine.⁴² Incorporation of the other biologically common transition metals was minimal (0.012 equiv Mn, 0.030 equiv Ni, and 0.032 equiv Zn).

Characterization of (An)aerobically Purified *Np* ADO by Mössbauer Spectroscopy. Confirmation of the dinuclear nature of the iron cofactor was sought by Mössbauer spectroscopy on the protein purified in the presence and absence of O₂. The 4.2-K/53-mT spectrum of the aerobically purified *Np* ADO (Figure S1A in the Supporting Information) can be adequately analyzed as a single quadrupole doublet with an isomer shift (δ) of 0.52 mm/s and quadrupole splitting (ΔE_Q) of 1.25 mm/s. These parameters are characteristic of high-spin ($S = 5/2$) Fe^{III} ions.⁴³ The spectrum of the aerobically isolated ADO containing 10 mM *n*-decanal (Figure 1A) is essentially identical. This observation implies either that




Figure 1. Mössbauer spectra of *Np* ADO isolated from *Ec* grown in minimal medium with ⁵⁷Fe supplementation. (A) Spectrum of the aerobically isolated ADO in the presence of *n*-decanal recorded at 4.2 K and a magnetic field of 53 mT (vertical bars) and simulation of the experimental spectrum with parameters provided in the text (purple line). (B) Spectrum of the sample in (A) recorded with a magnetic field of 6 T (vertical bars) applied parallel to the γ -beam and simulation thereof with parameters described in the text (purple line). (C) 4.2-K/53-mT spectrum of anaerobically isolated ADO in the presence of *n*-decanal (vertical bars); the 6% contribution from the Fe₂^{III/III} complex in this sample is shown in purple.

the substrate does not bind tightly to the Fe₂^{III/III} form of the enzyme or that the aldehyde binds in a manner that does not detectably alter its Mössbauer properties. The fact that the spectrum is a quadrupole doublet is consistent with the presence of *two* AF coupled high-spin Fe^{III} ions in a dinuclear cluster. Indeed, the spectrum collected with a strong (6 T) applied magnetic field (Figure 1B, vertical bars) can be simulated with $\delta = 0.52$ mm/s, $\Delta E_Q = -1.25$ mm/s, asymmetry parameter $\eta = 0.4$, and a diamagnetic ($S_{\text{total}} = 0$) ground state (Figure 1B, purple line).

The 4.2-K/53-mT Mössbauer spectrum of the anaerobically isolated *Np* ADO is dominated by a broad quadrupole doublet with (average) isomer shift ($\delta \sim 1.3$ mm/s) and quadrupole splitting parameter ($\Delta E_Q \sim 3.0$ mm/s) characteristic of high-

spin Fe^{II} ions in an octahedral O/N coordination environment (Figure S1B).⁴³ In addition, there is a minor (6%) contribution from the oxidized Fe₂^{III/III} cluster (purple subspectrum). Treatment of the aerobically isolated Fe₂^{III/III}-ADO with dithionite affords a reduced state with a spectrum that is very similar to that of the anaerobically purified protein. Addition of 16.6 mM 1-[¹³C]-octanal to the anaerobically isolated enzyme, yielding the Fe₂^{II/II}-ADO•1-[¹³C]-octanal complex that is shown below to react rapidly with O₂ to accumulate an intermediate, does not detectably alter its Mössbauer spectrum (Figure 1C).

SF-Abs Evidence for “Substrate Triggering” of O₂ Addition and Accumulation of an Intermediate. Upon mixing of the Fe₂^{II/II} form of *Np* ADO (prepared either by anaerobic purification or by dithionite reduction of the aerobically purified enzyme) with oxygen-saturated buffer at 5 °C, a broad absorption band centered at ~ 350 nm develops slowly (Figure 2A), signifying oxidation of the cofactor. The




Figure 2. SF-Abs experiment monitoring the reaction at 5 °C of Fe₂^{II/II}-ADO (0.35 mM) with O₂ (0.9 mM) (A) in the absence and (B) in the presence of 10 mM *n*-decanal. The arrows in (B) indicate the formation/decay of an intermediate absorbing at ~ 450 nm only in the presence of the aldehyde substrate.

absorption feature is similar to those associated with μ -oxo-Fe₂^{III/III} clusters in other ferritin-like diiron carboxylate proteins, in which it has been assigned to oxo-to-Fe^{III} charge-transfer (CT) transitions.^{44–46} By contrast to the slow, monotonic development of the ~ 350 nm band in the reaction of substrate-free Fe₂^{II/II}-ADO with excess O₂ (Figure S2), inclusion of an aldehyde substrate (e.g., *n*-decanal) with the Fe₂^{II/II}-ADO in the mix with O₂ results in a much faster reaction, and a new transient absorption feature at ~ 450 nm is observed (Figure 2B). The spectrum after completion has the same ~ 350 nm absorption band seen in the reaction in the absence of substrate, suggesting formation of a μ -oxo-Fe₂^{III/III} product cluster upon decay of the transient species. The short-chain C_{6–10} *n*-aldehydes, which were previously shown to be ADO substrates,^{4,47} all “trigger” reaction of the Fe₂^{II/II} cofactor with O₂, leading to development of the transient absorption feature (Figure S3). Corresponding *n*-alkyl compounds lacking a functional group or having acid or alcohol functional groups do not support accumulation of the absorbing species (Figure 3A, red, blue and purple traces). Thus, the aldehyde functional group is essential for “substrate triggering.” At a concentration of 4 mM, the shorter aldehydes (e.g., *n*-hexanal) are less effective at substrate triggering (Figure S3), as expected from the fact that the natural substrates are C₁₄, C₁₆, or C₁₈ saturated or monounsaturated fatty aldehydes provided by the acyl-ACP reductase.¹

Photolytic Lability of the Intermediate. In the SF-Abs experiments of Figure 2, multiwavelength detection was achieved by use of a photodiode array, which requires intense,




Figure 3. Characteristics of the formation and decay of the 450-nm-absorbing complex. (A) Requirement for the aldehyde functional group for substrate triggering of intermediate formation in reaction of $\text{Fe}_2^{\text{II/II}}\text{-ADO}$ with O_2 . A solution of 200 μM $\text{Fe}_2^{\text{II/II}}\text{-ADO}$, 2 mM substrate or analog [*n*-nonanal (green), *n*-octanoic acid (red), *n*-nonan-1-ol (blue), *n*-nonane (purple), or no substrate (black)] was mixed in the SF-Abs instrument at 5 °C with O_2 -saturated (1.8 mM) buffer (50 mM sodium HEPES, 10% glycerol, pH 7.5). The reaction was monitored at 450 nm with PDA detection. (B) Assessment of the intermediate's photolytic lability by SF-Abs spectroscopy monitored at 450 nm. A solution of 700 μM $\text{Fe}_2^{\text{II/II}}\text{-ADO}$ and 10 mM *n*-decanal was mixed with an equal volume of an O_2 -saturated buffer solution at 5 °C. $\Delta A_{450 \text{ nm}}$ -vs-time traces were recorded with PDA (green trace) or PMT (blue trace) detection. Regression analysis of the latter trace was carried out according to eq 6, giving rate constants mentioned in the text. Control experiments carried out in the absence of *n*-decanal with PDA and PMT detection are shown as black and gray traces, respectively.

polychromatic incident light. Under these conditions, the intermediate was observed to have a half-life ($t_{1/2}$) of ~ 0.7 min at 5 °C (Figure 3, green traces). However, upon use of monochromatic incident light and the photomultiplier detector (PMT), the intermediate species was found to decay ~ 10 -fold less rapidly ($t_{1/2} \sim 7$ min, blue trace with superimposed red fit, Figure 3B). This observation indicates that the intermediate is photolytically labile, as has been found for O_2 -derived complexes in other nonheme-diiron oxygenases,^{21,48,49} including the peroxo- $\text{Fe}_2^{\text{III/III}}$ complex in $\Delta^9\text{D}$,²⁷ and related model compounds.⁵⁰ Therefore, accurate kinetics for the intermediate were obtained by use of monochromatic incident light and photomultiplier detector.

Complex Kinetics of Formation of the Intermediate and Their Origin. Attempts to analyze the $\Delta A_{450 \text{ nm}}$ -versus-time traces from the reaction according to a sequence of two irreversible steps gave poor agreement between the fit and experimental data, suggesting the presence of two distinct formation phases with different rate constants, each associated with a decay phase of the same rate constant. The traces were thus fit by eq 6, which gives change in absorbance ($\Delta A_{450 \text{ nm}}$) as a function of time (t) from two converging reaction channels that form a common intermediate (I) and product (P) from two different forms of the reactant (R and R'). I forms from R and R' with apparent first-order rate constants k_1 and k_1' , respectively, and decays to P with rate constant k_2 . R (R'), I, and P have molar absorption coefficients at 450 nm of ϵ_R (assumed to be equal for R and R' because the $\text{Fe}_2^{\text{II/II}}$ complexes should be essentially transparent at this wavelength), ϵ_I , and ϵ_P , respectively. The product complex was also assumed to have practically zero absorption ($\epsilon_P \sim 0$) at 450 nm, which is justified by its spectrum (Figure 2B). This regression analysis applied to the trace in Figure 3B gave apparent first-order rate constants for formation of $k_1 = 0.75 \text{ s}^{-1}$ and $k_1' = 0.20 \text{ s}^{-1}$ and a rate constant for decay of $k_2 = 0.0017 \text{ s}^{-1}$. According to this

analysis, reactant form R predominates (64%) over R' (36%). Fitting the data so as to account for the nonzero absorption of P at 450 nm gave essentially the same results. The origin of the two formation phases and the distinction between the two reactant states (R and R') giving rise to them were probed in two additional SF-Abs experiments. In the first, variation of the concentration of the aldehyde substrate (decanal) revealed a titration behavior, in which the relative contribution of the slow formation phase to the total amplitude of the transient is diminished with increasing substrate (Figure S4). This result suggests that incomplete formation of the $\text{Fe}_2^{\text{II/II}}\text{-ADO}$ •substrate complex in the reactant solution could be one source of the kinetic complexity. To confirm this deduction, a second experiment, in which substrate-free $\text{Fe}_2^{\text{II/II}}\text{-ADO}$ was mixed with an O_2 -saturated solution containing the substrate, was carried out (Figure S5). The markedly slower formation of the intermediate in the reaction initiated in this manner proves that substrate acquisition by $\text{Fe}_2^{\text{II/II}}\text{-ADO}$ is slower than O_2 addition to the $\text{Fe}_2^{\text{II/II}}\text{-ADO}$ •decanal complex and provides support for the notion that subsaturation by the substrate could be one source of the kinetic complexity in intermediate formation in the experiment of Figure 3B. Interestingly, the reaction initiated by mixing with decanal and O_2 also exhibits kinetic complexity. This observation suggests that, in addition to the heterogeneity related to the presence of both substrate-free and substrate-bound forms, the free enzyme is itself heterogeneous, having two or more forms differing in their kinetics of substrate binding.

Characterization of the Intermediate by Mössbauer Spectroscopy. To obtain insight into the electronic structure of the intermediate, freeze-quench (FQ) samples were prepared for analysis by Mössbauer spectroscopy. The $\text{Fe}_2^{\text{II/II}}\text{-ADO}$ •octanal complex was mixed with a solution of O_2 -saturated buffer, and the reaction was quenched after 32 s, the time of maximum ΔA_{450} in a SF-Abs experiment under identical reaction conditions (Figure S6). The 4.2-K/53-mT Mössbauer spectrum of this sample (Figure 4A, vertical bars) has minor contributions from the $\text{Fe}_2^{\text{III/III}}$ complex (6%) present in the reactant solution (purple trace) and unreacted $\text{Fe}_2^{\text{II/II}}$ species (14%, light blue trace). After removal of these two components, the remaining spectral features (Figure 4B) can be analyzed as two partially resolved quadrupole doublets of equal intensity (40% each of the integrated intensity of the original, experimental spectrum) with parameters $\delta_1 = 0.48 \text{ mm/s}$, $\Delta E_{\text{Q}1} = 0.49 \text{ mm/s}$ (blue subspectrum) and $\delta_2 = 0.55 \text{ mm/s}$, $\Delta E_{\text{Q}2} = 1.23 \text{ mm/s}$ (green subspectrum) characteristic of high-spin Fe^{III} subsites (Table 1).⁴³ The 4.2-K/6.0-T spectrum (Figure 4C, vertical bars) can be simulated with values of δ and ΔE_{Q} determined from the 53-mT spectrum, $\eta_1 = 0$, $\eta_2 = 0.9$, and $S_{\text{total}} = 0$. The 53-mT spectrum can be simulated equally well with parameters $\delta_1 = 0.32 \text{ mm/s}$, $\Delta E_{\text{Q}1} = 0.80 \text{ mm/s}$ and $\delta_2 = 0.70 \text{ mm/s}$, $\Delta E_{\text{Q}2} = 0.93 \text{ mm/s}$, but this analysis is inconsistent with the 6.0-T spectrum. From the isomer shifts of the two subsites (0.48 and 0.55 mm/s), it can be concluded that the intermediate has a $\text{Fe}_2^{\text{III/III}}$ core, which is more oxidized by two electrons than the cofactor in the $\text{Fe}_2^{\text{II/II}}\text{-ADO}$ •octanal reactant complex. Because the transient complex forms in a reaction without an obvious source of additional electrons, the cosubstrate O_2 must be reduced by two electrons to the peroxide level, yielding, for example, peroxo- $\text{Fe}_2^{\text{III/III}}$ state B or $\text{Fe}_2^{\text{III/III}}$ -peroxyhemiacetal state C in Scheme 1. The transient species is, therefore, referred to hereafter as a "Fe₂^{III/III}-peroxide" complex, a designation intended to encompass both




Figure 4. Characterization of the $\text{Fe}_2^{\text{III/III}}$ -peroxide complex by FQ Mössbauer spectroscopy. (A) 4.2-K/53-mT Mössbauer spectrum of a FQ sample prepared by reacting $\text{Fe}_2^{\text{II/II}}$ -ADO•1- ^{13}C -octanal with O_2 -saturated buffer for 32 s (vertical bars). The contribution of unreacted $\text{Fe}_2^{\text{II/II}}$ -ADO•1- ^{13}C -octanal (14%) and the $\text{Fe}_2^{\text{III/III}}$ -ADO (6%) are shown as light blue and purple lines, respectively. (B) Removal of the contributions from the $\text{Fe}_2^{\text{II/II}}$ -ADO•1- ^{13}C -octanal and $\text{Fe}_2^{\text{III/III}}$ -ADO from (A) yields the spectrum of the intermediate (vertical bars), which can be simulated as two quadrupole doublets of equal intensity (blue and green shaded areas and red line) with parameters quoted in the text. (C) 4.2-K/6-T spectrum of the 32-s sample described above (vertical bars) and simulation of the subspectra of the $\text{Fe}_2^{\text{III/III}}$ -peroxide intermediate (blue and green shaded areas) as described in the text. The spectrum of the minor $\text{Fe}_2^{\text{III/III}}$ contribution is shown in purple, and the summation of the contributions from the intermediate and $\text{Fe}_2^{\text{III/III}}$ contaminant is shown as the red line plotted on top of the data.

possible structural formulations. Correlation of the concentration of the intermediate determined from analysis of the Mössbauer data (80% of 0.75 mM giving $[\text{Fe}_2] = 0.60 \text{ mM}$) and the maximum value of the absorbance change ($\Delta A_{450} = 0.72$) in the SF-Abs experiment carried out under identical conditions (Figure S6) provides an estimate of ϵ_{450} for the intermediate of $1,200 \text{ M}^{-1} \text{ cm}^{-1}$. This value is similar to those observed for known peroxo- $\text{Fe}_2^{\text{III/III}}$ complexes.^{31,51–56}

The SF-Abs experiments of Figures 2 and 3 suggest that the $\text{Fe}_2^{\text{III/III}}$ -peroxide complex accumulates only in the presence of aldehyde. This conclusion was verified by FQ Mössbauer spectroscopy (Figure S7). Upon mixing of the anaerobically isolated *Np* ADO with O_2 -saturated buffer in the absence of an aldehyde substrate, the $\text{Fe}_2^{\text{III/III}}$ -peroxide complex was found to accumulate to a minor extent ($\sim 4\%$), if at all, after a reaction time of 15 s (Figure S7B). The spectrum reflects the presence of $\sim 11\%$ of the oxidized ($\text{Fe}_2^{\text{III/III}}$) state, which is also present in the reactant solution for this experiment (Figure S7A), but is dominated by the contribution from unreacted $\text{Fe}_2^{\text{II/II}}$ complex. Thus, the $\text{Fe}_2^{\text{II/II}}$ -ADO is relatively unreactive toward O_2 , as deduced from the SF-Abs data. By contrast, in the presence of *n*-decanal, the $\text{Fe}_2^{\text{III/III}}$ -peroxide complex accumulated (Figure S7C), albeit to a lesser extent ($\sim 44\%$) than in the experiment of Figure 4. The SF-Abs and FQ Mössbauer data thus confirm

Table 1. Mössbauer and Optical Properties of $\text{Fe}_2^{\text{III/III}}$ -peroxide Complexes in *Np* ADO and Other Diiron-Carboxylate Proteins

peroxo species	δ (mm/s)	ΔE_Q (mm/s)	λ_{max} (nm)	ref
ADO	0.48	0.49	450	this work
	0.55	1.23		
sMMO	0.66	1.51	725	31, 66
AurF	0.55	0.74	500	70
	0.61	0.35		
Δ^9 -ACP desaturase	0.64	1.06	700	51
	0.68	1.90		
ToMOH	0.55	0.67	n.d.	69
oxy-hemerythrin	0.54	1.92	~ 500	86
	0.51	1.09		
deoxyhypusine hydroxylase	0.55	1.16	630	92
	0.58	0.88		
frog M ferritin	0.62	1.08	650	56
<i>Ec</i> D84E RNR- β	0.63	1.58	700	53, 82

substrate triggering of O_2 addition to form the $\text{Fe}_2^{\text{III/III}}$ -peroxide complex.

Chemical Competence of the Intermediate for Reduction by Me^{O} PMS. According to our working hypothesis for the mechanism of the ADO reaction (Scheme 1), the $\text{Fe}_2^{\text{III/III}}$ -peroxide complex is expected to be analogous to state **B** or state **C** (which could also be in equilibrium). Accordingly, production of formate and the C_{n-1} alkane from this complex should require a total of two electrons from the reducing system. The ability of the complex to oxidize Me^{O} PMS, the reductant shown by Marsh and co-workers⁴⁷ to support greater ADO activity than the originally employed heterologous (spinach) ferredoxin,^{1,3,4} was assessed in a sequential-mix SF-Abs experiment. The $\text{Fe}_2^{\text{III/III}}$ -peroxide complex was allowed to accumulate by a 32-s incubation (t_{delay}) at 5°C following a first mix of the $\text{Fe}_2^{\text{II/II}}$ -ADO•1- ^{13}C -octanal complex with an equal volume of O_2 -saturated buffer, and the solution was then mixed with an equal volume of a solution containing one equivalent of the dithionite-reduced form of Me^{O} PMS. The reaction was monitored by the development of the spectrum of the oxidized Me^{O} PMS⁵⁷ with $\lambda_{\text{max}} = 388 \text{ nm}$ ($\epsilon_{388} \sim 26,000 \text{ M}^{-1} \text{ cm}^{-1}$) (Figure 5A). The ΔA_{388} -vs-time trace reveals rapid oxidation, essentially complete within 1 s (Figure 5B, green trace). Corresponding traces from control experiments omitting ADO (black), having an initial delay time insufficient to permit accumulation of the absorbing complex ($t_{\text{delay}} = 0.01 \text{ s}$, blue trace), or substituting the $\text{Fe}_2^{\text{II/II}}$ -ADO with the aerobically isolated $\text{Fe}_2^{\text{III/III}}$ form (Figure S8) reveal much slower Me^{O} PMS oxidation. Similarly, with $t_{\text{delay}} = 500 \text{ s}$, a time sufficient to permit decay of the transient complex to $\sim 50\%$ of its maximum concentration (Figure 3B, blue trace), the fast phase of Me^{O} PMS oxidation is diminished accordingly (Figure 5B, red trace). These observations show that the transient complex rapidly oxidizes Me^{O} PMS, whereas neither the $\text{Fe}_2^{\text{II/II}}$ -ADO•1-




Figure 5. Sequential-mixing SF-Abs experiments establishing the ability of the $\text{Fe}_2^{\text{III/III}}$ -peroxide complex to oxidize MeOPMS . The $\text{Fe}_2^{\text{II/II}}$ -ADO•1- ^{13}C -octanal reactant solution was mixed with O_2 -saturated buffer, the solution was allowed to react for the delay time (t_{delay}) indicated in the figure, the aged reaction solution was then mixed with a solution of dithionite-reduced MeOPMS , and spectra were acquired following the second mix. (A) Selected spectra from the experiment with $t_{\text{delay}} = 32$ s, reflecting the rapid oxidation of reduced MeOPMS . (B) ΔA_{388} -vs-time traces for experiments with different delay times between the two mixes [$t_{\text{delay}} = 0.01$ s (blue), 32 s (green), 500 s (red)]. A control experiment in which ADO was omitted is shown in black. Dashed lines are fits of eq 7 to the data.

^{13}C -octanal reactant nor the aerobically isolated $\text{Fe}_2^{\text{III/III}}$ -ADO does so. The results thus establish the oxidizing nature of the complex, consistent with the hypotheses that it possesses a bound peroxide moiety and is an intermediate on the pathway to formate and alk(a/e)n.

Chemical Competence of the Intermediate for Formate Production and Reaction Stoichiometry. The expectation that MeOPMS -promoted decay of the $\text{Fe}_2^{\text{III/III}}$ -peroxide intermediate should produce formate (Scheme 1) was evaluated in a chemical-quench experiment (see Experimental Section) designed to mimic the sequential-mixing SF-Abs experiment of Figure 5. The quantity of ^{13}C -labeled formate from 1- ^{13}C -octanal, determined after conversion to the formyl-2-nitrophenylhydrazide [^{13}CHO]-2NPH derivative,³ was found in two independent trials to vary as an approximately linear function of the quantity of reduced MeOPMS added (Figure 6).

Decay of the intermediate in the absence of reductant in an Eppendorf tube under ambient light produces only a low yield of formate (<15%). The formate that is produced under these conditions could reflect electron flow from the unreacted $\text{Fe}_2^{\text{II/II}}$ -ADO complex (~14% observed in the FQ Mössbauer




Figure 6. Determination of the reductant/formate stoichiometry upon reduction of the $\text{Fe}_2^{\text{III/III}}$ -peroxide complex with MeOPMS . The dotted fit line gives a formate/reductant ratio of ~0.5, which is half of the theoretical yield.

experiment of Figure 1) to the $\text{Fe}_2^{\text{III/III}}$ -peroxide complex. With increasing MeOPMS added (again, under ambient light), the yield of formate increases. Because the phenazine is a two-electron reductant, the maximum theoretical yield of formate/ MeOPMS is unity. The observed deviation most likely reflects the “uncoupling” of phenazine oxidation from formate production in approximately half of the reaction events. This result is not surprising in view of the nonphysiological nature of the reducing system and the possibilities for (1) unproductive two-electron reduction of the $\text{Fe}_2^{\text{III/III}}$ -peroxide intermediate and (2) competition for reaction with the reductant from either the $\text{Fe}_2^{\text{III/III}}$ -ADO product emerging from initial turnover of the intermediate, residual O_2 , or both. We conclude that the complex is chemically competent for formate production upon its reduction by MeOPMS and is an authentic intermediate in the ADO reaction.

Mössbauer-Spectroscopic Characterization of the Product of Reduction of the $\text{Fe}_2^{\text{III/III}}$ -Peroxide Intermediate by MeOPMS . The state of the ADO cofactor after the MeOPMS -induced productive decay of the $\text{Fe}_2^{\text{III/III}}$ -peroxide complex was probed by Mössbauer spectroscopy. A sample was freeze-quenched 0.56 s after the preformed complex was mixed with the reduced form of MeOPMS . This incubation time is sufficient to allow for the reaction to near completion (Figure 5B, green trace). The 4.2-K/53-mT Mössbauer spectrum of this sample (Figure 7A), has residual contributions from $\text{Fe}_2^{\text{II/II}}$ -ADO (~13%) but is dominated by broad quadrupole doublets. Subtracting the spectrum of the sample quenched before reaction with MeOPMS (Figure 4A) from the spectrum of postreduction sample (Figure 7A) cancels out the contributions of unreacted $\text{Fe}_2^{\text{II/II}}$ -ADO•octanal complex and $\text{Fe}_2^{\text{III/III}}$ impurity and illustrates that decay of the features of the transient complex (~46%, pink trace in Figure 7B) is associated with the appearance of a broad doublet (downward features). By adding back the 46% contribution of the $\text{Fe}_2^{\text{III/III}}$ -peroxide spectrum, the spectrum of the reduction product is obtained (Figure 7C). The spectrum can be adequately analyzed as two quadrupole doublets with parameters $\delta_1 = 0.55$ mm/s, $\Delta E_{\text{Q1}} = 1.80$ mm/s, $\delta_2 = 0.51$ mm/s, and $\Delta E_{\text{Q2}} = 1.08$ mm/s. These parameters are typical of high-spin Fe^{III} ions^{43,44} and indicative of an AF coupled high-spin $\text{Fe}_2^{\text{III/III}}$ cluster. The spectroscopic signature of this cluster is distinct from that of the $\text{Fe}_2^{\text{II/II}}$ complex that forms upon the slow (30 min at 5 °C) decay of the $\text{Fe}_2^{\text{III/III}}$ -peroxide complex in the absence of MeOPMS (Figure 7D, vertical bars). This latter complex has a spectrum that is essentially identical to that of aerobically isolated ADO. The observation that reduction of the transient $\text{Fe}_2^{\text{III/III}}$ complex by MeOPMS generates a distinct state with the cofactor still in the $\text{Fe}_2^{\text{III/III}}$ oxidation state confirms that the oxidizing equivalents of the transient complex are in the form of a bound peroxide. The fact that the spectrum of the ADO after the productive, reductive decay of the $\text{Fe}_2^{\text{III/III}}$ -peroxide state is distinct from those of the aerobically isolated enzyme and the spontaneous (unproductive) decay product could be explained by the presence of bound formate, alkane or both in at least a fraction of the reduction product, but other explanations are also possible.

DISCUSSION

Definitive Evidence that ADO Employs a Diiron Cofactor. A better understanding of the structure and mechanism of ADO might enable rational enhancement of its in vivo efficiency in renewable energy applications. A key aspect




Figure 7. Mössbauer-spectroscopic characterization of the ADO cofactor following the productive decay of the $\text{Fe}_2^{\text{III/III}}$ -peroxide intermediate that is initiated by reduced MeO-PMS . (A) Experimental 4.2-K/53-mT spectrum after reaction of the preformed complex with dithionite-reduced MeO-PMS for 0.56 s at 5 °C. (B) Difference spectrum (vertical bars) prepared by subtracting the spectrum of the sample quenched before reduction (Figure 4A) from spectrum A. In this presentation, the features associated with the $\text{Fe}_2^{\text{III/III}}$ -peroxide complex point upward (the pink line is the experimental "reference spectrum" shown in Figure 4B scaled to 46% intensity) and the features associated with the product of the productive decay point downward. (C) "Reference spectrum" of the product of the reaction of the $\text{Fe}_2^{\text{III/III}}$ -peroxide intermediate with reduced MeO-PMS (vertical bars) and quadrupole doublet simulations with parameters quoted in the text (solid lines). (D) Spectrum of a sample prepared by allowing the $\text{Fe}_2^{\text{III/III}}$ -peroxide intermediate to decay at 5 °C for 30 min in the absence of reduced MeO-PMS .

of its structure is the identities of the metal ions in its functional cofactor. Although a diiron cofactor has generally been presumed, the multiturnover enzyme assays reported to date have left open the possibility that a minor fraction of the ADO preparations, and not the major diiron component, was responsible for the alk(a/e)ne and formate produced. We sought to unequivocally test the catalytic competence of the diiron cofactor by preparing ADO as highly enriched in diiron cluster as possible and examining the reaction of these preparations in single turnover experiments, with steps in the sequence resolved by controlling the delivery of the requisite electrons.

The ICP-AES metal analysis and Mössbauer spectra show that our preparations of *Np* ADO produced in *Ec* grown on minimal medium supplemented with iron contain a diiron cluster. When the enzyme is isolated aerobically, it contains an AF coupled $\text{Fe}_2^{\text{III/III}}$ cofactor, which can be chemically reduced to the $\text{Fe}_2^{\text{II/II}}$ form. Following isolation under anaerobic conditions, it contains a $\text{Fe}_2^{\text{II/II}}$ cluster with spectroscopic properties identical to those of the chemically reduced form. Although the detection of a particular metal ion in a recombinant protein is, at best, very poor evidence that that

metal is *functional*, the reactivity of the Fe_2 -ADO characterized here indicates that it is indeed the active state. The reaction of $\text{Fe}_2^{\text{II/II}}$ -ADO with O_2 is very slow and produces the μ -oxo- $\text{Fe}_2^{\text{III/III}}$ state without obvious accumulation of an intermediate species. The reaction of the $\text{Fe}_2^{\text{II/II}}$ -ADO•aldehyde complex with O_2 is much faster and yields a transient complex with distinct absorption and Mössbauer spectra. Accumulation of this intermediate is strictly dependent on the presence of an *n*-aldehyde substrate: all short-chain *n*-aldehydes tested (C_6 – C_{10}) support its accumulation, whereas the corresponding *n*-alkyl compounds terminated by other functional groups (*n*-alkanoic acids, primary *n*-alcohols, and *n*-alkanes) do not. Aldehydes with longer side chains trigger intermediate formation more efficiently. This trend is expected, because the physiological substrates are C_{14} , C_{16} , or C_{18} saturated or monounsaturated fatty aldehydes provided by the acyl-ACP reductase and because the longer-chain substrates should bind more tightly within the hydrophobic active-site channel.¹ Triggering of O_2 reactivity by the substrate or an effector protein is a common characteristic of the ferritin-like diiron-carboxylate oxidases and oxygenases, and its occurrence in our ADO preparations is consistent with the diiron form being functional. Importantly, the fact that ~80% of the $\text{Fe}_2^{\text{II/II}}$ cluster can, with proper substrate triggering, convert to an intermediate that reacts rapidly with the reducing system to afford ~0.6 formate/ADO rules out the possibility that a minor fraction of the recombinant protein, and not the predominant enzyme form, could be responsible for the activity. Thus, the spectroscopic and reactivity data together provide the most definitive evidence to date that cyanobacterial ADOs can use a diiron cofactor in catalysis.

Nature of the Diiron- O_2 Intermediate. The chemically competent intermediate state exhibits an absorption band at 450 nm ($\epsilon_{450} = 1,200 \text{ M}^{-1} \text{ cm}^{-1}$) and two resolved Mössbauer quadrupole doublets characteristic of an AF coupled $\text{Fe}_2^{\text{III/III}}$ cluster. Its spectroscopic properties, its formation by reaction of the $\text{Fe}_2^{\text{II/II}}$ -ADO•aldehyde complex with O_2 in the absence of reductant, and its capacity to oxidize the reduced form of MeO-PMS rapidly while converting to a distinct $\text{Fe}_2^{\text{III/III}}$ state, imply that it possesses a bound peroxide. Indeed, this two-electron reduction of O_2 to peroxide is the common first step in the reactions of the carboxylate-bridged diiron oxygenases and oxidases.¹⁸ Moreover, with the exception of the complex formed in ToMOH, all peroxo- $\text{Fe}_2^{\text{III/III}}$ complexes trapped in metalloproteins and inorganic models thereof share with the ADO intermediate the characteristics of absorption in the visible region and Mössbauer spectra reflecting AF-coupling of two Fe^{III} ions (see Table 1).

Possible Structures of the Intermediate and Positions in the Chemical Mechanism. Scheme 1 shows the mechanistic hypothesis initially proposed by us³ and subsequently adopted by Marsh and co-workers⁵⁸ upon their corrections^{59,60} to their claim of O_2 -independent ADO activity.^{47,61} The mechanism accommodates all published data on the reaction, including: (i) our discovery that the C1-derived coproduct of *Np* ADO is formate² (which was subsequently confirmed for the ADO orthologue from *Prochlorococcus marinus* (*Pm*)⁶¹) rather than the originally proposed carbon monoxide¹ or carbon dioxide;⁵ (ii) our demonstration that the O-atom incorporated into the formate coproduct is derived from O_2 ^{3,4} rather than from solvent;⁴⁷ and (iii) recent evidence from the Marsh group for a C2 methylene radical intermediate accompanying formate production.⁵⁸ The mechanism can

therefore guide consideration of the structure of the diiron intermediate and its position in the reaction sequence.

The reaction begins with binding of the aldehyde substrate to $\text{Fe}^{\text{II/II}}\text{-ADO}$, yielding the reactant $\text{Fe}_2^{\text{II/II}}\text{-ADO}\cdot\text{aldehyde}$ complex (state A in Scheme 1). Coordination of the substrate in this manner could displace one or more solvent ligand (not depicted), a common mechanism for substrate triggering of O_2 addition.^{33–35,62} There is extensive precedent for triggering in iron oxygenases upon binding of the substrate merely in the vicinity of, and not directly to, the cofactor.^{26,33–35,62–65} Aldehyde coordination is, therefore, only a reasonable conjecture that is consistent with both the substrate triggering demonstrated here and the presence of a μ -1,3 fatty acid ligand suggested by the X-ray crystallographic data on the *Pm* ADO (PDB accession codes 2OC5 and 4KVQ).¹

Analogously to the reactions of other members of the superfamily, addition of O_2 to A would produce a peroxo- $\text{Fe}_2^{\text{III/III}}$ intermediate, B, which could have the symmetrical $\mu\text{-}\eta^2\text{:}\eta^2\text{-peroxo-Fe}^{\text{III/III}}$ core structure depicted in Scheme 1 or a less symmetrical peroxide binding mode (e.g., end-on to a single iron). The peroxide in B is proposed to nucleophilically attack the aldehyde carbonyl, yielding the peroxyhemiacetal state C. This step would distinguish the ADO mechanism from those of related diiron oxygenases and oxidases studied to date, in which peroxo- $\text{Fe}_2^{\text{III/III}}$ complexes undergo redox-neutral or reductive O–O bond cleavage to yield high-valent states (in sMMO⁶⁶ and class Ia ribonucleotide reductase β subunit (RNR- β),^{67,68} respectively) or are attacked by a nucleophilic substrate to yield oxidized (most often hydroxylated) products (in ToMOH⁶⁹ and perhaps AurF^{54,70}). The proposed nucleophilicity of the ADO peroxide complex is thus reminiscent of the ambivalent electrophilicity/nucleophilicity of C4a-(hydro)peroxyflavin intermediates in flavin-dependent monooxygenases.⁷¹ The demonstrated protonolyses of an inorganic peroxo- $\text{Fe}_2^{\text{III/III}}$ complex⁷² and the corresponding intermediate in ferritin⁷³ to produce H_2O_2 suggest that such complexes can indeed also be basic/nucleophilic. Moreover, the nucleophilic attack by a peroxo-Fe intermediate on a carbonyl carbon has been proposed as a key step in the C–C-bond-cleaving reactions of the mixed-valent nonheme diiron enzyme, *myo*-inositol oxygenase (MIOX),^{74,75} the mononuclear nonheme-iron enzymes, 2-hydroxyethyl-1-phosphonate dioxygenase (HEPD)^{76,77} and methylphosphonate synthase (MPNS),⁷⁸ and the heme-dependent cytochrome P450 sterol 14a-demethylase (CYP51).^{7,79}

The next step in the proposed mechanism is the one-electron reductive cleavage of the O–O bond of the $\text{Fe}_2^{\text{III/III}}$ -peroxyhemiacetal complex in C by the reducing system, which yields a μ -oxo- $\text{Fe}_2^{\text{III/III}}$ cluster and *gem*-diolyl radical (D). The radical undergoes a β -scission of the C1–C2 bond, generating formate and a C2 methylene radical (E). Support for the formation of this radical was recently provided by the Marsh group, who found evidence for the expected ring-opening accompanying formate production from a C3–C4 cyclopropanated substrate analogue.⁵⁸ The C2 methylene radical then abstracts a hydrogen atom from a nearby amino acid (XH in Scheme 1, pink arrows) or a solvent ligand to the cofactor (green arrows), and quenching of the resultant oxidized species (amino acid radical or $\text{Fe}_2^{\text{III/IV}}$ cluster) by another electron from the reducing system and a proton gives the product state, F. The fact that the Mössbauer spectrum of the samples quenched immediately after the *productive* (reductive) decay of the $\text{Fe}_2^{\text{III/III}}$ -peroxide complex differs

significantly from the spectra of both the aerobically purified protein and the product of *spontaneous* decay of the intermediate is consistent with the formation of a downstream intermediate such as state F. Release of the products from state F and two-electron reduction of the cofactor by the reducing system completes the cycle.

The peroxo- $\text{Fe}_2^{\text{III/III}}$ state, B, and $\text{Fe}_2^{\text{III/III}}$ -peroxyhemiacetal state, C, in Scheme 1 both have the requisite characteristics of a III/III oxidation state and a bound peroxide equivalent and are, therefore, both viable representations of the ADO intermediate. Three characteristics of the intermediate favor a C-like structure but can, with additional rationalization, also accommodate a B-like structure. First, formation of the intermediate requires a bound aldehyde. The carbonyl group of the substrate would not be directly involved in formation of B but would be indispensable for formation of C. The observation that *n*-alkyl compounds lacking the carbonyl uniformly fail to support intermediate accumulation could thus be readily understood in terms of structure C. However, an alternative explanation is that the structural requirements for activating the $\text{Fe}_2^{\text{II/II}}$ cofactor for addition of O_2 to form B are just unusually stringent and fulfilled only by the aldehyde. Second, the intermediate persists for minutes in the absence of reductant but oxidizes the reduced form of MeO^{PMS} in hundreds of milliseconds. Progression of B to C in Scheme 1 is redox neutral, but conversion of C to D requires an electron. The promotion of its decay by reduced MeO^{PMS} (yielding formate) is thus most naturally rationalized by a C-like structure. However, B and C could interconvert in a rapid equilibrium favoring B. In this case, decay of B would be fast only under conditions favoring rapid reductive trapping of C. Alternatively, binding of reduced MeO^{PMS} to state B could induce its conversion to C and subsequent fast reduction to D. Indeed, the Marsh group provided evidence for the tight binding of a similar phenazine to the enzyme.⁶¹ Either of these idiosyncratic kinetic scenarios could rationalize both the stability of a complex similar to B in the absence of reductant and the acceleration of its decay by reduced MeO^{PMS} .

The third characteristic of the intermediate that is also most easily explained by a $\text{Fe}_2^{\text{III/III}}$ -peroxyhemiacetal rather than a μ -peroxo- $\text{Fe}_2^{\text{III/III}}$ formulation is its Mössbauer spectrum, which exhibits more pronounced site-differentiation than has often been seen among μ -peroxo- $\text{Fe}_2^{\text{III/III}}$ complexes analogous to B (Table 1). An inorganic complex originally prepared by Kitajima and the intermediate trapped in the D84E variant of *Ec* RNR- β are the most extensively studied among the known inorganic and protein complexes (respectively) of this type. Each has been shown to possess a *cis*- μ -1,2-peroxo- $\text{Fe}_2^{\text{III/III}}$ core structure.^{53,80–82} Each exhibits an absorption band at ~ 700 nm, assigned to a peroxide $\rightarrow \text{Fe}^{\text{III}}$ charge transfer transition,⁸¹ but only a single Mössbauer quadrupole doublet with a rather high isomer shift (>0.60 mm/s) for both Fe sites. Peroxide complexes in the wild-type RNR- β proteins from *Ec* and mouse,^{67,68} sMMO,^{31,66} ToMOH,⁶⁹ and frog M ferritin,⁵⁶ as well as several inorganic complexes,^{52,80,83,84} also have iron sites that are unresolved by Mössbauer and thought to contain the *cis*- μ -1,2-peroxo- $\text{Fe}_2^{\text{III/III}}$ core, though calculations on the intermediate from sMMO suggest that a $\mu\text{-}\eta^2\text{:}\eta^2\text{-peroxo}$ binding mode is also possible.⁸⁵

There are, however, several examples of protein^{51,54,86} and inorganic (hydro)peroxo- $\text{Fe}_2^{\text{III/III}}$ complexes^{87,88} that *do* exhibit site resolution in their Mössbauer spectra (Table 1). The peroxo- $\text{Fe}_2^{\text{III/III}}$ complexes in oxy-hemerythrin, the *N*-oxygenase

AurF, and $\Delta^9\text{D}$ desaturase are such examples. Differentiation of the two iron subsites in these complexes arises either from asymmetric coordination of the peroxide moiety (e.g., to a single Fe), from asymmetry imposed by the protein or solvent ligands, or from a combination of these factors. For example, in oxyhemerythrin, coordinative saturation of one iron with a μ -hydroxo, two μ -1,3-carboxylate, and three histidine ligands enforces addition of O_2 terminally only to the other Fe^{II} ion,⁸⁹ which has one fewer histidine and thus a single open coordination site. The structure of the (hydro)peroxo- $\text{Fe}_2^{\text{III/III}}$ intermediate in AurF is not known, but here again the protein provides one more histidine ligand to the Fe^{II} in site 1⁹⁰ and may therefore enforce O_2 addition either only to the site 2 Fe^{II} or in a manner that more intimately involves and profoundly affects site 2. On the other hand, for the peroxo- $\text{Fe}_2^{\text{III/III}}$ state in $\Delta^9\text{D}$, recent quantum mechanical calculations suggest that the asymmetry reflected by its Mössbauer spectrum arises in spite of an inherently symmetrical μ -1,2 mode of peroxide coordination by virtue of a bidentate-to-monodentate carboxylate shift at one of the Fe^{III} ions, which makes that site five-coordinate while leaving the other Fe^{III} six-coordinate.⁹¹

The amino-acid iron ligands of ADO are, at least in terms of their identities, inherently site-symmetrical (two His and four Glu residues), as they are in $\Delta^9\text{D}$. The site resolution in the Mössbauer spectrum of the $\text{Fe}_2^{\text{III/III}}$ -peroxide complex in ADO is thus most likely associated with either rearrangement of protein ligand(s) in the first coordination sphere (as proposed for $\Delta^9\text{D}$), asymmetrical coordination of the peroxide moiety, direct, asymmetric coordination of the substrate (or an intermediate derived from it), or a combination of these factors. As depicted in Scheme 1, the postulated $\text{Fe}_2^{\text{III/III}}$ -peroxyhemiacetal state C could potentially possess both an asymmetrically coordinated hemiacetal anion and an asymmetrically bound peroxide. In this respect, state C provides a potential rationale for the spectroscopic properties of the $\text{Fe}_2^{\text{III/III}}$ -peroxide intermediate in ADO. More detailed structural studies on the intermediate will be required to test our hypothesis that it is a unique type of peroxide (peroxyhemiacetal) adduct to its nonheme diiron cofactor.

CONCLUSIONS

Our results verify that cyanobacterial ADO can employ a *diiron* cofactor rather than a homo- or heterodinuclear cofactor involving one or more transition metal other than iron. Upon reaction of the $\text{Fe}_2^{\text{II/II}}$ -ADO•substrate complex with O_2 in the absence of the reducing system, an intermediate with a $\text{Fe}_2^{\text{III/III}}$ cluster and bound peroxide equivalent accumulates to ~80% yield. It is chemically competent to generate formate upon reacting with the reducing system, and this reaction generates a stable $\text{Fe}_2^{\text{III/III}}$ cluster that is spectroscopically distinct from the aerobically isolated “resting” state. The identification of the intermediate provides support for our previously proposed mechanism involving a novel attack of the peroxide of a peroxo- $\text{Fe}_2^{\text{III/III}}$ complex upon the substrate carbonyl and the reductant promoted decomposition of the resultant peroxyhemiacetal complex, leading to formate and alkane production via a free-radical β -scission mechanism.

ASSOCIATED CONTENT

Supporting Information

Figures S1–S8 containing spectra of stopped-flow absorption and Mössbauer experiments. This material is available free of charge via the Internet at <http://pubs.acs.org>.

AUTHOR INFORMATION

Corresponding Authors

sjb14@psu.edu
cdk10@psu.edu
jmb21@psu.edu

Present Addresses

[§]N.L.: Joule Unlimited Technologies, Inc., 18 Crosby Drive, Bedford, MA 01730.

^{||}H.N.: Department of Plant and Environmental Sciences, University of Copenhagen, Denmark.

Author Contributions

[†]Equal contributors: M.E.P. and N.L.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

This work was supported by the National Science Foundation (MCB-1122079 to C.K., S.J.B., and J.M.B.) and the National Institutes of Health (GM-055365 to J.M.B. and C.K.).

REFERENCES

- (1) Schirmer, A.; Rude, M. A.; Li, X. Z.; Popova, E.; del Cardayre, S. B. *Science* **2010**, *329*, 559.
- (2) Warui, D. M.; Li, N.; Nørgaard, H.; Krebs, C.; Bollinger, J. M., Jr.; Booker, S. J. *J. Am. Chem. Soc.* **2011**, *133*, 3316.
- (3) Li, N.; Nørgaard, H.; Warui, D. M.; Booker, S. J.; Krebs, C.; Bollinger, J. M., Jr. *J. Am. Chem. Soc.* **2011**, *133*, 6158.
- (4) Li, N.; Chang, W. C.; Warui, D. M.; Booker, S. J.; Krebs, C.; Bollinger, J. M., Jr. *Biochemistry* **2012**, *51*, 7908.
- (5) Reppas, N. B., and Ridley, C. P. Methods and compositions for the recombinant biosynthesis of *n*-alkanes U.S. Patent 7794969, Joule Unlimited Inc., 2010.
- (6) Ducat, D. C.; Way, J. C.; Silver, P. A. *Trends Biotechnol.* **2011**, *29*, 95.
- (7) Patra, T.; Manna, S.; Maiti, D. *Angew. Chem., Int. Ed.* **2011**, *50*, 12140.
- (8) Zhang, F. Z.; Rodriguez, S.; Keasling, J. D. *Curr. Opin. Biotechnol.* **2011**, *22*, 775.
- (9) Robertson, D. E.; Jacobson, S. A.; Morgan, F.; Berry, D.; Church, G. M.; Afeyan, N. B. *Photosynth. Res.* **2011**, *107*, 269.
- (10) Serrano-Ruiz, J. C.; Ramos-Fernández, E. V.; Sepúlveda-Escribano, A. *Energy Environ. Sci.* **2012**, *5*, 5638.
- (11) Schirmer, A.; Rude, M. A.; Brubaker, S. Methods and compositions for producing hydrocarbons, U.S. Patent 140696, LS9, 2009.
- (12) Akhtar, M. K.; Turner, N. J.; Jones, P. R. *Proc. Natl. Acad. Sci. U.S.A.* **2013**, *110*, 87.
- (13) Wallar, B. J.; Lipscomb, J. D. *Chem. Rev.* **1996**, *96*, 2625.
- (14) Merkx, M.; Kopp, D. A.; Sazinsky, M. H.; Blazyk, J. L.; Müller, J.; Lippard, S. J. *Angew. Chem., Int. Ed.* **2001**, *40*, 2782.
- (15) Stubbe, J. *Curr. Opin. Chem. Biol.* **2003**, *7*, 183.
- (16) Fox, B. G.; Lyle, K. S.; Rogge, C. E. *Acc. Chem. Res.* **2004**, *37*, 421.
- (17) Andrews, S. C. *Biochim. Biophys. Acta, Gen. Subj.* **2010**, *1800*, 691.
- (18) Krebs, C.; Bollinger, J. M., Jr.; Booker, S. J. *Curr. Opin. Chem. Biol.* **2011**, *15*, 291.
- (19) Nordlund, P.; Eklund, H. *Curr. Opin. Struct. Biol.* **1995**, *5*, 758.
- (20) Barynin, V. V.; Whittaker, M. M.; Antonyuk, S. V.; Lamzin, V. S.; Harrison, P. M.; Artymiuk, P. J.; Whittaker, J. W. *Structure* **2001**, *9*, 725.
- (21) Jiang, W.; Yun, D.; Saleh, L.; Barr, E. W.; Xing, G.; Hoffart, L. M.; Maslak, M.-A.; Krebs, C.; Bollinger, J. M., Jr. *Science* **2007**, *316*, 1188.
- (22) Boal, A. K.; Cotruvo, J. A., Jr.; Stubbe, J.; Rosenzweig, A. C. *Science* **2010**, *329*, 1526.

- (23) Cotruvo, J. A., Jr.; Stubbe, J. *Biochemistry* **2011**, *50*, 1672.
- (24) Price, J. C.; Barr, E. W.; Hoffart, L. M.; Krebs, C.; Bollinger, J. M., Jr. *Biochemistry* **2005**, *44*, 8138.
- (25) Bollinger, J. M., Jr.; Price, J. C.; Hoffart, L. M.; Barr, E. W.; Krebs, C. *Eur. J. Inorg. Chem.* **2005**, 4245.
- (26) Matthews, M. L.; Krest, C. M.; Barr, E. W.; Vaillancourt, F. H.; Walsh, C. T.; Green, M. T.; Krebs, C.; Bollinger, J. M., Jr. *Biochemistry* **2009**, *48*, 4331.
- (27) Broadwater, J. A.; Ai, J. Y.; Loehr, T. M.; Sanders-Loehr, J.; Fox, B. G. *Biochemistry* **1998**, *37*, 14664.
- (28) Paulsen, K. E.; Liu, Y.; Fox, B. G.; Lipscomb, J. D.; Münck, E.; Stankovich, M. T. *Biochemistry* **1994**, *33*, 713.
- (29) Liu, Y.; Nesheim, J. C.; Lee, S. K.; Lipscomb, J. D. *J. Biol. Chem.* **1995**, *270*, 24662.
- (30) Bailey, L. J.; McCoy, J. G.; Phillips, G. N.; Fox, B. G. *Proc. Natl. Acad. Sci. U.S.A.* **2008**, *105*, 19194.
- (31) Tinberg, C. E.; Lippard, S. J. *Acc. Chem. Res.* **2011**, *44*, 280.
- (32) Lee, S. J.; McCormick, M. S.; Lippard, S. J.; Cho, U.-S. *Nature* **2013**, *494*, 380.
- (33) Solomon, E. I.; Brunold, T. C.; Davis, M. I.; Kemsley, J. N.; Lee, S. K.; Lehnert, N.; Neese, F.; Skulan, A. J.; Yang, Y. S.; Zhou, J. *Chem. Rev.* **2000**, *100*, 235.
- (34) Neidig, M. L.; Brown, C. D.; Light, K. M.; Fujimori, D. G.; Nolan, E. M.; Price, J. C.; Barr, E. W.; Bollinger, J. M., Jr.; Krebs, C.; Walsh, C. T.; Solomon, E. I. *J. Am. Chem. Soc.* **2007**, *129*, 14224.
- (35) Pavel, E. G.; Zhou, J.; Busby, R. W.; Gunsior, M.; Townsend, C. A.; Solomon, E. I. *J. Am. Chem. Soc.* **1998**, *120*, 743.
- (36) Hitchman, M. L. *Measurement of dissolved oxygen*; John Wiley and Sons: New York, 1978.
- (37) Connors, K. A. *Chemical kinetics: the study of reaction rates in solution*; Wiley-VCH: New York, 1990.
- (38) Espenson, J. H. *Chemical kinetics and reaction mechanisms*; McGraw-Hill: New York, 1995.
- (39) Jiang, W.; Hoffart, L. M.; Krebs, C.; Bollinger, J. M., Jr. *Biochemistry* **2007**, *46*, 8709.
- (40) Whitteck, J. T.; Cicchillo, R. M.; van der Donk, W. A. *J. Am. Chem. Soc.* **2009**, *131*, 16225.
- (41) Price, J. C.; Barr, E. W.; Tirupati, B.; Bollinger, J. M., Jr.; Krebs, C. *Biochemistry* **2003**, *42*, 7497.
- (42) Fish, W. W. In *Methods in Enzymology*; James, F., Riordan, B. L. V., Eds.; Academic Press: New York, 1988; Vol. 158, p 357.
- (43) Münck, E. Aspects of ^{57}Fe Mössbauer spectroscopy. In *Physical Methods in Bioinorganic Chemistry*; Que, L., Jr., Ed.; University Science Books: Sausalito, CA, 2000; pp 287–319.
- (44) Kurtz, D. M. *Chem. Rev.* **1990**, *90*, 585.
- (45) Fox, B. G.; Shanklin, J.; Somerville, C.; Münck, E. *Proc. Natl. Acad. Sci. U.S.A.* **1993**, *90*, 2486.
- (46) Brown, C. A.; Remar, G. J.; Musselman, R. L.; Solomon, E. I. *Inorg. Chem.* **1995**, *34*, 688.
- (47) Eser, B. E.; Das, D.; Han, J.; Jones, P. R.; Marsh, E. N. G. *Biochemistry* **2011**, *50*, 10743.
- (48) Valentine, A. M.; Stahl, S. S.; Lippard, S. J. *J. Am. Chem. Soc.* **1999**, *121*, 3876.
- (49) Xing, G.; Diao, Y.; Hoffart, L. M.; Barr, E. W.; Prabhu, K. S.; Arner, R. J.; Reddy, C. C.; Krebs, C.; Bollinger, J. M., Jr. *Proc. Natl. Acad. Sci. U.S.A.* **2006**, *103*, 6130.
- (50) Hayashi, Y.; Kayatani, T.; Sugimoto, H.; Suzuki, M.; Inomata, K.; Uehara, A.; Mizutani, Y.; Kitagawa, T.; Maeda, Y. *J. Am. Chem. Soc.* **1995**, *117*, 11220.
- (51) Broadwater, J. A.; Achim, C.; Münck, E.; Fox, B. G. *Biochemistry* **1999**, *38*, 12197.
- (52) Dong, Y. H.; Zang, Y.; Shu, L. J.; Wilkinson, E. C.; Que, L., Jr.; Kauffmann, K.; Münck, E. *J. Am. Chem. Soc.* **1997**, *119*, 12683.
- (53) Bollinger, J. M., Jr.; Krebs, C.; Vicol, A.; Chen, S. X.; Ley, B. A.; Edmondson, D. E.; Huynh, B. H. *J. Am. Chem. Soc.* **1998**, *120*, 1094.
- (54) Korboukh, V. K.; Li, N.; Barr, E. W.; Bollinger, J. M., Jr.; Krebs, C. *J. Am. Chem. Soc.* **2009**, *131*, 13608.
- (55) Tinberg, C. E.; Lippard, S. J. *Biochemistry* **2009**, *48*, 12145.
- (56) Pereira, A. S.; Small, W.; Krebs, C.; Tavares, P.; Edmondson, D. E.; Theil, E. C.; Huynh, B. H. *Biochemistry* **1998**, *37*, 9871.
- (57) Zaugg, W. S. *J. Biol. Chem.* **1964**, *239*, 3964.
- (58) Paul, B.; Das, D.; Ellington, B.; Marsh, E. N. G. *J. Am. Chem. Soc.* **2013**, *135*, 5234–5237.
- (59) Eser, B. E.; Das, D.; Han, J.; Jones, P. R.; Marsh, E. N. G. *Biochemistry* **2012**, *51*, 5703.
- (60) Das, D.; Eser, B. E.; Han, J.; Sciore, A.; Marsh, E. N. G. *Angew. Chem., Int. Ed.* **2012**, *51*, 7881.
- (61) Das, D.; Eser, B. E.; Han, J.; Sciore, A.; Marsh, E. N. G. *Angew. Chem., Int. Ed.* **2011**, *50*, 7148.
- (62) Zhou, J.; Kelly, W. L.; Bachmann, B. O.; Gunsior, M.; Townsend, C. A.; Solomon, E. I. *J. Am. Chem. Soc.* **2001**, *123*, 7388.
- (63) Brazeau, B. J.; Lipscomb, J. D. *Biochemistry* **2000**, *39*, 13503.
- (64) Xue, G. Q.; Pokutsa, A.; Que, L., Jr. *J. Am. Chem. Soc.* **2011**, *133*, 16657.
- (65) Yang, Y. S.; Broadwater, J. A.; Pulver, S. C.; Fox, B. G.; Solomon, E. I. *J. Am. Chem. Soc.* **1999**, *121*, 2770.
- (66) Liu, K. E.; Valentine, A. M.; Wang, D. L.; Huynh, B. H.; Edmondson, D. E.; Salifoglou, A.; Lippard, S. J. *J. Am. Chem. Soc.* **1995**, *117*, 10174.
- (67) Tong, W. H.; Chen, S.; Lloyd, S. G.; Edmondson, D. E.; Huynh, B. H.; Stubbe, J. *J. Am. Chem. Soc.* **1996**, *118*, 2107.
- (68) Yun, D.; García-Serres, R.; Chicalese, B. M.; An, Y. H.; Huynh, B. H.; Bollinger, J. M., Jr. *Biochemistry* **2007**, *46*, 1925.
- (69) Murray, L. J.; Naik, S. G.; Ortillo, D. O.; García-Serres, R.; Lee, J. K.; Huynh, B. H.; Lippard, S. J. *J. Am. Chem. Soc.* **2007**, *129*, 14500.
- (70) Li, N.; Korboukh, V. K.; Krebs, C.; Bollinger, J. M., Jr. *Proc. Natl. Acad. Sci. U.S.A.* **2010**, *107*, 15722.
- (71) Ballou, D. P.; Entsch, B.; Cole, L. J. *Biochem. Biophys. Res. Commun.* **2005**, *338*, 590.
- (72) LeCloux, D. D.; Barrios, A. M.; Lippard, S. J. *Biorg. Med. Chem.* **1999**, *7*, 763.
- (73) Jameson, G. N. L.; Jin, W.; Krebs, C.; Perreira, A. S.; Tavares, P.; Liu, X. F.; Theil, E. C.; Huynh, B. H. *Biochemistry* **2002**, *41*, 13435.
- (74) Bollinger, J. M., Jr.; Diao, Y.; Matthews, M. L.; Xing, G.; Krebs, C. *Dalton Trans.* **2009**, 905.
- (75) Hirao, H.; Morokuma, K. *J. Am. Chem. Soc.* **2009**, *131*, 17206.
- (76) Cicchillo, R. M.; Zhang, H. J.; Blodgett, J. A. V.; Whitteck, J. T.; Li, G. Y.; Nair, S. K.; van der Donk, W. A.; Metcalf, W. W. *Nature* **2009**, *459*, 871.
- (77) Hirao, H.; Morokuma, K. *J. Am. Chem. Soc.* **2010**, *132*, 17901.
- (78) Metcalf, W. W.; Griffin, B. M.; Cicchillo, R. M.; Gao, J. T.; Janga, S. C.; Cooke, H. A.; Circello, B. T.; Evans, B. S.; Martens-Habbena, W.; Stahl, D. A.; van der Donk, W. A. *Science* **2012**, *337*, 1104.
- (79) Roberts, E. S.; Vaz, A. D.; Coon, M. J. *Proc. Natl. Acad. Sci. U.S.A.* **1991**, *88*, 8963.
- (80) Kim, K.; Lippard, S. J. *J. Am. Chem. Soc.* **1996**, *118*, 4914.
- (81) Brunold, T. C.; Tamura, N.; Kitajima, N.; Moro-oka, Y.; Solomon, E. I. *J. Am. Chem. Soc.* **1998**, *120*, 5674.
- (82) Moënne-Loccoz, P.; Baldwin, J.; Ley, B. A.; Loehr, T. M.; Bollinger, J. M., Jr. *Biochemistry* **1998**, *37*, 14659.
- (83) Ménage, S.; Brennan, B. A.; Juarez-Garcia, C.; Münck, E.; Que, L., Jr. *J. Am. Chem. Soc.* **1990**, *112*, 6423.
- (84) Zhang, X.; Furutachi, H.; Fujinami, S.; Nagatomo, S.; Maeda, Y.; Watanabe, Y.; Kitagawa, T.; Suzuki, M. *J. Am. Chem. Soc.* **2005**, *127*, 826.
- (85) Rinaldo, D.; Philipp, D. M.; Lippard, S. J.; Friesner, R. A. *J. Am. Chem. Soc.* **2007**, *129*, 3135.
- (86) Okamura, M. Y.; Klotz, I. M.; Johnson, C. E.; Winter, M. R. C.; Williams, R. J. P. *Biochemistry* **1969**, *8*, 1951.
- (87) Ookubo, T.; Sugimoto, H.; Nagayama, T.; Masuda, H.; Sato, T.; Tanaka, K.; Maeda, Y.; Okawa, H.; Hayashi, Y.; Uehara, A.; Suzuki, M. *J. Am. Chem. Soc.* **1996**, *118*, 701.
- (88) LeCloux, D. D.; Barrios, A. M.; Mizoguchi, T. J.; Lippard, S. J. *J. Am. Chem. Soc.* **1998**, *120*, 9001.
- (89) Holmes, M. A.; Letrong, I.; Turley, S.; Sieker, L. C.; Stenkamp, R. E. *J. Mol. Biol.* **1991**, *218*, 583.

(90) Choi, Y. S.; Zhang, H. J.; Brunzelle, J. S.; Nair, S. K.; Zhao, H. M. *Proc. Natl. Acad. Sci. U.S.A.* **2008**, *105*, 6858.

(91) Srnec, M.; Rokob, T. A.; Schwartz, J. K.; Kwak, Y.; Rulíšek, L.; Solomon, E. I. *Inorg. Chem.* **2012**, *51*, 2806.

(92) Vu, V. V.; Emerson, J. P.; Martinho, M.; Kim, Y. S.; Münck, E.; Park, M. H.; Que, L., Jr. *Proc. Natl. Acad. Sci. U.S.A.* **2009**, *106*, 14814.