

Robust Affinity Standards for Cu(I) Biochemistry


Pritha Bagchi,[†] M. Thomas Morgan,[†] John Bacsa,[‡] and Christoph J. Fahrni^{*,†}

[†]School of Chemistry and Biochemistry, Petit Institute for Bioengineering and Bioscience, Georgia Institute of Technology, 901 Atlantic Drive, Atlanta, Georgia 30332, United States

[‡]X-ray Crystallography Center, Department of Chemistry, Emory University, 1515 Dickey Drive, Atlanta, Georgia 30322, United States

S Supporting Information

ABSTRACT: The measurement of reliable Cu(I) protein binding affinities requires competing reference ligands with similar binding strengths; however, the literature on such reference ligands is not only sparse but often conflicting. To address this deficiency, we have created and characterized a series of water-soluble monovalent copper ligands, MCL-1, MCL-2, and MCL-3, that form well-defined, air-stable, and colorless complexes with Cu(I) in aqueous solution. X-ray structural data, electrochemical measurements, and an extensive network of equilibrium titrations showed that all three ligands form discrete Cu(I) complexes with 1:1 stoichiometry and are capable of buffering Cu(I) concentrations between 10^{-10} and 10^{-17} M. As most Cu(I) protein affinities have been obtained from competition experiments with bathocuproine disulfonate or 2,2'-bicinchoninic acid, we further calibrated their Cu(I) stability constants against the MCL series. To demonstrate the application of these reagents, we determined the Cu(I) binding affinity of CusF ($\log K = 14.3 \pm 0.1$), a periplasmic metalloprotein required for the detoxification of elevated copper levels in *Escherichia coli*. Altogether, this interconnected set of affinity standards establishes a reliable foundation that will facilitate the precise determination of Cu(I) binding affinities of proteins and small-molecule ligands.



INTRODUCTION

Copper is an essential trace element that is central to a broad range of biological processes, including cellular respiration, connective tissue formation, pigment synthesis, antioxidant defense, and photosynthesis in plants and bacteria.¹ Cellular copper levels are controlled by an intricate network of membrane transporters and metallochaperones, which escort the metal ion to specific subcellular locations for incorporation into metalloproteins or export through secretory pathways.² To rationalize the directivity of these trafficking routes, a detailed knowledge of the copper binding affinities of all involved ligands is essential.^{3,4} Although a host of copper-transporting proteins have been characterized, the reported affinity constants are often conflicting.^{4–11} These discrepancies were likely caused by the difficulties associated with determining the stability constants for complexes of Cu(I), the predominant oxidation state of copper in the reducing intracellular environment. Aqueous Cu(I) is not only highly reactive toward dioxygen but also prone to disproportionation at acidic pH or precipitation as Cu₂O at neutral or basic pH. Recently, Xiao and co-workers pointed out common pitfalls in stability constant measurements and offered a set of guidelines for the determination of Cu(I) protein binding affinities.^{8,9} Foremost, the choice of suitable equilibrium conditions is key for obtaining reliable binding affinities, irrespective of the analytical method applied. As a general rule, the degree of complex formation should not exceed 90% at equimolar concentrations of metal ion and


protein to avoid large errors during data analysis.¹² Because of the high affinities of Cu(I)-transporting proteins,¹⁰ direct addition of Cu(I) would result in nearly quantitative binding. For this reason, the stability constants of high-affinity ligands are best obtained through competition experiments with an affinity standard whose stability constant and solution chemistry have been well-characterized. At present, only a few such reference ligands exist for Cu(I), and similar to the reports on protein affinities, the published values are often conflicting. Because of the limited stability of Cu(I) in aqueous solution, the majority of Cu(I) stability constants were obtained from experimental Cu(II) affinities and the standard reduction potentials of the free and ligand-bound Cu(II/I) couples using the Nernst equation. Although this approach eliminates the challenges associated with the redox instability of aqueous Cu(I), there remain uncertainties due to the ionic strength dependence of the reduction potentials and the often large structural differences between the mono- and divalent forms of the copper–ligand complexes, which may lead to irreversible redox processes.

To address the scarcity of reliable Cu(I) affinity standards, we report here a set of three new water-soluble monovalent copper ligands, MCL-1, MCL-2, and MCL-3 (1–3; Chart 1), which we have rigorously characterized in terms of their Cu(I)

Received: August 26, 2013

Published: December 3, 2013

Chart 1



binding affinities and coordination chemistry in aqueous buffer. Rather than relying solely on the Nernst relationship to determine the Cu(I) affinities, we also identified the dibasic ligand PEMEA¹³ (4) as a pH-modulated intermediary to anchor the Cu(I) affinities of the strongly binding MCL series of chelators to that of the much weaker ligand acetonitrile, whose Cu(I) complex stability constants have been determined via a kinetic method.¹⁴ As the majority of Cu(I) protein affinities have been obtained from competition experiments with the chromogenic ligands bathocuproine disulfonate (BCS, 5) and 2,2'-bicinchoninic acid (BCA, 6), we additionally determined their Cu(I) stability constants by competition titrations with the MCL series of compounds. The colorless MCL series of affinity standards complement the chromogenic reagents by allowing direct monitoring of the degree of Cu(I) saturation of the target ligand in spectrophotometric or fluorimetric titrations. To illustrate this capability, we measured the Cu(I) affinity of CusF, a tryptophan-containing periplasmic protein important to copper resistance in *Escherichia coli*. Altogether, we have established a set of reliable Cu(I) affinity standards that should be broadly applicable for the determination of Cu(I) affinities in aqueous buffer at physiological pH.

EXPERIMENTAL SECTION

Synthesis. The procedures and analytical data for the synthesis of ligands 1–3, their Cu(I) complexes, and all intermediates are provided in the Supporting Information. The ligands BCS (disodium bathocuproine disulfonate hydrate, 97%) and BCA (2,2'-bicinchoninic acid, free acid, 98%) were purchased from Acros Organics and MP Biomedicals, respectively; PEMEA and DHEAMP were synthesized according to published procedures^{13,15} and purified by recrystallization as the phosphate (PEMEA) and hydrochloride (DHEAMP) salts.

Protonation Constants. All of the protonation constants ($\log K_{\text{H}}$) were derived from measurements of the hydronium ion concentration ($-\log[\text{H}_3\text{O}^+]$ or $\text{p}[\text{H}]$), not the activity.¹⁶ For this purpose, a combination glass electrode was calibrated by titrating a 5 mM solution of HCl in 0.1 M KCl (ionic background) with a standardized solution of 0.1 M KOH at 25 °C. The end point, electrode potential, and slope were determined using Gran's method as implemented in the GLEE software package.¹⁷ The protonation constants of PEMEA were measured by spectrophotometric titrations at 25 °C with $\text{p}[\text{H}]$ ranging between 1.9 and 8.2 (0.1 M KCl), followed by nonlinear least-squares fitting over the entire spectral range using the SPECFIT software package.¹⁸ The $\log K_{\text{H}}$ of BCA was analogously determined by varying the $\text{p}[\text{H}]$ between 3.7 and 5.5. All other protonation constants were obtained through potentiometric titrations by stepwise addition (motorized buret) of a standardized solution of 0.1 M KOH

to a solution of the ligand in 0.1 M KCl at 25 °C (ligand concentrations were 1 mM for BCS and 5 mM for MCL-1, MCL-2, and DHEAMP·2HCl). In the cases of BCS, MCL-1, and MCL-2, at least one full molar equivalent of HCl was added to protonate the basic sites prior to titration with KOH. The protonation constants were then determined from the potentiometric data by means of nonlinear least-squares fitting using the Hyperquad software package.¹⁹

Electrochemistry. Cyclic voltammograms were acquired in 10 mM PIPBS buffer (pH 5) containing 0.1 M KClO_4 as the electrolyte using a CH-Instruments potentiostat (model 600A). Experiments were carried out under an atmosphere of argon in a single-compartment cell with a glassy carbon working electrode, a Pt counter electrode, and an aqueous Ag/AgCl reference electrode (1 M KCl). The half-wave potentials ($E_{1/2}$) were referenced to ferrocenium (0.40 V vs SHE)^{20,21} or ferriin (1.112 V vs SHE)^{13,22} as external standards. Measurements were typically performed at a scan rate of 20–50 mV s^{-1} .


Stability Constants. Titrations involving Cu(I) were carried out under an atmosphere of argon with thorough exclusion of oxygen. All of the solutions were degassed immediately before use, and reagents were supplied from freshly prepared stock solutions with a gastight syringe. All spectrophotometric competition equilibrium titrations were carried out in a 1 cm path length quartz cuvette that was sealed with a PTFE-lined rubber septum. Equilibration times were determined by means of separate kinetic experiments. Stability constants were obtained from the titration data by nonlinear least-squares fitting over the relevant spectral range using the SPECFIT software package.¹⁸ Detailed experimental conditions for each equilibrium titration are provided in the Supporting Information.

X-ray Structural Determinations. Crystals were coated with Paratone N oil, suspended in a small fiber loop, and placed in a cooled nitrogen gas stream at 173 K on a Bruker D8 APEX II CCD sealed-tube diffractometer with graphite-monochromatized Mo $K\alpha$ radiation (0.71073 Å). Data collection, indexing, and initial cell refinements were carried out using APEX II software.²³ Frame integration and final cell refinements were done using the SAINT software.²⁴ All of the structures were solved using direct methods and difference Fourier techniques (SHELXTL, version 6.12).²⁵ Hydrogen atoms were placed in their expected chemical positions and were included in the final cycles of least-squares analysis with isotropic U_{ij} values related to the atom ridden upon. All non-hydrogen atoms were refined anisotropically except for the disordered oxygen and fluorine atoms of the ClO_4 and PF_6 anions, respectively. Additional details of data collection and structure refinement are given in Tables S1–S8 in the Supporting Information. Supplementary crystallographic data [CCDC 948257 (13a), 948431 (14a), and 949729 (16)] can be obtained free of charge from the Cambridge Crystallographic Data Centre via <http://www.ccdc.cam.ac.uk/Community/Requestastructure/Pages/DataRequest.aspx>.

Expression and Purification of CusF. *E. coli* BL21(DE3) cells hosting the pASK-IBA3 plasmid with the gene encoding for CusF residues 6–88 were grown at 37 °C to an optical density of 0.6–1.1 (600 nm) in LB medium containing 100 mg/L ampicillin.²⁶ Protein expression was induced with anhydrotetracycline (0.8 mg/L), and the cells were harvested after 5 h by centrifugation. The cell pellet was resuspended in MOPS buffer (50 mM MOPS, 0.15 M NaCl, pH 7), and the protein was extracted by three freeze–thaw cycles. After agitation at 4 °C for 1 h, the cell suspension was centrifuged at 6000g for 15 min at 4 °C, and the supernatant was concentrated with a Microsep 3K centrifugal device (Pall Life Sciences). The protein extract was purified by two sequential gel filtrations (GE Äkta) using a Superdex 75 column equilibrated with MOPS buffer. Fractions containing pure protein were confirmed by Laemmli SDS-PAGE, pooled, and concentrated. The protein was stored at –20 °C in the elution buffer for up to 2 days. Trace elemental analysis by TXRF (Bruker S2 Picofox spectrometer) revealed that purified CusF was present in the apo form (Figure S27 in the Supporting Information).

Cu(I) Affinity of CusF. A. Fluorescence Equilibrium Titration with MCL-3. A 20 μM solution of holo-CusF in MOPS buffer (50 mM MOPS, 0.15 M NaCl, pH 7.0) was titrated with MCL-3 (up to a

Scheme 1. Synthesis of the Water-Soluble Cu(I) Ligands MCL-1 (1), MCL-2 (2), and MCL-3 (3)



concentration of 280 μM). After addition of each aliquot, a fluorescence emission spectrum was acquired over the range of 305–500 nm with excitation at 290 nm. The precise concentration of CusF was determined from a molar-ratio titration of the apo protein with Cu(I), which was generated in situ by reduction of CuSO_4 with sodium ascorbate. To ensure quantitative loading with Cu(I), the holo protein was prepared from apo-CusF by addition of 1.0 molar equiv of Cu(I). The Cu(I) stability constant of CusF was then determined by nonlinear least-squares fitting of the fluorescence emission traces over the entire spectral range using the SPECFIT¹⁸ software package.

B. Spectrophotometric Equilibrium Titration with BCA. The equilibrium titrations were carried out by the following two approaches: either (1) the Cu(I) complex of BCA was first formed in situ by addition of $[\text{Cu}(\text{I})(\text{CH}_3\text{CN})_4]\text{PF}_6$ (16 μM) to a solution of BCA (50 μM) in MOPS buffer at pH 7 (50 mM MOPS, 0.15 M NaCl, 25 $^\circ\text{C}$) and then titrated with apo-CusF (0–72 μM) at pH 7 or (2) apo-CusF (65 μM) was titrated with $[\text{Cu}(\text{I})(\text{CH}_3\text{CN})_4]\text{PF}_6$ (0–87 μM) in the presence of 80 μM BCA as a competing ligand. In either case, the Cu(I) stability constant of CusF was obtained by nonlinear least-squares fitting of the UV–vis traces over the 450–700 nm range using SPECFIT.¹⁸

RESULTS AND DISCUSSION

Ligand Design. For biochemical applications, Cu(I) affinity standards should be water-soluble, discourage heteroleptic complex formation, and ideally form a redox-stable Cu(I) complex even under aerobic conditions. The latter requires a substantially positive reduction potential of the ligand-bound Cu(II/I) couple and discourages the use of redox-labile donors such as thiols or phosphines in the ligand design. Various aliphatic thioether-based macrocyclic or tripodal ligands with S_4 or NS_3 donor sets form stable 1:1 Cu(I) complexes under aerobic conditions and offer the additional advantage of transparency in the visible to near-UV range, thus minimizing spectral overlap with a competing ligand during spectrophotometric or fluorimetric titrations. While linear or macrocyclic ligands with an aliphatic S_4 or NS_3 donor set provide affinities in the range of $\log K_{\text{Cu}(\text{I})} = 12\text{--}14$,^{27,28} N-centered tripodal ligands such as tris(2-methylthioethyl)amine (TMMEA) provide stability constants 2–3 orders of magnitude larger and LCu(II/I) reduction potentials approximately 300 mV higher than those of their macrocyclic counterparts.¹³ Building upon the TMMEA architecture, we designed the water-soluble derivative MCL-1 by replacing the terminal methyl groups with 3-sulfopropyl moieties (Chart 1). In the structurally related ligand MCL-2, each of the three N–S bridges was elongated by a methylene spacer, a change that was found to lower the apparent affinity at pH 7 by approximately 5 orders of

magnitude relative to MCL-1 (vide infra). Finally, the thiocrown ligand MCL-3 was devised to bridge the affinity gap between the two tripodal ligands. While the parent macrocycle [16]ane S_4 exhibits a $\log K_{\text{Cu}(\text{I})}$ of 12.0,²⁷ we anticipated that the conformational preorganization imposed by the bulky solubilizing groups should substantially increase the affinity. A similar effect has been reported for [14]ane S_4 , whose Ni(II) affinity in nitromethane increased 49-fold (1.7 logarithmic units) upon geminal substitution with two pairs of methyl groups.²⁹

Ligand Synthesis. Both MCL-1 and MCL-2 were prepared in good yield from the corresponding tris(ω -chloroalkyl)amines 7 and 8 via reaction with excess sodium 3-mercaptopropanesulfonate in the presence of sodium hydroxide followed by recrystallization from methanol–water mixtures (Scheme 1). The NaCl byproduct was efficiently removed under these conditions as a result of the negative temperature dependence of its solubility in methanol.³⁰ After rigorous drying, both ligands were obtained as the anhydrous trisodium salts. The macrocyclic ligand MCL-3 was constructed from the versatile thietane building block 9, which was previously utilized in our syntheses of water-soluble Cu(I)-selective fluorescent probes.³¹ As outlined in Scheme 1, double ring opening with 1,3-diiodopropane gave diiodide 10, which was cyclized with 1,3-propanedithiol under Kellogg conditions³² and deprotected with aqueous HCl to give thiocrown 12. Alkylation of the corresponding lithium alkoxide with 1,3-propanesultone followed by precipitation with ethanolic NaI provided MCL-3 as the tetrasodium salt.

Synthesis and X-ray Crystal Structures of Cu(I) Complexes. Ligands MCL-1, MCL-2, and MCL-3 were reacted with stoichiometric quantities of $[\text{Cu}(\text{I})(\text{CH}_3\text{CN})_4]\text{PF}_6$ to produce the respective Cu(I) complexes 13–15. All three complexes were isolated as hexafluorophosphate adducts containing equal numbers of sodium cations and sulfonate groups, corresponding to the formulas of $[\text{Na}_3\text{LCu}(\text{I})]\text{PF}_6$ for MCL-1 and MCL-2 and $[\text{Na}_4\text{LCu}(\text{I})]\text{PF}_6$ for MCL-3. Crystallization of the MCL-2–Cu(I) complex (14) from ethanol–water yielded the hydrate $\text{Na}_6(\text{Cu}(\text{I})\text{L})_2(\text{PF}_6)_2 \cdot 15\text{H}_2\text{O}$ (14a) as colorless tablets suitable for X-ray diffraction. Under the same conditions, the PF_6^- adduct of MCL-1–Cu(I) produced only very thin acicular crystals; however, in the presence of excess NaClO_4 , the complex formed the perchlorate adduct $[\text{Na}_7(\text{Cu}(\text{I})\text{L})_2](\text{ClO}_4)_3 \cdot 4\text{H}_2\text{O}$ (13a), which crystallized as colorless triclinic prisms. Because crystallization of MCL-3–Cu(I) (15) under a wide variety of

conditions yielded only fibrous acicular crystals or powders, we selected compound **12** (Scheme 1) as a truncated analogue of MCL-3 (hereafter abbreviated as *tr*MCL-3) for crystal structure determination. The corresponding Cu(I) complex crystallized from ethanol as clear prisms with the formula [(*tr*MCL-3)–Cu(I)]PF₆·EtOH (**16**). To gauge the molecular preorganization of the tetrasubstituted thiocrown, we also determined the structure of free *tr*MCL-3, which was crystallized from DMSO–water. Lists of selected bond distances and angles for these structures are compiled in Tables 1 and 2. Additional data concerning the X-ray structural determinations are provided in the Supporting Information.

Table 1. Selected Bond Distances (Å) and Angles (deg) Describing the Cu(I) Coordination Geometries in the Crystal Structures of [Na₇(Cu(I)MCL-1)₂](ClO₄)₃·4H₂O (13a**) and [Na₆(Cu(I)MCL-2)₂](PF₆)₂·15H₂O (**14a**)**

	13a	14a (I)	14a (II)
Cu1–S1	2.2486(7)	2.2868(6)	2.2874(6)
Cu1–S2	2.2886(7)	2.2870(6)	2.2941(6)
Cu1–S3	2.2680(7)	2.2889(6)	2.2910(6)
Cu1–N1	2.1610(17)	2.1261(17)	2.1242(17)
S1–Cu1–N1	91.91(6)	98.78(5)	98.85(5)
S2–Cu1–N1	89.88(5)	97.86(5)	98.90(5)
S3–Cu1–N1	91.32(5)	98.51(5)	97.98(5)
S1–Cu1–S2	121.41(2)	117.60(2)	117.33(2)
S1–Cu1–S3	125.49(2)	118.36(2)	118.53(2)
S2–Cu1–S3	112.99(3)	117.77(2)	117.60(2)

Table 2. Selected Bond Distances (Å) and Angles (deg) for the Two Cationic Units in the Crystal Structure of [Cu(I)–(*tr*MCL-3)]PF₆·EtOH (16**)**

	16 (I)	16 (II)
Cu1–S1	2.2558(2)	2.2729(3)
S2–Cu1	2.2577(2)	2.2576(2)
S3–Cu1	2.3024(2)	2.3339(2)
S4–Cu1	2.3108(2)	2.3088(3)
S1–Cu1–S3	118.896(9)	116.703(10)
S1–Cu1–S4	110.019(9)	104.482(9)
S1–Cu1–S2	106.648(9)	106.517(9)
S2–Cu1–S3	106.772(9)	109.358(9)
S2–Cu1–S4	117.871(9)	123.669(10)
S3–Cu1–S4	97.026(8)	96.580(9)

Complexes with Tripodal Ligands. The Cu(I) adducts of the tripodal ligands MCL-1 and MCL-2 crystallized as discrete monomeric complexes in which the Cu(I) center is coordinated by the apical nitrogen and all three thioether donors (Figure 1). In both structures, the sulfopropyl arms adopt a helical conformation and do not participate in copper coordination. Instead, the anionic sulfonate groups are bridged by sodium cations and water molecules, forming a hydrophilic layer that links two Cu(I) complexes of opposing helicity within a single unit cell (Figure S1 in the Supporting Information). In the case of MCL-1 (**13a**), the two complexes contained within a unit cell are exact mirror images, whereas for MCL-2 (**14a**), slight structural differences produce a pair of pseudoenantiomeric geometries (I) and (II) (Table 1).

As illustrated with the ORTEP plot in Figure 1A, the Cu(I) complex of MCL-1 (**13a**) adopts a nearly trigonal-pyramidal coordination geometry with an S–Cu–S bond angle sum of




Figure 1. ORTEP drawings and atom numbering schemes for the crystal structures of the Cu(I) complexes of (A) MCL-1 (**13a**) and (B) MCL-2 [**14a (I)**]. Ellipsoids shown represent 50% probability. Hydrogen atoms and counterions have been omitted for clarity. A packing diagram for each structure is provided in Figure S1 in the Supporting Information.

359.9°. Despite the open coordination space below the basal plane of the trigonal pyramid, no additional donors are present within reasonable bonding distance in the crystal structure, rendering the coordination geometry of MCL-1–Cu(I) essentially identical to that previously reported for Cu(I)–TMMEA.¹³ In contrast to MCL-1, the longer N–S bridges of MCL-2 might be expected to accommodate a near-tetrahedral Cu(I)-center; however, the coordination geometry of complex **14a** is still strongly distorted toward a trigonal-pyramidal arrangement (Figure 1B). The sum of the S–Cu–S bond angles (353.7°) is substantially greater than that expected for an ideal tetrahedral geometry (328.4°) and approaches the value for a trigonal arrangement as in MCL-1–Cu(I). For comparison, the Cu(I) complex of the structurally related ligand tris(3-(benzylthio)propan-1-yl)amine (**18**) containing benzyl groups in place of the sulfopropyl moieties crystallized in four distinct conformers (*Z* = 4) with S–Cu–S bond angle sums ranging from 331.7 to 348.3°,³³ suggesting that the modest structural differences between **13a** and **14a** are not associated with a significant energetic penalty.

As is evident from Table 1, a comparison of the bond distances reveals a similar picture compared to the trends in

bond angles. The Cu–N and average Cu–S bond lengths of 2.161(2) and 2.27(2) Å, respectively, observed for **13a** are almost identical to the values of 2.17(1) and 2.26(1) Å previously reported for TMMEA–Cu(I),¹³ indicating that the pendant sulfopropyl groups have little effect on the Cu(I)–S interactions. The average Cu–N bond in **14a** is shorter by 0.036 Å compared with that in **13a**, a difference that may be attributed to the greater basicity of the nitrogen in MCL-2 compared with MCL-1 (vide infra); however, the magnitude of the effect is rather modest in view of the variation of the Cu–N bond lengths observed for the Cu(I) complex of ligand **18**, which differed by up to 0.030 Å within the same unit cell.³³

Macrocyclic Ligand and Complex. The macrocyclic ligand *tr*MCL-3 and Cu(I) form a discrete complex (**16**) with 1:1 stoichiometry (Figure 2A). Similar to the crystal structure of




Figure 2. ORTEP drawings and atom numbering schemes for the crystal structures of (A) [Cu(I)–(*tr*MCL-3)]PF₆ [**16** (**1**)] and (B) the free ligand *tr*MCL-3 (**12**). Ellipsoids shown represent 50% probability. Hydrogen atoms have been omitted for clarity.

complex **14a**, the unit cell contains two conformers with notable structural differences (Table 2). The average S–Cu bond distance in complex **16** (2.287 ± 0.030 Å, averaged over both conformers) is essentially identical to that in the MCL-2 complex **14a** (2.289 Å), and the overall coordination geometry is best described as flattened tetrahedral, with transannular S–Cu–S angles of 116.7–123.7° (vs 109.5° for an ideal tetrahedral arrangement). Overall, the crystallographic features of complex **16** are consistent with strong coordination between *tr*MCL-3 and Cu(I). Furthermore, all of the hydroxyl groups point away from the Cu(I) center, suggesting that complex **16** is a viable structural analogue for the Cu(I)–MCL-3 complex (**15**).

The crystal structure of free *tr*MCL-3 (Figure 2B) indicates significant preorganization toward Cu(I) coordination, especially when compared with the previously reported crystal structure of the unsubstituted parent ligand [16]aneS₄ (Figure 3).³⁴ In the latter, two of the thioether donors are in exo positions and point away from the macrocyclic cavity with a S–




Figure 3. Space-filling models illustrating the molecular preorganization of *tr*MCL-3 (**12**) vs [16]aneS₄³⁴ in comparison to the geometry of [Cu(I)–(*tr*MCL-3)]PF₆ (**16**) (representations were generated with Mercury CSD³⁵ from crystal structure atomic coordinates).

S separation of 8.48 Å. By contrast, the corresponding S–S distance in *tr*MCL-3 is only 6.13 Å, thus bringing the thioether donors much closer to the 4.03 Å transannular separation observed for the Cu(I) complex **16**. To underscore the importance of this preorganization, we note that no discrete monomeric [16]aneS₄–Cu(I) complex has been previously observed in the solid state; in fact, the crystal structure of a [16]aneS₄–copper(I) iodide adduct revealed a coordination polymer with the thiocrown in an exodentate conformation.³⁶ A dihydroxy derivative of [16]aneS₄ also failed to yield a discrete Cu(I) complex, and the Cu(I)–perchlorate adduct showed evidence of oxidation to Cu(II) even under an argon atmosphere.³⁷ In stark contrast, both *tr*MCL-3 and MCL-3 itself yield Cu(I) complexes that are air-stable even in solution. On the basis of these results, it appears that the preorganization imposed by the placement of the bulky solubilizing groups in MCL-3 not only increases the Cu(I) affinity relative to [16]aneS₄ (vide infra) but may actually be essential for the selective formation of a discrete 1:1 Cu(I) complex.

Protonation Constants. As the metal ion affinities of ligands with one or more basic donor atoms are pH-dependent, we determined for each compound the corresponding protonation constants (log *K*_H). A compilation of these results is provided in Table 3. All of the log *K*_H values were determined on the basis of the hydronium ion concentration (–log[H₃O⁺] or p[H]). Following the recommendation of Martell and Smith,³⁸ the tabulated concentration constants must be corrected upward by 0.11 (*I* = 0.1) to derive the corresponding mixed-mode protonation constants where hydronium ions are

Table 3. Protonation Constants for Ligands 1–6 and DHEAMP¹⁵ (17) in Aqueous Solution at 25 °C (0.1 M KCl)^a

ligand	log <i>K</i> _{H1}	log <i>K</i> _{H2}
MCL-1 (1)	7.00(2)	
MCL-2 (2)	8.98(1)	
MCL-3 (3)	–	
PEMEA (4)	7.24(4)	3.23(4)
BCS (5)	5.70(2)	
BCA (6)	3.80(2)	
DHEAMP (17)	6.94(1)	

^aFor each reported value, the standard deviation of the mean of three independent titrations is listed in parentheses. The digit refers to the significant figure furthest to the right [e.g., 7.00(2) means 7.00 ± 0.02].

Scheme 2




Table 4. Reduction Potentials and Averaged Thermodynamic and Conditional Stability Constants for Cu(I)/Cu(II) Complexes of Ligands 1–6 in Aqueous Solution at 25 °C (0.1 M KCl)

ligand	E^f (V vs SHE) ^a	$\log K_{\text{Cu(II)L}}$	$\log K_{\text{Cu(I)L}}$	$\log K'_{\text{Cu(I)L}}$ ^b	pCu ^c
MCL-1 (1)	0.716(1)	6.42(2)	16.33(2)	16.0	16.9
MCL-2 (2)	n/a	–	13.08(13)	11.0	11.9
MCL-3 (3)	0.729(3)	3.47(4)	13.80(3)	13.8	14.8
PEMEA (4)	0.595(2)	7.85(8)	15.71(2)	15.2	16.2
BCS (5) ^f	0.626(1)	12.42(7) ^d	20.81(2) ^d	20.8 ^e	16.6
BCA (6)	n/a	–	17.66(3) ^d	17.7 ^e	13.5

^aHalf-wave potential at pH 5.0. ^bApparent affinity at pH 7.0 (0.1 M KClO₄); ^cpCu = $-\log [\text{Cu(I)}_{\text{aq}}]$ at pH 7.0 for ligand and Cu(I) concentrations of 10 and 1 μM , respectively. ^d $\log \beta_2$ of the 2:1 complex. ^e $\log \beta'_2$. ^fMixture of the *m*–*m'*, *m*–*p'*, and *p*–*p'* isomers (60:32:8).

expressed in terms of their activities (pH) (see the Supporting Information). Because neutral PEMEA and BCA were not soluble at millimolar concentrations in aqueous solution, the protonation constants for these ligands were determined by spectrophotometric titrations at micromolar concentrations (Figures S2–S5 in the Supporting Information). The protonation constant of BCS was measured at 1 mM. All of the other compounds were titrated at 5 mM, with subsequent data fitting using Hyperquad.¹⁹

Potentiometric titrations of MCL-1 at 0.1 M ionic strength yielded a protonation constant of 7.00 ± 0.02 for the aliphatic amino group of the ligand. The corresponding mixed-mode protonation constant of 7.11 is lower than those reported for TMMEA (8.36) and its ethyl analogue TEMEA (8.32) in aqueous solution¹³ but closely matches the value of 7.1 for TEMEA extrapolated from data acquired in 80% water–methanol.³⁹ The differences between the extrapolated and measured protonation constants reported in the literature might be due to hydrophobic effects. Elongation of the N–S bridge in going from MCL-1 to MCL-2 increases the basicity of the amino group by 2 orders of magnitude to 8.98, a trend that

was also observed for tris(3-methylthiopropyl)amine (TMMPA) in 80% methanol when compared with TEMEA under the same conditions.³⁹ Finally, the previously reported values for the protonation constants of BCS (5.7),⁹ BCA (3.74 ± 0.04 ; 0.2 M ionic strength),⁴⁰ and PEMEA ($\log K_{\text{H1}} = 7.33$ and $\log K_{\text{H2}} = 3.26$, reported as mixed-mode protonation constants)¹³ corroborated well with the values determined in this work (Table 3).

Stability Constants of Cu(I) Complexes. The simplest method for the determination of complex stability constants involves the direct titration of a ligand with the metal ion; however, this approach yields reliable values only if the fractional complex formation resides below 90%,¹² a condition that is difficult to satisfy with tight-binding ligands. To arrive at a set of reliable thermodynamic stability constants for the Cu(I) complexes of ligands 1–6, we instead performed a series of equilibrium titrations, each involving a pair of ligands that compete for Cu(I) binding (Scheme 2). Because the concentration of aqueous Cu(I) is very low under these conditions, precipitation of Cu₂O and disproportionation to Cu(II) and Cu(0) are thermodynamically unfavorable and therefore

not of concern. Even the reactivity toward dioxygen is greatly reduced in the presence of a Cu(I)-stabilizing ligand, though all titrations were still carried out under deoxygenated conditions as a standard precautionary measure. For ligands that formed Cu(I) complexes with a reversible or quasi-reversible redox reaction, we derived the corresponding Cu(I) stability constants from the half-wave potentials E^f and the Cu(II) complex stability constant using the Nernst relationship (eq 1):

$$E^f = E_{\text{aq}}^{\circ} - \frac{2.303RT}{nF} \log \left(\frac{K_{\text{Cu(II)L}}}{K_{\text{Cu(I)L}}} \right) \quad (1)$$

where E_{aq}° represents the reduction potential for the aqueous Cu(II/I) couple expressed in terms of concentrations. To account for the differential activity at 0.1 M ionic strength, the IUPAC standard potential of $E^{\circ} = 0.153 \text{ V}^{20}$ was adjusted to $E_{\text{aq}}^{\circ} = 0.13 \text{ V}$ as recommended by Rorabacher (also see the Supporting Information).²⁸ In combination with the titration data, we established an interconnected network of thermodynamic equilibria from which we were able to arrive at a set of consistent Cu(I) stability constants for all of the involved ligands (Scheme 2). To outline the relative dependence and cross-validations within this extensive data set, the following sections describe the thermodynamic equilibria in more detail. A compilation of the pertinent thermodynamic data is given in Table 4. Stability constants which were determined based on multiple thermodynamic equilibria are reported as the cumulative average.

Ligand MCL-1. Slow-scan cyclic voltammetry studies of MCL-1 at pH 5 in the presence of excess Cu(II) revealed a quasi-reversible one-electron redox process at $E^f = 0.716 \text{ V}$ vs SHE with a peak separation of 74 mV (Figure 4A), thus




Figure 4. Cyclic voltammograms for (A) MCL-1 (150 μM) and (B) MCL-3 (150 μM) in the presence of CuSO_4 (1 mM) at pH 5.0 (10 mM PIPBS, 0.1 M KClO_4 ; glassy carbon electrode, scan rate 20 mV/s).

offering the opportunity to determine the corresponding Cu(I) stability constant from the Nernst relationship (eq 1). The Cu(II) stability constant of MCL-1 was obtained by spectrophotometric titration of the ligand with CuSO_4 under the same conditions (pH 5.0, 10 mM PIPBS, 0.1 M KClO_4 ; Figure S9 in the Supporting Information). Nonlinear least-squares fitting over the entire spectral range yielded $\log K_{\text{Cu(II)L}}$

$= 6.42 \pm 0.02$. On the basis of the above data, we derived a Cu(I) stability constant of $\log K_{\text{Cu(I)L}} = 16.33 \pm 0.02$. Taking into account the protonation constant of MCL-1, we calculated an apparent affinity of $\log K_{\text{Cu(I)L}}^{\text{app}} = 16.0$ at pH 7.0.

2,2'-Bicinchoninic Acid (BCA). In the presence of Cu(I), this ligand forms a purple 2:1 complex with an absorption maximum at 562 nm ($\epsilon = 7900 \text{ M}^{-1} \text{ cm}^{-1}$).⁴¹ Cyclic voltammetry experiments revealed irreversible redox behavior, thus precluding the application of the Nernst relationship to determine its Cu(I) stability constant. Taking advantage of the optical transparency of Cu(I)–MCL-1 in the visible range, we determined the stability constant of BCA by a spectrophotometric competition titration between the two ligands (Figure 5A). For this purpose, equimolar quantities of BCA and the




Figure 5. Spectrophotometric titrations of (A) 2,2'-bicinchoninic acid (BCA, **6**; 100 μM) at pH 5.0 (10 mM PIPBS, 0.1 M KClO_4) and (B) bathocuproine disulfonate (BCS, **5**; 50 μM) at pH 7.0 (10 mM PIPES, 0.1 M KClO_4) with MCL-1 in the presence of Cu(I).

Cu(I) complex of MCL-1 were pre-equilibrated in 10 mM PIPBS (pH 5.0, 0.1 M KClO_4). Under these conditions, less than 50% of the total BCA is bound to Cu(I), thus excluding the presence of a significant amount of the 1:1 complex. Titration with MCL-1 up to 4 molar equiv followed by nonlinear least-squares fitting yielded a stability constant of $\log \beta_2 = 17.67 \pm 0.03$. This value is 0.5 logarithmic units higher than the Cu(I) affinity estimated by Xiao et al.,⁹ however, the difference is consistent with their selected value of E_{aq}° ⁴² which is 34 mV higher than the ionic-strength-adjusted potential of 0.13 V used throughout this work.

Ligand MCL-2. Despite its structural similarity with MCL-1, we were not able to detect the formation of a complex with Cu(II) even at high-millimolar concentrations. Consistent with this observation, cyclic voltammetry experiments with the Cu(I) complex of MCL-2 showed an irreversible redox process, suggesting rapid dissociation of the copper ion upon oxidation. For this reason, we determined the Cu(I) affinity of MCL-2 only by spectrophotometric competition titrations with BCA

(Figure S16 in the Supporting Information). The data fit well to a 1:1 binding model and yielded $\log K_{\text{Cu(I)L}} = 13.08 \pm 0.13$, corresponding to an apparent affinity of 11.0 at pH 7.0. Compared with that of MCL-1, the Cu(I) affinity of MCL-2 is remarkably lower, a result that would not be predicted on the basis of the crystal structure data, which indicated similarly strong Cu(I) coordination for the two ligands. The dramatically lower Cu(I) affinity of MCL-2 is presumably due to a greater entropic penalty for conformational restriction of the longer N–S bridges upon chelation. While six-membered chelate rings generally result in less stable metal complexes compared with the analogous five-membered rings, Cu(I) complexes often exhibit little or no such effect.⁴³

Bathocuproine Disulfonate (BCS, Disodium Salt). Similar to BCA, this heterocyclic ligand and Cu(I) form a colored 2:1 complex with an absorption maximum at 483 nm ($\epsilon = 13\,300 \text{ M}^{-1} \text{ cm}^{-1}$).⁴⁴ Commercially available BCS is supplied as a mixture of three regioisomers bearing the sulfonate groups at either the meta or para positions of the 4- and 7-aryl rings (Chart 1).⁴⁵ Taking advantage of the resonance assignments previously established by multidimensional NMR techniques,⁴⁵ we determined for our batch a 60:32:8 composition of the *m*–*m'*, *m*–*p'*, and *p*–*p'* isomers, respectively. The following data should therefore be interpreted as a statistical average for this composition. Competition titrations with MCL-1 performed in the same fashion as described for BCA yielded a stability constant of $\log \beta_2 = 20.80 \pm 0.03$ (Figure S5B and Figure S17 in the Supporting Information). Slow-scan cyclic voltammetry experiments with BCS in the presence of either $[\text{Cu(I)}\text{-(CH}_3\text{CN)}_4]\text{PF}_6$ or CuSO_4 revealed a quasi-reversible one-electron redox process with $E^f = 0.626 \text{ V}$ vs SHE and a peak separation of 73 mV (Figure S7 in the Supporting Information), allowing an independent determination of the Cu(I) complex stability constant via eq 1. Although the Cu(II) complex stability constant of BCS was previously assessed via a potentiometric method requiring millimolar ligand concentrations,⁹ we relied instead on a spectrophotometric titration at micromolar concentrations to avoid interference from the reported self-association of BCS in aqueous solution.⁴⁶ For this purpose, we identified *N,N*-bis(2-hydroxyethyl)-*N*-(2-pyridylmethyl)amine (DHEAMP, 17) as suitable competing ligand, which according to Damu et al.¹⁵ forms a 1:1 Cu(II) complex with a stability constant of $\log K_{\text{Cu(II)L}} = 9.2 \pm 0.1$. On the basis of a molar ratio titration of DHEAMP with Cu(II) at low pH combined with the experimental protonation constant of 6.94 ± 0.01 (lit. 6.92 ± 0.01), we obtained a $\log K_{\text{Cu(II)L}} = 9.21 \pm 0.02$, in good agreement with the above literature value (Figure S11 in the Supporting Information). Finally, a competition titration of BCS with CuSO_4 in the presence of DHEAMP yielded a stability constant of $\log \beta_2 = 12.42 \pm 0.07$ for the 2:1 Cu(II) complex of BCS (Figure S12 in the Supporting Information). Together with the half-wave potential of the Cu(II/I) complex, we calculated a Cu(I) stability constant of $\log \beta_2 = 20.81 \pm 0.08$, which is in excellent agreement with the data obtained by the above competition titration with MCL-1. As in the case of BCA, the above result is higher than the previously reported $\log \beta_2$ of 19.9;⁹ here the deviation is somewhat larger than the 0.6 logarithmic units expected for a 34 mV difference in E_{aq}° .

Ligand MCL-3. In the presence of Cu(II), this ligand forms a complex with an absorption maximum at 577 nm and a strong charge-transfer band centered around 443 nm. A direct spectrophotometric molar ratio titration at pH 5.0 yielded a

Cu(II) stability constant of $\log K_{\text{Cu(II)L}} = 3.47 \pm 0.04$ (Figure S10 in the Supporting Information). Because MCL-3 does not contain any basic sites, its stability constants are invariant toward pH in aqueous solution. Slow-scan cyclic voltammetry studies of MCL-3 in the presence of Cu(II) showed a one-electron process at $E^f = 0.729 \text{ V}$ vs SHE (Figure 4B); however, with a peak separation of 100 mV even at scan rates as low as 2 mV/s (Figure S8 in the Supporting Information), the redox process cannot be considered reversible and is therefore not suitable for the derivation of a Cu(I) stability constant using eq 1. Instead, we performed spectrophotometric competition titrations with BCS, from which we derived a Cu(I) stability constant of 13.80 ± 0.04 (Figure S19 in the Supporting Information). Furthermore, we were able to corroborate this value by a competition titration with BCA, which yielded a stability constant of 13.78 ± 0.06 (Figure S20 in the Supporting Information). It is noteworthy that the Cu(I) affinity of MCL-3 is significantly higher than that of the parent macrocycle [16]aneS₄ ($\log K_{\text{Cu(I)L}} = 12.0$),²⁷ an observation that is likely due to the conformational preorganization of the free ligand. In fact, the 65-fold gain in Cu(I) affinity is very similar to the previously reported 49-fold increase in Ni(II) affinity upon methylation of [14]aneS₄ as mentioned above,²⁹ despite the different coordination preferences of Cu(I) and Ni(II).

Although the combination of competition titrations and thermodynamic cycles (eq 1) yielded a consistent set of cross-validated stability constants, the entire data set hinges on a single value, the standard reduction potential of the aqueous Cu(II/I) couple used in eq 1. When considering the spread of E_{aq}° values applied in the literature,^{20,28,42} the calculated Cu(I) stability constant may vary by up to 0.6 logarithmic units. To address this uncertainty, we sought to measure at least one of the stability constants by a method that is not dependent on the value of E_{aq}° . At present, there are only few ligands for which Cu(I) stability constants have been reported without the use of electrochemical methods. At first, we explored the utility of cyanide as competing ligand in spectrophotometric titrations with BCS; however, careful data analysis indicated the presence of heteroleptic ternary complexes, which prevented us from deriving a reliable stability constant. We next explored the suitability of acetonitrile, which not only may act as a competing ligand but also can stabilize Cu(I) toward disproportionation.⁴³ In view of the rather low stability constants of $\log \beta_1 = 2.63$, $\log \beta_2 = 4.02$, and $\log \beta_3 = 4.29$,¹⁴ acetonitrile cannot compete for Cu(I) binding with either BCS or BCA. We therefore tried to identify a spectrophotometrically active dibasic Cu(I) ligand whose apparent affinity could be adjusted over several orders of magnitude simply by changing the pH. At low pH, acetonitrile would be able to compete for Cu(I) binding, while at neutral pH the higher apparent affinity would match those of BCS or BCA.

With a reported Cu(I) stability constant of $\log K_{\text{Cu(I)L}} = 15.76$ and protonation constants of 3.26 ± 0.08 and 7.33 ± 0.15 , the dibasic pyridine derivative PEMEA (4) emerged as the most promising candidate.¹³ Slow-scan cyclic voltammetry measurements in the presence of Cu(II) revealed a quasi-reversible one-electron redox process with $E^f = 0.595 \text{ V}$ vs SHE and a peak separation of 68 mV (Figure S6 in the Supporting Information), thus allowing independent verification of the PEMEA–Cu(I) stability constant through a thermodynamic cycle. Spectrophotometric titration of PEMEA with Cu(II) at pH 5.0 gave rise to a new charge-transfer absorption band centered around 350 nm due to formation of the corresponding

1:1 complex. Nonlinear least-squares fitting over the spectral range of 300–450 nm provided a Cu(II) stability constant of $\log K_{\text{Cu(II)L}} = 7.85 \pm 0.08$ (Figure S13 in the Supporting Information). Combined with the electrochemical data, we calculated a Cu(I) stability constant of $\log K_{\text{Cu(I)L}} = 15.71 \pm 0.08$, which agrees well with the value reported by Rorabacher and co-workers (15.76).¹³ As anticipated, acetonitrile was able to compete for Cu(I) binding in the presence of PEMEA at low pH. On the basis of a molar ratio titration of PEMEA with $[\text{Cu(I)(CH}_3\text{CN)}_4]\text{PF}_6$ in aqueous buffer at pH 2.0 (14 mM HClO_4 , 100 μM sodium ascorbate, 3.9% CH_3CN , 0.1 M KClO_4 , 25 $^\circ\text{C}$), we determined a Cu(I) stability constant of $\log K_{\text{Cu(I)L}} = 15.75 \pm 0.02$ (Figure S21 in the Supporting Information). This value is in excellent agreement with the data obtained using eq 1 with $E_{\text{aq}}^\circ = 0.13$ V vs SHE. Finally, we used PEMEA to perform two additional competition titrations with MCL-1 (Figure S22 in the Supporting Information) and BCA (Figure S23 in the Supporting Information) to further cross-validate the above equilibrium network. On the basis of spectrophotometric titrations followed by nonlinear least-squares fitting, we determined Cu(I) stability constants of $\log \beta_2 = 17.63 \pm 0.05$ for BCA and $\log K_{\text{Cu(I)L}} = 16.33 \pm 0.07$ for MCL-1. Both results agree well with the values from the initial series of competition titrations (17.67 ± 0.03 and 16.33 ± 0.02 for BCA and MCL-1, respectively).

On the basis of all of the above data, we present in Table 4 the recommended values of the Cu(I) complex stability constants for the MCL series as well as for BCA, BCS, and PEMEA. All of the standard deviations are based on independent repeats with $n = 3$ or higher. When the uncertainty associated with the measurement of formal potentials is additionally taken into account, the listed stability constants are expected to be accurate within ± 0.1 logarithmic unit. In addition to the Cu(I) stability constant ($\log K_{\text{Cu(I)L}}$) for each ligand, we have also listed the corresponding conditional or apparent stability constant ($\log K_{\text{Cu(I)L}}'$) at neutral pH. Originally proposed by Schwarzenbach,⁴⁷ the apparent affinity takes into account the protonation equilibria of the ligand and metal ion at a given pH (see the Supporting Information). Because of competing protonation equilibria, the stability constants of ligands with basic donor atoms are strongly pH-dependent. For example, at neutral pH the apparent Cu(I) affinity of MCL-2 ($\text{p}K_{\text{a}} = 9.09$, $I = 0.1$) is lowered by 2 orders of magnitude, whereas the less basic ligand MCL-1 ($\text{p}K_{\text{a}} = 7.11$, $I = 0.1$) exhibits a much smaller change of 0.33 logarithmic units (Table 4). Because BCS and BCA form Cu(I) complexes with a 2:1 stoichiometry, their stability constants cannot be directly compared with those of the MCL series of ligands. For this reason, we have also listed in Table 4 the corresponding pCu values, which equate to $-\log [\text{Cu(I)}_{\text{aq}}]$ for a solution containing total ligand and Cu(I) concentrations of 10 and 1 μM , respectively, at pH 7.0 ($I = 0.1$).⁴⁸ Under these conditions, BCS and BCA exhibit similar chelating abilities compared to MCL-1 and MCL-3, respectively. Because of the 2:1 complex stoichiometry, the solution equilibrium depends not only on the ligand/Cu(I) ratio but also the total ligand concentration (Figure 6A). For example, if the BCS and Cu(I) concentrations are increased 10-fold, the pCu(I) value increases from 16.6 to 17.6.

On the basis of the apparent affinities at pH 7, the MCL series is suitable for buffering aqueous Cu(I) concentrations from 10^{-10} to 10^{-17} M, a range that is well-matched with the affinities of many Cu(I) metalloproteins.^{9,11} The three ligands




Figure 6. Dynamic range for buffering of free Cu(I)_{aq} with ligands exhibiting (A) 2:1 and (B) 1:1 complex stoichiometries. The fractional saturation of the corresponding ligand was calculated as a function of $-\log [\text{Cu(I)}_{\text{aq}}]$ for a buffer with constant total ligand concentration and varying $[\text{Cu(I)}]_{\text{total}}$ at pH 7.0 ($I = 0.1$).

exhibit complementary buffer ranges with almost equal spacing (Figure 6B). As the corresponding Cu(I) complexes are all crystalline and air-stable, Cu(I) buffer solutions can be readily prepared without the need of an additional Cu(I) source. Furthermore, the ligands and the corresponding Cu(I) complexes are optically transparent down to 300 nm (Figure S14 in the Supporting Information), permitting interference-free monitoring of Cu(I) binding to targets with an intrinsic spectroscopic signature, thus complementing the chromogenic reagents BCS and BCA.

The MCL series may also be applied for the calibration of Cu(I)-responsive fluorescent probes, the majority of which have been characterized via competition with low-affinity monodentate ligands amenable to heteroleptic complex formation. Since most fluorescent probes reported to date exhibit dissociation constants in the picomolar range,⁴⁹ MCL-2 would be well-suited as a reference standard. Compared with CH_3CN and two other low-affinity ligands recently proposed as quantitative probes for Cu(I),⁵⁰ MCL-2 forms a Cu(I) complex with improved redox stability and reduced reactivity toward dioxygen. The Cu(I) complex of MCL-2 would also be well-suited for the metalation of copper proteins, which are typically isolated in their apo forms upon overexpression.⁹ Similarly, the Cu(I) complexes of MCL-1 and MCL-3 offer superior redox stability compared with those of PEMEA, BCS, and BCA.

To illustrate the application of the MCL series as affinity standards, we describe in the last section the determination of a Cu(I)-protein complex stability constant based on fluorimetric titrations. For this purpose, we chose a tryptophan-containing protein whose affinity has been previously characterized by means of competition with a chromogenic ligand.

Determination of the Cu(I) Stability Constant of CusF.

As a component of the CusCFBA efflux complex,⁵¹ the periplasmic protein CusF plays an important role in the

detoxification of elevated levels of copper and silver in *E. coli*.⁵² Besides coordination to a Met₂His motif, the protein utilizes a tryptophan residue to stabilize Cu(I) via a strong cation- π interaction.^{6,7,53} In the apo protein, the tryptophan residue fluoresces at 343 nm, and the fluorescence is quenched upon binding of Cu(I), thus offering a convenient spectroscopic handle to monitor the degree of Cu(I) binding in an equilibrium titration with a competing ligand. On the basis of preliminary titrations, we found that MCL-3 matched best the Cu(I) affinity of CusF. As illustrated in Figure 7, gradual




Figure 7. Fluorescence titration of CusF (20 μ M) with MCL-3 at pH 7.0 (50 mM MOPS, 0.15 M NaCl, 25 $^{\circ}$ C). Inset: Change of the fluorescence intensity at 343 nm (excitation at 290 nm).

addition of MCL-3 to a solution of holo-CusF in MOPS buffer (pH 7.0) restored the tryptophan fluorescence. Nonlinear least-squares fitting of the emission response revealed a CusF-Cu(I) stability constant of $\log K_{\text{Cu(I)L}} = 14.29 \pm 0.11$.

This value is 0.5 logarithmic units higher compared to an earlier report in which the Cu(I) stability constant of CusF was determined relative to BCA [$K_{\text{d}}(\text{CusF}) \times \beta_2(\text{BCA}) = 7.3 \times 10^3$, or $\log K(\text{CusF}) = 13.8$ assuming $\log \beta_2(\text{BCA}) = 17.7$]⁶ but agrees well with the data from a more recent independent re-evaluation using the same competitor ligand [$\log K(\text{CusF}) = 14.3$ assuming $\log \beta_2(\text{BCA}) = 17.7$].⁷ Because of this discrepancy, we also re-examined the Cu(I) affinity of CusF by spectrophotometric equilibrium titrations using BCA as a competing ligand. Regardless of whether the Cu(I)-BCA complex was preformed and then titrated with apo-CusF (Figure S25 in the Supporting Information) or a mixture of apo-CusF and BCA was titrated with Cu(I) supplied from a stock solution in acetonitrile (Figure S26 in the Supporting Information), the stability constant converged at a value of $\log K_{\text{Cu(I)L}} = 14.21 \pm 0.03$, thus confirming within experimental error the above result from the competition titration with MCL-3.

CONCLUSIONS

To address the pressing need for robust affinity standards in Cu(I) biochemistry, we have created and characterized a series of three water-soluble ligands that form well-defined, colorless, and air-stable Cu(I) complexes in aqueous solution. All three ligands and their Cu(I) complexes can be isolated and recrystallized in multigram quantities, thus allowing for the preparation of precisely defined Cu(I) buffer solutions simply by mixing the free ligand and corresponding Cu(I) complex in the appropriate ratios. Grounded in an extensive network of equilibrium titrations and electrochemical measurements, we determined reliable acidity and Cu(I) stability constants for not only the colorless MCL series of ligands but also two

chromogenic reagents, bathocuproine disulfonate (BCS) and 2,2'-bicinechonic acid (BCA), which are frequently used for the determination of Cu(I) protein stability constants. Altogether, these ligands represent a cohesive set of optically transparent and chromogenic affinity standards that are capable of buffering Cu(I) concentrations from 10^{-10} to 10^{-18} M. Despite the high binding affinities, rapid equilibration was observed for all of the ligand competition experiments, presumably as a result of an associative Cu(I)-ligand exchange mechanism. The fast exchange kinetics also raises the possibility of conducting competition experiments by isothermal titration calorimetry,⁵⁴ which would allow the measurement of reliable Cu(I) binding affinities of proteins and ligands regardless of their spectroscopic signatures. Furthermore, the Cu(I) complex of the lowest-affinity ligand, MCL-2, may be used for the quantitative metalation of Cu(I)-binding proteins, thus offering an alternative Cu(I) source in place of the air- and moisture-sensitive tetrakis(acetonitrile)copper(I) salts. Conversely, the chelation of copper with high-affinity ligands might offer avenues for the treatment of diseases associated with copper overload such as Alzheimer's disease and Wilson's disease.⁵⁵ Most importantly, we anticipate that these reagents will facilitate the reliable measurement of Cu(I) binding affinities of metalloproteins and other biologically relevant Cu(I) ligands.

ASSOCIATED CONTENT

Supporting Information

Synthetic procedures, compound characterization, crystal structure data, and details of the determination of protonation constants and Cu(I) stability constants. This material is available free of charge via the Internet at <http://pubs.acs.org>.

AUTHOR INFORMATION

Corresponding Author

fahrni@chemistry.gatech.edu

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

We thank Prof. M. M. McEvoy for the generous gift of the CusF expression vector and K. Odegaard for assistance with the TXRF measurements. Financial support from the National Institutes of Health (R01GM067169) is gratefully acknowledged.

REFERENCES

- (1) Linder, M. C. *Biochemistry of Copper*; Springer: New York, 1991.
- (2) Kim, B.-E.; Nevitt, T.; Thiele, D. J. *Nat. Chem. Biol.* **2008**, *4*, 176.
- (3) Lutsenko, S. *Curr. Opin. Chem. Biol.* **2010**, *14*, 211.
- (4) Huffman, D. L.; O'Halloran, T. V. *J. Biol. Chem.* **2000**, *275*, 18611. Xiao, Z. G.; Wedd, A. G. *Chem. Commun.* **2002**, 588. Badarau, A.; Dennison, C. *Proc. Natl. Acad. Sci. U.S.A.* **2011**, *108*, 13007.
- (5) Niemiec, M. S.; Weise, C. F.; Wittung-Stafshede, P. *PLoS One* **2012**, *7*, No. e36102. Fu, Y.; Tsui, H.-C. T.; Bruce, K. E.; Sham, L.-T.; Higgins, K. A.; Lisher, J. P.; Kazmierczak, K. M.; Maroney, M. J.; Dann, C. E.; Winkler, M. E.; Giedroc, D. P. *Nat. Chem. Biol.* **2013**, *9*, 177.
- (6) Banci, L.; Bertini, I.; Ciofi-Baffoni, S.; Kozyreva, T.; Zovo, K.; Palumaa, P. *Nature* **2010**, *465*, 645.
- (7) Jensen, P. Y.; Bonander, N.; Møller, L. B.; Farver, O. *Biochim. Biophys. Acta* **1999**, *1434*, 103. Wernimont, A. K.; Yatsunyk, L. A.; Rosenzweig, A. C. *J. Biol. Chem.* **2004**, *279*, 12269. Kittleson, J. T.; Loftin, I. R.; Hausrath, A. C.; Engelhardt, K. P.; Rensing, C.; McEvoy, M. M. *Biochemistry* **2006**, *45*, 11096. Yatsunyk, L. A.; Rosenzweig, A. C. *J. Biol. Chem.* **2007**, *282*, 8622.

- (6) Xue, Y.; Davis, A. V.; Balakrishnan, G.; Stasser, J. P.; Staehlin, B. M.; Focia, P.; Spiro, T. G.; Penner-Hahn, J. E.; O'Halloran, T. V. *Nat. Chem. Biol.* **2008**, *4*, 107.
- (7) Loftin, I. R.; Blackburn, N. J.; McEvoy, M. M. *J. Biol. Inorg. Chem.* **2009**, *14*, 905.
- (8) Xiao, Z. G.; Wedd, A. G. *Nat. Prod. Rep.* **2010**, *27*, 768.
- (9) Xiao, Z.; Brose, J.; Schimo, S.; Ackland, S. M.; La Fontaine, S.; Wedd, A. G. *J. Biol. Chem.* **2011**, *286*, 11047.
- (10) Badarau, A.; Dennison, C. *J. Am. Chem. Soc.* **2011**, *133*, 2983.
- (11) Allen, S.; Badarau, A.; Dennison, C. *Biochemistry* **2012**, *51*, 1439.
- (12) Hansen, L. D.; Fellingham, G. W.; Russell, D. J. *Anal. Biochem.* **2011**, *409*, 220.
- (13) Ambundo, E. A.; Deydier, M. V.; Grall, A. J.; Aguera-Vega, N.; Dressel, L. T.; Cooper, T. H.; Heeg, M. J.; Ochrymowycz, L. A.; Rorabacher, D. B. *Inorg. Chem.* **1999**, *38*, 4233.
- (14) Kamau, P.; Jordan, R. B. *Inorg. Chem.* **2001**, *40*, 3879.
- (15) Damu, K. V.; Shaikjee, M. S.; Michael, J. P.; Howard, A. S.; Hancock, R. D. *Inorg. Chem.* **1986**, *25*, 3879.
- (16) Martell, A. E.; Motekaitis, R. J. *The Determination and Use of Stability Constants*; Wiley: New York, 1992.
- (17) Gans, P.; O'Sullivan, B. *Talanta* **2000**, *51*, 33.
- (18) *SPECFIT Global Analysis System*; Spectrum Software Associates: Marlborough, MA, 2001.
- (19) Gans, P.; Sabatini, A.; Vacca, A. *Talanta* **1996**, *43*, 1739.
- (20) Milazzo, G.; Caroli, S. *Tables of Standard Electrode Potentials*; Wiley: New York, 1978.
- (21) Koepf, H.-M.; Wendt, H.; Strehlow, H. Z. *Elektrochem.* **1960**, *64*, 483.
- (22) Yee, E. L.; Cave, R. J.; Guyer, K. L.; Tyma, P. D.; Weaver, M. J. *J. Am. Chem. Soc.* **1979**, *101*, 1131.
- (23) *APEX II*, version 2005; Bruker AXS, Inc.: Madison, WI, 2005.
- (24) *SAINTE*, version 6.45A; Bruker AXS, Inc.: Madison, WI, 2003.
- (25) Sheldrick, G. M. *Acta Crystallogr.* **2008**, *A64*, 112.
- (26) Loftin, I. R.; Franke, S.; Roberts, S. A.; Weichsel, A.; Héroux, A.; Montfort, W. R.; Rensing, C.; McEvoy, M. M. *Biochemistry* **2005**, *44*, 10533.
- (27) Bernardo, M. M.; Schroeder, R. R.; Rorabacher, D. B. *Inorg. Chem.* **1991**, *30*, 1241.
- (28) Bernardo, M. M.; Heeg, M. J.; Schroeder, R. R.; Ochrymowycz, L. A.; Rorabacher, D. B. *Inorg. Chem.* **1992**, *31*, 191.
- (29) Desper, J. M.; Gellman, S. H.; Wolf, R. E., Jr.; Cooper, S. R. *J. Am. Chem. Soc.* **1991**, *113*, 8663.
- (30) Pinho, S. P.; Macedo, E. A. *J. Chem. Eng. Data* **2005**, *50*, 29.
- (31) Morgan, M. T.; Bagchi, P.; Fahrni, C. J. *J. Am. Chem. Soc.* **2011**, *133*, 15906. Morgan, M. T.; Bagchi, P.; Fahrni, C. J. *Dalton Trans.* **2013**, *42*, 3240.
- (32) Buter, J.; Kellogg, R. M. *J. Org. Chem.* **1981**, *46*, 4481.
- (33) Blumenkemper, M.; Schröder, H.; Pape, T.; Hahn, F. E. *Inorg. Chim. Acta* **2011**, *366*, 76.
- (34) Blake, A. J.; Gould, R. O.; Halcrow, M. A.; Schroeder, M. *Acta Crystallogr.* **1993**, *B49*, 773.
- (35) Macrae, C. F.; Bruno, I. J.; Chisholm, J. A.; Edgington, P. R.; McCabe, P.; Pidcock, E.; Rodriguez-Monge, L.; Taylor, R.; van de Streek, J.; Wood, P. A. *J. Appl. Crystallogr.* **2008**, *41*, 466.
- (36) Brooks, N. R.; Blake, A. J.; Champness, N. R.; Cooke, P. A.; Hubberstey, P.; Proserpio, D. M.; Wilson, C.; Schröder, M. *J. Chem. Soc., Dalton Trans.* **2001**, 456.
- (37) Munakata, M.; Wu, L. P.; Yamamoto, M.; Kuroda-Sowa, T.; Maekawa, M. *J. Chem. Soc., Dalton Trans.* **1995**, 3215.
- (38) Martell, A. E.; Smith, R. M. *Critical Stability Constants*; Plenum Press: New York, 1974.
- (39) Cooper, T. H.; Mayer, M. J.; Leung, K. H.; Ochrymowycz, L. A.; Rorabacher, D. B. *Inorg. Chem.* **1992**, *31*, 3796.
- (40) Gershuns, A. L.; Koval, V. L. *Izv. Vyssh. Uchebn. Zaved., Khim. Khim. T.* **1975**, *18*, 1543.
- (41) Xiao, Z.; Donnelly, P. S.; Zimmermann, M.; Wedd, A. G. *Inorg. Chem.* **2008**, *47*, 4338.
- (42) Hawkins, C. J.; Perrin, D. D. *J. Chem. Soc.* **1963**, 2996.
- (43) Balakrishnan, K. P.; Kaden, T. A.; Siegfried, L.; Zuberbühler, A. D. *Helv. Chim. Acta* **1984**, *67*, 1060.
- (44) Xiao, Z.; Loughlin, F.; George, G. N.; Howlett, G. J.; Wedd, A. G. *J. Am. Chem. Soc.* **2004**, *126*, 3081.
- (45) De Pascali, S. A.; Migoni, D.; Papadia, P.; Muscella, A.; Marsigliante, S.; Ciccarese, A.; Fanizzi, F. P. *Dalton Trans.* **2006**, 5077.
- (46) Yao, S.; Cherny, R. A.; Bush, A. I.; Masters, C. L.; Barnham, K. J. *J. Pept. Sci.* **2004**, *10*, 210.
- (47) Schwarzenbach, G. *Complexometric Titrations*; Interscience: New York, 1957.
- (48) Harris, W. H.; Raymond, K. N. *J. Am. Chem. Soc.* **1979**, *101*, 6534.
- (49) Fahrni, C. J. *Curr. Opin. Chem. Biol.* **2013**, *17*, 656. Morgan, M. T.; Bagchi, P.; Fahrni, C. J. *Fluorescent Probes for Monovalent Copper*. In *Metals in Cells*; Culotta, V., Scott, R. S., Eds.; Wiley: Chichester, U.K., 2013; pp 65–83. Hyman, L. M.; Franz, K. J. *Coord. Chem. Rev.* **2012**, *256*, 2333.
- (50) Xiao, Z.; Gottschlich, L.; van der Meulen, R.; Udagedara, S. R.; Wedd, A. G. *Metalomics* **2013**, *5*, 501.
- (51) Franke, S.; Grass, G.; Rensing, C.; Nies, D. H. *J. Bacteriol.* **2003**, *185*, 3804.
- (52) Mealman, T. D.; Blackburn, N. J.; McEvoy, M. M. *Curr. Top. Membr.* **2012**, *69*, 163.
- (53) Chakravorty, D. K.; Wang, B.; Ucisik, M. N.; Merz, K. M. *J. Am. Chem. Soc.* **2011**, *133*, 19330.
- (54) Trapaidze, A.; Hureau, C.; Bal, W.; Winterhalter, M.; Faller, P. J. *Biol. Inorg. Chem.* **2012**, *17*, 37.
- (55) Delangle, P.; Mintz, E. *Dalton Trans.* **2012**, *41*, 6359. Pujol, A. M.; Cuillel, M.; Jullien, A.-S.; Lebrun, C.; Cassio, D.; Mintz, E.; Gateau, C.; Delangle, P. *Angew. Chem., Int. Ed. Engl.* **2012**, *51*, 7445.