

Silvio Th. Mariani, General Manager, Futehally Chemicals Ltd., Karachi, Pakistan, Chemical Engineering, Federal Technical Inst., Zurich, Switzerland.

Dean Ralph Meurer, Laboratory Technician, Honeymead Products Co., Mankato, Minn.

James Albert Miller, Research Scientist, Wyeth Laboratories, Inc., Philadelphia, Pennsylvania.

Donald N. Mitcham, Physicist, Spectroscopy Group S. Utiliz. Res. & Dev. Div., USDA, New Orleans, Louisiana.

George Arthur Muck, Research Chemist, Dean Foods Co., Rockford, Illinois.

Arthur J. Murphy, Surfactant Techn. Service and Sales, Witco Chem. Co., Chicago, Ill.

Bryan William Nichols, Biochemist, Unilever Research Laboratories, Sharnbrook, Bedfordshire, England.

Homer Clyde Odom, Jr., Research Chemist, Pan American Tung Res. & Dev. League, Hattiesburg, Mississippi.

Vasanth Madhav Pai, Works Manager (Soap) Godrej Soaps Private Ltd., Bombay, India.

Wilbur A. Parker, Research Chemist, Corn Products Co., Bayonne, New Jersey.

Richard H. Plantholt, Research Chemist, Emery Industries, Inc., Cincinnati, Ohio.

Richard G. Powell, Research Chemist, N. Utiliz. Res. & Dev. Div., ARS, USDA, Peoria, Ill.

Henry Rakoff, Associate Professor—Chemistry Department, Texas A&M University, College Station, Texas.

Humber Joseph Ranauto, Group Leader—Product. Dev. Dept., Atlas Chemical Industries, Wilmington, Delaware.

Albert M. Reynaud, Superintendent of Production & Control, Gulf Soap Corp., Arabi, Louisiana.

Robert Allen Reynolds, Manager, G. H. T. Laboratories, Brawley, California.

Allen Dinwoody Robinson, Professor of Chemistry, University of Manitoba, Winnipeg, Manitoba, Canada.

Antoni Rogozinski, Plant Lab. & Quality Control Chf. Techn., Patrick Cudahy, Inc., Cudahy, Wis.

Clyde B. Rowntree, Chief Chem., Wellman Combing Co., Johnsonville, South Carolina.

Antoni Rutkowski, Assistant Director, Head of Food Technology, Institute of General Chemistry, Warszawa, Poland.

Francesco Sanguinetti, Associate Professor, University of Rome, Pisa, Italy.

Cornelius Theodorus Schnetler, Chemist, H. Lewis & Co., Ltd., Transvaal, S. Africa.

Joseph Harry Schwartz, Research Chemist, E. Regional Res. Lab., USDA, Philadelphia, Pennsylvania.

Fred William Shapiro, Lab. Technician, Best Foods Div., Corn Products, Brooklyn, New York.

David R. Shenkenberg, Techn. Director, Jewett & Sherman Co., Milwaukee, Wisconsin.

Marion Edmonds Smith, Neurochemist, Veterans Administration Hospital, Instructor, Stanford University, Palo Alto, California.

Ronald B. Stapley, Chemist, Carnation Research Laboratories, Van Nuys, California.

Victor V. Studer, Manager Food and Res. Section, Thomas J. Lipton, Inc., Englewood Cliffs, New Jersey.

Walter R. Trent, Chemical Director, John T. Stanley Co., Inc., Menlo Park, New Jersey.

Lloyd Woodbury Trevoy, Research Chemist, Chemcell Limited, Edmonton, Alberta, Canada.

Ph. J. Van Der Kelen, Laboratoire Ph. J. Van Der Kelen, Bruxelles, Belgium.

James Velasco, Research Chemist, USDA, Washington, D. C.

Frank Philip Wetterau, Research Chemist, Atlas Chem. Industries, Inc., Wilmington, Delaware.

Dr. Henry Karl Wren, Director of Research, Baltimore Paint & Chemical Corp., Baltimore, Maryland.

#### Active Junior

Roger C. Crum, Jr., Research Assistant, Animal Science Dept., University of Florida, Gainesville, Florida.

Arthur W. Kruski, Research Assistant, Burnside's Research Laboratory, Urbana, Ill.

Benny K. H. Tjho, Graduate Student, MIT, Cambridge, Mass.

Lee-Shin Tsai, Graduate Student, University of California, Davis, Calif.

#### Individual Associates

William R. Barnes, Sales Engineer, Manufacturers Engineering and Equipment Corp., Houston, Texas.

John R. Barone, Regional Sales Manager, Industrial Filter & Pump Manufacturing Co., Cicero, Ill.

Lloyd Roy Custer, Jr., Chemist, Foods Division, Anderson, Clayton & Co., Sherman, Texas.

William R. Cutts, Sr., Laboratory Technician, Honeymead Products Co., Mankato, Minnesota.

Lawrence Deleve Johns, Chemist, Best Food Company, Bayonne, New Jersey.

Andrew M. Lubienski, Technical Sales, Miranol Chem. Co., Irvington, New Jersey.

Katsunori Mukai, Associate Professor of Nutrition, University of Osaka Prefecture, Sakai, Japan.

C. R. Rathbone, General Manager, Ranchers Cotton Oil, Fresno, California.

Richard Crawford Slover, Marketing Adviser, Humble Oil & Refining Co., Houston, Texas.

Max Edward Westby, Department Foreman, HumKo Products, Chem. Div., Memphis, Tennessee.

#### Corporation Associate

Intoco, Inc., Basking Ridge, New Jersey.

### • Referee Applications

SECOND NOTICE. Ronald Mayo Fox of Texas Testing Laboratories, Inc., P. O. Box 2144, Dallas, Texas 75221 has applied for a Referee Certificate on Oil Cake and Meal, Cottonseed and Soybean Oils and Cottonseed. Interested parties wishing to comment on this certification should communicate with the Chairman of the Examination Board. Please write to R. T. Doughtie, Jr., Chairman of the Examination Board, P. O. Box 17469, Memphis, Tenn. 38117.

SECOND NOTICE. Hans J. Schulze of New Jersey Feed Laboratory, 910 Pennsylvania Avenue, Trenton, New Jersey has applied for a Referee Certificate on Oil Cake and Meal. Interested parties wishing to comment on this certification should communicate with the Chairman of the Examination Board. Please write to R. T. Doughtie, Jr. Chairman of the Examination Board, P. O. Box 17469, Memphis, Tennessee 38117.

FIRST NOTICE: Charles R. Norris of Barrow-Agee Laboratories, Inc.; P.O. Box 858, Shreveport, Louisiana, 71102 has applied for a Referee Certificate on Cottonseed, Oil Cake and Meal, Protein Concentrates, Cottonseed Oil and Soybean Oil. Interested parties wishing to comment on this certification should communicate with the Chairman of the Examination Board. Please write to R. T. Doughtie, Jr., Chairman of the Examination Board, P.O. Box 17469, Memphis, Tennessee 38117.

**LABORATORY SERVICES**

Analytical Services  
Organic Synthesis  
Testing & Evaluation  
Surface Chemistry  
Physical Chemistry

**FOSTER D. SNELL, INC.**

A SUBSIDIARY OF BOOZ-ALLEN APPLIED RESEARCH, INC.



**FDS**

Member - American Council of  
Independent Laboratories

**CHEMISTS  
ENGINEERS  
BIOLOGISTS**

29 West 15th St.  
N. Y. N. Y. 10011  
(212) 924-8800

**LAW & COMPANY  
CHEMISTS**

Consulting and Analytical  
Atlanta, Ga.    Montgomery, Ala.    Wilmington, N.C.

(Continued from page 309A)

Urbana). *J. Dairy Science* **48**, 381-390 (1965). n-Butyrate-1-C<sup>14</sup> was injected intravenously into four adult goats. Observations were made on the subsequent incorporation of C<sup>14</sup> into liver and muscle glycogen and into various organic acids of blood, taken either from the jugular vein or simultaneously from both the carotid artery and ruminal vein. Though several acids were labeled with C<sup>14</sup>, β-hydroxybutyrate exhibited the highest specific activity. Ruminal vein blood contained over twice as much β-hydroxybutyrate as did blood from the carotid artery. The concentration of n-butyrate in ruminal vein blood was frequently lower than that of iso-butyrate, even though the ratio of these two acids in the rumen fluid was approximately 7:1. This suggests that β-hydroxybutyrate is a normal intermediate in the metabolism of n-butyrate absorbed from the rumen and that the rumen epithelium is a major site of this conversion. The lactate of ruminal vein blood was slightly greater in concentration but several times higher in specific activity than lactate of arterial blood. Glucose from both liver and muscle glycogen was labeled with C<sup>14</sup> predominantly in Carbons 3 and 4, the ratio of labeling between these two carbons being approximately 1:1.

INHERITANCE OF LINOLEIC AND OLEIC ACIDS IN MAIZE. C. G. Poneleit and D. E. Alexander (Dept. of Agronomy, Univ. of Illinois, Urbana, Ill.). *Science* **147**, 1585-1586 (1965). Gas-liquid chromatographic analysis of methyl esters of fatty acids of individual maize seeds of parental and segregating populations suggests that desaturation at the Δ<sup>12-13</sup> position in oleic acid is under simple Mendelian control. High linoleic acid content is recessive to low.

**Gas Chromatograph NOISE  
a problem?**



- Reduce Noise
- Increase Sensitivity


**Cause**

- ... Instability in balance circuit
- ... Limit of Balance Resistors

**Remedy**

Install the Gyra E-307  
Electrometer . . . increases  
sensitivity beyond  
your expectations

Send for Bulletin 791



**Gyra ELECTRONICS CORPORATION**

Phone: Area 312 354-4644.

P. O. Box 184-W-6

LaGrange, Illinois 60526

THE EFFECT OF A MEAT-ENRICHED DIET ON THE DEVELOPMENT OF EXPERIMENTAL ATHEROSCLEROSIS IN RABBITS. J. Polcak, F. Melichar, D. Sevelova, I. Dvorak and M. Skalova (Second Med. Clinic, Med. Faculty, J. E. Purkyne Univ., Brno, Czechoslovakia). *J. Atheroscler. Res.* **5**, 174-180 (1965). The effect of a standard diet enriched with 20 g meat on experimental cholesterol-induced atherosclerosis in rabbits was followed-up in a number of experiments of 4 and 8 months' duration. The serum-levels of cholesterol, phospholipids and total lipids and the cholesterol content of the tissues were significantly lower when given a meat-enriched standard diet than with a standard diet alone. Also the atherosclerotic changes in the aorta were less marked. The authors conclude that addition of meat to the diet has an inhibiting effect on the development of experimental atherosclerosis in rabbits.

FATTY ACIDS IN BLUE-GREEN ALGAL MAT COMMUNITIES. P. L. Parker and R. F. Leo (Inst. of Marine Sci., Univ. of Texas, Port Aransas). *Science* **148**, 373-374 (1965). The concentrations of the major fatty acids were determined for the varied layers of an algal mat community. The living mat contains substantial amounts of the unsaturated acids, while the underlying layers become progressively depleted in unsaturated molecules. A similar progressive increase in the ratio of saturated to unsaturated acids was detected in a sediment core from a hypersaline lagoon.

DIETARY EFFECT OF MAGNESIUM ON CHOLESTEROL-INDUCED ATHEROSCLEROSIS OF RABBITS. M. Nakamura, S. Torii, M. Hiramatsu, J. Hirano, A. Sumiyoshi and K. Tanaka (Res. Inst. of Angiocardiology and Dept. of Pathol., Kyushu Univ. Med. School, Fukuoka, Japan). *J. Atheroscler. Res.* **5**, 145-158 (1965). The dietary effect of various amounts of magnesium on cholesterol-induced atherosclerosis of rabbits was investigated. There was a significant enhancement of the lipid deposition in the aorta of rabbits fed a magnesium-deficient diet containing cholesterol. However, this study failed to demonstrate any retarding effect from the feeding of magnesium excess diets on aorta lipid deposition.

FATTY LIVERS OF WEANLING RATS FED A LOW PROTEIN THREONINE-DEFICIENT DIET. I. EFFECT OF VARIOUS DIET FATS. L. Morris, D. Arata and D. C. Cederquist (Dept. of Foods and Nutr., Michigan State Univ., East Lansing, Mich.). *J. Nutr.* **85**, 362-366 (1965). Weanling albino rats were fed a diet containing 9% casein and 30% fat with and without supplements of threonine. The fat source in the diet was provided by corn oil, olive oil, cottonseed oil, hydrogenated vegetable oil, or corn oil hydrogenated to an iodine value of 74. Animals were maintained with their respective diets for 2 to 4 weeks. Feeding weanling rats a threonine-deficient diet containing 30% corn oil resulted in the appearance of fatty livers. Replacing the corn oil in this diet with either cottonseed oil or hydrogenated vegetable oil caused a significant reduction in liver fat concentration. Substituting olive oil for corn oil slightly increased liver fat levels. In every instance, the addition of threonine lowered liver fat levels. When hydrogenated corn oil was substituted for corn oil in threonine-deficient diets, liver fat concentrations decreased markedly after 2 weeks; this effect persisted after 4 weeks. Since all diets within a series were isocaloric and since no significant differences in food intake or growth were observed among any of the deficient groups, the "protective" action of some diet fats is apparently not mediated through a more equitable balance between the amino acid and calorie more.

COMPARISONS OF CASEIN AND SOY PROTEINS UPON MINERAL BALANCE AND VITAMIN D<sub>2</sub> REQUIREMENT OF THE BABY PIG. E. R. Miller, D. E. Ullrey, C. L. Zutaut, J. A. Hoefler (Depts. of Animal Husbandry and Biochem., Michigan State Univ., East Lansing, Mich.). *J. Nutr.* **85**, 347-354 (1965). Comparisons were made of the effects of dietary protein source (casein versus isolated soy protein) upon growth, serum mineral level, skeletal development, mineral balance and vitamin D<sub>2</sub> requirement of the baby pig. The data indicate that the vitamin D<sub>2</sub> requirement of the baby pig receiving purified diets containing isolated soybean protein is greater than the 100 IU/kg requirement level of casein diets and may be several times this level depending upon the criteria selected, the level of isolated soybean protein in the diet and the method of isolation of the soybean protein.

LIPID CLASSES OF BOVINE SPERMATOZOA. L. D. Miller and D. T. Mayer (Dept. of Agricultural Chem.) and C. P. Merilan (Dept. of Dairy Husbandry, Univ. of Missouri, Columbia). *J. Dairy Sci.* **48**, 395-397 (1965). A modified silicic acid technique was developed for separation of the lipids of bovine

(Continued on page 320A)

(Continued from page 320A)

spermatozoa into seven distinct classes: saturated hydrocarbons, cholesterol esters, free cholesterol, triglycerides, diglycerides, monoglycerides, and phospholipids. The average lipid content of lyophilized spermatozoa was 11.05%. Monoglycerides and diglycerides were found in exceptionally large quantities, but no unsaturated hydrocarbons of free fatty acids could be detected.

EFFECT OF DIETARY CHOLESTEROL UPON SERUM LIPIDS IN RURAL GUATEMALAN INDIAN CHILDREN. J. Mendez (Inst. of Nutr. of Central America and Panama (INCAP), Guatemala, C. A.). *Am. J. of Clin. Nutr.* 16, 304-308 (1965). School children from a rural Indian community in the highlands of Guatemala were divided into three groups. Group 1 (the egg group) received 2 boiled eggs and 2 glasses of Incaparina daily. The cholesterol content of this supplement was 600 mg. Group 2 (the cholesterol group) received 600 mg. of crystalline cholesterol daily, plus 2 glasses of Incaparina containing 15 ml. of cottonseed oil each. Group 3 (the control group) received only the Incaparina with the same amount of oil. A significant increase in serum cholesterol levels was observed in group 1 at the end of four weeks; no increase was observed in the other groups. Although the increase in serum cholesterol levels observed in group 1 was significant, the increase was small, indicating that probably this Indian group has an intrinsically low responsiveness to dietary manipulations.

EFFECT OF PRENATAL DIET ON SERUM CHOLESTERYL ESTER FATTY ACIDS IN NEWBORN AND ADULT RATS. A. Lopez-Santolino, O. N. Miller, and J. E. Muldrey (Dept. of Biochem., Tulane Univ. School of Med., New Orleans, La.). *Proc. Soc. Exp. Biol. Med.* 118, 829-834 (1965). Experiments were conducted to study the effect of prenatal diet on the serum CEFA pattern of newborn rats and their mothers. Evidence was obtained confirming the existence of a typical "fetal" CEFA pattern in the newborn rats characterized by a high oleate: linoleate (O/L) ratio (1.5 to 5.0), and of a typical "adult" CEFA pattern in the mother characterized by an O/L ratio of less than unity (0.2 to 0.6). These characteristic "adult" and "fetal" CEFA patterns were found to be maintained in spite

of moderate alterations in the fatty acid composition of the prenatal diets, but could not be maintained in the face of the exaggerated imbalance of diets supplying only single fatty acids. Based on the data obtained, it was postulated that homeostatic mechanisms exist which function so as to maintain the characteristic adult and fetal CEFA patterns in the face of stresses consisting of variations in the dietary fatty acid composition.

EFFECT OF DIETARY FAT ON PLASMA AND LIVER LIPIDS OF PROPYLTHIOURACIL-TREATED RATS. Ching-Tong Liu (Dept. of Physiology and Biophys., Univ. of Tenn., Med. Units, Memphis, Tenn.). *J. Nutr.* 85, 426-428 (1965). The administration of 0.03% propylthiouracil to rats resulted in an increase in plasma cholesterol and phospholipid levels. No change was observed in liver cholesterol and total lipids. The induced hypercholesterolemia was not lowered by feeding a supplement of 20% cottonseed oil. However, further increased concentrations of plasma cholesterol and phospholipid were observed in the hypothyroid rats maintained with a diet containing 20% hydrogenated coconut oil. In the normal animals, the plasma cholesterol concentrations were augmented by the addition of dietary fat. The liver cholesterol and total lipids increased in all animals when their diets were mixed with unsaturated fatty acids.

A COMPARISON OF THE DISPOSITION OF INTRAVENOUSLY-INJECTED, ALBUMIN-BOUND STEARIC ACID-1-C<sup>14</sup> AND PALMITIC ACID-1-C<sup>14</sup> IN THE RAT. P. Kapiloff, W. J. Lossow, and I. L. Chaikoff (Dept. of Physiol., Univ. of Calif., Berkeley). *Proc. Soc. Exp. Biol. Med.*, 118, 800-803 (1965). Stearic acid-1-C<sup>14</sup> or palmitic acid-1-C<sup>14</sup> was injected intravenously into rats in the form of albumin-bound free fatty acids. At various intervals thereafter livers, hearts and samples of adipose tissue, skeletal muscle and plasma were taken for determination of total C<sup>14</sup> and its distribution among glycerides, phospholipids, cholesterol esters and free fatty acids. At all intervals the ratio (C<sup>14</sup> in phospholipid/C<sup>14</sup> in glycerides) was greater after injection of C<sup>14</sup>-labeled stearic acid than after that of labeled palmitic acid. This indicates that preferential incorporation of stearic acid into phospholipids is not limited to incorporation during absorption from the intestines into lymph.

## ARMSTRONG SCRAPED SHELL CRYSTALLIZERS

FOR: Low Temperature Crystallizing, Crystallizing Organics or Viscous Fluids, Cooling Viscous Fluids up to 80,000 cp., Materials with High Boiling Point Rise

WE WILL CUSTOM FIT OUR EQUIPMENT TO YOUR PROCESS CONDITIONS

### TYPICAL FLUIDS

FATTY ACIDS  
ORGANIC INTERMEDIATES  
PARA-XYLENE  
PARADICHLOROBENZENE

Variety of Shell Sizes & Internal Designs

Pilot Units for Rent

Process Know How & Refrigeration Experience Available

Contact:


**RICHARD M. ARMSTRONG  
COMPANY**

Box 566-J

West Chester, Penna. 19380




2 Stands of 12 Shells Each, 6" Inner Pipe, All Steel, Drive Cabinet Removed. De-waxing Lube Oils.



## -LAYER CHROMATOGRAPHY

- Plexiglas Applicator with variable adjustments, \$18.00. One of the finest instruments on the market at the lowest price anywhere.
- Introductory Kit for TLC . . . includes applicator, plates, adsorbents . . . all necessary materials for \$29.50.
- ADSORBOSIL Adsorbents, Silica Gels and Porous Glass. Ultra-high purity at very reasonable prices.
- Wide varieties of accessories and special apparatus . . . Write for our free catalog and GAS-CHROM Newsletter.



**APPLIED SCIENCE LABORATORIES, Inc.**  
P.O. Box 140, State College 3, Penna.  
phone: 814 - 238 - 2406

**RETINOYL BETA-GLUCURONIC ACID: A MAJOR METABOLITE OF VITAMIN A IN RAT BILE.** P. E. Dumagin, Jr., E. H. Meadows, Jr., J. A. Olson (Dept. of Biochem., Univ. of Florida College of Med., Gainesville). *Science* 148, 86-87 (1965). The major metabolite in rat bile of injected  $C^{14}$ -retinoic acid was purified by ion-exchange and silicic acid chromatography; it has the spectrum of methyl retinoate, releases retinoic acid upon basic hydrolysis or by treatment with  $\beta$ -glucuronidase, and contains glucuronic acid. The metabolite was characterized by treatment with diazomethane followed by hexamethyldisilazane, or with periodate followed by semicarbazide, and the products were chromatographed. The metabolite has been tentatively identified as retinoyl  $\beta$ -glucuronide.

**INCORPORATION OF PLASMA CHOLESTEROL-4- $C^{14}$  INTO EGG YOLK CHOLESTEROL.** W. E. Connor, J. W. Osborne and W. L. Marion (Dept. of Internal Medicine and Radiation Research Lab., Univ. of Iowa, College of Medicine, Iowa City, Iowa). *Proc. Soc. Exp. Biol. Med.* 118, 710-713 (1965). These data indicate that the laying hen transfers cholesterol-4- $C^{14}$  from the blood across the ovarian membranes into developing yolks. Eggs laid after administration of cholesterol-4- $C^{14}$  intravenously contained up to 45.8% of the administered isotope confined completely to the cholesterol fraction of the yolk. Many individual eggs contained from 5-10  $\mu$ c of cholesterol-4- $C^{14}$ . The specific radioactivity of yolk cholesterol always had a well defined temporal relationship to the specific radioactivity of cholesterol in the blood. While exact quantitative data are not yet available, the results of these experiments suggest that most, if not all, of the cholesterol in the egg yolk originates from the blood.

**RELATIONSHIP BETWEEN CHOLESTEROL AND VITAMIN A METABOLISM IN RATS FED AT DIFFERENT LEVELS OF VITAMIN A.** S. V. Bring, C. A. Ricard, and M. V. Zaehring (Home Econ. Research, College of Agr., Univ. of Idaho, Moscow, Idaho). *J. Nutr.* 85, 400-406 (1965). The relationship between cholesterol and vitamin A metabolism was studied in 3 replications using a total of 225 weanling male rats. Following vitamin A depletion, each animal was assigned at random to an experimental group, receiving one treatment from each of the following 3 variables: 0.0 or 0.5% cholesterol mixed in the basal purified diet; 10, 150 or 1000 IU vitamin A acetate daily administered orally; and fed 7, 18 or 28 days. Increasing the dietary vitamin A decreased the serum and liver total cholesterol concentrations when cholesterol was fed. The rats given 10 IU vitamin A daily had significantly higher ( $P < 0.01$ )

liver total cholesterol values than those given either 150 or 1000 IU. Cholesterol-fed rats given 10 IU also had significantly higher ( $P < 0.01$ ) mean serum cholesterol values than those given 1000 IU. Dietary cholesterol significantly lowered ( $P < 0.05$ ) mean liver vitamin A concentrations on a dry weight basis. Liver vitamin A and free cholesterol were found to be positively correlated ( $P < 0.01$ ) on either a percentage or per liver basis.

**PARALYSIS INDUCED BY FEEDING SYNTHETIC GLYCERIDES TO CHICKS.** S. Brambila and F. W. Hill (Dept. of Poultry Husbandry, Univ. of Calif., Davis). *Proc. Soc. Exp. Biol. Med.* 118, 845-847 (1965). Feeding mixed triglycerides synthesized from soybean oil fatty acids as the sole non-protein energy source is a simplified diet produced severe growth retardation, incoordination, and a crippling leg paralysis in young chicks. The possibility that these effects may be due to the lack of an unidentified essential nutrient is discussed.

**LIPOPROTEIN IMMUNIZATION AND INDUCED ATHEROSCLEROSIS IN RABBITS.** J. M. Bailey and R. Tomar (Dept. Biochem., School of Med., George Washington Univ., Washington, D. C.). *J. Atheroscler. Res.* 5, 203-214 (1965). Groups of rabbits were immunized with  $\beta$ -lipoproteins isolated from serum of normal and cholesterol-fed chickens. Precipitin titers in excess of 5000:1 were developed against antigen protein from cholesterol-fed birds. Plasma lipid levels increased in immunized animals and increase paralleled development of precipitin titers. Rabbit serum was analyzed by chromatography on DEAE cellulose ion exchange resin. Cholesterol in normal serum was found almost exclusively in proteins having the electrophoretic characteristics of  $\alpha$ -globulins. Following immunization, significant quantities of cholesterol were found in  $\gamma$ -globulins in rabbits on both normal and on cholesterol-supplemented diets. Immunized and control animals were fed a diet supplemented with 2% cholesterol for 12 weeks. Development of atherosclerosis in thoracic aorta was not significantly different in normal and immunized rabbits. Lipid levels in serum and tissues also were not significantly reduced by the prior immunization procedure. When immunization was carried out subsequent to cholesterol feeding, intensity of atherosclerotic plaques did not decrease in 6 weeks following transfer to normal diet whereas serum lipids returned to normal at same rate in control and immunized animals during this period.

**FATTY ACID AMIDES OF ETHANOLAMIDE IN MAMMALIAN TISSUES.** N. R. Bachur, Karel Masek, K. L. Melmon and S. Udenfriend

(Nat'l Heart Institute, Nat'l Inst. of Health, Bethesda, Maryland). *J. Biol. Chem.* 240, 1019-24 (1965). A method for the isolation, purification, and quantitative assay of fatty acid ethanolamides from natural sources is described. In rats and guinea pigs from which food has been withheld, the amide is found in brain, liver, and skeletal muscle, the level being highest and most constant in brain. Verification of the structure of the isolated palmitoylethanolamide was obtained through gas-liquid chromatography, infrared spectroscopy, and hydrolysis and analysis of the component fatty acid and amine. Evidence was presented to rule out the artificial production of palmitoylethanolamide.

EFFECT OF SIMPLE AND COMPLEX CARBOHYDRATES UPON TOTAL LIPIDS, NONPHOSPHOLIPIDS, AND DIFFERENT FRACTIONS OF PHOSPHOLIPIDS OF SERUM IN YOUNG MEN AND WOMEN. M. A. Antar and M. A. Ohlson (Dept. of Internal Medicine, State Univ. of Iowa, Iowa City, Iowa). *J. Nutr.* 85, 329-337 (1965). Eight young, healthy persons, were fed experimental diets for 4 dietary periods of 4 weeks each. Total fats, proteins, and carbohydrates constituted 40, 16 and 44% of total calories of the basic diet, respectively. The ratio of complex to simple carbohydrates was 1:4 in periods 1 and 3 and this was reversed in periods 2 and 4. Serum total lipids, nonphospholipids, and different fractions of phospholipids (ethanolamine phosphatides, inositides, lecithins, lysolecithins, and sphingomyelins) were determined using silicic acid column chromatography. Serum total lipids, phospholipids, and nonphospholipids were found to be significantly reduced with the high cereal diet and increased with the high sugar diet when the total calories and fats were held constant.

EVIDENCE FOR A CONTRIBUTION BY THE INTESTINAL WALL TO THE SERUM CHOLESTEROL OF THE RAT. C. A. Lindsey, Jr., and J. D. Wilson (Dept. of Internal Medicine, Univ. of Texas Southwestern Med. School, Dallas, Texas). *J. Lipid Res.* 6, 173-181 (1965). Rats were fed cholesterol in order to block hepatic cholesterol synthesis, and their intestinal lymph ducts were cannulated. Experiments with these rats showed that cholesterol synthesized in the intestinal wall enters into the circulating cholesterol pool. The quantitative significance of this source of serum cholesterol has not been established.

EFFECT OF VERY LOW-DENSITY LIPOPROTEINS ON LIPID TRANSFER IN INCUBATED SERUM. A. V. Nichols and L. Smith (Donner Lab., Lawrence Radiation Lab., Univ. of Calif., Berkeley, Calif.). *J. Lipid Res.* 6, 206-210 (1965). Incubation of human serum solutions low in glyceride content for 16 hr resulted in a net increase of cholesterol esters in all three major human serum lipoprotein classes, namely the very low-density lipoproteins of d 1.006 (VLDL), low-density lipoproteins of d 1.006-1.063 (LDL) and high-density lipoproteins of d 1.063-1.21 (HDL). Transfer of glycerides from the VLDL to the LDL and HDL was also observed during incubation. Reductions in lipoprotein unesterified cholesterol and phospholipids occurred during incubation and indicated the presence of serum fatty acid transferase activity in these solutions. When serum was incubated in the presence of high concentrations of VLDL lipoproteins, the increase in cholesterol esters was found only in the VLDL fraction. The VLDL apparently took up newly-formed cholesterol esters produced by serum fatty acid transferase activity, and accepted some cholesterol esters initially bound to the LDL and HDL. When the transferase was inhibited with sodium *p*-hydroxymercuribenzoate, some of the cholesterol esters initially bound to the LDL and HDL were still transferred to the VLDL. The HDL class was then the principal contributor of the cholesterol esters. These observations indicate that reciprocal transfer of cholesterol esters for glycerides in human serum lipoproteins can occur.

LIPIDS OF THE SPLEEN IN GAUCHER'S DISEASE. W. D. Suomi and B. W. Agranoff (Dept. of Biol. Chem. and Mental Health Res. Inst., Univ. of Mich., Ann Arbor, Mich.). *J. Lipid Res.* 6, 211-219 (1965). Thin-layer chromatography (TLC) was used to analyze lipids of eight spleens of patients with Gaucher's disease. Four non-Gaucher spleens were also analyzed. Phospholipid concentrations are not increased in Gaucher spleens, while several classes of neutral lipids are moderately increased.

ESTABLISHMENT OF ERYTHRO CONFIGURATION OF CERAMIDES FROM BEEF BRAIN AND CHICKEN LIVER. V. Groom and M. Sribney (Depts. of Biochem. and Psychiatry, Yale Univ. School of Medicine, New Haven, Connecticut). *J. Lipid Res.* 6, 220-221 (1965). The sphingosine moiety of ceramides isolated from beef brain and chicken liver has been characterized as having the erythro stereochemical configuration. This is in agreement with the configuration of sphingosine as found in cerebrosides and sphingomyelin.

IN VITRO ESTERIFICATION OF CHOLESTEROL BY PANCREATIC JUICE AND BY ACETONE POWDER EXTRACTS OF SMALL INTESTINE: THE EFFECT OF UNSATURATED FATTY ACID UPON ESTERIFICATION WITH SATURATED FATTY ACIDS. S. N. Shah, W. J. Lossow, and I. L. Chaikoff (Dept. of Physiology, Univ. of Calif., Berkeley, Calif.). *J. Lipid Res.* 6, 228-232 (1965). The esterification of cholesterol by rat pancreatic juice and by extracts of acetone powders of rat small intestine was studied with palmitic, stearic, oleic, linoleic, and linolenic acids. When each fatty acid was tested individually, the extent of esterification was highest with oleic acid and lowest with the two saturated fatty acids. The esterification of cholesterol with palmitic and stearic acids by the intestinal extracts and with palmitic acid by pancreatic juice was greatly increased by the addition of either oleic or linoleic acid to the incubation mixture. It is suggested that the difference between esterification of cholesterol with saturated fatty acids and esterification with unsaturated fatty acids, as well as the enhancement of the esterification with saturated fatty acids by the addition of an unsaturated fatty acid, is accounted for by micellar solubilization of cholesterol and of saturated fatty acid in the presence of an unsaturated fatty acid and bile salts.

PARTICLE SIZE AND COMPOSITION OF DOG LYMPH CHYLOMICRONS. A. Yokoyama and D. B. Zilversmit (Dept. of Physiology, Univ. of Tennessee Medical Units, Memphis, Tenn.). *J. Lipid Res.* 6, 241-246 (1965). Chyle from the thoracic duct of dogs fed cream or corn oil was fractionated into chylomicrons of different particle size: 140  $\mu$ , 140-200  $\mu$ , and 200  $\mu$ . The composition of chylomicrons was studied before and after washing with saline solutions. In general the percentages of protein, phospholipid, cholesterol, and cholesterol ester were greater in the small chylomicrons than in the larger ones. Washing the chylomicrons did not appear to affect their composition. Chylomicrons collected at different times during fat feeding showed significant differences in phospholipid and cholesterol concentrations. Storage of lymph at 4C was accompanied by alterations in phospholipid and cholesterol concentrations of the chylomicrons. The composition of chylomicrons is consistent with the hypothesis that at some stage during their formation the nontriglyceride components are present on the surface of the oil droplet.

FAT/OIL

facts faster!



Steinlite

MODEL  
300-LOS

FAT AND OIL  
TESTER

FRED STEIN  
LABORATORIES, INC.

Atchison, Kansas

Steinlite determines either content in just 10 to 15 minutes!

Steinlite offers profitable advantages for these products:

- Frankfurter Emulsions
- Luncheon Meat
- Corn Chips
- Bologna Emulsions
- Deviled Ham
- Pork Sausage
- Flax
- Ground Beef
- Fried Noodles
- Copra
- Potato Chips
- Ground Pork
- Soy Beans
- Trimmings
- Peanuts
- Corn Meal
- Sesame Seed
- Dog Food
- Cottonseed
- Cabbage Seed
- Fish Meal
- Corn Germ
- Castor Beans
- Pumpkin Seed
- Mink Food
- Mafura Beans
- Safflower Seed
- Sunflower Seed
- Rice Bran
- Chocolate

Tell Us More!

Write today - Tell us about your product - We'll tell YOU all about STEINLITE, as it applies to your product. No obligation on your part! Just address inquiries to our Fat and Oil Department.

## • Drying Oils and Paints

ELECTRICAL INSULATING VARNISH. C. K. Sung, T. W. Park and B. S. Lee. *Bull. Sci. Res. Inst.* 4(1), 20-5 (1959). The varnish is based on 5 pts. of a drying oil mixture, containing 85% tung oil and 15% linseed oil, with 4 pts. of an oil-soluble phenolic resin. The applied coating is stoved at 160C for 1 hr. Information on breakdown potential, flexibility, relative hardness and acid, alkali, oil and moisture resistance is given. (Rev. Current Lit. Paint Allied Ind., No. 268)

AUSTRALIAN PRODUCTION STATISTICS—PAINTS AND OTHER SURFACE COATINGS, MAY, 1963. Anon. *Austral. Paint J.* 9(6), 34 (1963). The data are classified by type of product and states in which they were produced (Victoria, Queensland, etc.). Similar information is given each month. (Rev. Current Lit. Paint Allied Ind., No. 268).

SURFACE COATINGS IN THE CHEMICAL COMPLEX. W. P. Georgans. *Austral. Paint J.* 9(6), 22-4 (1963). A discussion of the relationship between the chemical and paint industries, with statistics on the Australian production and consumption of different types of paint and the growth in the consumption of TiO<sub>2</sub>. (Rev. Current Lit. Paint Allied Ind., No. 268).

POLYMERIZATION OF DRYING OILS. H. Wexler (Continental Can Co., Chicago, Ill., 60620). *Chem. Rev.* 64, 591-611 (1964). An excellent review.

USE OF OXIDATION IN THE SYNTHESIS OF DRYING OILS FROM TALL OIL MODIFIED WITH CYCLOPENTADIENE. A. A. Subbotin. *Lakokras. Mat.* 1963 (6), 18-21 (1963). Tall oil was reacted with cyclopentadiene, esterified and oxidised with air. The resulting drying oil was found to have good wetting properties in relation to the common pigments and be suitable as a thinner for thick oil-bound paints. (Rev. Current Lit. Paint Allied Ind., No. 268).

EFFECT OF WATER ON PAINT FILMS. II. [ADHESION]. III. WATER VAPOUR PERMEABILITY. IV. BLISTERING OF EXTERIOR HOUSE PAINTS. G. Gärdenäs and E. Wählin. *Färg och Lack* 10 (2), 20-36; (3), 71-4; (4), 83-5 (1964). Alkyd paints, oil paints and polyvinyl acetate latex paints have been studied by measuring the force required to pull off steel cylinders, which have been glued on to the paint film. Alkyd paints retain most of the adhesion even after 10 days' treatment in a moisture cabinet, which is not the case with oil paints and polyvinyl acetate latex paints. Most of the adhesion is restored after drying the panels in air. Interoat adhesion and cohesion in some cases is lower than the adhesion to the substrate. III. The water vapour permeability has been studied using the Payne cup method, the paint films being applied to Cellophane. Alkyd paints show very low permeability, 1.1-1.3 mg./cm.<sup>2</sup> day. Oil paints 2.0-2.5 mg./cm.<sup>2</sup> day and polyvinyl acetate latex paints 20-25 mg./cm.<sup>2</sup> day. IV. The blistering tendency of oil paints, alkyd paints and polyvinyl acetate latex paints has been investigated by subjecting wood panels to 100% relative humidity on the unprotected side while keeping the painted side at 65% relative humidity. Three out of four oil paints did show blistering, whereas with the alkyd and polyvinyl acetate latex paints no blistering was observed during the 10 days' testing period. (Rev. Current Lit. Paint Allied Ind., No. 269).

FORMATION OF TRIMER ACIDS IN THE THERMAL POLYMERIZATION OF LINSEED OIL. E. Fedeli, P. Capella, A. F. Valentini and G. Jacini (Fats and Oils Exper. Stat., Milan, Italy). *Riv. Ital. Sostanze Grasse* 41, 647-51 (1964). The presence of trimer acids in the thermal polymerization of linseed oil has been proved by chromatographic analysis. The kinetics of formation of trimer acids indicates that these are not formed by further polymerization of dimer acids. The interrelation between polyunsaturates and the formation of dimer and trimer acids is discussed.

GENERAL DEVELOPMENT AND PRESENT TRENDS FOR PAINTS. A. Cailliez. *Peint Pig. Vernis* 40, (4), 189-96 (1964). Difficulties of introducing new products and developments during the last war and the following decade are discussed. Present-day problems and trends are considered in relation to certain recent

statistics. The probable future for different media, solvents, latex and pigments is discussed and zinc-rich paints singled out for special mention. (Rev. Current Lit. Paint Allied Ind. No. 271).

STATISTICS—READY MIXED PAINTS AND RAW MATERIALS. Anon. (Central Bureau voor de Statistiek (Netherlands)). *Verf-kroniek* 37 (4), 128-9 (1964). Import and export figures for 1963 between the Netherlands and other countries for different types of paints, lacquers, printing inks, etc., and their raw materials are set out in detail. (Rev. Current Lit. Paint Allied Ind. No. 271).

NEW CONJUGATED SAFFLOWER OIL. L. Cummings and W. Rathjen. *Am. Paint J.* 48 (49), 16, 18 (1964). Safflower oil containing 18-19% conjugation is now available and this may be used to prepare alkyds suitable for modification with methyl methacrylate/styrene mixtures to give films of improved properties. (Rev. Current Lit. Paint Allied Ind. No. 272).

UNDERSTANDING PAINT, XI-XVI. W. R. Fuller. *Am. Paint J.* 48, (1964), No. 33, 86 (11 pp.); No. 54, 61 (12 pp.); No. 55, 90 (10 pp.); No. 56, 60 (4 pp.); 49, No. 1, 59 (13 pp.); No. 2, 66 (10 pp.). A comprehensive survey of the paint industry, covering both the properties, chemistry and uses of the different raw materials and the formulation and uses of different types of paints. (Rev. Current Lit. Paint Allied Ind., No. 272).

UTILISATION OF COTTONSEED OIL FOR PREPARATION OF OLEORESIN VARNISHES. M. Aslam. *Pakist. J. Sci. Ind. Res.*, 6, 188-91 (1963). Cottonseed oil has been successfully used for the preparation of tack-free coating compositions by a simple open kettle cooking process. Best results were obtained by judicious control of (a) cooking temp., (b) amount of catalyst, (c) extent of cooking and (d) proportion of resin. These varnishes compare favourably with those prepared from linseed oil. (Rev. Current Lit. Paint Allied Ind. No. 272).

MARKET FOR DRYING OILS. Anon. *Tropical Products Inst. Report No. 5*, 1963, 56 pp.—A very good account of the use of the major drying oils in paints, etc., is given. Many tables of statistics, production, prices, uses and consumption of the drying oils are included, also sales of paints and varnishes in the U.K. (Rev. Current Lit. Paint Allied Ind. No. 272).

## • Detergents

STUDIES ON SODIUM ALKYL BENZENE SULFONATES. IV. SULFONATION OF HIGHER MONOALKYL BENZENES IN RELATION TO STRUCTURAL FACTOR OF THE ALKYL CHAIN. Yasushi Kimura, Syuhei Tanimori, and Terunosuke Shimo (Lion Fat and Oil Co., Tokyo). *Yukagaku* 14, 63-6 (1965). The relative rate of sulfonation of linear dodecylbenzenes and *n*-hexylbenzene at 18C with 20% oleum has been determined by a unique method of applying gas chromatography. The relative reaction rate of linear dodecylbenzenes decreases as the position of aromatic nucleus is shifted toward the center of the alkyl group. The rate of 6-phenyl isomer is about 40% less than that of 1-phenyl isomer. The rate of *n*-hexylbenzene, having the half-length of alkyl chain, is about 60% more than that of *n*-dodecylbenzene. Also, there is more pronounced selectivity in the sulfonation of higher monoalkylbenzene owing to the steric hindrance of the alkyl chain. Selective sulfonation was eliminated by heating at 50C and a new product with lower boiling point was obtained.

PREPARATION OF UNSATURATED HIGHER ALCOHOL UNDER HIGHER HYDROGEN PRESSURE (300-430 kg./cm.<sup>2</sup>) USING ZN-AL-O CATALYST. Isao Ikeda, Motoaki Tanaka, and Saburo Komori (Osaka Univ.). *Yukagaku* 14, 58-62 (1965). High yields (90-95%) of unsaturated alcohol were obtained from methyl esters of rice bran oil at 330C with pressure at 300-430 kg./cm.<sup>2</sup> by use of 3-5% Zn-Al-O catalyst. Hydrogen at high pressure gave a higher yield of alcohol than at low pressure. Reduction product can be used immediately as the raw material for surface-active agent manufacture without purification on account of its high alcohol content.

APPLICATION OF SURFACTANTS ON WOOL INDUSTRY. III. CONSIDERATION ON WOOL SHRINKAGE IN DETERGENT SOLUTION. Chikaaki Sakai and Saburo Komori. *Yukagaku* 14, 66-71 (1965). In order to explain the phenomenon of wool shrinkage in detergent solution, the change of physical properties of a single wool fiber was investigated and it was observed that the lubrication caused by the sorption of detergent on wool surface seemed to be the main factor. Aqueous solutions of surface-active agents having aromatic group in their molecule, such as alkylbenzenesulfonate and butyl naphthalenesulfonate, increased


**H A H N L A B O R A T O R I E S**  
Consulting and Analytical  
Chemists

1111 Flora St. P.O. Box 1163 Columbia, S.C.

the friction of fiber and retarded its shrinkage. In the presence of fatty alcohol, the shrinkage of wool was accelerated from its lubrication effect.

**SURFACE TENSION OF AQUEOUS SOLUTION OF POLYVINYL ALCOHOL ESTERIFIED WITH FATTY ACID IN THE HOMOGENEOUS SYSTEM.** Sadao Hayashi, Chiyoko Nakano, and Takuhiko Motoyama (High Polymer Chemical Industries, Ltd., Osaka). *Yukagaku* 14, 24-6 (1965). Polyvinyl alcohol was esterified with formic, acetic, and propionic acids, respectively, in water containing hydrochloric acid as a catalyst. The surface tensions of these esterified products were greater in order of formic acid < acetic acid < propionic acid ester.

**STUDIES ON ANTISTATIC PROPERTY OF SURFACE ACTIVE AGENTS.** Takashi Yamamoto, Ikuo Katsura and Seiji Sumida (Nippon Oils and Fats Co., Amagasaki-shi, Japan). *Yukagaku* 13, 480-6 (1964). Surface electrical resistivity of fibers treated with surfactant solutions and specific conductivity of the solutions were measured as well as the effect of added electrolyte on the orientation of surfactants on the fibers. Surface electrical resistivity was decreased as the concentration of surfactant increased. In the vicinity of the critical micelle concentration of surfactant it increased then decreased at higher concentrations. Inorganic salts added to surfactant solutions did not increase electrical conductivity of the fibers but contributed to the change of the micelle state of surfactant solutions. Addition of sodium chloride to the solutions of cationic surfactants resulted in an increase sorption of surfactant by the fibers. Sodium hydroxide was not effective. Antistatic treatment is enhanced by use of cationic surfactants together with electrolytes having the tendency to salt out cationic surfactants and to make micelle molecular weight larger. Antistatic effect of fibers treated with organic solvent was less than that treated in water because of low molecular weight of micelle. Piles of Nylon, Dacron, cotton and glass fibers treated with surfactants were dispersed in benzene-water and the orientation to fibers of the surfactant was studied. Piles treated with sodium dioctyl phosphate, which is the only antistatic agent of anionic surfactants, dispersed in benzene or at the interface. Orientation of surfactant to fibers may explain the effectiveness of cationic surfactants and of sodium dioctyl phosphate.




*Analyses and Research*  
**REID H. LEONARD, Ph.D.**  
*Consulting Biochemist*  
 P. O. BOX 2334  
**PENSACOLA, FLORIDA 32503**

**APPLICATION OF THIN-LAYER CHROMATOGRAPHY TO OIL CHEMISTRY. I. CHROMATOGRAPHY OF SYNTHETIC SURFACTANTS BY SILICA PLATES.** Toru Takagi and Kazuo Fukuzumi (Nagoya Univ., Nagoya, Japan). *Yukagaku* 13, 520-2 (1964). Seventeen kinds of synthetic surfactants were chromatographed on silica-coated plates. Chloroform-methanol-water system was selected for developing. The surfactants gave sharp spots when free components were developed but not the salts. Anionic surfactants gave peculiar fluorescence by ultraviolet light after spraying of sulfuric acid and heating.  $\alpha$ -Sulfo-fatty acid was separated from other sulfonic acids and sulfates. Polyoxyethylene surfactants were separated into major components and polyethylene oxide. Cationic surfactants were detected by Dragendorff reagent and platinum chloride-potassium iodide reagent. These procedures are useful to detect impurities in synthetic surfactants and to check progress of the reaction in the manufacturing process.

**STUDIES ON THE SURFACE TENSION OF LOWER POLYOXYETHYLENE ETHERS OF LAURYL-, CETYL-, OLEYL-ALCOHOL AND NONYLPHENOL.** Shoichiro Watanabe (Kitasato Univ., Tokyo). *Yukagaku* 13, 603-7 (1964). The surface activities of polyoxyethylene ethers having definite length of oxyethylene chain have been investigated. Mono-, di-, tri-, tetra- and penta-oxyethylene ethers of lauryl, cetyl and oleyl alcohols and nonylphenol were synthesized. Those were dissolved or suspended in water and the surface tension of the solutions was measured. With the exception of cetyl ethers and mono- and di-oxyethylene lauryl ethers, the surface tension-concentration curves of the solutions showed the same characteristics as those of usual surfactants.

# super kleeer

## NICKEL HYDROGENATION CATALYST



**CAT-CHEM**

**Filterability**  
excellent clarity without added filter aids.

**Activity**  
superior, even under the most adverse hardening conditions.

**Selectivity**  
improved over a wide range of commercial hardening conditions.

**Re-useability**  
the re-use quality of this catalyst affords many additional uses, thereby reducing operating costs.

**Resistance to Poisoning**  
this rugged catalyst can be used for fatty acids and other inedible products, reducing hardening times along with excellent filterability.

**Predictable Performance**  
the hydrogenator can be assured of uniform reproduceable results.

**Proved Process Efficiency**  
the superior quality of this catalyst has been verified in many commercial operations, improving over-all hydrogenation performance.

After meeting all exacting requirements  
containers are sealed for shipment.  
Your drums arrive with "QUALITY SEALED IN"

For further information, technical data and samples write:


## CAT-CHEM CORPORATION

68 ELIZABETH AVENUE, ELIZABETH, NEW JERSEY • 201-351-4555

75 years of skill and experience combined to give the ultimate in product, performance and service.

For processing  
winterized oils

# NIAGARA<sup>®</sup> FILTERS OFFER 3-WAY ECONOMY



**SAVE ON INSTALLATION COSTS:** The fully jacketed design of NIAGARA filters eliminates the need for refrigerated rooms. **SAVE LABOR COSTS:** Design utilizes a simple system of push buttons and valves that enable one man to operate an entire bank of NIAGARA filters. **SAVE FILTER MEDIA:** Leaf design permits easy cake removal with no scraping. This lengthens cloth life.

NIAGARA filters are available in 3 basic types . . . vertical pressure leaf, horizontal pressure leaf and hor-

izontal plate filter. Investigate NIAGARA filters for all your filtration problems. See the advantages of greater flow; lower maintenance and operating costs; positive solids removal and totally enclosed construction. For full details on custom-built, cost-saving NIAGARA filters, write today. Address Dept. JO-665.



## NIAGARA FILTERS

A DIVISION OF AMETEK, INC.  
EAST MOLINE, ILLINOIS

(Niagara Filters Europe: Kwakelpad 28, Alkmaar, Holland)

Namely, there was little change in the surface tension at higher concentration, while it increased rapidly as the concentration decreased in the range below critical micelle concentration. The surface tension of the solutions of paired ethers containing lauryl-oleyl, oleyl-nonylphenyl, and nonylphenyl-lauryl, having tetraoxyethylene chain, respectively, was in a range between those of individuals of the pair.

**DETERGENTS VIA NEW ROUTE.** C. Y. Shen and C. F. Callis (Monsanto Co.). *Soap Chem. Specialties* 41(2), 64, 68, 70, 109-12, 114 (1965). The alkaline hydrolysis of trimetaphosphate opens the way to new technology in processing conventional spray-dried detergents as well as novel types of products. Detergents made by this route usually contain more sodium tripolyphosphate and less degradation products than those made directly from STPP. The apparent viscosity of a STMP slurry is less than that of a corresponding STPP slurry. Slurries of high solids concentration and high density can be made from TMPP, allowing processes and formulation for a much lower drying load than is required by present spray tower practice. STMP provides a uniform source of phosphate with chemical properties that are easier to control than are the properties of STPP. STMP shows no tendency to lump in the crutching operation.

**DETERGENT MARKET IN WESTERN EUROPE.** G. L. Hollis (Imperial Chemical Industries, Ltd.). *Soap Chem. Specialties* 41(2), 55-8, 90 (1965). This is the first part of an article discussing the rise of synthetics and the markets in Western Europe, the introduction of biodegradable detergents in the United Kingdom and Western Europe, and the future of syndet materials.

**DETERGENT BRIQUETTE.** D. F. Percival and W. A. Sweeney (California Research Corp.). *U.S.* 3,172,859. A water-soluble briquette comprises an intimate admixture compacted in the presence of 5 to 20% of water at a pressure in the range of 250 to 700 pounds per square inch. The briquette consists of, by weight, 15-80% water-soluble normal primary alkyl sulfonate detergent salt selected from the group consisting of the alkali metal, alkaline earth metal and ammonium salts, the alkyl group containing 10-20 carbon atoms and averaging 12-18 carbon atoms; 25-75% water-soluble inorganic condensed sodium phosphate detergent builder; and the 5-20% of water which was present during the compacting step.

**DETERGENT COMPOSITIONS COMPRISING INORGANIC ESTERS OF EPOXYHYDROCARBON POLYMERS.** D. R. Jackson (Wyandotte Chemicals Corp.). *U.S.* 3,173,877. The described detergent composition consists of about 36% by weight of a disodium polyoxypropylene glycol disulfate in which the molecular weight of the polyoxypropylene base is at least 1250, about 55% of sodium tripolyphosphate, and about 9% of sodium carboxymethylcellulose.

**EFFECT OF THE MOLECULAR STRUCTURE OF SURFACTANTS ON THEIR AEROBIC BIOLOGIC DEGRADATION.** H. Kölbl, P. Kurzen-dörfer and M. Zahiruddin (Univ. of Berlin, Germany). *Ten-side* 1, 7-18 (1964). The effect of specific structural groups on the biologic degradability of anionic surfactants of the alkyl aryl sulfonate type has been investigated. Among the surfactants studied were: *n*-alkyl sulfonates, *p*-*n*-alkyl benzene sulfonates, *p*-*n*-alkyl diphenyl sulfonates, 2,6-*n*-alkyl naphthalene sulfonates, *p*-2,2-bis(*n*-alkyl)-ethyl benzene sulfonates and *p*-*N*-methyl-*N*-acyl aniline sulfonates. All biological tests were conducted by Henkel's closed flask test, with cultures of *E. coli* ( $10^8$  to  $10^4$  organisms/ml.), at an initial surfactant concentration of  $10^{-5}$  mol/l and initial oxygen concentration of 8-10 mg/l. During the course of the test surfactant and  $O_2$  concentrations, pH,  $KMnO_4$  titration and microorganism concentration were all measured. Alkyl sulfonates are the most degradable among the surfactants tested, since they not only disappear completely in 5-6 days, but are completely oxidized in a period of 14-20 days, regardless of chain length. Alkyl benzene sulfonates, alkyl diphenyl- and alkyl naphthalene sulfonates are also completely degraded in 7-15 days but do not oxidize completely (the max.  $O_2$  consumption is only 20-40% of theoretical). The optimum chain length of the alkyl group is in most cases 10-14 C atoms. The *p*-2,2-bis(*n*-alkyl)-ethyl benzene sulfonates are very little degraded, with only the  $C_{12}$  chain being completely destroyed in 35-40 days. The sodium *p*-*N*-methyl-*N*-acyl aniline sulfonates disappear completely in 8-14 days (optimum chain length 12-16 C atoms) but are not completely oxidized.

**BIOLOGIC BEHAVIOR OF SURFACTANTS.** A. DeJong (Unilever Res. Lab., Vlaardingen, The Netherlands). *Riv. Ital. Sostanze Grasse* 41, 547-9 (1964). Fish appear to be very sensitive to the presence of surfactants in their environment.  $LD_{50}$  values of 3-5 ppm are reported for linear alkyl aryl sulfonates, while

(Continued on page 327A)



## AOCS Short Course

(Continued from page 296A)

of Lipid Mixtures on Liquid-Solid Chromatographic Columns—Gary Nelson

- 1) Structure Determination of Unsaturated Acids by Ozonolysis. 2) Quantitative Thin-Layer Chromatography of Lipids—Orville S. Privett

Analytical Fractionation of Complex Lipid Mixtures (Column Chromatography, Thin-Layer Chromatography and Associated Techniques)—George Rouser

Mass Spectrometry Directly Coupled to Gas Chromatography. Demonstration of Techniques—A. Struck

- 1) Analysis of Steroids by Thin-Layer and Gas Chromatography. 2) Membrane Systems and the Use of Molecular Models—F. A. Vandenheuvel

Plant Inositol Lipids. Importance and Alteration of Salt Forms—Evelyn Weber

For further information contact the Director, C. Freeman Allen, Department of Chemistry, Pomona College, Claremont, California.

tetrapropylene benzene sulfonate and fatty alcohol sulfates are far less toxic. Bacteria and other microorganisms in water seem to be able to survive rather high surfactant concentrations. Concentrations above 5 ppm in irrigation water may affect plant growth, especially when the organic matter content of the soil is low. Mice, rats, rabbits and dogs consumed between 50 and 30,000 ppm surfactants in their daily diet for up to two years without any discernible effects on their health. Men have consumed water with 100 ppm surfactants for several weeks without any sign of chronic harm to their health. There is at present very little chance that the surfactant concentration in the diet of man will exceed 2 ppm.

THE DIRECT CONDENSATION OF FATTY ACIDS AND TAURINE. E. Elbel (Farbwerke Hoechst A.G., Frankfurt/Main, Germany). *Tenside* 1, 26-28 (1964). The condensation of fatty acid chlorides with N-methyl taurine in the presence of NaOH leaves considerable amounts of NaCl in the end product. Attempts to avoid this by using a direct condensation reaction with fatty acids in the past were only partly successful, mainly because of side reactions leading to the formation of very hygroscopic by-products, such as the monosodium salt of N-methyl ditaurine. This difficulty can be overcome by the use of salts of volatile fatty acids as neutralization media in the direct condensation process. Sodium formate and acetate have both been successfully used in the preparation of detergents of the Hostapon T type, the resulting acid being then driven off by distillation. The end product of this process exhibits very little hygroscopicity.

MEASUREMENT OF THE SURFACE TENSION OF AQUEOUS SURFACTANT SOLUTIONS. G. Schwen (Ludwigshafen/Rhein, Germany). *Tenside* 1, 46-50 (1964). The surface tension of aqueous solutions of surface active agents depends not only on chemical composition, concentration, electrolyte level and temperature, but also on time. Experimental methods are therefore required to take the time variable also into consideration. A new quasistatic method for measuring surface tension, called the capillary pressure method, is described. The results of three series of experiments are reported, in which surface tension was measured for four different surfactants, varying concentrations of a non-ionic surfactant and varying amounts of electrolyte. The time dependency of surface tension was followed up to 25 min., at which time it had generally reached a constant, minimum value.

SURFACTANTS IN TEXTILE INDUSTRY EFFLUENTS. M. Kehren (Holzschlag, Germany). *Tenside* 1, 109-11 (1964). The situation of the textile industry in Germany is reviewed with respect to the German law on detergent biodegradability.

THE ADVENT OF SURFACTANTS. A. Chwala (Vienna, Austria). *Tenside* 1, 41-45 (1964). A historical review of the development of surface active agents.

THE ANALYSIS OF SURFACTANTS, I. D. Hummel (Univ. of Cologne, Germany). *Tenside* 1, 50-9 (1964). This paper gives a comprehensive survey on qualitative and quantitative methods of surfactant analysis, with special reference to spectroscopic and ion exchange methods. Besides extensive literature survey there are reported experimental results on the identification of surfactants which have not so far been published. In Part 1 of this paper the author deals with qualitative methods for

## Rensselaer Polytechnic Plans June Short Course in Color Technology

Through the Department of Chemistry, Rensselaer Polytechnic Institute will offer for the first time, a special comprehensive short course in The Principles of Color Technology, from June 21-25, 1965. This course is intended to provide both theory and practice in the description, specification and measurement of color. It will be of particular interest to industrial personnel responsible for color matching and color control.

Address application for admission to Prof. A. A. K. Booth, Director, Division of Special Programs, Rensselaer Polytechnic Institute, Troy, New York 12181.

## • Obituaries

H. E. Moore (1934), of New Orleans, Louisiana, died recently, it was learned at the AOCS headquarters in Chicago.

testing surface activity and the charge of surface active ions as well as analytical methods for anionic and cationic surfactants.

III. *Ibid.*, 116-25. The third part of this paper deals with methods for the quantitative analysis of surfactants. These include gravimetric, volumetric, colorimetric and absorption-spectroscopic methods as well as special methods for the quantitative analysis of mixtures. More recently developed methods, such as I.R. and U.V. spectroscopy, ion exchange, as well as new titrimetric methods are given special attention. It is the object of these critical discussions to make the choice of a suitable analytical method easier.

AMINE OXIDE SURFACTANTS. H. Lindner (Berlin-Lichterfelde, Germany). *Tenside* 1, 112-5 (1964). The reaction of tertiary fatty amines (e.g. lauryl dimethyl amine) with hydrogen peroxide solutions is discussed. This is a reversible reaction, which can also cause the decomposition of the amine oxide under some conditions, e.g. at elevated temperatures. Different views on the molecular structure of these compounds are discussed, as well as the surface active properties of their aqueous solutions. The excellent cleaning and foaming properties of amine oxides, especially at low temperatures, make them well suited for use in detergents, dishwashing liquids, hair shampoos and textile finishing.

THE NATURE OF HYDROTROPY AND ITS SIGNIFICANCE IN INDUSTRIAL CHEMISTRY. H. Ratts (Inst. für Textilchemie, Stuttgart, Germany). *Tenside* 2, 1-6 (1965). Examples of the various branches of industrial chemistry in which hydrotropic substances can be used are given from the fields of organic technology, dyestuff synthesis, dyeing and printing, pharmaceuticals and cosmetics, cellulose production, tanning, prevention of calcium soap formation and swelling of high molecular weight substances. The nature of hydrotropy consists in the formation of adducts between the materials to be dissolved and the hydrotropic substance. The tendency to adduct formation and the property of the adduct to dissolve in water are closely dependent on the chemical constitution of the hydrotropic agent. This connection is illustrated by numerous examples.

ATTEMPTS AT SEPARATING SUCROSE MONOESTERS BY TLC AND DETERMINATION OF CAPILLARY PROPERTIES OF POSITIONAL ISOMERS. W. Wachs and K. Gerhardt (Univ. of Berlin/Charlottenburg, Germany). *Tenside* 2, 6-10 (1965). With the aid of thin layer chromatography, industrial sucrose-fatty acid esters were separated into their mono-, di- and triesters. It was also possible to separate the fraction of sucrose monoesters obtained into three groups (altogether seven individual compounds) of positional isomers. The various industrial com-

**THE POPE TESTING LABORATORIES**  
Analytical Chemists

2618½ Main

P.O. Box 903

Dallas, Tex.