

Gdansk, Poland, Host to International Rapeseed Symposium

The Food Technology and Chemistry Committee of the Polish Academy of Science and the Department of Fat Technology of the Gdansk Technical University will host the International Symposium for the Chemistry and Technology of Rapeseed Oil and other Cruciferae Oils from Sept. 19-23, 1967.

Three days will be given to discussions; two days to visiting fats industrial works in the Gdansk region and the tourist region of the Kaszub lake district. The discussions will be held at the Technical University, Gdansk, Majakowskiego 11.

Provisional applications for participation should be submitted not later than Dec. 31, 1966, and final application for participation should be sent not later than April 1, 1967. Also due before April 1, 1967 are two copies of abstracts (in English) of not more than 100 words.

The program will include research work regarding the chemistry and technology of rapeseed oil and of other cruciferae oils. There will be plenary and sectional meetings. The 45-minute plenary meetings will be devoted to lectures prepared by top specialists. Informational news from original works will be part of the 15-minute sectional meetings.

The Symposium will be composed of the following five sections: 1) cruciferae seeds as industrial raw material; 2) chemistry, biochemistry, analysis; 3) technology of edible fats; 4) technology of nonedible fats and their derivatives; 5) food and fodder value of products from cruciferae.

The main language of the symposium will be English. All papers will be in English, and English will be preferred in the discussions. French, German, and Russian will be admitted. There will be no simultaneous translations into other languages.

Summaries of papers will be printed before the beginning of the Symposium. The Polish Academy of Science will publish full texts of papers in the language in which they are presented. The printing deadline for submitting full texts of papers with illustrations in a form ready for publication is the time of the Symposium.

There is a fee of \$5.00 for participation in the scientific part of the Symposium. Further information will be provided upon request by Prof. Dr. H. Niewiadomski, Katedra Technologii Tluszczow, Politechnika Gdanska, Gdansk-6 (Poland), Majakowskiego 11.

• Industry Items

DREW CHEMICAL CORPORATION, New York, N. Y., has sold its Wilmington, Calif., plant to a group of investors who will continue the operations under its former name, Vegetable Oil Products Company, Inc. W. L. Dickinson will continue as president.

All SHARPLES centrifuge operations of the PENNSALT CHEMICALS CORPORATION have been transferred from Philadelphia to the company's new \$8 million equipment division plant in Warminster, Pa.

According to A. R. Greenlaw (1950), plant general manager, the 285,000 sq ft facility will increase the company's centrifuge production capacity. It houses administrative, research, engineering and manufacturing operations for the complete line of Sharples and Fletcher centrifuges. It is also headquarters for Stokes tableting presses, punches and dies.

EMERY INDUSTRIES, INC., of Cincinnati has broken ground on a new technical center. The cost of the new structure will be approximately \$1 million and will be located at Emery's Ivorydale location. The center will house engineering, technical sales service, and product development personnel.

(Continued from page 594A)

paraffins and mixtures thereof, into detergent grade polycyclic compounds boiling in the range of 570-725F comprises reacting the bottoms fraction with aluminum chloride at a temperature between 40-150C.

ENTROPY-DETERMINED COMPLEX FORMATION; THERMODYNAMIC DATA ON COMPLEX FORMATION BY THE TRIPOLYPHOSPHATE ION. G. Anderegg (Tech. Hochschule, Zürich, Switzerland). *Helv. Chim. Acta* 48, 1712-7 (1965). The formation of the 1:1 complexes of 10 divalent metal ions with tripolyphosphate has been studied from a thermodynamic standpoint. The stability constants and the heats of formation of these complexes have been determined. In all cases, the complex formation is endothermic and entropically favored. The 1:2 complexes are known to be considerably less stable than 1:1 complexes and hence were not investigated.

SURFACTANTS DERIVED FROM P-CUMYL PHENOL. P. Peri (Edison S.p.A., Mantua, Italy). *Riv. Ital. Sostanze Grasse* 43, 166-74 (1966). Ethylene oxide adducts of p-cumyl phenol (a by-product in the preparation of phenol and acetone from cumene) have been prepared. Non-ionic adducts with 2 to 12 mols of ethylene oxide as well as their corresponding sulphates have been prepared and evaluated. Both classes of compounds thus obtained had very low foaming power, with the non-ionic adducts being generally somewhat better in detergency than their sulfated equivalents.

APPLICATION OF TLC TO THE ANALYSIS OF CATIONIC SURFACTANTS. M. R. Gasco and G. Gatti (Univ. of Turin, Italy). *Riv. Ital. Sostanze Grasse* 43, 175-6 (1966). The conditions for thin layer chromatographic analysis of several cationic surfactants are described and the R_F values in various solvent combinations are reported.

SURFACE ACTIVE AGENTS IN THE ELECTROPLATING INDUSTRY. W. Strauss (Deutsche Hydrierwerke GmbH., Düsseldorf, Germany). *Tenside* 3, 144-50 (1966). Surface active agents are used in

SODIUM METHYLATE
(Sodium Methoxide)

Spec 101. Packaged in airtight steel drums of 10, 25, 50 and 200 pounds net.

Free flowing white powder.
Formula NaOCH_3 .

Bulk density 4.15 pounds per gallon.
Molecular weight 54.03.

Write today for full information.

HARSHAW
THE HARSHAW CHEMICAL COMPANY
CLEVELAND, OHIO 44106