


Chicago, Illinois, Scene of the 44th Fall Meeting of AOCS and 10th Biennial Congress of ISF September 27-October 1, 1970



All Roads Lead to Chicago

There are certain spots on the globe that were meant by destiny to occupy key positions no matter what the change in civilization or the new twists in man's developments with their attendant results on his habits, and one of these is Chicago.

Situated at the bottom of Lake Michigan in the richest section of the Mississippi Valley and within a few hundred miles of both the nation's center of population, as well as its geographical center, Chicago occupies a strategic location unparalleled anywhere.

This Fall, the spotlight is on Chicago, as it plays host to the 44th Annual Fall Meeting of the American Oil Chemists' Society in joint session with the 10th Biennial Congress of the International Society for Fat Research, at the famed Conrad Hilton Hotel September 27 through October 1st.

With world food problems being brought more into focus, and with the reformulation of soaps, detergents and cosmetics moving at a rapid pace, this World Congress should be the most important marketing event in the oils and fats industries in 1970.

Technical Program

Bob Reiners and his excellent staff have put together the most comprehensive technical program ever prepared for a meeting of this type. Papers will be presented on the scientific, technological and economical aspects of oils, other lipids and associated substances. The program will consist of four daily plenary lectures by invited speakers, a broad array of symposia and many reports of original research divided into topical sessions. As an example of the excellent quality of the symposia being offered, we are advised that Reinhard Marcuse of the Swedish Institute for Food Preservation Research at Göttenborg, Sweden, has assembled an imposing group of speakers for his symposium on "Metal Catalyzed Lipid Oxidation." Abstracts of the complete technical program will appear in the July issue of your AOCS Journal.

In addition, two excellent short courses have been scheduled in conjunction with the World Congress; one immediately before the Congress and one immediately after. They are: "Processing and Quality Control of Fats and Oils" and "Behavior of Membrane Lipids at Surfaces." These courses are expected to appeal to a broad spectrum of participants, from both industry and the academic world as well. Do not overlook the opportunity to attend one or both of these fine courses.

There will also be an International Conference on the "Science, Technology and Marketing of Rapeseed and Rapeseed Products," at the Le Chantecler Hotel, Ste. Adele,

Quebec, Canada, September 20-23. This Conference is being organized by Bernd Weinberg of the Edible Oils Section, Department of Industry, 112 Kent Street, Ottawa 4, Ontario, and requests for additional information should be addressed to him.

Plant Visits

The Chicago Area provides an important concentration of industrial and academic facilities devoted to the various facets of the fat, oil and lipid sciences. Visits to a number of these outstanding plants and laboratories will be available to registrants.

International Fats and Oils Exposition

The first truly "international" Fats and Oils Exposition will be held in conjunction with the World Fat Congress. Here is an opportunity for all registrants to examine and discuss the latest innovations in laboratory and plant equipment, processes and techniques, ingredients and additives, and the latest in literature.

Anyone wishing to inquire about display space should contact W. R. Deutscher, Exhibits Chairman, at Somes-Nick & Co., 407 South Dearborn Street, Chicago, Illinois 60605.

Social Events

When the business meetings are over, it is time to relax; to talk with old friends and make new ones. It is time to have fun.

On Sunday evening, there will be the traditional Reception and Cocktail Party. "Christine" at the electric organ will provide entertainment, while colleagues from every corner of the globe will gather to celebrate the opening of this first ISF Congress to be held in the United States.

Monday evening is left free to enjoy as you wish. Registrants will be provided with complete information on Chicago's renowned restaurants and supper clubs, as well as the other forms of entertainment available within easy distance of the hotel.


On Tuesday evening, we will be treated to a special concert by the world-famous Swedish Glee Club with the singing magic of their 60 beautiful voices.

Highlighting the social schedule of the Congress will be the banquet on Wednesday evening. There will be dinner in the Conrad Hilton's Grand Ballroom, dancing to Lou Breeze and his orchestra, and superb entertainment starring the well known comedian and television personality, Don Rice.

In addition to all this, Toni Trinchese, Ladies' Chairman, assures us that she has a special, exciting program planned, just for the ladies.

CONFIDENTIAL:

our custom catalyst service



When you come to Chemetron's Catalysts Division with a request for a custom catalyst, a highly productive dialog begins—behind closed doors.

You tell us what you want and when you want it. This information is restricted to the scientists and engineers who'll work on your catalyst-development project.

Would you want it any other way? Neither do we. You'll get full utilization of the Catalysts Division's know-how. Of our years of experience in catalyst research. Of our technical service which covers every facet of catalyst technology. Of our stringent standards of quality control and manufacturing integrity.

We appreciate the confidence you've placed in us. The proprietary data on the nature of your catalyst and your process will be treated with respect and discretion. We'll put it in writing.

That's why our Custom Catalyst Service is so successful. Arrange a rendezvous with us on your next catalyst project.

(Special note: The special catalyst you need may be one of our "T" series of experimental catalysts. In stock, in a variety of forms and sizes, these catalysts can be supplied in sample quantities for your evaluation. Write for a complete listing of our "T" series.)

Catalysts Division, Chemetron Corporation, P.O. Box 337, Louisville, Ky. 40201 Phone: (502) 366-9541

*Girdler-Südchemie Katalysator, GmbH, Munich
Nissan Chemetron Catalysts, Ltd., Tokyo*

CHEMETRON CORPORATION

Catalysts Division

CHEMETRON
TRADEMARK

• Four Corners . . .

(Continued from page 213A)

Technology of New Oil-bearing Materials

Exploratory work has been completed on the characteristics and processing of hitherto uninvestigated oil-bearing materials in India such as muskmelonseed, sweet orange seed, palm oil fruits and spent silk worm pupae. Pilot plant processing has been carried out on muskmelonseed and sweet orange seed. A centrifugal method of recovering palm oil using indigenous equipment has been developed. Spent silkworm pupae, produced to the extent of 7,000 tons per year, have been found to contain 25% of fat. Its extracted meal is high in protein content (70%), which is found suitable as substitute for fishmeal in poultry feed and as a sumptuous feed for sheep when mixed with molasses.

Regional Research Laboratory, Hyderabad

The Regional Research Laboratory is the only national laboratory of India which functions under the auspices of the Council of Scientific and Industrial Research, and in which research on oilseeds and oils, as well as on surface coatings, is conducted. Much attention is devoted to castor oil on which the following work is currently in progress: preparation of monoglycerides of over 90% purity by a solvent-glycerolysis technique, preparation and evaluation of direct ethylene oxide condensates of castor oil, development of a continuous method for alkali-fusion of castor oil to sebacic acid and 2-octanol and development of a one-stage process to obtain tristearin from castor oil.

In the area of hydrogenation, a study is in progress regarding the effect of modifying agents on the nickel-catalyzed hydrogenation of peanut oil, with particular reference to the course of the reaction. The presence of a trace of phosphoric acid greatly speeds up the saturation reaction without affecting its course. Considerable acyl migration has been shown to occur during commercial fat hydrogenation, and the presence of linolic moieties in the 1 and 3 positions facilitates hydrogenation.

A process for continuous conversion of fatty oils to fatty alcohols is under development in which comparatively low pressure of only 70 atm are used. The first seed oil of the Malpighiaceae family, *Hiptage benghalensis*, has shown a remarkable resemblance in fatty acid composition to castor oil, which belongs to the Euphorbiaceae family. In contrast to stearic oleate oil, *Garcinia indica*, the seed fat of *Garcinia xanthochymus* is found to be a palmitic oleate oil with some hexadecenoic acid.

Indo-Soviet Symposium on Natural Products

The second Indo-Soviet Symposium on the Chemistry of Natural Products (including Pharmacology) was organized by the Indo-Soviet Joint-Committee for Scientific Collaboration, under the auspices of the National Institute of Sciences of India, in New Delhi on February 1 to 6, 1970. It included the following papers relating to lipids.

Synthetic Studies on Optically Active Phosphoinositides, by N. A. Preobrazhensky from the Institute of Fine Chemical Technology, Moscow, USSR.

Two unusual Natural Fatty Materials (Kamalaseed Oil, i.e., *Mallotus philippinensis* and Seed Oil of Snake Gourd, i.e., *Trichosanthes anguina*) by M. R. Subbaram and co-workers, from the Regional Research Laboratory, Hyderabad-9, India.


The symposium was convened by S. Rangaswami of the Department of Chemistry, University of Delhi; abstracts of papers are available from S. Rajaraman, National Institute of Sciences of India, Bahadur Shah Zafar Marg, New Delhi-1.

Japan Teruzo Asahara

General Meeting

At a recent meeting in Tokyo, several oil chemists, who conducted research on fats and oils and on flavor chemistry

at Rutgers University, New Brunswick, New Jersey, gathered for the following portrait. S. S. Chang, AOCS President, who worked with them while they were in the United States, received the picture from the chemists.



Top row (left to right) Yoshiyuki Kawase, Nisshin Oil Mills, Ltd.; Tsukasa Kawada, Kao Soap Company, Ltd.; Yukinobu Murase, Asahi Denka Kogyo Company. Front row (left to right) Choichiro Hirai, Nihon University; Kosaku Yasuda, Nisshin Oil Mills; Yoshiyuki Toyama, Toyo University; Akio Kato, Government Chemical Industrial Research Institute.

Activities of the JOCS

—A biodegradation test method for nonionics is being studied by the Detergent Committee of JOCS. An interim report presented by the Technical and Materials Division of the Soap and Detergent Association will be considered as the prototype procedure for this study.

—A Preparatory Committee for the JOCS Joint Meeting with AOCS in 1972 was formed. The meeting is expected to be very successful, considering the intensive cooperation expected from both societies in the next two years. As part of this preparation many participants from Japan are expected to attend the coming AOCS-ISF Joint Meeting.

Statistics

Production of Soaps and Detergents in Japan for the Calendar Year of 1968 and 1969 (unit metric tons).

	1968	1969	ratio (%)
Soaps	154,263	152,633	99
Detergents	558,453	634,775	114
Total	712,716	787,408	110

Supply Plan of Fats and Oils in Fiscal Year of 1970 (unit thousand metric tons).

	Edible	Industrial	Export	Total
Import	837	324	23	1,184
Domestic production	208	77	24	309
Total	1,045	401	47	1,493

Annual Meeting of the Chemical Society of Japan

The 23rd Annual Meeting of the Chemical Society of Japan was held starting on April 3. Almost 10,000 registrants joined this meeting and over 2,000 papers were reported. Thirty-seven lectures by guest speakers included Solvation Effects on the Conformational Equilibria of Carbohydrate Substances, by R. U. Lemieux of the University of Alberta, Canada; Progress in the Chemistry of

Continuous Fatty Acid Hydrogenation successfully put on stream

LURGI
NEWS

Over 20 years experience in the construction of discontinuous hydrogenation plants have led to the development of a process for continuous hydrogenation.

Our initial plant is running since autumn 1969.

Capacity: 50 t/24 h

Crudestock: distilled and undistilled fatty acids of animal and vegetable origin.

Operating pressure: 25 – 28 atm. g.


Operating temperature: up to 220 °C

The results achieved on our trial plant have been surpassed in operation:

Catalyst savings up to 20% over discontinuous operation,

low hydrogen consumption,

iodine number reduction below 0.5.



We shall be glad to consult you:

US agents:

WINFIELD EQUIPMENT CO. INC.
19615, LAKE ROAD
CLEVELAND, OHIO 44116
Tel. EDISON 3-2460

**LURGI GESELLSCHAFT FÜR WÄRME-
UND CHEMOTECHEKNIK MBH**
6 FRANKFURT (MAIN) GERMANY

Subsidiaries in: Amsterdam · Brussels · Johannesburg · Madrid · Melbourne · Mexico D. F. · Milano · New York · Paris · Stockholm · Tokyo · Toronto · Vienna · Zurich

Antibiotics from Higher Plant by R. Tschesche of Universität Bonn, West Germany; and Effect of Solvents on the Rates and Routes of Oxidation Reactions, by G. E. Zaikov of the Academy of Science, USSR. Dr. Zaikov presented another paper, Contribution From Solvation to the Action of Inhibitors, at a panel discussion co-sponsored by the Japan Oil Chemists' Society (JOCS) and other three societies, on April 15 in Tokyo.

JOCS held its 16th Annual Meeting on April 4th and papers were presented as one of the sections of the 23rd Annual Meeting of the Society of Japan.

Spain Eduardo Vioque

The Spanish Society of Cosmetic Chemists, Sixth Congress of the International Federation of Societies of Cosmetic Chemists (I.F.S.C.C.)

The Spanish Society of Cosmetic Chemists, organizer of the next Congress of the I.F.S.C.C. to be held in Barcelona, September 20-25, 1970, invites all members of the Societies of Cosmetic Chemists who integrate the International Federation and all interested persons to present original unpublished papers. The papers may include any of the following subjects: (a) Color in cosmetics. (b) Physical chemistry in cosmetic. (c) Biochemistry and dermatology in cosmetic. (d) Practical aspects of the preparation of cosmetics. (e) Regulation in cosmetic standardization of raw materials.

In order to reward the best paper presented to this Congress, the I.F.S.C.C. has offered a prize of \$1000. Additional information may be obtained from the Sociedad Espanola de Quimicos Cosméticos, Mallorca 279, Barcelona (9) Spain.

The World Olive Oil Year

The World Olive Oil Year began on November 1, 1969 and will last until the next campaign in 1970. The idea for the celebration came from the International Olive Oil Council (COI) which will commemorate the 10th anniversary of its foundation. The COI and the OEI (Ibero-American Office of Education) are the only two international organizations with official residence in Madrid.

The COI was created in 1959 with the purpose of carrying out the goals of the International Convention of Olive Oil in 1956-1958. Lucien Denis, director of the COI, has devoted much energy to defend and promote the olive tree culture and its products, not only from an economic point of view, but also with regard to propaganda and scientific knowledge.

The President of the Spanish delegation is F. J. Villaure, general vice director of the International Organization of the Foreign Office. The same delegation has representatives from several cabinets (Agriculture, Finance and Commerce) and other organizations related to the production and commerce of olive oil and table oils, such as the Olive Tree Syndicate, the Institute for Foreign Propaganda of Olive Oil Products (IPEPO) and the Fats Institute.

During the International Olive Oil Year at Lucca, Italy, an International Congress on the Biological Value of Olive Oil was held October 10-12, 1969. Several other meetings have been held throughout the year in Madrid.

Among the many other activities planned is the celebration of World Day of the Olive Tree, similar to Harvesting Day for grapes, etc.

The National Meeting of Food Quality Control

The Second National Meeting on Food Quality Control took place in Madrid on July 7 and 8, 1969, as a contribution of the Spanish Association for Food Quality Control and the National Institute of Food Science and Technology (INCYTA) to the International Food Fair (AIDA 1969).

Martinez Moreno, president of the INCYTA, served as

chairman of the first section. The two main subjects were: Quality Control in the Oil Industry and its Application to Olives, by M. J. Fernandez and A. Garrido, both of the Fats Institute of Seville, and Quality Control in Margarine Industries, by E. Ruiz from Agra S.A., Bilbao.

The second section, with E. Primo, vice president of INCYTA, as the chairman, was dedicated to: Quality Control in Canned Goods, by L. Duran, from the Institute of Agrochemistry and Technology of Foods, and Industrial Aspects of Quality Control in Canned Foods, by V. Corell from CODESA.

Official Inauguration of Pilot Plants of the Fats Institute of Seville

One of the events included in the World Olive Oil Year was the inauguration of pilot plants, with the assistance of high personnel from the Spanish government and international organizations. Among those present were: the Spanish Minister of Commerce, the President of the High Council of Scientific Research, the Vice President of the Spanish Deputies' Assembly and the Director of the International Council of Olive Oil.

The Director of the Fats Institute of Seville indicated the following three main concerns of the pilot plants:

(a) The quality of olive oil; the first and fundamental reason for its presence in the market as well as the quality of table olives.

(b) The mechanization and automatization of olive oil extraction; not only from an economical point of view, but also from a social one, because the harvesting of olives is very difficult and therefore causes many problems in the olive tree regions.

(c) The training; instruction of technicians is one target of the Fats Institute. Some experience in this field has already been acquired, since short courses for those in the olive oil and table oil industries have been offered since 1960.

The Director of COI, Lucien Denis, said: "The inauguration today is of great importance for the solution of many problems in the olive oil industry. Important problems are due to the peculiar characteristics of olive oil and to its competition from other edible oils, whose prices can not be compared to that of olive oil. It is quite necessary for the olive industry to gain benefits from improvements obtained by the technicians directed to harvest, transport and store the olives, as well as those directed to extract and stock the oil."

During the ceremony, the Medalla Marqués de Acapulco was delivered to Manuel Ortega, Agriculture engineer, by the Assembly of Members of the Fats Institute of Seville. The award is for his relevant contributions of knowledge concerning the olive oil industry.

Fourteenth Biennial Meeting of the Real Sociedad Espanola de Fisica y Quimica

The Fourteenth Biennial Meeting of the Real Sociedad Espanola de Fisica y Quimica was held in Seville, September 28 to October 4, in the Faculty of Sciences of the University. Papers dealing with many different aspects of Chemistry were discussed. Among the papers of interest in the field of fats and oils were:

Chemical Transformations of Olives During Pickling: The Desoxyoleanolic Acid and Other Products in the Residual Lye, by F. J. Ribelles and J. M. Viguera.

Reaction Mechanism of the Alkaline Metal Catalyzed Transesterification, by M. P. Jordan and R. Martinez.

Secondary Products in the Acyloinic Condensation of Alifatic Fatty Acids, by M. Martinez and R. Martinez.

Ester Formation When Passing Oils Through Ionexchange Resins Which Have Been Regenerated in the Presence of Ethanol, by M. A. Albi, A. Vioque, and T. Albi.


Air Pollution by Organochlorine Pesticides IV: Interference of Polychlorides Biphenyl in the Gas Liquid Chroma-

(Continued on page 233A)

Beta carotene plus TENOX[®] antioxidants improve snacks.



**EASTMAN
FOOD PRODUCTS**



There's nothing like a rancid odor to turn off your potential customers. It's easy to help inhibit oxidative rancidity with our TENOX[®] Antioxidants. They're ready to help keep your snack foods fresh-smelling and appealing. Add beta carotene, and you've got a golden color with plenty of vitamin A.

We also supply vitamins A and E in a number of convenient dry and liquid forms and concentrates. And if you need an emulsifier to solve starch gel problems,

there's our wide range of distilled monoesters—including MYVEROL[®], MYVAPLEX[®], MYVATEX[®], and MYVACET[®] products.

That's our way of saying we've got a broad range of products for a host of modern foods. We'd like to prove just how broad our line is. Write for a brochure, or let our food experts go to work on your special requirements.

DPI Division,
EASTMAN CHEMICAL
PRODUCTS, INC., Kingsport,
Tennessee 37662.

ABSTRACTS

R. A. REINERS, Editor. ABSTRACTORS: N. E. Bednarczyk, J. E. Covey, J. G. Endres, J. Iavicoli, Sin'itiro Kawamura, F. A. Kummerow, E. G. Perkins, J. A. Thompson and R. W. Walker

• Fats and Oils

FATTY ACID COMPOSITION OF LIPID EXTRACTS OF A THERMOPHILIC BACILLUS SPECIES. H. H. Caron (Dept. Animal Sci., Agr. Exp. Sta., Auburn Univ., Auburn, Ala. 36830). *J. Bacteriol.* 101, 145-151 (1970). Fatty acids of C_{16} and C_{17} accounted for over 80% of the fatty acids of a thermophilic *Bacillus*. Under most conditions branched chain acids were more abundant than *n*-fatty acids. The unsaturated fatty acid content varied inversely with growth temperature but was never more than 14%. Increasing growth temperature from 40 to 60°C resulted in a three to four-fold increase in the ratio of *n*- to branched-chain C_{16} . Two *n*-hexadecenoic acids were found, their relative abundance was influenced by temperature.

LIPID COMPOSITION OF BACILLUS CEREUS DURING GROWTH AND SPOREULATION. D. R. Lang and D. G. Lundgren (Dept. Bacteriology and Botany, Biol. Res. Labs., Syracuse Univ., Syracuse, N.Y. 13210). *J. Bacteriol.* 101, 483-89 (1970). The lipid composition of *B. cereus* during growth and sporulation consisted of 30-40% neutral lipids and 60-70% phospholipid. The phospholipids were phosphatidylethanolamine and glycerol and diphosphatidyl glycerol. Also present was diglycosyl diglyceride and the alanine ester of phosphatidyl glycerol. Diphosphatidyl glycerol, which is difficult to extract in the vegetative and stationary growth phases, becomes more easily extractable during spore maturation when its cellular level increases. Phosphatidyl glycerol had a high turnover rate and accounted for 70% of the phospholipid synthesized during sporulation. Phosphatidyl ethanolamine, the major phospholipid, had a low turnover rate.

LIPIDS OF BACTERIODES MELANINOGENICUS. V. Rizza, Anne N. Tucker and D. C. White (Dept. Biochem., Univ. of Kentucky Med. Center, Lexington, Ky. 40506). *J. Bacteriol.* 101, 84-91 (1970). Four percent of the chloroform-methanol extractable lipids of *B. melaninogenicus* was neutral lipid which contained the vitamin K_2 isoprenologues K_2-35 , K_2-40 and K_2-45 . The rest of the extract consisted of phosphatidic acid, phosphatidyl-serine and -ethanolamine as well as ceramide phosphorylethanolamine, ceramide phosphorylglycerol and ceramide phosphorylglycerol phosphate.

THE EFFECTS OF PHOSPHOLIPIDS AND AMINO ACIDS ON THE STABILITY OF RAPESEED OIL. K. Babuchowski *et al.* *Zeszyty Nauk. Wyższej Szkoły Rolniczej Olsztynie* 25(689), 415-21 (1969). The phospholipids of rapeseed oil, in the form of technical lecithin, appreciably prolong the induction period when added to refined rapeseed oil at the level of 1.5%. Addition of 0.36 mg of acidic and basic amino acids per 100 g of oil along with the phospholipids had no effect on the stability of the oil. The important factor which lowers the stability of the oil following degumming and neutralization is removal of the phospholipids. (Rev. Franc. Corps Gras)

ISOMERIZATION OF METHYL ESTERS OF OLEIC AND ERUCIC ACIDS DURING HYDROGENATION. K. Modzelewska. *Prace Inst. Lab. Badawczych Przemysłu Spożywczego* 19(3), 417-43 (1969). The rates of hydrogen uptake by oleic and erucic acid molecules hydrogenated separately are nearly the same. The rate constants for formation of saturated acids are 3.7×10^{-3} mole/min and 3.8×10^{-3} mole/min, respectively. The formation of saturated acids follows zero order kinetics, which shows that the process depends only on diffusion. Although the degree of geometrical isomerization is nearly the same for both acids, the rate of isomerization is much higher for erucic acid. At equilibrium, the ratio of trans to cis isomers is 2:1 for both acids. (Rev. Franc. Corps Gras)

RECOVERY OF THE FAT FROM COCOA SHELLS BY EXTRACTION WITH ETHANOL. J. Salmonowicz *et al.* *Tłuszcze Jadalne* 14(1), 7-18 (1970). The amount of shell in the cocoa bean varies from 12 to 14%. The shell contains about 7% fat, 2.6%

theobromine, small amounts of caffeine, tannins, natural pigments and very active antioxidants. Cocoa shell fat, obtained by ethanolic extraction, is similar to cocoa butter in physical and chemical characteristics. As such, it can be refined and used as a cocoa butter substitute. Recovery of the fat from the miscella by crystallization, in addition to eliminating distillation, offers the possibility of recovery of other by-products. (Rev. Franc. Corps Gras)

DISTILLATION OF OLEIC ACID IN A STREAM OF HYDROGEN. G. M. Pavlov *et al.* *Izv. Vysshikh Uchebn. Zavedenii, Pishchevaya Tekhnol.* 1970, 71-73. Distillation of oleic acid can be carried out at atmospheric pressure in a stream of hydrogen. The rate of distillation depends on the temperature and on the rate of hydrogen flow. The best results are obtained at 230°C and a hydrogen flow of 5.5 l/min. (Rev. Franc. Corps Gras)

LIPID CONTENT OF THE BALTIC SPRAT. T. Krassowska *et al.* *Przemysł Spożywczy* 23(7-8), 340-41 (1969). *Spratus balticus* attains a maximum lipid content of about 17% in November and December. It contains only 6-7% fat in May and June. Seasonal variations occur only in the contents of moisture and fat. The amounts of protein and ash remain the same throughout the year. An inverse linear correlation exists between moisture and fat contents. Above 75% water, the sprat is too lean and not good for smoking. (Rev. Franc. Corps Gras)

STABILITY OF AN OIL BLEACHED WITH A CATION-EXCHANGED ACTIVATED EARTH. R. Guillaumin, J. F. Pertuisot and M-F. Bosquet (Lab., Inst. des Corps Gras, Paris). *Rev. Franc. Corps Gras* 17, 21-24 (1970). An oil may be adequately bleached while forming few conjugated double bond systems by using an activated earth with the protons replaced by Ca or Mg ions. In the present study, a peanut oil was bleached with 0.5% of an activated Montmorillonite earth containing Ca ions at 80°C for 30 minutes. This oil together with a control bleached with the same earth before exchange, was stored in sealed bottles in the light at room temperature (18-21°C). After 100 days, there were no significant differences (P.V., uv absorption, Kreis test) between the two oils. Organoleptic evaluations during the first five weeks also showed no differences.

SOVIET SUNFLOWER TECHNOLOGY. B. Solomon (Inst. des Corps Gras, Paris). *Rev. Franc. Corps Gras* 17, 25-37 (1970). The more important Russian articles on sunflower technology published since 1951 are reviewed. The subjects covered include treatment of the sunflower seeds, *i.e.*, cleaning, decortication, breaking, pressing, and extraction; storage of the crude oil; refining, including degumming, winterization, and neutralization; and hydrogenation of the oil. Flow diagrams for the winterizing operation and for the continuous hydration of vegetable oil phosphatides are included.

NUCLEAR MAGNETIC RESONANCE STUDIES OF INTERACTIONS OF PHOSPHOLIPIDS WITH CYCLIC ANTIBIOTICS. E. G. Finer, H. Hauser and D. Chapman (Unilever Res. Lab. Colworth/Welwyn, The Frythe, Welwyn, Hertfordshire, G.B.). *Chem. Phys. Lipids* 3, 386-93 (1969). The cyclic antibiotics alamethicin, valinomycin and gramicidin S dihydrochloride interact with aqueous dispersions of phospholipids (ox brain phosphatidylserine and egg yolk lecithin). A mixture of each of these antibiotics with the phospholipid dispersed in 0.2 M phosphate buffer (pH 8.0) forms tightly bound phospholipid-antibiotic aggregates, in which the N.M.R. spectrum shows that some sort of cooperative hydrophobic interaction has caused a reduction in the intramolecular motion of the alkyl chains of the phospholipids. In the absence of salt, the interaction of egg yolk lecithin with alamethicin and valinomycin is similar to that observed in phosphate buffer, whereas its interaction with gramicidin S dihydrochloride results in a different type of complex. The resultant particles are smaller and the phospholipid alkyl chains are less tightly bound, resulting in greater molecular motion of the phospholipid compared with egg yolk lecithin/alamethicin (or valinomycin) mixtures. The use of other techniques has shown that the multilamellar structure of the phospholipids is broken down on the addition of each of the antibiotics. These findings may be relevant to the mode of action of some cyclic antibiotics, to the mechanism of induced ion transport across


(Continued on page 223A)

HAHN LABORATORIES

Consulting and Analytical
Chemists

1111 Flora St. P.O. Box 1177 Columbia, S.C. 29202

SYSTEMS FOR OIL SEED PROCESSING



**COMPLETE
TURN-KEY
INSTALLATIONS**

**DESIGN
ENGINEERING
MANUFACTURING
GENERAL CONTRACTING
ERECTION & PROCESS ENGINEERS**

**BULK STORAGE
CONVEYING SYSTEMS
ELECTRICAL CONTROL SYSTEMS
PROCESS CONTROL SYSTEMS
LOADING & UNLOADING SYSTEMS**

Our great store of research, technical knowledge and experience accumulated in 60 years of building extraction equipment and systems can be brought to bear on your most stubborn processing problems. We'll welcome your inquiry and respond promptly. Or, ask for our Solvent Extraction Catalog, 08-10-DG.

SOLVENT EXTRACTION DIVISION











F074

THE FRENCH OIL MILL MACHINERY CO. • PIQUA, OHIO 45356

FILTREX

THE DIRECT SOLVENT THAT ELIMINATES

A FEW OF OUR FILTREX PLANT INSTALLATIONS



EXTRACTION PROCESS COSTLY PREPRESSING

FILTRATION EXTRACTION

FILTREX effectively combines the steps of extraction, filtration and washing. High extraction rates are obtained at low solvent ratios with correspondingly high miscella concentrations. Thorough washing with effective filtration insures low residual oil and solvent in the marc. Efficient solvent removal in the Marc Desolventizing System results in meal commercially free of solvent. FILTREX is designed for compactness of layout, accurately controlled with provisions for safety and ease of operation.

FILTREX plants now in operation on rice bran, soybeans, cottonseed, rapeseed, flaxseed, sunflower seed and various press cakes producing highest quality oils and meals.

Request detailed information about FILTREX processing of your particular material and plan to visit a FILTREX installation at your earliest convenience.

ADVANTAGES OF W & S FILTREX PLANTS

- Eliminates pre-pressing.
- Eliminates "fines" problem.
- Lowest solvent requirements.
- Lowest utilities consumption.
- Compact installation indoors or outdoors.
- Operated easily by one man.
- Highest quality oil and meal.

For NEW plants or plant CONVERSION, FILTREX is the best investment for processing ANY vegetable-oil bearing material . . . extraction efficiencies to 99.5%.

WURSTER & SANGER
DO BRASIL
COMMERCIO E INDUSTRIA LTDA.
Caixa Postal 7707
Sao Paulo, Brasil

Rep: Arturo Samudio & Cia. Ltda.
Calle 37 No. 44-40 Barranquilla, Colombia

LICENSEE:
Stork Smulders N.V.
Zeelantlaan 27-29
Utrecht, Holland



Rep: The E. J. Nell Company
P. O. Box 612 Manila, Philippines

WURSTER & SANGER, INC., Dept. 9, 164 W. 144th Street,
Chicago (Riverdale), Ill. 60627, CABLE: WURSANCHEM
A Division of Jacobs Engineering Co., Pasadena, Calif.—Linden, N.J.

(Continued from page 220A)

natural membranes observed with many cyclic antibiotics and also to the question of lipid-protein interactions and membrane organization in general.

NUCLEAR MAGNETIC RESONANCE SPECTRA OF SOME OCTADECADIENOIC ACIDS AND OF SOME METHYL CIS, CIS- AND TRANS, TRANS-OCTADECADIENOATES. F. D. Gunstone, M. Lie Ken Jie and R. T. Wall (Dept. of Chem., Univ., St. Andrews, Scotland). *Chem. Phys. Lipids* 3, 297-303 (1969). The NMR spectra of a number of octadecadienoic acids and of the derived cis,cis- and trans,trans-octadecadienoates have been studied using a 100 MHz instrument. Characteristic spectral features associated with the number of methylene groups between the two unsaturated centres are described. In diunsaturated acids and esters of the general formula $\text{CH}_2 \cdot (\text{CH}_2)_n \text{X} \cdot (\text{CH}_2)_m \cdot \text{X} \cdot (\text{CH}_2)_p \cdot \text{COOR}$ where X represents the unsaturated centre, those compounds can be differentiated by their NMR spectra.

SYNTHESIS OF GALACTOSYL GLYCERIDES AND RELATED LIPIDS. H. P. Wehrli and Y. Pomeranz (Kansas St. Univ., Dept. of Grain Sci. and Ind. and Crops Res. Div. Agr. Res. Ser., U.S. Dept. Agr.). *Chem. Phys. Lipids* 3, 357-370 (1969). Synthesis of natural monogalactosyl glycerides is reported. The synthesis involves acylation of the primary hydroxyl groups of 2,5-methylene-D-mannitol, cleavage of the mannitol moiety between C-3 and C-4 by lead tetraacetate, reduction of the resulting aldehyde, attachment of galactose by the Koenig-Knoor reaction, hydrolysis of the acetal, acylation of the hydroxyl group and hydrazinolysis of the acetylated glycolipids. A simplified procedure, in which the optical activity at C-2 of the glycerol moiety is lost, is also described.

ON THE NOMENCLATURE OF ASYMMETRICALLY SUBSTITUTED MYOINOSITOL DERIVATIVES WITH PARTICULAR REFERENCE TO PHOSPHATIDYLINOSITOL. B. A. Klyashchitskii, V. I. Shvets and N. A. Proebrazhenskii (The M. V. Lomonosov Instit. of Fine Chemical Tech., Moscow, USSR). *Chem. Phys. Lipids* 3, 393-400 (1969). A nomenclature based on the system of stereospecific numbering is proposed for optically active asymmetrically substituted myoinositols.

A METHOD FOR THE DIFFERENTIAL ANALYSIS OF MIXTURES OF ESTERIFIED AND FREE FATTY ACIDS. R. L. Glass and Susan W. Christopherson (Dept. Biochem., Univ. Minn., St. Paul, Minn. 55101). *Chem. Phys. Lipids* 3, 405-08 (1969). A procedure is described which permits the preferential methanolysis of esterified fatty acids in the presence of free fatty acids and the subsequent esterification of the free fatty acids. Analysis, by gas chromatography, of the mixture before and after esterification of the free fatty acids gives an accurate determination of the esterified and free fatty acids in the presence of each other.

THE PARTIAL SYNTHESIS OF SOME NATURALLY OCCURRING GLYCOSPHINGOLIPIDS WITH SPECIAL REFERENCE TO O- β -D-GALACTOSYL-(1 \rightarrow 4)-O- β -D-GALACTOSYL-(1 \rightarrow 1)-CERAMIDE. J. B. Hay and G. M. Gray (Lister Inst. of Preventive Med., London, S.W. 1). *Chem. Phys. Lipids* 3, 59-69 (1969). A procedure is described for the preparation of some naturally occurring glycosphingolipids by partial synthesis from natural ceramide (N-acyl-sphingosine). The ceramide was obtained in quantitative yield from sphingomyelin by the action of phospholipase C. It was converted to 3-O-benzoylceramide which was then condensed with the chosen acetobromosugar in the presence of mercuric cyanide. Details of the preparations of O- β -D-glucosyl-(1 \rightarrow 1)-ceramide, O- β -D-lactosyl-(1 \rightarrow 1)-ceramide and O- β -D-galactosyl-(1 \rightarrow 4)-O- β -D-galactosyl-(1 \rightarrow 1)-ceramide are given.

GAS PHASE ANALYTICAL SEPARATION AND STRUCTURAL STUDY OF CERAMIDES. G. Casparrini, E. C. Horning and M. G. Horning (Inst. Lipid Res., Baylor Univ. College of Med., Houston, Texas 77025). *Chem. Phys. Lipid* 3, 1-10 (1969). Ceramides may be subjected to direct separation and structural study by gas phase analytical methods. Suitable derivatives may be prepared by reaction with silylating reagents (leading to trimethylsilyl ether derivatives), and separations may be carried out with 1% SE-30 columns by temperature programming at 2°C/min to 340-350°C. The structure of individual ceramides is indicated by their mass spectra; both the acyl group and the sphingosine base moiety can be recognized in this way. A STUDY OF PALMITIC-STEARIC TRIGLYCERIDES AND THEIR BINARY MIXTURES BY DIFFERENTIAL THERMAL ANALYSIS (DTA). I. PURE TRIGLYCERIDES. R. Perron, J. Petit and A. Mathieu

(Continued on page 226A)

Speculative Open Interest in Soybean Futures

D. M. BARTHOLOMEW, Commodity Analyst, Merrill Lynch, Pierce, Fenner & Smith Inc.

Open interest is one of the important indicators used in analysis of technical aspects of futures markets. To some degree it is related to fundamental conditions, i.e., large production may be expected to reflect large open interest if at the same time substantial price movement is possible. Conversely, large production may result in small open interest if price fluctuation is anticipated to be small due to government price support mechanisms, for example.

Beyond this relationship to fundamentals, however, open interest is primarily to technical factor. By definition, open interest is a measure of the size of unliquidated long and short positions in the futures market. For soybeans and grains it is expressed in number of bushels. For most other commodities, including soybean oil and soybean meal, it is expressed in number of contracts. The figures for the end of trade on a given day are released on the following market day. Sometimes this report may influence trading activity depending on the magnitude of change in open interest.

For the open interest to show an increase there must be both new long and new short positions established. Likewise for the open interest to show a decrease there must be both liquidation of longs and covering of shorts. The open interest would not change if, for example, there were new purchases made from sellers who were liquidating a long position. Likewise, open interest would not change if new short positions were sold to previous shorts who were covering their position. This explains one reason why volume of trade is usually larger than open interest change.

Volume of trade is another technical factor which is usually considered in conjunction with open interest. Volume figures are released with the open interest. Another obvious reason why volume is much larger than open interest changes at the end of the day is because of the participation of "day traders." These are speculators who initiate and liquidate positions several times during the day and seldom carry a position over night. It is the "day traders" who provide the necessary liquidity to futures markets so that commercial trade interests may always find a buyer when they are ready to sell, or a seller when they are ready to buy. (Volume is of no significance in this study, but is mentioned by way of explanation.)

Open interest changes at the end of the day are watched for a signal of change in trader confidence in the market, but one day's results must not be afforded too much significance unless the change is usually large. On the other hand, several days' change in the same direction merits considerable attention. Such a situation represents collective attitude which cannot be ignored.

Once a month the government releases a report which shows types of traders holding open positions in the market. This is analyzed by the Commodity Exchange Authority (CEA) and represents a profile of the open interest in the regulated commodity futures. It is usually released about the eleventh of the month for the close of business on the last day of the preceding month. It shows long and short positions of large speculators, large hedgers, large spreaders and small speculators and hedgers combined. By observing changes in these categories from the previous month, some judgment may be formed as to change in market sentiment by the various participants. Of course, price action must also be related to these changes. Furthermore, market action must be observed from the first of the month until the date the report is released. The market profile may have changed significantly during the intervening time.

The purpose of this study is to see if a relationship exists between the market position of large speculators and subsequent price action. Large speculators are considered to be representative of the market sentiment of speculators in general because small speculators frequently

(Continued on page 231A)




Fig. 1. 1968-69 Soybean Oil




Fig. 2. 1968-69 Soybean Meal



Enter the ageless age

With Sustane[®]—the proven antioxidant that keeps your foods and food tastes alive longer.

Nothing turns off a customer faster than rancid foods. And, she'll never forget the brand name of the offender.

She doesn't know or care about "shelf life"—that's *your* problem. And that's where we lend a helping hand—with Sustane BHA—knowing freshness and long shelf life are musts for food processors and packagers.

UOP pioneered and developed BHA, and trademarked it Sustane. For cooking oils and other food products, Sustane survives the rigors of processing, baking and frying. Its effectiveness extends into plastic films, wax coatings and other modern food packaging materials. The result: Better protected and safer products—for you and your customers.

For information on the many Sustane formulations that can introduce your products to the Ageless Age, write: UOP Chemicals, East Rutherford, N.J. 07073, U.S.A.
Or phone 201/438-7800.

UOP Chemical is a division of Universal Oil Products Company, a half-billion dollar company busy making life better through research and product innovation across the world.



UOP CHEMICALS

BHA is quality protection;
SUSTANE is quality BHA.

(Continued from page 223A)

(Groupe de Lab. du C.N.R.S., 2 à 8, Rue Henry Dunant, 94, Thiais, France). *Chem. Phys. Lipids* 3, 11-28 (1969). The cryothermograms of the six pure triglycerides derived from palmitic and stearic acids have been determined by DTA. As well as revealing itself to be an excellent method of distinguishing between the positional isomers of the mixed triglycerides, DTA also showed that cooling triglycerides at a rate of 1.2C/min yielded an α_L form (except PSP), which on subsequent heating gave the β_L form with the symmetrical triglycerides, and the β'_L form with the non-symmetrical triglycerides. In the process of cooling melted triglycerides, various forms may appear depending on the rate of cooling. The symmetrical triglycerides, with the exception of SSS, give stable forms more easily than the non-symmetrical triglycerides. Further, a sub- β'_L form can be detected in the case of PSP, PSS and SPP.

LIPIDS OF STREPTOMYCES SIOYAENSIS. V: ON THE 2-HYDROXY-13-METHYL-TETRADECANOIC ACID FROM PHOSPHATIDYLETHANOLAMINE. J. Kawanami, A. Kimura, Y. Nakagawa and H. Otsuka (Shionogi Res. Lab., Shionogi and Co., Ltd., Fukushima-ku, Osaka, Japan). *Chem. Phys. Lipids* 3, 29-38 (1969). Phosphatidylethanolamine from *St. sioyaensis* afforded a double spot on a thin-layer chromatogram, typical of most glycosphingolipids from animal tissue. They were phosphatidylethanolamines one of which had only non-hydroxylated fatty acids and the other hydroxy fatty acids in addition to non-hydroxylated fatty acids, respectively. The distribution of the fatty acids was studied by hydrolysis with snake venom phospholipase A. Hydroxy fatty acids were located in the β -position of the glycerol moiety, differing from the results for *Brucella abortus* phospholipids in which location in the α -position has been reported. The main hydroxy fatty acid was purified by preparative gas-liquid chromatography. The structure of the hydroxy fatty acid was analyzed by oxidation with lead tetraacetate, proton magnetic resonance and mass

spectrometry, etc. From these results, it was shown that the main acid was 2-hydroxy-13-methyltetradecanoic acid.

FATTY ACIDS. PART 19. CONVERSION OF ALKENOIC ACIDS TO ALKYNIC ACIDS BY BROMINATION-DEHYDROBROMINATION. F. D. Gunstone and G. M. Hornby (Dept. Chem., Univ. St. Andrews, North Haugh, St. Andrews, Scotland). *Chem. Phys. Lipids* 3, 91-7 (1969). Alkynoic acids (including octadec-10-ynoic, hendec-10-ynoic, and 12-hydroxy-octadec-9-ynoic) can be prepared from the cis alkenoic acids by bromination followed by dehydrobromination with sodium in liquid ammonia or with DBU (1,5-diazabicyclo(5.4.0)undec-5-ene). With other bases extensive migration of the unsaturated centre was observed and no satisfactory procedure for converting trans alkenoic acids to alkynoic acids without migration was discovered. Both types of alkenoic acids could be converted to ene-bromides, sometimes in high yield, with DBU and DBN (1,5-diazabicyclo(4.3.0)non-5-ene).

PHOSPHOLIPIDS OF MARINE INVERTEBRATES. V. E. Vaskovsky and E. Y. Kostetsky (Inst. Biol. Active Substances, Siberian Dept. of the Acad. of Sci. of the URRS, Vladivostok 22, USSR). *Chem. Phys. Lipids* 3, 102-5 (1969). The quantitative and qualitative phospholipid composition is reported for all the main phyla of the marine animals. No simple correlation was found between the phospholipid composition and taxonomic system of marine animals. Unusual phospholipids were shown in a great number of invertebrates.

CHARACTERIZATION OF THE STRUCTURE OF A 4-METHYL- $\Delta^{8,24}$ -CHOLESTADIEN-3 β -OL ISOLATED FROM RAT SKIN. A. Sanghvi (Dept. of Biochem., College of Med. Sci., Univ. of Minnesota, Minneapolis, Minn. 55455). *J. Lipid Res.* 11, 124-30 (1970). A new sterol has been isolated from the skin of rats treated with triparanol. Its chromatographic behavior on silicic acid-Celite columns and in gas-liquid chromatographic systems indicated it to be a 4-methyl- $\Delta^{8,24}$ -cholestadien-3 β -ol. The specific rotation, the delayed color reaction with Liebermann-Burchard reagent and the nuclear magnetic resonance (NMR) data support the $\Delta^{8(9)}$ -unsaturation. Previous workers have shown that triparanol treatment results in an accumulation of Δ^{24} -unsaturated sterols in animal tissues. Consonant with this observation, the infrared, NMR and mass spectrometric data confirm the presence of a C-24(25) unsaturated side chain in this sterol.

Short Course on Processing and Quality Control of Fats and Oils

Are you up-to-date in the principles, practices and latest innovations in the processing of edible oils? If you "do your thing" in processing (are you a dial twister?) or quality control (will that customer accept our last tank of oil?), you will be happy to hear that plans for the next AOCS Short Course on Processing and Quality Control of Fats and Oils are well underway. This very popular Short Course, last presented in 1966, will be held September 23 through 25 at Michigan State University, the week before the Joint AOCS-ISF Meeting in Chicago.

The objectives of this Short Course will be to provide each participant with:

- Review and/or new information on the chemistry and physics of edible oils which are pertinent to an understanding of their processing;
- fundamental principles and commercial practices in all of the major edible oil unit operations; and
- principles and practices in statistical quality control and involuntary operations.

These topics will be covered by recognized industrial experts representing food, consulting and equipment companies. Wherever possible, latest innovations in particular fields will be presented, with emphasis on continuous processing. An evening session devoted to the impact of federal regulations on refinery operations is also planned. This course will be of particular value to technical people who are new to the edible oil industry and will also serve as an excellent refresher for our most experienced people.

Co-chairmen for this Short Course are Leroy Dugan (arrangements), Michigan State University, and Bob Hlavacek (program), Hunt Wesson Foods. Registration, including meals and lodging will be \$140.00 for the three-day course. Early reservations may be directed to Dr. Dugan, Michigan State University, East Lansing, Michigan 48823. A semi-detailed program will be available in the July issue of the journal. Please watch for this announcement.

GEL PERMEATION CHROMATOGRAPHY OF NEUTRAL HYDROXY LIPIDS ON SEPHADEX LH-20. M. Calderon and W. J. Baumann (Univ. of Minn., The Hormel Inst., Austin, Minn. 55912). *J. Lipid Res.* 11, 167-69 (1970). Gel-permeation chromatography on Sephadex LH-20, using ethanol as eluent, permits the resolution of neutral hydroxy lipids according to molecular size. The influence of molecular shape, functional groups, chain lengths and degree of unsaturation, as well as the effect of the eluent on the elution pattern are discussed. The usefulness of the method for the separation of classes of hydroxy lipids, which cannot be resolved by other chromatographic procedures, is demonstrated. Examples include the separations of 1,2- and 1,3-diglycerides from long-chain alcohols and of alkyl ethanediol monoethers from cholesterol.

FLUORIMETRIC DETERMINATION OF SPHINGOSINE AND ITS APPLICATION TO NATURAL MIXTURES OF GLYCOSPHINGOLIPIDS. L. Coles and G. M. Gray (The Lister Inst. of Preventive Med., Chelsea Bridge Road, London, S. W. 1, Eng.). *J. Lipid Res.* 11, 164-66 (1970). A sensitive estimation of sphingosine, by measurement of the fluorescence of a complex formed with 1-naphthylamino-4-sulfonic acid, is described. The practical range is 5-35 nmoles sphingosine. The method is used to estimate, in terms of sphingosine, amounts of ceramide and glycosphingolipids. The isolation of microamounts (5-30 μ g) of individual glycosphingolipids from a mixture and their quantitative estimation is described. The percentage composition of a glycosphingolipid mixture from the kidneys of adult C57/BL male mice is given.

GAS-LIQUID CHROMATOGRAPHY-MASS SPECTROMETRY OF SYNTHETIC CERAMIDES CONTAINING 2-HYDROXY ACIDS. S. Hammarstrom, B. Samuelsson and Karin Samuelsson (Dept. Med. Chem., Royal Vet. College, Dept. Neurol., Karolinska sjukhuset, Stockholm, Sweden). *J. Lipid Res.*, 11, 150-57 (1970). Ceramides containing either sphingosine or sphinganine and one of the 2-hydroxy acids, 14h:0, 16h:0, 18h:0, 20h:0, 22h:0, 24h:0, and 26h:0 were prepared and separated by gas chromatography as the 1,3,2'-tri-O-trimethylsilyl derivatives. Mass spectrometric analyses of these derivatives showed that the

(Continued on page 228A)

AOCS-ISF Announces Computer Symposium for Chemists

A computer symposium designed especially for chemists includes descriptions of computer systems and applications in various chemical fields. The symposium is part of the AOCS-ISF World Congress to be held September 27-October 1, 1970, at the Conrad Hilton Hotel in Chicago, Illinois, USA.


Computer systems from programmable calculators and desk top computers to time-sharing with a large computer complex and including dedicated laboratory computer systems will be described for chemists by authorities in these fields. Applications in spectroscopy, chromatography, fatty acids, foods, pesticides, lipids and other areas will be presented by individual researchers from Australia, Europe and North America. Data acquisition, instrument interfacing and laboratory automation will be discussed.

The symposium is designed for the chemist who is contemplating the use of computers or who may be using one computer system and needs organized, factual information presented in a concise manner to enable him to choose the best solutions for his individual requirements. The World Congress is sponsored jointly by the American Oil Chemists' Society and the International Society for Fat Research.

Abstracts of the symposium papers are available prior to the meeting. Those interested in attending should contact the symposium chairman, Leonard H. Ponder, Research Chemist, American Enka Corporation, Enka, North Carolina 28728, USA.

The program will include:

- Introduction to Modern Computers: Development, Terminology, Uses, James Brown, Managing Editor, Auerbach Standard EDP Reports.
- An Off-Line System for Data Acquisition and Analysis, R. C. Master, Engineering Manager, Analytical Chemistry, Nuclear Data.
- Time Sharing Today, J. G. Rude, Chairman of the Board, Call-A-Computer.
- The Desk Top Computer: I. General Capabilities; II. Applications in a Clinical Chemistry Laboratory, W. R. Dito, Director of Laboratories, Pontiac General Hospital.
- Application of a Desk Top Computer to Chemistry, A. P. Damoglou, Research Scientist, CSIRO (Australia).
- A Computer Approach Toward Automation of a Chemical Services Laboratory: I. Instrument Support, E. L. Schneider, Manager, Ralston Purina Company, and Arvid Munson.
- A Computer Approach Toward Automation of a Chemical Services Laboratory: II. Management Information, Arvid Munson, Manager, Ralston Purina Company, and E. L. Schneider.
- A Computerized Search System for Infrared Spectral Data, B. M. Vasta, Chief, Chemical Compound Information, FDA.
- Use of the Computer as a Technical Service Tool in Pesticide Formulations, B. F. Fay, Laboratory Supervisor, Atlas Chemical Industries.
- Computer Processing of Gas Chromatographic Data: Analysis of Fatty Acid Mixtures Using an Off-Line Magnetic Tape System, G. J. Nelson, Lawrence Radiation Laboratory.
- Collection, Storage and Processing Gas Chromatographic Data by Digital Integrator-Paper Tape Punch Computer System, H. E. Pattee, Research Chemist, Market Quality Research, USDA, and J. A. Singleton, Chemist.
- Digital Registration on Magnetic Tape of Data From 15 Gas Chromatographs, F. Woutman, Chemist, AKZO Research Laboratories, (The Netherlands).
- Applications of Nonlinear Programming to Detergent Formulations, E. C. Steinle, Project Scientist, Union Carbide Corporation, C. D. Hendrix, Research Scientist, and R. R. Fields, Chemist.
- Improved Software for GC Automation Via the Time Share Computer, G. D. Dupre, J. M. Gill and J. R. Hubbard, Vidar Corporation.



Meetings

AOCS National Meetings

Sept. 27-Oct. 1, 1970—Chicago, Conrad Hilton Hotel.

May 2-6, 1971—Houston, Shamrock Hotel.

Oct. 2-6, 1971—Atlantic City, Chalfonte-Haddon Hall Hotel.

AOCS Section Meetings

Northeast Section—June 2, 1970, Whyte's Restaurant, New York.

Other Organizations

June 7-12, 1970—Fourth ISA Process Analytical Instrumentation Short Course, Temple Buell College, Denver, Colorado.

June 9-12, 1970—14th International Conference on the Biochemistry of Lipids, Lund, Sweden.

June 21-26, 1970—73rd Annual Meeting of the American Society for Testing and Materials, Royal York Hotel, Toronto, Canada.

June 23-25, 1970—Fourth International Sunflower Conference, Sheraton-Peabody Hotel, Memphis, Tenn.

June 22-27, 1970—14th International Congress of Esthetics and Cosmetology, Amsterdam, The Netherlands.

July 7-9, 1970—International Association of Seed Crushers, the Royal Garden Hotel, London, England.

July 26-August 1, 1970—5th International Water Pollution Research Conference, San Francisco, California.

Aug. 9-14, 1970—Third International Congress of Food Science and Technology, Washington, D.C.

Aug. 23-25, 1970—41st Annual Meeting of the National Soybean Processors Association, Fairmont Hotel, San Francisco, Calif.

Sept. 20-23, 1970—International Conference on the Science, Technology and Marketing of Rapeseed and Rapeseed Products, Chantecler Hotel, St. Adele, Quebec.

Oct. 11-14, 1970—Ninth Annual Meeting, ASTM Committee E-19 on Chromatography, Brown Palace Hotel Denver, Colorado.

Oct. 14-17, 1970—International Symposium on Computer Applications in Engineering Sciences, Istanbul Technical University, Istanbul, Turkey.

Oct. 12-15, 1970—84th Annual Meeting of the Association of Official Analytical Chemists, Marriott Motor Hotel, Twin Bridges, Washington, D.C.

Oct. 26-29, 1970—ISA 25th Annual Conference and Exhibit on Instrumentation, Systems and Automatic Control, Civic Center, Philadelphia, Pa.

* Oct. 26-28, 1970—17th Spectroscopy Symposium and Exhibition of Instrumentation, Skyline Hotel, Ottawa, Ontario, Canada.

* Oct. 26-30, 1970—Fourth Materials Research Symposium, National Bureau of Standards, U.S. Department of Commerce, Gaithersburg, Md.

* Nov. 2-4, 1970—Technicon International Congress on Automated Analysis, New York Hilton Hotel, New York.

*Additions to previous calendar.

(Continued from page 226A)

ions formed on electron impact can be used to determine unequivocally the structures of the long-chain base and the fatty acid residue in the ceramide. Proposed structures of ions and the mechanisms of reaction of their formation are supported by mass spectra of homologous derivatives, by deuterium labeling experiments and by high-resolution mass spectrometry.

SPECTROPHOTOMETRIC DETERMINATION OF PROTEIN AND FAT IN MILK SIMULTANEOUSLY. S. Nakai and Anh Chi Le (Dept. of Food Sci., Univ. of British Columbia, Vancouver 8, B.C., Canada). *J. Dairy Sci.* 53, 276-78 (1970). A clear solution was obtained by adding 5 ml of 97% acetic acid to 0.05 ml of whole milk. The protein was calculated from the absorbance at 280 m μ . Turbidity depending on fat content was developed thereafter by adding 2.5 ml of a solution containing 20% urea and 0.2% imidazole. The fat was determined from absorbance at 400 m μ with a round cuvette. This method does not require prehomogenization of milk for turbidity measurement, because of a dispersing effect by the combination of reagents. Application of this method to other food products is suggested.

THE GLYCERIDE STRUCTURE OF SAPIUM SEBIFERUM SEED OIL. W. W. Christie (Hannah Dairy Res. Inst., Ayr, Great Britain). *Biochim. Biophys. Acta* 187, 1-5 (1969). *S. sebiferum* seed oil is known to contain some tetraester triglycerides. The oil was separated by preparative thin-layer chromatography into normal triglycerides (76.9%) and estolide (23.1%) components which were each subjected to stereospecific analysis procedures. In the normal triglycerides, saturated and monoenoic fatty acids were in greatest abundance in the 1-position, while linoleic and linolenic acids (18:3 ω 3) were in greatest abundance in the 2- and 3-positions, respectively. The estolide fatty acids were entirely in the 3-position of the tetraester fraction. The fatty acid compositions of the 1- and 2-positions of the two components of the oil were very similar.

THE EFFECT OF TECHNOLOGICAL PROCESSING ON THE TOCOPHEROL CONTENT OF RAPESEED OIL. A. Rutkowski and L. Mzyk (Univ. of Agr., Olsztyn, Poland). *Riv. Ital. Sostanze Grasse* 46, 614-6 (1969). The changes in α and γ tocopherol content during industrial extraction and refining of rapeseed oil have been investigated. The results show that refining causes a decrease of about 75% of the tocopherol content initially present in the crude oil, with the largest losses occurring during alkali refining and during bleaching. The ratio of α to γ tocopherol does not change significantly during the refining processes. Thus relative losses of α and γ tocopherols are the same. The concentration of tocopherols in the deodorization condensate is higher than in other waste products of the refining processes. In all these waste products the ratio of α to γ tocopherol is equal to that in the crude oil.

ISOLATION AND CHARACTERIZATION OF THE HYDROPEROXIDES OF METHYL OLEATE, II. I.R. CHARACTERIZATION. M. Piretti (Univ. of Bologna, Bologna, Italy). *Riv. Ital. Sostanze Grasse* 46, 591-601 (1969). The I.R. spectra of methyl oleate hydroperoxide, reduced methyl oleate hydroperoxide and methyl oleate have been obtained in the interval of 2 to 15 μ . Within the range of applicability of Lambert and Beer's law, the extinction coefficients of methyl oleate hydroperoxide have been calculated at $\lambda = 2.82 \mu$ (free OH) and at $\lambda = 2.91 \mu$ (bound OH) in CCl₄ and at $\lambda = 2.85 \mu$ (free OH) in CS₂. The experimental results suggest the possible existence of intramolecular hydrogen bonds in methyl oleate hydroperoxide and in its reduction product.

DEGRADATION OF LINOLEIC ACID DURING FRYING. L. Kilgore and M. Bailey (Mississippi State Univ.). *J. Am. Dietetic Assoc.* 56, 130-2 (1970). The decrease in linoleic acid content of fats used for frying has been studied on samples of safflower oil, cottonseed oil, corn oil and a commercial shortening advertised as being highly unsaturated. The percentage of

linoleic acid (based on total fatty acids) for the fresh fats was: safflower, 72%; corn, 57.2%; cottonseed, 55.5%; and shortening, 30.2%. After the fats had been used for intermittent frying periods totaling 7½ hrs., during which 10 lbs. of potatoes were fried, the percentages of linoleic were: safflower, 69.2%; corn, 51.6%; cottonseed, 49.0%; and shortening, 26.7%. The linoleic acid content of the fat extracted from the tenth pound of potatoes fried was equal to slightly lower than the content in the oil at the end of the 7½ hrs.

NEW FRACTIONATION PROCEDURES IN FATS AND OILS TECHNOLOGY. E. Bernardini and M. Bernardini (Costruzioni Meccaniche Bernardini, Pomezia, Italy). *Riv. Ital. Sostanze Grasse* 46, 607-13 (1969). A novel solvent fractionation process for fats and oils is described, consisting mainly of a solvent phase crystallization with a final filtration, yielding three fractions: one with relatively high I.V., another with relatively low I.V. and an intermediate recycle fraction. Operational data and process information are given.

FATTY ACIDS IN NEWER BRANDS OF MARGARINE. P. Miljanich and R. Ostwald (Univ. of Calif., Berkeley, Calif.). *J. Am. Dietetic Assoc.* 56, 29-30 (1970). The composition of some newer margarine products has been determined. This information is as yet unavailable on the labels of these products and should be useful to dietitians and consumers concerned with their dietary intake of polyunsaturated fatty acids.

FURTHER INVESTIGATIONS ON LEGUMINOSAE SEED OILS. V. Averna, G. Lotti and F. P. Tartaglia (Univ. of Palermo, Palermo, Italy). *Riv. Ital. Sostanze Grasse* 46, 602-6 (1969). The oils extracted from the seeds of herbaceous and arboreal Leguminosae were examined. The data reported, on a total of 15 botanical species, include chemical analysis, fatty acid composition, U.V. and I.R. absorption characteristics.

PREPARATION OF PURE FATTY ACIDS. E. Fedeli, F. Camurati and A. Lanzani (Exper. Stat. for Fats and Oils, Milan, Italy). *Riv. Ital. Sostanze Grasse* 46, 514-9 (1969). The results of experiments on the preparation of high purity fatty acids are discussed. The methods used have been mainly physical (distillation, crystallization, chromatography) to avoid the isomerization which often accompanies chemical treatment of fatty acids.

PREPARATION OF COLORED FATTY COMPOSITION. G. W. Brankamp (Procter & Gamble Co.). *U.S. 3,489,573*. Fats and oils are provided with heat-stable color by incorporating in them water-soluble dyes in combination with particular polyglycerol esters, such as decaglycerol trilinoleate and tetraglycerol monooleate.

HYDROGENATION OF UNSATURATED ALIPHATIC COMPOUNDS. L. P. van't Hof (Lever Bros. Co.). *U.S. 3,489,778*. Unsaturated aliphatic compounds, especially soybean oil, are catalytically hydrogenated by contacting with hydrogen in the presence of a solution of a platinum compound, for instance chloroplatinic acid, and a stannous halide, for instance stannous chloride, in a solvent consisting of a lower dialkyl ether, dialkyl ketone or aliphatic carboxylic acid or its ester.

FATTY EMULSIONS AND THEIR PREPARATION. D. P. J. Moran (Lever Bros. Co.). *U.S. 3,490,919*. Emulsions of edible fats are described, having a semi-solid plastic fat as a continuous phase and a stabilized dispersion of liquid oil in an aqueous medium as the disperse phase.

COOKED SWEET CORN FLAVORING COMPOSITION FOR VEGETABLE OILS. D. Melnick and H. L. Zmachinski (Corn Products Co.). *U.S. 3,490,921*. A flavoring composition which, when added to a liquid vegetable oil imparts to it the flavor of freshly cooked sweet corn, comprises a combination of 1.0 to 4.0 parts by wt. of ethyl vanillin and 1 part by wt. of ethyl n-butyrate. The flavoring composition should be added to the vegetable oil in an amount of from about 30 parts per billion to about 400 parts per billion. In addition, if it is desired to add a buttery flavor to the flavored oil, diacetyl, in an amount of at least 90 parts per billion, may be added.

GLYCERIDE OIL TREATMENT. R. A. Reiners and F. J. Birkhaug (Corn Products Co.). *U.S. 3,491,132*. A method is described for reducing the free fatty acid levels of glyceride oils so that they may be later efficiently refined, the method consisting of combining a cyclodextrin with oil and water, breaking the mixture so formed and obtaining the oil with the desirably reduced levels of free fatty acids. The cyclodextrin is recovered by decomposing the clathrate of the cyclodextrin and the fatty acid.

POPE TESTING LABORATORIES, INC.

Analytical Chemists

2618½ Main

P.O. Box 903

Dallas, Tex.

HARD BUTTER COMPOSITIONS AND METHOD OF PRODUCTION. J. Harwood (SCM Corp.). *U.S. 3,492,130*. Compositions have been found which comprise mixtures of certain symmetrical and asymmetrical mono-unsaturated triglycerides and asymmetrical di-unsaturated triglycerides of domestic origin and which are temperable to a stable triple chain length beta crystalline form. The term 'symmetrical glyceride' refers to positional isomer symmetry of the unsaturated acyl groups rather than the kind or type of saturated acyl groups present. When tempered, the compositions are compatible with cocoa butter and can be used as extenders or substitutes for it. Cocoa butter itself is temperable to the stable triple chain length beta crystalline form, but naturally occurring domestic glycerides generally temper to a double chain length or beta prime crystalline form or a mixture of beta prime and beta crystalline forms and when in such forms are not compatible with cocoa butter.

PROCESS FOR VACUUM DISTILLING RANDOMLY INTERESTERIFIED TRIGLYCERIDES TO PRODUCE NOVEL TRIGLYCERIDE FRACTIONS. P. Seiden (Procter & Gamble Co.). *U.S. 3,494,944*. Hardened lauric acid oils are randomly rearranged, or randomly interesterified, with hardened non-lauric acid oils containing predominantly C₁₆ and higher saturated fatty acids. The hardened randomized oils are distilled to produce fractions and residues useful in hard butter and margarine oil formulations.

STABILIZATION OF OXIRANE CONTAINING FATTY DERIVATIVES. R. J. Sims (Swift & Co.). *U.S. 3,497,531*. Fatty, oxirane-containing compounds are stabilized against loss of oxirane oxygen by the presence of phenolic fat antioxidants such as propyl gallate.


STABILIZATION OF FATS AND OILS WITH EDTA AND RELATED COMPOUNDS. W. J. Lennon (Geigy Chemical Corp.). *U.S. 3,497,535*. Fats and oils which are subject to oxidative deterioration are stabilized by means of an effective amount of an antioxidant and from about 0.05 to about 1000 parts per million by wt. of esters of EDTA as chelating agents.

• Biochemistry and Nutrition

THE BIOSYNTHESIS OF $\Delta^{9,12,15,18}$ -TETRACOSATETRAENOIC AND OF $\Delta^{6,9,12,15,18}$ -TETRACOSAPENTAENOIC ACIDS BY RAT TESTES. R. B. Bridges and J. G. Coniglio (Dept. of Biochem., Vanderbilt Univ., Nashville, Tenn. 37203). *J. Biol. Chem.* 245, 46-9 (1970). Two 24-carbon polyenes previously reported in rat testes have been isolated and characterized to be $\Delta^{9,12,15,18}$ -tetracosatetraenoic and $\Delta^{6,9,12,15,18}$ -tetracosapentaenoic acids. The compounds were separated and purified by gas-liquid chromatography, and their structures were established by use of gas-liquid chromatography before and after hydrogenation, by ultraviolet spectroscopy of the alkaline isomerized derivatives and by identification of fragments resulting from cleavage at double bonds by oxidative and reductive ozonolyses. The biosynthesis of the 24-carbon polyenes was studied after intratesticular injections of either 1^{14}C -linoleate or 1^{14}C -arachidonate. The two metabolic products, individually, were cleaved by oxidative ozonolysis, and the location of the ^{14}C in the molecule was determined by gas-liquid radiochromatography of the fragments. The distribution of the ^{14}C was consistent with the hypothesis that the 24-carbon polyenoic acids were biosynthesized by elongation and further desaturation of linoleic acid. Final proof was obtained by chemical carbon by carbon degradation and measurement of the specific activity of individual carbon atoms. After 1^{14}C -linoleate injection, $\Delta^{6,9,12,15,18}$ -tetracosapentaenoic acid from testicular tissue was labeled primarily in the 7th carbon. After 1^{14}C -arachidonate injection, testicular $\Delta^{9,12,15,18}$ -tetracosatetraenoic was labeled primarily in the 5th carbon. The suggested pathway of biosynthesis of the 24-carbon tetraenoic and pentaenoic acids is by a 2-carbon elongation of docosatetraenoic and of docosapentaenoic acids.

ENZYMATIC ALKYLENATION OF PHOSPHOLIPID FATTY ACID CHAINS BY EXTRACTS OF MYCOBACTERIUM PHLEI. Y. Akamatsu and J. H. Law (Dept. Biochem., Univ. Chicago, Chicago, Ill. 60637). *J. Biol. Chem.* 245, 701-8 (1970). The enzymatic synthesis of

ARMSTRONG-CHEMTEC BUILDS FATS CRYSTALLIZERS



For use crystallizing: Tallow fatty acids, tall oil heads cuts and mid cuts, many neutral fats from vegetable, animal and marine sources, and cooling viscous materials.

Bring us your fats crystallization problems.

RICHARD M. ARMSTRONG CO.
Box 566-J
West Chester,
Pennsylvania 19380

CHEMTEC N.V.
Box 52-J
Soestdijk,
Holland

CHEMTEC N.V.
Box 3-J, Willowyard Rd.
Beith, Ayrshire,
Scotland

ANNOUNCEMENT

1970-71

SMALLEY CHECK SAMPLE PROGRAM

The Smalley Committee annually offers a number of Check Sample Series in various analytical categories. Interested analysts should write to Smalley Committee, AOCS, 35 E. Wacker Drive, Chicago, Illinois 60601, prior to July 15, 1970 for order forms and complete information, which will be distributed before each series begins.

The following Check Sample Series (the number of samples being shown in parenthesis) are offered:

Cottonseed (10)	Oilseed Meals (15)	Cottonseed Oil (4)
Soybeans (10)	Edible Fats (5)	Soybean Oil (4)
Peanuts (7)	Drying Oils (6)	Copra (4)
Safflower Seed (7)	Tallow & Grease (5)	N.I.O.P. Fats & Oils (5)
Gas Chromatography (fatty acid composition) (6)		
Cellulose Yield (cotton linters) (10)		

Additional series will be offered should sufficient interest be indicated. Please advise the Smalley Committee of series you feel would be of value.

R. T. Doughtie, Jr., Chairman
Smalley Committee

ABSTRACTS: BIOCHEMISTRY AND NUTRITION

tuberculostearic acid (10-methylstearic acid) was catalyzed by extracts of *Mycobacterium phlei*. This process involved two reactions of the olefinic fatty acid chain of phospholipids. The chain was first alkylated at the 10-carbon to give a methylene group, which was subsequently reduced to a methyl group. The first reaction could be measured by using S-adenosyl-methionine-methyl-¹⁴C. The enzyme was found in the supernatant fraction when extracts of cells broken down by sonic oscillation were subjected to centrifugation at 100,000 g. S-Adenosyl-L-methionine was the only effective donor of the 1-carbon unit. Phosphatidylglycerol, phosphatidylinositol and phosphatidylethanolamine were substrates for the reaction, and both 16- and 18-carbon chains were alkylated although only the Δ⁹-olefinic chains appeared to be converted.

THE ENZYMIC SYNTHESIS OF FATTY ACID METHYL ESTERS BY CARBOXYL GROUP ALKYLATION. *Ibid.*, 709-13. The reaction was studied by incubation of fatty acid with S-adenosylmethionine-methyl-¹⁴C and isolation of the labeled ester. This was characterized by thin-layer and gas-liquid chromatography and by isolation of labeled methanol from alkaline hydrolysates of the ester. Of several substrates tested only S-adenosylmethionine was an effective methyl donor ($K_m = 2.5 \times 10^{-6}M$) and oleic acid was the most effective fatty acid acceptor ($K_m = 1.3 \times 10^{-3}M$). Some methyl ester was formed when phospholipids were added to the incubation mixture, presumably because lipase action liberated fatty acids which could serve as substrates.

RAPESEED PRESSCAKE. XV. DEGRADATION OF SULFUR COMPOUNDS DURING PROCESSING OF THE PRESSCAKE. A. Rutkowski and H. Kozłowska (Dept. of Food Technol., Agricultural Univ. of Olsztyn, Olsztyn, Poland). *Oléagineux* 24, 687-90 (1969). The 5-vinyl-2-oxazolidinethione thiocyanates produced by hydrolysis of the goitrogenic thioglucosides are heat-labile. Heat treatment (100-125°C for 1 hour) of the rapeseed presscake lead to a marked drop in their concentration. This treatment did not affect the isothiocyanates, whose level can be reduced by steam stripping. The conditions used in the desolventizing-toasting process enable a presscake of satisfactory quality to be obtained.

EFFECT OF DIETARY LIPIDS ON THE RATES OF SYNTHESIS OF NUCLEIC ACIDS AND OF DEVELOPMENT OF ADIPOSE TISSUE. J. Raulin (Unité U 56 de l'INSERM, Hôpital Parrot, 78 rue de General Leclerc, 94-Bicêtre). *Rev. Franc. Corps Gras* 16, 767-70 (1969). In the rat, the size of the adipose cells increased both with the quantity of lipids fed and also with the hardness of the fat. In animals of the same age, having the same amount of adipose tissue, the number of adipocytes was greater when the diet was richer in polyunsaturated fatty acids (from sunflower oil) and less when the dietary lipids were more saturated (i.e., lard). The amount of DNA in the perigenital tissue was greater when the diet contained sunflower oil and lower when the diet contained lard. Using radioactive precursors, the author found that the specific activity of the adipose subcellular particulates (nucleus, mitochondria) depended on the nature of the dietary lipid. In general, the rate of synthesis (or breakdown) and the development of adipose tissue appeared to be closely related to the composition of dietary lipid.

INFLUENCE OF DIETHYLSTILBESTROL ON THE TURKEY WITH SPECIAL REFERENCE TO HISTOLOGICAL CHANGES IN THE AORTA. L. M. Krista, J. H. Sautter and P. E. Waibel (Depts. of Animal Sci., and Vet. Pathol., Univ. of Minnesota, St. Paul, Minn. 55101). *Poultry Sci.* 48, 1961-68 (1969). Physiological and metabolic changes were induced by DES treatment. Even though distinct morphological differences are hard to establish, the levels of DES could be related to changes in body weight, blood pressure, carcass composition, general conformation and secondary sex characteristics. The lowest level of DES did not influence body weight, blood pressure or aortic rupture, but an increase in body fat and liver size was observed. The medium level of DES resulted in a significant reduction in weight gain at 12 weeks of age, a significant decrease in blood pressure at eight weeks of age and a significant increase in aortic rupture, plaque formation and degeneration. The highest level of DES had an obvious toxic effect on the birds as indicated by a debilitated appearance and depressed weight gains by six weeks of age. Morphological changes due to

(Continued on page 232A)

(Continued from page 224A)




Fig. 3. 1968-69 Soybeans


lack confidence in personal judgment and tend to duplicate the action of large speculators. This may be due to an assumption that large speculators became large because of successful market judgment more than 50% of the time. At any rate it is impossible to segregate small speculator positions since these are lumped together with small hedges in one figure.

This study deals with the soybean, soybean oil and meal markets during the period July 1965 to June 1969. It was observed that during this period large speculators' positions fluctuated from heavily long to heavily short.

No attempt was made to show price action during the month. Only prices used were the closing futures prices for the spot month on the last day of each month, which is the same date on which the CEA position report is compiled. Certainly there were price fluctuations, some very drastic ones, during the month but these are frequently related to the closing out of a futures contract in its final days of trade which may not accurately reflect reasonable market value. The charts for 1968/69 illustrate the large speculator position profile plotted against price at the end of each month. This is expressed as a ratio of long vs short or short vs long depending on which was largest.

Conclusion: The most obvious conclusion is that price goes up as large speculators move more heavily to the long side and go down as they move to the short side. This is not intended to indicate that speculator action causes price change, or that price change causes speculator action. It is valid to conclude, however, that when large speculators have greater short positions than long there will be price improvement soon. It is also demonstrated that prices will weaken when large speculators are heavily long but this reaction may be delayed when there exists a substantially bullish situation. It necessarily follows that speculative position changes occur in relation to price changes, for they seldom hang onto a losing position.

COLEMAN



Refinery control lab determines nitrogen automatically on round-the-clock basis

Using the automated Coleman Nitrogen Analyzer as basic equipment, the quality control laboratory of The Atlantic Refining Company (Philadelphia) determines nitrogen content of petroleum products down to 0.02% nitrogen.

Operated by control chemists on all three shifts, the instrument provides a complete nitrogen analysis in as little as 8 minutes. It is used for critical quality control work with gas oils, lube oil additives, greases and synthetic detergents.

The instrument is fully as efficient in the control laboratory environment as it has proven under carefully-controlled research laboratory conditions.

Write for the specific method developed by Atlantic Refining's Product Quality Department for nitrogen determination in refinery fluids.

Send for Bulletin H-291

Write to Coleman Instruments Division, The Perkin-Elmer Corporation, 42 Madison St., Maywood, Illinois 60153

PERKIN-ELMER

(Continued from page 230A)

DES included increased reorientation of fibers (presumably smooth muscle cells) from a parallel position to a radial position. This reorientation of fibers and subsequent thinning of the aortic wall and degeneration of the elastic tissue of the media are factors that may lead to the rupture of the wall. Plaque size increased with age. Many of the vacuoles observed did not stain with Oil Red O. Dosage of DES represents an important variable in relation to results obtained and should be carefully considered in experimental work.

ESTIMATES OF CHANGES IN PLASMA CHOLESTEROL AND PROTEIN IN RELATION TO CERTAIN REPRODUCTIVE TRAITS IN FEMALE BREEDER TURKEYS. T. K. Mukherjee, G. W. Friars and J. D. Summers (Dept. of Poultry Sci., Univ. of Guelph, Guelph, Ontario, Canada). *Poultry Sci.* 48, 2081-86 (1969). When 534 turkey hens from three generations were shifted from approximately an 11 to a 14 hour photoperiod per day for 14 days, average increases of 72.91 mgm/100 ml and 0.87 gm/100 ml were observed for cholesterol and total protein of blood plasma respectively. The birds that hatched later in each generation showed a higher increase in plasma cholesterol levels than the earlier hatched birds within each strain. No such trend was observed in the levels of plasma protein. The measures of increments in the plasma cholesterol and protein levels due to increased photoperiod were used as independent variables in predicting measures of subsequent reproductive performance. It was noted, in general, that the traits associated with reproduction are not dependent on the increments in the levels of these two biochemical components. Heritability estimates obtained by the regression of progeny performance on dam's record for the increments in the plasma cholesterol and protein levels were 0.24 ± 0.40 and 0.24 ± 0.54 respectively. Estimates of heritability for cholesterol and protein levels measured after 14 days of extended photoperiod were 0.28 ± 0.30 and 0.26 ± 0.42 respectively. Significant ($P < .05$) positive correlations were observed between the increments in cholesterol and protein levels in the different subclasses under study.

A RAPID ASSAY FOR LIPOPROTEIN LIPASE. M. C. Schotz, Arlene S. Garfinkel, R. J. Huebotter, and J. E. Stewart (Radioisotope Res., Veterans Admin. Center, Los Angeles, Cal. 90073). *J. Lipid Res.* 11, 68-9 (1970). A rapid assay for lipoprotein lipase activity employing a ^{14}C -labeled substrate is described. The method is very sensitive and suitable for routine use.

SIZE AND LIPID COMPOSITION OF CHYLOMICRONS OF DIFFERENT SVEDBERG UNITS OF FLOTATION. R. Fraser (Dept. of Expt. Pathol., J. Curtin School of Med. Res., Australian Nat. Univ., Canberra, Australia). *J. Lipid Res.* 11, 60-5 (1970). Chylomicrons from thoracic duct lymph of rabbits which were fed corn oil were separated in a preparative ultracentrifuge into subfractions of different S_v values in order to compare their size, as determined by electron microscopy, with that expected from ultracentrifugation data. The lipid composition of the chylomicrons of different S_v values was also correlated with their morphology in order to elucidate their structure. Although the diameter distribution of chylomicrons from sub-

fractions of lower S_v ranges corresponded approximately to the expected size distribution, that of the higher S_v ranges contained many small particles. The findings were consistent with the hypothesis that, irrespective of the S_v range of chylomicrons, the core is comprised of triglycerides, while phospholipids is spread as a monomolecular layer on the surface of the particles.

COMPOSITION OF HUMAN SERUM SPHINGOMYELINS. E. L. Hirvisalo and O. Renkonen (Dept. of Biochem., Univ. Helsinki, Helsinki, Finland). *J. Lipid Res.* 11, 54-9 (1970). Serum sphingomyelins were analyzed by argentation chromatography of the corresponding ceramide diacetates. Six subfractions were obtained. Three of them contained 4-sphingenes in combination with saturated, *trans*-, or *cis*-monoenoic fatty acids; the remaining three contained sphingadienine, also in combination with saturated, *trans*-, or *cis*-monoenoic fatty acids. Palmitic acid was the principal fatty acid combined with 4-sphingenes, while nervonic acid was the principal fatty acid combined with sphingadienine. About 4% of the total fatty acids of sphingomyelin were *trans*-monoenoic. They were comprised of many positional isomers of straight-chain C_{22-24} compounds. The *cis*-monoenoic acids made up 33% of the total acids and consisted of almost pure nervonic acid. The rest of the acids were saturated. The 4-sphingenes contained small amounts of iso- C_{18} and anteiso- C_{19} compounds in addition to the straight-chain C_{16-22} bases.

STIMULATION OF LIPOLYSIS IN ADIPOSE TISSUE IN VITRO BY INHIBITORS OF LIPID MOBILIZATION. F. P. Kupiecki and Diana I. Schneider (Dept. of Diabetes Res., The Upjohn Co., Kalamazoo, Michigan 49001). *J. Lipid Res.* 11, 38-41 (1970). 5-Methylpyrazole-3-carboxylic acid (U-19425) and nicotinic acid, which apparently inhibit lipolysis *in vivo* as indicated by low plasma FFA and glycerol concentrations, stimulate lipolysis *in vitro* in adipose tissue removed from fasted rats 30-90 min after treatment. This stimulation is not the result of low initial levels of FFA in adipose tissue. An increased rate of lipolysis is not induced *in vitro* by preincubating tissue of untreated rats with U-19425.

A METHOD FOR THE QUANTITATIVE DETERMINATION OF NEUTRAL GLYCOSPHINGOLIPIDS IN URINE SEDIMENT. R. J. Desnick, C. C. Sweeley and W. Krivit (Dight Inst. for Human Genetics, Univ. of Minnesota, Minneapolis, Minn. 55455). *J. Lipid Res.* 11, 31-7 (1970). A method is described for the isolation and quantitation of six neutral glycosyl ceramides from human urinary sediment. Total lipids were extracted from sediments of 24-hr. urine collections, and the glycosyl ceramides were isolated by silicic acid column chromatography followed by thin-layer chromatography. Methanolysis of the individual glycosyl ceramides yielded methyl glycosides which were quantitated as the trimethylsilyl ethers by gas-liquid chromatography. By this technique, the submicromolar concentrations of six glycosyl ceramides in normal subjects and in individuals with Fabry's disease, an hereditary glycosphingolipid storage disease, were determined. Trihexosyl ceramide (galactosyl-galactosylglucosyl ceramide) and a digalactosyl ceramide accumulated in the urinary sediment of patients with Fabry's disease.

BIOSYNTHESIS OF FATTY ACIDS IN MAMMARY TISSUE. I. PURIFICATION AND PROPERTIES OF FATTY ACID SYNTHETASE FROM LACTATING GOAT MAMMARY TISSUE. A. K. N. Nandedkar, E. W. Schirmer, T. I. Pynadath and S. Kumar (Dept of Chem., Georgetown Univ., Washington, D.C.). *Arch. Biochem. Biophys.* 134, 554-62 (1969). Fatty acid synthetase of a high degree of purity has been prepared from the particle-free supernatant fraction of lactating goat mammary tissue. Ultracentrifugation and polyacrylamide gel electrophoresis revealed heterogeneity of the preparation. However, the enzyme complex was sufficiently free of malonyl-CoA decarboxylase activity to enable the demonstration of the requirement for the 'primer' acyl-CoA in addition to malonyl-CoA and NADPH. The enzyme appeared to utilize butyryl-CoA more efficiently than acetyl-CoA as 'primer.' The nature and extent of the fatty acids synthesized using these two 'primers' were characteristically different and were pronouncedly influenced by the concentration of malonyl-CoA.

II. SYNTHESIS OF BUTYRATE IN LACTATING RABBIT MAMMARY SUPERNATANT FRACTION BY THE REVERSAL OF β -OXIDATION. A. K. N. Nandedkar and S. Kumar. *Ibid.* 563-71. An enzyme fraction has been obtained from lactating rabbit mammary particle-free supernatant fraction which synthesizes butyrate from acetyl-CoA by the reversal of β -oxidation. This fraction

(Continued on page 234A)



Plan Now for
AOCS 44TH
FALL
MEETING
Chicago
Conrad Hilton
Sept. 27-Oct. 1, 1970

• Four Corners . . .

(Continued from page 218A)

- tographic Determinations of Insecticides, by G. Baluja, J. M. Franco and M. E. Pereira.
- Densitometry on Tin Layer Chromatography, by E. Vioque, H. Murillo and M. P. Maza.
- Biochemistry of Development: Biosynthesis of Fatty Acids in the Metamorphosis of *Ceratitis capitata*, by J. M. Odriozola, A. Pineiro, A. Ribera and A. Martin.
- Biochemistry of Development: Evolution of Positional Fatty Acid Distribution in the Phosphatidylethanolamine During the Metamorphosis of *Ceratitis capitata*, by L. J. M. Fernandez-Sousa, A. Martin and A. Ribera.
- Changes in the Glycerides of Olives During Ripening, by A. Vazquez and M. Mancha.
- Mixed Monolayers of Cholic Acid and Lecithin, by J. Llopis, A. Albert, J. L. Saiz and D. Alonso.
- The Fatty Acid Distribution in the Cocoa Butter Glycerides, by F. Mazuelos and J. L. Aguila.
- Studies on the Superficial Monomolecular Layers IX: Compression Isotherms of Soybitan Esters, by E. Otero and O. Valls.
- Studies on the Superficial Monomolecular Layers X: Association of Cholesterol and Sorbitan Cholesterol. E. Otero, O. Valls and M. Castillo.
- Rheology of Sodium Dodecyl-oxydibencene-disulfonate, by J. Alba and C. Gomez.
- Retention of Miscelle in Olive Press Cake of High Humidity, by V. Flores, F. Lopez and J. Pereda.
- Contribution to the Cinetic study of Fat Hydrolysis, by J. Rodrigo and B. Lopez.

Syndicalist National Week of Research Industry

The Syndicalist National Week of Research Industry meeting will be held April 20-25, 1970, and has been promoted by the Syndicalist Organization with the cooperation of official organizations of investigation.

The purpose of the meeting is to demonstrate the need for and the convenience of investigation for both industrial people and scientific research institutions. The general topics to be discussed will be: (a) Investigation as a factor of development. Spanish investigation and international cooperation. (b) Participation of syndicates in industrial investigation and development. (c) The industrial investigation and its economical conditioning. The official research centers and their projection towards the industry. (d) Lecture and seminar on possible assistance for investigation.

Sweden Reinhard Marcuse

Scandinavian Forum for Lipid Research and Technology (Lipidforum)

An association of Scandinavian lipid scientists and technicians and other persons interested in lipids has been established. The General Secretary is Reinhard Marcuse, Swedish Institute of Food Preservation Research, Göteborg. The Membership which is acquired by simple application is free of fee and the number of members is about 200.

The association is taking over the organization of Scandinavian Symposia on Lipids which formerly were arranged by the Scandinavian Board of Research about each third year. The next symposium will be held in Denmark next year. The chairman of the organizing committee of this symposium is also chairman of the board of Lipidforum. It is at present director Anders Herløw, Grindstedvaerket, Braband, Denmark. The board consists of representatives of the Scandinavian countries.

Lipidforum is intended to promote professional information and education and to become a center for contact with colleagues and organizations in the Scandinavian countries as well as abroad. The activity is now being

developed in accordance with the results of an inquiry and will comprise seminars, symposia and congresses. To begin with, information is distributed twice a year to the members. Communications or notes on events of interest to be forwarded in this way are very welcome and should be sent to the office. Mailing address: Fack, S-400 21 Göteborg 16, Sweden.

International Summer School on Glycolipids in Membranes at the University of Helsinki, Finland

On behalf of the Finnish Biochemical Society, the 16th FEBS (Federation of European Biochemical Societies) Summer School on Glycolipids in Membranes has been arranged by Ossi Renkonen to take place August 10 to 14. Twenty-eight papers will be presented. Those interested in attending the School are invited to write O. Renkonen, Department of Biochemistry, Laboratory of Lipid Research, University of Helsinki, Haartmaninkatu 3B, Helsinki 29, Finland.

The Proceedings of the Fifth Scandinavian Symposium on Lipids

The Fifth Scandinavian Symposium on Lipids took place at Tyringe, Sweden, during June 10 to 13, 1969, and was attended by about 150 persons. The proceedings are now available for distribution to all attendants. They can be ordered from R. Ohlson, AKO, Dept. o. Research, S-292 00 Karlshamn, Sweden at Sw. crws. 20. They are published in Scandinavian languages, a few in English. The publication comprises 40 papers including five plenary lectures by: K. G. Jensen (The Role of Fat in the Scandinavian Countries); G. Andersson (The Development of Swedish Oil Seed Breeding); K. Larsson (Molecular Packing in Fats); R. Marcuse (Metal Catalyzed Lipid Oxidation); and S. Friberg (Equilibria in Liquid-Water System and Their Influence Upon the Properties of Products). Further plenary lectures were given by E. Stenhagen (The Use of Masspectrometry for Lipid Analysis) and B. Samuelsson (New Possibilities for Determining the Effect of Dietary Fat Composition).

Scandinavian Symposium on Lipids to be Arranged in 1971 in Denmark

The Sixth Scandinavian Symposium on Lipids will take place in June 1971 in Denmark, probably at Grenaa, Jutland. On behalf of the Danish Institute for Lipid Research, Ole Tolboe, of the Jutish Institute of Technology at Aarhus, has taken over the chairmanship of the organizing committee.

The Food and Nutrition Group in Göteborg

CLiNG, The Food and Nutrition Group in Göteborg, was formed February 12, 1969, and has an office at The Swedish Institute for Food Preservation Research, Fack, S-400 21 Göteborg 16, Sweden. The purpose of this group is to create better possibilities for worthwhile contacts between research workers from different disciplines but with a common interest in food and nutrition. Hopefully new team projects in research and education will arise as a result of the formation of CLiNG. New contacts between scientists have already been taken. It is also likely that existing resources in the Göteborg area in personnel, know-how and equipment will be more efficiently used. Suitable areas for cooperation in research concern lipids and proteins respectively. A considerable amount of research in these two areas is presently being carried out in Göteborg, but a need for coordination and cooperation over the faculty boundaries is strongly felt among the participating scientists.

An inventory of research workers and research projects in the area of Göteborg has been made. Two interdisciplinary seminars have been held, one concerning "Fats

(Continued on page 234A)

(Continued from page 233A)

in Foods," where the consumption of fat was discussed from mercantile and organoleptic points of view. Chemical, medical and technical aspects on fat were also discussed. The second seminar was about medical aspects of protein deficiency and new sources of protein. A seminar about enzymes will be held in September 1970.

Other immediate activities include plans for a joint educational program in connection with the new Swedish doctors degree.

Scandinavian Symposium on Surface Chemistry at Tylösand, Sweden

The Fourth Scandinavian Meeting on Surface Chemistry will take place at Tylösand in Sweden August 27-29, 1970. The structure of lipids is one topic in the program. Among the invited speakers are Egon Matijevic, director of Institute of Colloid and Surface Science, Potsdam, N.Y. and James L. Ferguson, associate director of Liquid Crystal Institute, Kent, Ohio.

For information contact the Institute of Surface Chemistry, Drott. Kristinas v. 45 S11428 Stockholm Ö, Sweden.

Different Forms of Obesity Studied by Group in Göteborg

At the annual meeting of the Swedish Medical Association in December 1969, results of research on two forms of obesity in man have been presented by a group working in association with Per Björntorp at the First Medical Service of Sahlgrenska Sjukhuset in Göteborg. One form is due to an increased diameter (without any considerable change of number) of the adipose tissue fat cells (Hypertrophic obesity), the other shows mainly an increased number of the fat cells (Hyperplastic obesity). Hypertrophic obesity is more closely connected to well-known metabolic disturbances found in obesity as increased insulin production and possibly diabetes mellitus. Increased levels of blood lipids and cardiovascular diseases may be more frequent as well. The hyperplastic form, probably inherited to at least some extent, is presumably less dangerous.

Group Flight From Copenhagen to the World Congress in Chicago

A group flight for participants of the World Congress is being arranged by SAS, departing from and returning to Copenhagen. This flight should be particularly suitable for participants from Scandinavia and other Northern European countries. Two alternative schedules of two or three weeks in Chicago are being offered, i.e., departing from Copenhagen September 19 or 26.

Detailed information can be obtained from R. Marcuse, Lipidforum, c/o SIK, Faek S-400 21 Goteborg 16, Sweden, or from Wallace J. Quick, President, Chemtex Products, Inc., 400 E. Randolph Street, Chicago, Illinois 60601, U.S.A.

(Continued from page 232A)

has been shown to contain acetoacetyl-CoA thiolase, NADH-dependent acetoacetyl-CoA reductase, and crotonyl-CoA hydratase. The reduction of crotonyl-CoA appears to be brought about by the fatty acid synthetase using NADPH or NADH. The two reducing reactions are believed to overcome the thermodynamically unfavorable reaction involved in the condensation of two molecules of acetyl-CoA.

CHEMICAL, PHYSICAL AND BIOLOGIC PROPERTIES OF BUTTER AND GHEE. R. I. Tannous and A. Merat (American Univ. of Beirut, Beirut, Lebanon). *J. Am. Dietetic Assoc.* 55, 267-72 (1969). Fermenting milk into yoghurt increases the total acidity and biacetyl value of butter and ghee compared with samples made from milk. Other chemical and physical properties were generally similar. During fermentation also, lauric, myristic and palmitic acids increased, while a decrease in oleic acid was observed. Comparison of butter properties with those of ghee show a lower biacetyl value for ghee and a distinctive flavor difference from butter. These are attributed to the effect of the heat treatment involved in converting butter into ghee. In animal experiments, feeding either ghee or corn oil at a 20% level in the diet for three weeks caused no appreciable increase in blood cholesterol. The recovery of ghee from the gastrointestinal tract of rats was lower than that of corn oil only when measured at three hours after ingestion, but not after six hours.

THE EFFECT OF CHOLANIC ACID ON STEROL AND FATTY ACID SYNTHESIS. G. R. Jansen, M. E. Zanetti, C. F. Hutchison, F. J. Andriuli and E. E. Howe (Merck Inst., Rahway, N.J.). *Arch. Biochem. Biophys.* 134, 185-95 (1969). The effects of cholanolic acid on incorporation of glucose-U-¹⁴C into digitonin-precipitable sterol (DPS) and fatty acid in liver and extrahepatic tissues of mice and rats have been investigated. Cholanolic acid lowers plasma triglycerides in both species, but lowers cholesterol only in mice. Incorporation of glucose-U-¹⁴C into liver fatty acid was lowered by cholanolic acid in both mice and rats, as was the weight percentage of this fraction. Incorporation into liver DPS in both species was reduced after 1-day treatment with cholanolic acid, but was considerably elevated when the cholanolic acid was fed for a 7-day period. Cholanolic acid caused the weight percentage of DPS in the liver to decrease in mice but to increase in rats. Incorporation of glucose-U-¹⁴C into extrahepatic fatty acid or DPS was not influenced in rats or mice by feeding cholanolic acid in the diet for up to 7 days. When fed in the diet for 6 weeks, cholanolic acid caused the percentage of body fat as well as the weight of the epididymal fat pads to be reduced approximately 35%. Under these conditions incorporation of glucose-U-¹⁴C into fatty acid in epididymal fat was stimulated by a three-fold factor.

LYSOLECITHIN METABOLISM OF THYMUS AND BURSA CELLS OF THE CHICKEN. J. Kruger, E. Ferber and H. Fischer (Max Planck Inst. fur Immunbiologie, Freiburg, Germany). *Proc. Soc. Exp. Biol. Med.* 132, 543-47 (1969). The purpose of the present investigation was to determine whether lysophosphatide-metabolizing enzymes in lymphocytic cells might be an important component of certain tissue-damaging immune reactions. Thymus and bursa cells were used for technical reasons. The two cell populations differed in total lipid content according to differences in cell size but there was no difference with regard to phospholipid composition. In the cell homogenate the "apparent" reaction rates and Michaelis constants of the following enzymes were determined: ligase, acyltransferase, acyl-CoA-hydrolase, lysophospholipase and phospholipase A, the only difference between both cell homogenates was shown for the transferase activity. The investigation of the incorporation of radio labeled substrates used in the enzyme assays indicated that the enzymes determined in the cell homogenates (with the exception of phospholipase A) are also active in the intact cells, but to a much lesser extent.

FACTORS AFFECTING WHOLE- AND PART-LACTATION MILK YIELD AND FAT PERCENTAGE IN A HERD OF HOLSTEIN CATTLE. R. H. Miller and N. W. Hooven (Animal Husbandry Res. Div., USDA, Beltsville, Md. 20705). *J. Dairy Sci.* 52, 1588-1600 (1969). Relationships among milk and fat-corrected milk yield, fat percentage, weight change, body weight, and days open were studied in 1004 lactations collected over a 14-year period in a herd of Holstein cattle. Least-squares analyses of 31 to 60-day, 121 to 150-day, and of 181 to 210-day part lactations and total lactation were performed on an intra-sire basis. The regressions of milk and fat-corrected milk on weight change

(Continued on page 236A)

SOUTHWESTERN

LABORATORIES

Fort Worth, Tex.

Analytical Chemists
Inspection & Testing
Engineers

Agricultural Products
Feeds—Seeds—Oils
& Related Products

817-332-5181, P.O. Box-1379, 2900 Cullen St., 76101

(Continued from page 212A)




FIG. 2. The De Laval Immediumfilter. Immediumfilter is the registered trademark of Industriele Maatschappij Activit N. V., Amsterdam N., Netherlands.

enters the bottom of the filter, passes thru gravel and fine sand layer, held in place by grid, and exits at the top. The whole sand bed serves as a dirt retention zone. The top fine grain sand is the final polishing zone. This unit offers large dirt holding capacity, long filter cycles, and may be readily cleaned by flow of raw water and air thru an expanded bed to dislodge the dirt.

Solvent Extraction Plants

At solvent extraction plants, there is frequently a degumming or miscella refining operation. Since the miscella soapstock and filter bleach clay are usually added to the meal in the desolventizer-toaster of solvent extractor, miscella refineries do not have the pollution problems associated with conventional refinery installations.

Average waste water flows were sampled at two solvent plants, with miscella refinery and conventional degumming operation. Typical analyses of composite waste water appear in Table IV. The refinery operation definitely increases impurities in the waste water. There is increase in organic and suspended solids, fat content and BOD figures.

TABLE IV
Chemical Analyses of Waste Water: Values in ppm

Components	Solvent plant miscella refinery	Solvent plant degumming operation
Total solids	1820	417
Suspended solids	570	127
Organic solids	740	203
Fat	232	Trace
BOD	462	67
pH	6.3	7
After chemical treatment and centrifugation		
Fat	None	None
Suspended solids	None	None
BOD	<50	<15

At a seed mill, with solvent extraction plant, the usage of fresh water is high, 50-70 gal/min for a 750 ton/day installation. This represents 75,000-100,000 gal of water per day. When such a plant is situated in a western area of the United States, where fresh water supply is limited, the treatment of solvent plant waste waters has real merit. Water can be recovered for reuse as cooling water to reduce fresh water demands.

It has been found that the waste water from a solvent plant can be chemically treated with coagulant in 90-100 ppm dosage, followed by polyelectrolyte at 1 ppm, and centrifugation in a PX Separator, to give a clear effluent. By treatment and centrifugation of this waste water, a clear water phase with no fat or suspended solids is recovered, and there is 78-90% BOD reduction. The precipitated solids, that may be concentrated and removed by centrifuge, are principally protein. This concentrated protein slurry may be added to desolventizer-toaster. In the area where this study was made, the well water had higher total solids content than the clarified process water.

REFERENCES

1. Alikonis, J. J., and J. V. Ziemba, "Waste Treatment," Food Engineering, July, 1967.
2. Butrico, F. A., "New Standards for Streams and What These Mean to Industry," AOCs Fall Meeting, 1966.
3. Dietz, J. C., "Joint Treatment of Municipal and Industrial Wastes," 16th Annual Dairy Engineering Conference, 1968.
4. Dow Chemical Co., Bulletin Separan.
5. Parsons, W. A., "Chemical Treatment of Sewage and Industrial Wastes," Bulletin No. 215, National Lime Association, 1965.
6. Roderick, R. E., J. Water Pollution Control Federation, 34, (1962).
7. Water Pollution Control Federation, Regulation of Sewer Use, Manual of Practice No. 3, 1968.
8. Water Resources Commission, Rules and Classifications and Standards of Quality and Purity for Waters of New York State, 1961.

[Received February 28, 1969]

Newest Detergency Concepts to be Heard at Chicago Congress

Eminent scientists will speak at a Symposium on "Basic Aspects of Detergency" on September 29 at the 10th World Congress in Chicago. Recent concern for pollution, the environment and ecology has focused continuing attention on the influence of various detergents. There have been recommendations from various sources to eliminate certain detergent components which allegedly contribute to pollution. It is, however, essential to know and understand the chemistry and physics of cleaning quite thoroughly if cleaners are to be developed which are both effective and environmentally safe. The planned Symposium is to provide a review and up-dating of key properties involved in detergency and will be basic to many aspects of the cleaning process.

The Symposium has been organized by M. E. Ginn of the Masury-Columbia Company (formerly of Armour-Dial Inc.). Titles and speakers will include:

1. "Recent Advances in Detergency Theory," by A. M. Schwartz, Gillette Research Institute.

2. "Gel Filtration of Surfactants," by Toshio Nakagawa, Shionogi Research Laboratory, Shionogi & Co., Ltd.
3. "The Association of Surfactants Into Liquid Crystal Systems and Its Influence on Solubilization and Emulsification Phenomena," by Stig Friberg, Swedish Institute for Surface Chemistry.
4. "Interactions of Colloidal Particles With Complex Ions and Polymers," Egon Matijevic, Institute of Colloid and Surface Science, Clarkson College of Technology.
5. "Statistical Analysis of Detergency Tests With a Natural Soil," J. R. Trowbridge, Colgate-Palmolive Research Center.

The Symposium is part of a Joint Meeting, September 28-October 1, 1970 between the International Society for Fat Research and the American Oil Chemists' Society at the Conrad Hilton Hotel in Chicago. The international flavor of the meeting should enhance an exchange of differing points of view.

(Continued from page 234A)

After 39 years with the company, W. R. CHASE, executive vice president and member of the board of directors of The Procter & Gamble Company, retired as both an officer and director on April 30. Mr. Chase, a native of Brookline, Mass., joined Procter & Gamble in 1931. He held various management positions before being elected vice president-advertising in 1955. He became vice president-soap products division and a director of the company in 1957 and was elected an executive vice president of P&G in 1960. Mr. Chase is a trustee of the Cincinnati Summer Opera Association. He is a past president of the Cincinnati Harvard and Harvard Business School Clubs, and was a national vice president of the alumni. He is a past president of the Cincinnati Planned Parenthood Association, and was a director of the national organization. He is an honorary vice president for life of the National Association of Manufacturers. J. W. HANLEY, vice president-group executive, assumed overall responsibility for the company's major staff departments which reported to Mr. Chase.

R. D. VOLD of Culver City, professor of chemistry at the University of Southern California, received the 11th Annual Tolman Medal from the Southern California Section of the American Chemical Society in recognition of his distinguished contributions to the field of chemistry over a period of 35 years. The Richard C. Tolman Medal bears the name of a distinguished scientist and profound scholar who had a deep concern for his fellow human beings. Selection of Dr. Vold as recipient of the 1970 Tolman Medal acknowledges his studies of colloidal systems, gels, emulsions, and surface films. It praises him for exemplifying the academic scientist with the integration of his talents for teaching, research and administration, and for his service to the chemistry profession nationally and abroad. For a number of years Dr. Vold was a research chemist with Procter & Gamble. After a brief stay at Stanford as a research associate, he was appointed an assistant professor of chemistry at USC in 1941. By 1947, Dr. Vold had advanced in academic rank to the position of professor and served as department chairman in 1950-53. As a Fulbright Fellow, he carried on research studies at the University of Utrecht, Holland, 1953-54, then spent two years as a visiting professor of physical chemistry at the Indian Institute of Science, Bangalore, India. In 1965 he returned to India to conduct a Summer Chemical Institute at the University of Jadavpur.

SIDNEY LOEB has joined Water Pollution Research & Applications, Inc., Washington, as a member of the firm's Scientific Advisory Council. In this capacity he will contribute to the firm's consulting activities especially as they apply to reverse osmosis, a process receiving increasing interest in desalting as well as in the treatment of wastewaters and process solutions. Dr. Loeb, who is often referred to as the "father" of reverse osmosis directed the project at the University of California, Los Angeles, where much of the pioneering work on the process was carried out. Recently he assisted Israel, under UNESCO auspices, to become independently established in reverse osmosis technology.

H. F. P. HOEPERMANS, has been named Director of Marketing-Europe of Stepan Chemical Company, Northfield, Illinois which has announced the opening of European headquarters in Brussels, Belgium. The office will direct the sales of all Stepan Chemical divisions' products; however, most emphasis will be placed on the sale of surface active agents. Mr. Hoepermans joined Stepan International Department, Northfield, Illinois in 1965. Prior to his new appointment he had been assigned to the firm's Commercial Development Group. Mr. Hoepermans is a native of the Netherlands and attended the Dutch Institute for Foreign Trade, Nijenrode. He was an exchange scholar at DePauw University and in 1969 received his MBA from the University of Chicago Graduate School of Business.

were consistently large, negative and linear. The regressions of yield on average lactation body weight were significant and curvilinear; production increased up to average or above-average weight, then declined. Effects of the number of days open were small, accounting for less than 2% of the variance in all cases. Calendar year and age were the factors most closely associated with variation in fat percentage.

THE MECHANISM OF INTRODUCTION OF ALKYL GROUPS AT CARBON 24 OF STEROLS. III. THE SECOND ONE-CARBON TRANSFER AND REDUCTION. R. T. Van Allen, H. Chikamatsu, N. J. de Souza, J. P. John and W. R. Nes (Dept. Chem., Univ. Mississippi 38677). *J. Biol. Chem.* 244, 6645-55 (1969). Seeds of the "stone pine", *Pinus pinea* have been shown to contain 24 Δ^5 -stigmasten-3 β -ol as well as smaller amounts of 24 Δ^5 -ergosten-3 β -ol and trans- $\Delta^{5,24(28)}$ -stigmastadien-3 β -ol. Germination of the seeds in the presence of 2-¹⁴C-mevalonate led to labeling of the steroids in a high yield. Isolation and reincubation of the labeled 24-ethylidene compounds, trans- $\Delta^{5,24(28)}$ -stigmastadien-3 β -ol, produced the labeled 24-ethyl derivative, 24 Δ^5 -stigmasten-3 β -ol. The 24-substituted C₁ analogue of the latter, 24 Δ^5 -ergosten-3 β -ol, was obtained labeled when the substrate contained the 24-methylene group (synthetic 28-¹⁴C- $\Delta^{5,24(28)}$ -ergostadien-3 β -ol). From germination in the presence of the 28-¹⁴C- $\Delta^{5,24(28)}$ -ergostadien-3 β -ol the 24-ethylidene derivative, 28-¹⁴C-trans- $\Delta^{5,24(28)}$ -stigmastadien-3 β -ol, was also formed. The results demonstrate that a second transfer of a C₁ group to the $\Delta^{24(28)}$ bond occurs with the formation of a 24-ethylidene group in the biosynthesis of 24-substituted C₂ steroids. They also show that the $\Delta^{24(28)}$ bond can undergo reduction to give the saturated C-24(28) bond in both the 24-substituted C₁ and 24-substituted C₂ cases.

EFFECT OF DIETARY FATS ON SOME CHEMICAL AND FUNCTIONAL PROPERTIES OF EGGS. R. D. Pankey and W. J. Stadelman (Animal Sciences Dept., Purdue Univ., Lafayette, Ind. 47907). *J. Food Sci.* 34, 312-17 (1969). Fatty acid composition of total yolk lipids (triglycerides, cephalin, and lecithin fractions of lipovitellin and lipovitellenin) was influenced by 10% vegetable oil diet supplement (corn, soybean, olive, safflower or hydrogenated coconut oil). The fatty acid composition of the total yolk lipids was influenced by all dietary fats. The major change was in the linoleic acid at the expense of oleic acid with corn, soybean and safflower oil. Olive oil increased the oleic acid and hydrogenated coconut oil increased lauric, myristic and myristoleic acids. The fatty acid composition of the fractions of the lipo-proteins was influenced by the dietary fats and varied between fractions.

CUTICULAR LIPIDS OF ADULTS AND PUPARIA OF THE AUSTRALIAN SHEEP BLOWFLY LUCILIA CUPRINA (WIED.). B. S. Goodrich (Div. of Animal Health, C.S.I.R.O., McMaster Lab., Glebe, N.S.W., 2037, Australia). *J. Lipid Res.* 11, 1 (1970). The presence of a strong contact component in the sex and ovipositing behavior of the sheep blowfly *Lucilia cuprina* Wied. prompted an investigation into the chemical composition of the cuticular wax of the adult male and female flies as well as that of the blowfly puparia. Thin-layer chromatography indicated that the lipids in all the waxes examined comprise hydrocarbons, nonglyceryl esters, triglycerides, free fatty acids and hydroxy compounds, probably diglycerides and monoglycerides. Phospholipids were not detected. Straight- and branched-chain saturated compounds, the latter often predominating, are present in the hydrocarbon, free fatty acid, and ester fractions. Unsaturated molecules were absent. The hydrocarbons resemble those of the cricket to some extent, but the absence of unsaturated compounds is in striking contrast to both the cricket and the cockroach. Pheromones may be present in the low molecular weight fatty acids obtained on brief extraction of the insects.

ISOLATION OF PLASMA LIPOPROTEINS BY ZONAL ULTRACENTRIFUGATION IN THE B14 AND B15 TITANIUM ROTORS. H. G. Wilcox and M. Heimberg (Dept. of Pharm., Vanderbilt Univ., School of Med., Nashville, Tenn. 37203). *J. Lipid Res.* 11, 7-22 (1970). Lipoproteins were isolated from plasma of man, dog, rabbit, rat and chicken by ultracentrifugation in continuous density gradients using the B14 titanium and B15 titanium zonal rotors. Both the VLDL and the LDL of human plasma were separated easily from the HDL and from the other more plentiful plasma proteins by centrifugation for only 1 or 2 hr in the B14 or B15 rotor, respectively. Satisfactory separation of the HDL from the more dense plasma proteins was not achieved with these rotors. The human LDL achieved isopycnic equilibrium (d 1.04) on prolonged