

109. Azulenes. Part I. Synthesis of Vetivazulene.

By ROBERT R. COATS and J. W. COOK.

Condensation of ethyl 4-isopropylcyclopentanone-2-carboxylate (II) with Δ^{β} -hexen- δ -one (III) by the Michael–Claisen–Robinson reaction gave a keto-ester (IV), which was converted by decarboxylation and then reduction into a mixture of dicyclic ketones represented by the structure (V). A pure constituent of this mixture was isolated, and its structure established. Vetivazulene (VIII) has been synthesised from the mixture (V) by ring-enlargement with diazomethane, followed by dehydrogenation with selenium.

THE elucidation by St. Pfau and Plattner (*Helv. Chim. Acta*, 1936, **19**, 858; 1937, **20**, 224, 469) of the type of structure present in the blue hydrocarbons obtained by dehydrogenation of certain essential oils made it of interest to study the synthesis and chemical properties of members of this group of coloured hydrocarbons. Interest in these compounds was reinforced by the elegant analytical proof, published while the present work was in progress (St. Pfau and Plattner, *ibid.*, 1940, **23**, 768), that the dicyclic hydroazulene system is in fact present in β -vetivone, a constituent of vetiver oil (*idem, ibid.*, 1939, **22**, 640; compare Sabetay and Traband, *Bull. Soc. chim.*, 1942, **8**, 152).

It was decided in the first instance to undertake the synthesis of vetivazulene (VIII), a product of dehydrogenation of Java vetiver-oil. Our synthesis was well under way when it found that vetivazulene had already been synthesised by another method (*idem, ibid.*, 1939, **22**, 202), which, however, did not completely establish the structure. This prior work made it possible to devote less attention to the rather troublesome later stages of our synthesis than would otherwise have been necessary.

The type of method which we adopted was that developed with such success by Robinson and his collaborators (*e.g.*, Rapson and Robinson, J., 1935, 1285; du Feu, McQuillin, and Robinson, J., 1937, 53) for the synthesis of polycyclic ketones. Attempted condensation of sodio-3-isopropylcyclopentanone with Δ^{β} -hexen- δ -one (III) gave unsatisfactory results on account of the large proportion of 3-isopropylcyclopentanone converted into diisopropylcyclopentylidenecyclopentanone (probably I) by sodamide in cold ether. Δ^{β} -Hexen- δ -one (III) condensed smoothly, however, with the sodio-derivative of ethyl 4-isopropylcyclopentanone-2-carboxylate (II). The resulting mixture of esters (IV) underwent successive hydrolysis and decarboxylation with boiling alcoholic potash. The ready hydrolysis of the quaternary ester and the lability of the resulting acid are unusual and must be ascribed to stereochemical influences. Hydrogenation with palladium of the unsaturated ketones so formed gave a mixture of stereoisomeric ketones (V), from which one pure component was isolated through its semicarbazone. Oxidation of the mixed ketones with chromic acid gave not only the β -(2-acetyl-4-isopropylcyclopentyl)butyric acid (VI) formed by similar treatment of the stereochemically pure ketone (V), but also a stereoisomeric keto-acid (VI).

The m. p. of the semicarbazone of the pure ketone (V) is not very different from that of the semicarbazone

Propylcyclohexanol (100 g.) was added to the vigorously stirred solution at such a rate that the temperature was maintained at 50–55° without external heat (4–5 hours). The temperature was then raised to 100° for $\frac{1}{2}$ hour, and the excess of water and nitric acid removed at 100° under reduced pressure. The residual β -isopropyladipic acid was esterified with ethyl alcohol and hydrogen chloride, and the ester distilled (135 g.; b. p. 145–165°/15 mm.).

For the Dieckmann reaction, sodium (15 g.) was atomised in dry toluene (200 c.c.), and ethyl β -isopropyladipate added at such a rate that the reaction could be controlled readily. The mixture was refluxed for 80 minutes (oil-bath at 120–130°), with occasional shaking, and cooled to 0°, ice-cold 10% hydrochloric acid (1 l.) added, and the whole shaken until no more solid remained. The separated toluene solution was washed with sodium carbonate solution and water, dried, and distilled. The resulting ethyl 4-isopropylcyclopentanone-2-carboxylate (64 g.; 65% yield) had b. p. 128–135°/15 mm., and gave an intense purple colour with alcoholic ferric chloride.

Hydrolysis of this ester (14 g.) by $4\frac{1}{2}$ hours' boiling with 3*N*-hydrochloric acid (140 c.c.), followed by steam distillation, gave 3-isopropylcyclopentanone (8.5 g.), b. p. 75–80°/15 mm. Its semicarbazone had m. p. 192° (cf. Bouveault and Blanc, *Compt. rend.*, 1908, **146**, 233; **147**, 1314; Wallach, *Annalen*, 1912, **388**, 59) and its phenylhydrazone formed colourless plates, m. p. 75–76.5°.

$\Delta\beta$ -Hexen- δ -one (III).—Oxidation of the carbinol resulting from the interaction of ethylmagnesium bromide and crotonaldehyde could not be effected by Oppenauer's method, aluminium isopropoxide, acetone, and benzene being used, or by chromic acid (after protection of the double bond by addition of bromine), and chromic acid oxidation of the corresponding δ -chloro- $\Delta\beta$ -hexene (cf. Courtot and Pierron, *Bull. Soc. chim.*, 1929, **45**, 290) gave the required ketone in very poor yield. Recourse was therefore had to Blaise's method (*ibid.*, 1905, **33**, 39, 45), whereby allylzinc halide was condensed with propionitrile, and the ethyl allyl ketone isomerised by boiling dilute sulphuric acid. An attempt to improve this method by substituting allylmagnesium bromide (cf. Gilman and McGlumphy, *ibid.*, 1928, **43**, 1322) was unsuccessful. Allyl bromide was used in place of the allyl iodide used by Blaise (*loc. cit.*).

Interaction of Sodio-3-isopropylcyclopentanone with $\Delta\beta$ -Hexen- δ -one.—A slow current of nitrogen was passed for 4 hours through a stirred suspension of powdered sodamide (3.2 g.) in 3-isopropylcyclopentanone (18 g.) and dry ether (60 c.c.), $\Delta\beta$ -hexen- δ -one (8 g.) added slowly, stirring continued at room temperature for 4 hours, the mixture refluxed for 6 hours, and the product decomposed with hydrochloric acid. An ethereal extract was evaporated, unchanged ketones removed in steam, and the residue distilled at 10 mm., giving fractions (1) 7 g., b. p. 150–170°, (2) 5.1 g., b. p. 170–190°, and (3) 4 g., b. p. 190–220°. Fraction (2) gave a solid which crystallised from aqueous alcohol in white plates, m. p. 75–76°. This was evidently not a condensation product but was formed by the action of sodamide on 3-isopropylcyclopentanone, for the same ketone (probably 1) (5 g.) was isolated when the cyclic ketone (14 g.) was stirred for an hour at –20° with sodamide (3.9 g.) in ether (50 c.c.). This unsaturated ketone was hydrogenated over palladium-black in acetone to a saturated ketone which gave a semicarbazone, m. p. 161.5° (from acetone) (Found: C, 69.8; H, 10.85. $C_{19}H_{31}ON_3$ requires C, 69.6; H, 10.7%), and an oxime, m. p. 142–143.5° (from alcohol) (Found: C, 76.3; H, 11.7. Calc. for $C_{18}H_{29}ON$: C, 76.5; H, 11.65%). Wallach and Worlitzer (*Nachr. K. Ges. Wiss. Göttingen*, 1915, **1**, 17) obtained the bimolecular condensation product of 3-isopropylcyclopentanone by treatment with cold sodium ethoxide. They gave the m. p. as 77–78°, and obtained by hydrogenation a saturated ketone of which the oxime had m. p. 136°.

Condensation of Ethyl 4-isopropylcyclopentanone-2-carboxylate (II) with $\Delta\beta$ -Hexen- δ -one (III).—To the thick paste of sodio-compound obtained by adding the keto-ester (130 g.) to a solution of sodium ethoxide (12 g. of sodium in 650 c.c. of absolute alcohol distilled over quicklime) was added a solution of $\Delta\beta$ -hexen- δ -one (50 g.) in absolute alcohol (200 c.c.). The whole was kept at room temperature for 60 hours (most of the solid had dissolved after 12 hours), the solution then heated, much of the alcohol (650 c.c.) distilled, and the residual solution poured into ice and water and extracted with ether. The aqueous alkaline solution was acidified, and the precipitated oil separated into fractions soluble in sodium carbonate (this was essentially isopropyladipic acid and gave 12 g. of ethyl ester when esterified) and in sodium hydroxide (this fraction gave an intense purple colour with ferric chloride and was apparently unchanged keto-ester). The ethereal extract of neutral product, which gave no colour with alcoholic ferric chloride, was distilled at 12 mm. into fractions (i) 14 g., b. p. 120–170°, and (ii) 70 g., b. p. 170–190°.

Fraction (ii), consisting of the dicyclic keto-ester (IV), was hydrolysed by 6 hours' boiling with alcohol (400 c.c.) and potassium hydroxide (50 g. in 35 c.c. of water), potassium carbonate separating. Some of the alcohol was distilled off, and the solution diluted with water and extracted with ether. The alkaline solution gave very little product on acidification. The ethereal extract was distilled and gave a yellowish liquid (43 g.), b. p. 160–170°/14 mm. This was essentially the unsaturated ketone formed by decarboxylation of (IV), but it yielded no crystalline derivatives.

Hydrogenation of this material, dissolved in alcohol (250 c.c.), over palladium-black (0.5 g.) was complete in 25 hours (hydrogen absorption, 4.2 l.). Treatment of an aqueous-alcoholic solution of the resulting saturated ketones (V) with semicarbazide, followed by crystallisation from alcohol, gave 7.2 g. of pure semicarbazone, which formed colourless leaflets, m. p. 204–205° (Found: C, 68.4; H, 10.2; N, 15.7. $C_{17}H_{27}ON_3$ requires C, 68.0; H, 10.3; N, 15.85%). The oxime, prepared from another portion of the mixed saturated ketones by heating with hydroxylamine hydrochloride in pyridine, formed fine white needles, m. p. 154.5° (Found: C, 75.1; H, 10.9. $C_{14}H_{25}ON$ requires C, 75.3; H, 11.3%).

The liquor from which the pure semicarbazone had been isolated gave only resinous material on dilution with water. This consisted of a semicarbazone mixture, from which the ketones were regenerated by steam-distillation with 20% sulphuric acid. The resulting ketone mixture was used in some experiments described below. The poor yield of pure semicarbazone hindered the subsequent stages of the synthesis, and the yield was not improved in other experiments in which the conditions of interaction of (II) and (III) were somewhat modified.

Stereochemically Pure 4:7-Dimethyl-2-isopropyl-5-hydrindanone (V).—The pure semicarbazone, m. p. 204°, was hydrolysed by steam-distillation with 20% sulphuric acid (200 c.c.). The regenerated ketone was extracted from the steam-distillate with ether, and the extract dried (sodium sulphate) and evaporated on the water-bath. The pure ketone, distilled at 0.3 mm. from an air-bath at 100°, formed a colourless liquid with a feeble peppermint odour, and had d_4^{16} 0.9419 and n_D^{16} 1.4778; whence $[R_D]_D = 62.52$ (calc., 62.70) (Found: C, 80.4; H, 11.7. $C_{14}H_{24}O$ requires C, 80.7; H, 11.6%). Reconversion of this ketone into the same semicarbazone from which it was prepared showed that the sulphuric acid used for hydrolysis did not effect configurational change. A crystalline benzylidene derivative was not obtained from this ketone.

Dehydrogenation of the pure ketone (1 g.) with palladium-black (0.1 g.) was effected by heating for 9 hours at 290–300° in an evacuated sealed tube. The resulting 4:7-dimethyl-2-isopropyl-5-indanol (IX) (0.25 g.), isolated by digestion with light petroleum, in which it was sparingly soluble, crystallised from hexane in colourless needles, m. p. 133–134° (Found: C, 81.9; H, 9.5. Calc. for $C_{14}H_{20}O$: C, 82.3; H, 9.9%) (St. Pfau and Plattner, *Helv. Chim. Acta*, 1940, **23**, 792, give m. p. 129–130°). This dehydrogenation product had the character of a cryptophenol (cf. Farinholt, Harden, and Twiss, *J. Amer. Chem. Soc.*, 1933, **55**, 3383). It gave a 3:5-dinitrobenzoate, which crystallised from alcohol in fine, colourless needles, m. p. 145–146° (Found: C, 63.6; H, 5.8; N, 7.4. $C_{21}H_{22}O_6N_2$ requires C, 63.3; H, 5.6; N, 7.0%).

β -(2-Acetyl-4-isopropylcyclopentyl)butyric Acid (VI).—A solution of the pure ketone (V) (0.5 g.) in glacial acetic acid was treated with a concentrated aqueous solution of chromic acid (0.2 g.). The solution was warmed to 75° and then allowed to cool slowly. The acidic oxidation product gave a semicarbazone ((0.15 g.) which crystallised from alcohol in

short colourless needles, m. p. 181—182° (Found: C, 60.9; H, 9.2; N, 14.3. $C_{15}H_{17}O_3N_3$ requires C, 60.5; H, 9.1; N, 14.1%).

Hydrolysis of this semicarbazone (1.5 g.) by aqueous sulphuric acid in boiling alcohol, followed by alkaline hydrolysis of the ester, gave colourless slender needles, m. p. 64—65° (from hexane), consisting of β -(2-acetyl-4-isopropylcyclopentyl)-butyric acid (VI) (Found: C, 69.9; H, 10.2. $C_{11}H_{24}O_3$ requires C, 69.9; H, 10.1%). Its *p*-phenylphenacyl ester formed colourless leaflets (from alcohol), m. p. 75—76° (Found: C, 77.8; H, 7.9. $C_{22}H_{24}O_4$ requires C, 77.8; H, 7.7%).

Examination of Crude Mixture of Stereoisomeric Dicyclic Ketones (V).—The material recovered by steam-distillation with 20% sulphuric acid of the mixture present in liquors from which the semicarbazone, m. p. 204°, had been isolated (see above) was treated with chromic acid with the object of oxidising any hydroxylic compounds present. For this purpose a solution of chromic acid (20 g.) in 80% acetic acid (50 c.c.) was added to a solution of the oil in glacial acetic acid (250 c.c.). The temperature rose slowly to 45°. After being kept overnight, the solution was poured into water and extracted with ether. By shaking the extract with sodium carbonate solution there was removed some acidic oxidation product, isolated as a yellow gum (8 g.). This gave a solid mixture of semicarbazones (6 g.), which was separated by fractional crystallisation from alcohol into the semicarbazone (2.6 g.), m. p. 181°, of the substituted butyric acid (VI) already described, and a more soluble semicarbazone, m. p. 151—155°, of an isomeric keto-acid. This crystallised from alcohol in colourless leaflets (2 g.) (Found: C, 60.8; H, 9.2%) and was hydrolysed to a keto-acid which formed a gum. This gave a *p*-phenylphenacyl ester, which crystallised from methyl alcohol in colourless microscopic leaflets, m. p. 66—67° (Found: C, 77.3; H, 7.9%). The m. p. was slightly but definitely depressed by the stereoisomeric ester, m. p. 75°. As one of the three substituents in the cyclopentane ring is an acetyl group, it is unlikely that there would be more than two optically inactive stable forms of the keto-acid (VI). In the case of the dicyclic ketone (V) the number of possible stereoisomerides is considerable.

The neutral fraction recovered from this oxidation experiment formed a yellow oil (25 g.), b. p. 145—155°/12 mm. Crystalline semicarbazones could not be obtained from it, and it was resistant to hydrogenation over platinum-black. This material was used for subsequent stages of the synthesis.

Action of Diazomethane on 4:7-Dimethyl-2-isopropyl-5-hydrindanone (V).—(i) In view of the finding of Adamson and Kenner (*loc. cit.*) that ethereal diazomethane reacts exceedingly slowly with 2-methylcyclohexanone we adopted in the first instance the experimental conditions which enabled these workers to obtain 2- and 3-methylcycloheptanone. Anhydrous potassium carbonate (0.1 g.) was added to a solution of the pure dicyclic ketone (V) (2 g.) in absolute alcohol (2 c.c.). Nitrosomethylurethane (1.4 g.) was added during 3 days, the mixture being kept at 0°. After 2 further days the alcohol was removed from the filtered solution. The resulting oil, which, like the original ketone, did not react with sodium bisulphite, was treated with semicarbazide hydrochloride and sodium acetate in cold aqueous alcoholic solution. The resulting crystalline product was not homogeneous, but by crystallisation from alcohol there was readily isolated a semicarbazone in well-formed colourless needles, m. p. 167—168° (Found: C, 67.9, 68.1, 68.7; H, 10.1, 10.2, 10.15; N, 15.6, 15.9. $C_{15}H_{19}ON_3$ requires C, 68.0; H, 10.3; N, 15.85%. $C_{14}H_{19}ON_3$ requires C, 68.8; H, 10.5; N, 15.0%). These analyses, carried out on two different specimens of the semicarbazone, suffice to show that isomerisation rather than ring-enlargement had taken place.

(ii) Methyl alcohol (9 c.c.) was added to a solution of the pure ketone (V) (2 g.) in ethereal diazomethane (from 3 c.c. of nitrosomethylurethane, 35 c.c. of ether, and 3.6 c.c. of 25% methyl-alcoholic potassium hydroxide). Slow liberation of nitrogen set in and after 48 hours this had almost ceased and the solution was nearly colourless. The oil remaining after removal of the solvents gave a semicarbazone (2.7 g.), m. p. 145—150°. A pure component could not be obtained from this by fractional crystallisation. An unsuccessful attempt was made to obtain a crystalline dibenzylidene derivative from the ketone mixture formed from this by hydrolysis with 20% sulphuric acid (tetrahydro- β -vetivone gives such a derivative).

(iii) The neutral oil recovered from the oxidation of the mixture of ketones (V) (20 g.) was treated with diazomethane for 5 days at room temperature, as described under (ii). Ketonic material (11 g.) was separated from the product by means of Girard's reagent T. Treatment of some of this (5 g.) with semicarbazide gave a resinous mixture of semicarbazones, an alcoholic solution of which deposited crystals (0.5 g.) on long standing. These consisted of the semicarbazone, m. p. 167—168°, described under (i).

Vetivazulene (VIII).—Although tetrahydro- β -vetivone (VII) or a stereoisomeride was not isolated from the foregoing experiments with diazomethane, the presence in the mixed products of a compound of this structure was shown by dehydrogenation. The ketonic material obtained as described in the previous paragraph (5 c.c.) was heated with selenium (10 g.) at 280—310° for 24 hours. The product was extracted with ether and distilled at 10—11 mm. from a bath slowly heated from 155° to 190°, the final fractions being taken off at 0.1 mm. The violet distillate, which was partly crystalline, was treated with hexane and gave crystals of 4:7-dimethyl-2-isopropyl-5-indanol (IX) (0.75 g.). The violet hexane liquors were diluted with hexane, and the solution passed through a column of alumina. The violet zone was separated and eluted with alcohol, the solvent mostly evaporated, and the concentrated solution treated with *s*-trinitrobenzene (100 mg.). The crystalline product, recrystallised from alcohol, formed reddish-brown needles, m. p. 150—151.5°. This was the complex of vetivazulene (VIII), for which St. Pfau and Plattner (*Helv. Chim. Acta*, 1936, **19**, 871; 1939, **22**, 207) give m. p. 151.5° (Found: C, 61.5; H, 5.3; N, 10.3. Calc. for $C_{15}H_{19}, C_6H_5O_2N_3$: C, 61.3; H, 5.15; N, 10.2%).

The analyses were all micro-analyses carried out by Mr. J. M. L. Cameron. We thank the Carnegie Trust for a scholarship which enabled one of us (R. R. C.) to take part in this work.