

## 79. *Spectroscopic Evidence for the N-H-N Bond in Ethyleneimine.*

By H. W. THOMPSON and G. P. HARRIS.

The infra-red absorption spectrum of solutions of ethyleneimine in carbon tetrachloride in the region of  $3\mu$  has been measured, and the results suggest the occurrence of association through N-H-N bonds. Other experimental results support this conclusion.

SEVERAL workers have discussed the existence of N-H-N bonds. For example, Hunter and his collaborators (*J.*, 1941, 1, 777, 820) have shown, by consideration of the physical properties and molecular weights, that this type of bond probably occurs in pyrazoles, indazoles, amidines, and related compounds. Gordy (*J. Chem. Physics*, 1939, 7, 170) also found a displacement of the infra-red absorption band of aniline at  $2.9\mu$  towards shorter wave-lengths when solutions in inert organic solvents were used, and he attributed this effect to the same cause.

During a recent measurement of the infra-red absorption spectrum of ethyleneimine it was noticed that the absorption band at  $3\mu$  due to the NH group is appreciably displaced in wave-length on passing from the vapour to the liquid, whereas other bands are not much affected. The relationships have therefore been examined more closely using solutions in carbon tetrachloride, the spectrum being measured between about  $2.9\mu$  and  $3.5\mu$ . The results provide strong evidence for the occurrence of hydrogen bonds in this case.

### EXPERIMENTAL.

Ethyleneimine, prepared from ethanolamine by Wenker's method (*J. Amer. Chem. Soc.*, 1935, 57, 2328), was dried over sodium, redistilled, and finally refractionated in a vacuum. Carbon tetrachloride for use as solvent was dried and redistilled.

A Hilger D 88 spectrometer with quartz prism was used, with a Hilger-Schwarz compensated vacuum thermocouple



arguments support this conclusion. Thus, ethyleneimine boils at  $55.5^{\circ}$ , *C*-methylethyleneimine at  $67^{\circ}$ , but *N*-methylethyleneimine at  $27.5^{\circ}$ . Replacement of the hydrogen atom of the N-H link removes the possibility of a hydrogen bridge, and when there is no association the boiling point is reduced. Measurements (with W. G. Leeds) on the absorption spectrum in the region of  $1\mu$  also suggest the formation of hydrogen bonds. The N-H absorption band of liquid ethyleneimine at  $0.9\mu$  is very broad, with a weak component on the higher-frequency side. With dilute solutions in carbon tetrachloride or chloroform the broad band is less pronounced and the higher-frequency component sharpens. Another interesting observation is that the displaced line in the Raman spectrum of ethyleneimine due to the N-H vibration frequency appears particularly broad.

THE PHYSICAL CHEMISTRY LABORATORY, OXFORD.

[Received, March 29th, 1944.]

---