

Nature of previously reported thermally unstable products derived from the reaction of $[\text{Co}_2(\text{CO})_8]$ with PhSSPh, EtSSEt or PhSeSePh and of the reactions of these products with alkynes or isocyanides

John E. Davies, Martin J. Mays,*† Paul R. Raithby, Vallipuram Sarveswaran and Gregory P. Shields

Department of Chemistry, Lensfield Road, Cambridge, UK CB2 1EW

Thermally unstable red complexes previously isolated from the reaction of $[\text{Co}_2(\text{CO})_8]$ with REER (E = S, R = Ph or Et; E = Se, R = Ph) and assigned the structure $[\text{Co}_3(\mu_3\text{-ER})(\text{CO})_9]$ have been reformulated as the trinuclear complexes $[\text{Co}_3(\mu_3\text{-E})\text{R}(\text{CO})_8]$ (E = S, R = Ph **1a** or Et **1b**; E = Se, R = Ph **1c**) on the basis of their spectroscopic properties and the nature of their reactions with alkynes and isocyanides. Thus the complexes $[\text{Co}_3(\mu_3\text{-E})\text{R}(\text{CO})_8]$ **1a–1c** react with $\text{PhC}\equiv\text{CH}$ to give $[\text{Co}_3(\mu_3\text{-E})\{\text{PhCCHC}(\text{O})\text{R}\}(\text{CO})_7]$ **2a–2c** respectively (E = S, R = Ph **2a** or Et **2b**; E = Se, R = Ph **2c**). It is proposed that **2a–2c** are formed *via* migration of the R group in **1a–1c** from the metal to a carbonyl carbon atom to give an acyl group. Subsequent insertion of $\text{PhC}\equiv\text{CH}$ into the metal–acyl group bond is followed by co-ordination of the acyl oxygen to give a five-membered metallacyclic ring. The reaction of $\text{Bu}^t\text{N}\equiv\text{C}$ with **1a–1c** gave the complexes $[\text{Co}_3(\mu_3\text{-E})(\mu\text{-Bu}^t\text{N}=\text{CR})(\text{CO})_7]$ **3a–3c** respectively (E = S, R = Ph **3a** or Et **3b**; E = Se, R = Ph **3c**) and a similar reaction pathway is proposed. The structures of complexes **2a** and **3a** have been determined by single-crystal X-ray diffraction studies.

The reaction of $[\text{Co}_2(\text{CO})_8]$ with dialkyl and diaryl disulfides, RSSR, was first studied more than thirty years ago by Klumpp and co-workers.^{1,2} The products obtained were found to depend on the nature of the R groups and, for R = Ph, the two thermally stable products of the reaction were originally formulated as $[\text{Co}_6\text{S}(\text{SPh})_5(\text{CO})_{10}]$ and $[\text{Co}_3(\text{SPh})_5(\text{CO})_4]$; a further air-sensitive and thermally unstable red product was formulated as the SPh-capped tricobalt complex $[\text{Co}_3(\mu_3\text{-SPh})(\text{CO})_9]$.¹ For R = *p*-tolyl an analogous unstable red complex, formulated as $[\text{Co}_3(\mu_3\text{-SC}_6\text{H}_4\text{Me-}p)(\text{CO})_9]$ and obtained in less than 1% yield, was the only reaction product,¹ whereas the reaction between EtSSEt and $[\text{Co}_2(\text{CO})_8]$ ² led to three stable complexes formulated as $[\text{Co}_3(\mu_3\text{-S})(\mu_3\text{-SEt})(\text{CO})_6]$, $[\text{Co}_4(\text{SEt})_7(\text{CO})_5]$ and $[\text{Co}_4(\text{SEt})_3(\text{CO})_7]$ but no thermally unstable red product was reported.

Several of the stable cluster complexes obtained in the above reactions were later structurally characterised by X-ray analysis and reformulated in the light of the structural evidence,^{3,4} but the nature of the thermally unstable red complex obtained in the reactions with R = Ph or *p*-tolyl and formulated as $[\text{Co}_3(\mu_3\text{-SR})(\text{CO})_9]$ remained unclear. Recently the reaction of $[\text{Co}_2(\text{CO})_8]$ with bis(perfluorophenyl) disulfide, $\text{C}_6\text{F}_5\text{SSC}_6\text{F}_5$, has been studied and the product structurally characterised as the trinuclear complex $[\text{Co}_3(\mu_3\text{-S})(\text{C}_6\text{F}_5)(\text{CO})_8]$.⁵ We now present spectroscopic and chemical evidence that all the thermally unstable red products obtained in the reaction of RSSR with $[\text{Co}_2(\text{CO})_8]$ and originally formulated as $[\text{Co}_3(\mu_3\text{-SR})(\text{CO})_9]$ are analogous to the above trinuclear pentafluorophenyl complex and should be reformulated as $[\text{Co}_3(\mu_3\text{-S})\text{R}(\text{CO})_8]$. We show also that a similar unstable red complex can be obtained in the reaction of EtSSEt with $[\text{Co}_2(\text{CO})_8]$, even though such a product was not reported in the original investigation. Only traces of the originally reported complexes $[\text{Co}_4(\text{SEt})_7(\text{CO})_5]$ and $[\text{Co}_4(\text{SEt})_3(\text{CO})_7]$, later reformulated as $[\text{Co}_3(\text{SEt})_5(\text{CO})_4]$ ³ and $[\text{Co}_6\text{S}(\text{SEt})_4(\text{CO})_{11}]$ ⁴ respectively, were obtained under the milder reaction condition employed in the present study. Finally we show that diphenyl diselenide gives a corresponding product when it is treated with $[\text{Co}_2(\text{CO})_8]$. The chemical evi-

dence which supports these conclusions is provided by the nature of the reactions of the unstable red products with $\text{PhC}\equiv\text{CH}$ and with Bu^tNC .

Results and Discussion

(a) Reaction of $[\text{Co}_2(\text{CO})_8]$ with PhSSPh, EtSSEt or PhSeSePh

These reactions when carried out in hexane solution at room temperature give red products which, on the basis of their spectroscopic properties and further reactions with alkynes or isocyanides (see below), may be formulated as the trinuclear complexes $[\text{Co}_3(\mu_3\text{-E})\text{R}(\text{CO})_8]$ (E = S, R = Ph **1a** or Et **1b**; E = Se, R = Ph **1c**) (Scheme 1). Thus the overall pattern of five $\nu(\text{CO})$ absorption bands seen in the IR spectrum of each of the complexes **1a–1c** (Table 1) is closely similar to that of the analogous complex $[\text{Co}_3(\mu_3\text{-S})(\text{C}_6\text{F}_5)(\text{CO})_8]$.⁵ In the ¹H NMR spectra of **1b** the ethyl protons give rise to two separate multiplets (δ 2.85 and 2.60) due to the diastereotopic CH_2 protons and to a doublet of doublets (δ 1.58) due to the CH_3 protons. The diastereotopic nature of the CH_2 group can be understood in terms of the asymmetry of the $\text{Co}_3(\mu_3\text{-S})(\text{CO})_8$ moiety to which the Et group is attached. The structure originally proposed for **1a**, $[\text{Co}_3(\mu_3\text{-SPh})(\text{CO})_9]$, is not similarly asymmetric and if **1b** had a structure analogous to **1a** it would not give rise to a diastereotopic CH_2 group.

Partial decomposition on attempted recrystallisation precluded analytically pure samples of complex **1a–1c** being obtained but the FAB mass spectra showed peaks at *m/z* values corresponding to the respective molecular ions and to fragment ions obtained by stepwise loss of carbonyl ligands from the molecular ions.

(b) Reaction of $[\text{Co}_3(\mu_3\text{-E})\text{R}(\text{CO})_8]$ (E = S, R = Ph **1a** or Et **1b**; E = Se, R = Ph **1c**) with $\text{PhC}\equiv\text{CH}$

The complexes **1a–1c** were treated in hexane with 1 equivalent of $\text{PhC}\equiv\text{CH}$ to give $[\text{Co}_3(\mu_3\text{-E})\{\text{PhCCHC}(\text{O})\text{R}\}(\text{CO})_7]$ (E = S, R = Ph **2a** or Et **2b**; E = Se, R = Ph **2c**) as red crystalline solids. The products have been characterised by microanalysis, mass spectrometry and IR, ¹H and ¹³C-¹H NMR spectroscopy (see

† E-Mail: mjm14@cam.ac.uk

Table 2 Selected bond lengths (Å) and angles (°) for $[\text{Co}_3(\mu_3\text{-S})\{\text{PhCCHC}(\text{O})\text{Ph}\}(\text{CO})_7]$ **2a** with estimated standard deviation in parentheses

| | | | |
|-------------------|------------|-------------------|------------|
| Co(1)–C(17) | 1.926(4) | Co(1)–O(15) | 1.953(3) |
| Co(1)–S | 2.1316(14) | Co(1)–Co(3) | 2.5163(11) |
| Co(1)–Co(2) | 2.5342(9) | Co(2)–S | 2.1650(14) |
| Co(2)–Co(3) | 2.5582(11) | Co(3)–S | 2.149(2) |
| C(8)–C(14) | 1.471(6) | C(14)–O(15) | 1.276(5) |
| C(14)–C(16) | 1.414(6) | C(16)–C(17) | 1.369(6) |
| C(17)–C(18) | 1.476(6) | | |
| C(17)–Co(1)–O(15) | 84.3(2) | C(17)–Co(1)–S | 111.82(13) |
| O(15)–Co(1)–S | 152.19(10) | C(17)–Co(1)–Co(3) | 100.50(3) |
| C(15)–Co(1)–Co(3) | 101.76(9) | S–Co(1)–Co(3) | 54.31(4) |
| C(17)–Co(1)–Co(2) | 160.71(13) | C(15)–Co(1)–Co(2) | 103.56(9) |
| S–Co(1)–Co(2) | 54.47(4) | Co(3)–Co(1)–Co(2) | 60.87(3) |
| S–Co(2)–Co(1) | 53.25(4) | S–Co(2)–Co(3) | 53.33(4) |
| Co(1)–Co(2)–Co(3) | 59.22(3) | S–Co(3)–Co(1) | 53.68(4) |
| S–Co(3)–Co(2) | 53.92(4) | Co(1)–Co(3)–Co(2) | 59.91(3) |
| Co(1)–S–Co(3) | 72.01(5) | Co(1)–S–Co(2) | 72.28(5) |
| Co(3)–S–Co(2) | 72.75(5) | O(15)–C(14)–C(16) | 117.0(4) |
| O(15)–C(14)–C(8) | 118.4(4) | C(14)–O(15)–Co(1) | 112.4(3) |
| C(17)–C(16)–C(14) | 116.3(4) | C(16)–C(17)–C(18) | 121.5(4) |
| C(16)–C(17)–Co(1) | 110.0(3) | C(18)–C(17)–Co(1) | 128.5(3) |





Fig. 1 Molecular structure of $[\text{Co}_3(\mu_3\text{-S})\{\text{PhCCHC}(\text{O})\text{Ph}\}(\text{CO})_7]$ **2a** including the atom numbering scheme



Scheme 2 Possible reaction pathway for the formation of complexes **2a–2c** from **1a–1c**

is slightly twisted (torsion angle 8.2°). The dihedral angle between the mean plane through these atoms and the Co_3 triangular core is 170.6° ; the plane through the atoms C(13), C(12), N lies closer to the plane through the Co_3 triangle

Table 3 Selected bond lengths (Å) and angles (°) for $[\text{Co}_3(\mu_3\text{-S})(\mu\text{-Bu}^t\text{NCPh})(\text{CO})_7]$ **3a** with estimated standard deviation in parentheses

| | | | |
|-------------------|-----------|-------------------|-----------|
| Co(1)–N | 2.010(8) | Co(1)–S | 2.155(3) |
| Co(1)–Co(2) | 2.411(2) | Co(1)–Co(3) | 2.575(2) |
| Co(2)–C(12) | 1.964(9) | Co(2)–S | 2.160(3) |
| Co(2)–Co(3) | 2.545(2) | Co(3)–S | 2.183(3) |
| N–C(12) | 1.285(12) | | |
| N–Co(1)–S | 95.0(2) | N–Co(1)–Co(2) | 74.9(2) |
| S–Co(1)–Co(2) | 56.13(9) | N–Co(1)–Co(3) | 135.3(2) |
| S–Co(1)–Co(3) | 54.08(9) | Co(2)–Co(1)–Co(3) | 61.28(6) |
| C(12)–Co(2)–S | 95.8(3) | C(12)–Co(2)–Co(1) | 71.9(3) |
| S–Co(2)–Co(1) | 55.94(9) | C(12)–Co(2)–C(3) | 134.2(3) |
| S–Co(2)–Co(3) | 54.54(8) | Co(1)–Co(2)–Co(3) | 62.55(6) |
| S–Co(3)–Co(2) | 53.71(9) | S–Co(3)–Co(1) | 53.09(8) |
| Co(2)–Co(3)–Co(1) | 56.17(5) | Co(1)–S–Co(2) | 67.93(9) |
| Co(1)–S–Co(3) | 72.83(10) | Co(2)–S–Co(3) | 71.75(10) |
| C(12)–N–C(11) | 130.0(8) | C(12)–N–Co(1) | 101.6(6) |
| C(11)–N–Co(1) | 128.4(7) | N–C(12)–C(13) | 130.6(9) |
| N–C(12)–Co(2) | 111.1(7) | | |




Fig. 2 Molecular structure of $[\text{Co}_3(\mu_3\text{-S})(\mu\text{-Bu}^t\text{NCPh})(\text{CO})_7]$ **3a** including the atom numbering scheme

(dihedral angle 5.7°) than that through C(12), N, C(11) (10.3°). The planar co-ordination around C(12) and N indicates sp^2 hybridisation which is consistent with the C(12)–N bond distance of 1.285(12) Å (within the range of C–N double-bond distances).¹¹


A plausible reaction pathway for the formation of complexes **3a–3c** from **1a–1c** is illustrated in Scheme 3. It is proposed that a carbonyl ligand on the metal atom to which the R group is attached is substituted by the Bu^tNC ligand. Migration of the R group from the cobalt to the carbon atom of the isocyanide group is then followed by co-ordination of the nitrogen atom of the resulting imino group to the other metal centre. This must be accompanied by migration of a CO ligand from one cobalt atom to the other to maintain the requirements of the EAN rule for each centre. Complexes similar to **3a–3c** have been synthesized previously *via* the reaction of $[\text{Co}_2(\text{CO})_8]$ with the thioamides $\text{R}'\text{C}(\text{S})\text{NHR}''$ ($\text{R}' = \text{Me}$ or Ph , $\text{R}'' = \text{C}_6\text{H}_{11}$).⁸

Experimental

Unless otherwise stated all reactions were carried out under a nitrogen atmosphere. Solvents were distilled from the appropriate drying agent prior to use.¹² Products are presented in order of decreasing R_f values. Unless otherwise stated, all reagents were obtained from commercial suppliers and used without further purification. The instrumentation used to obtain the spectroscopic data has been described previously.¹³

Reactions of $[\text{Co}_2(\text{CO})_8]$

With PhSSPh. To a solution of $[\text{Co}_2(\text{CO})_8]$ (1.0 g, 2.924



Scheme 3 Possible reaction pathway for the formation of complexes **3a–3c** from **1a–1c**

mmol) in hexane (100 cm³) was added PhSSPh (0.213 g, 0.974 mmol) and the resulting solution stirred at room temperature for 2 h. After removal of solvent the residue was taken up in the minimum volume of dichloromethane and adsorbed onto silica. The silica was pumped dry and added to the top of a silica chromatography column. Elution with hexane–dichloromethane (4:1) gave red [Co₃(μ₃-S)Ph(CO)₈] **1a**. ¹³C NMR (CDCl₃, 20 °C, ¹H composite pulse decoupled): δ 208.4 (CO), 208.2 (CO) and 133.6–125.1 (m, Ph). FAB mass spectrum: *m/z* 510 (*M*⁺) and *M*⁺ – *n*CO (*n* = 1–6). The yield of unrecrystallised product was 1.0 g.

With EtSSEt. To a solution of [Co₂(CO)₈] (1.0 g, 2.924 mmol) in hexane (100 cm³) was added EtSSEt (0.120 cm³, 0.974 mmol) in three equal portions at 10 min intervals and the resulting solution was then stirred at room temperature for 5 h. Work-up as for complex **1a** gave red [Co₃(μ₃-S)Et(CO)₈] **1b** followed by trace amounts of the known complexes [Co₃(SEt)₅(CO)₄]³ and [Co₆S(SEt)₄(CO)₁₁].⁴ The identity of both the known complexes was confirmed by IR spectroscopy. Complex **1b**: FAB mass spectrum *m/z* 462 (*M*⁺) and *M*⁺ – *n*CO (*n* = 1–4). The yield of unrecrystallised product **1b** was 0.5 g.

With PhSeSePh. To a solution of [Co₂(CO)₈] (1.0 g, 2.924 mmol) in hexane (100 cm³) was added PhSeSePh (0.304 g, 0.974 mmol) and the resulting solution stirred at room temperature for 2 h. Work-up as for **1a** gave red [Co₃(μ₃-Se)Ph(CO)₈] **1c**. ¹³C NMR (CDCl₃, 20 °C, ¹H composite pulse decoupled): δ 134.7–124.2 (m, Ph). FAB mass spectrum: *m/z* 558 (*M*⁺) and *M*⁺ – *n*CO (*n* = 1–8). The yield of unrecrystallised product was 0.65 g.

Reactions of PhC≡CH

With [Co₃(μ-S)Ph(CO)₈] 1a. To a solution of complex **1a** (0.30 g, 0.588 mmol) in dichloromethane (50 cm³) was added PhC≡CH (0.06 cm³, 0.57 mmol) and the resulting solution stirred at room temperature for 4 h. The solvent was then removed on a rotary evaporator, the residue taken up in the minimum volume of dichloromethane and adsorbed onto silica. The silica was pumped dry and added to the top of a silica chromatography column. Elution with hexane gave brown [Co₃(μ₃-S){PhCCHC(O)Ph}(CO)₇] **2a** (0.072 g, 20%) (Found: C, 43.03; H, 1.85. C₂₂H₁₁Co₃O₈S requires C, 43.14; H, 1.80%). ¹³C NMR (CDCl₃, 20 °C, ¹H composite pulse decoupled): δ 142.4–137.9 (m, Ph) and 131.2–125.2 [m, μ-PhCC(H)C(O)]. FAB mass spectrum: *m/z* 613 (*M*⁺) and *M*⁺ – *n*CO (*n* = 1–7).

With [Co₃(μ-S)Et(CO)₈] 1b. To a solution of complex **1b** (0.30 g, 0.649 mmol) in hexane (50 cm³) was added PhC≡CH (0.07 cm³, 0.64 mmol) and the resulting solution stirred at

room temperature for 3 h. Work-up as for **2a** gave brown [Co₃(μ₃-S){PhCCHC(O)Et}(CO)₇] **2b** (0.15 g, 40%) (Found: C, 38.21; H, 1.85. C₁₈H₁₁Co₃O₈S requires C, 38.30; H, 1.95%). FAB mass spectrum *m/z* 565 (*M*⁺) and *M*⁺ – *n*CO (*n* = 1–7).

With [Co₃(μ-Se)Ph(CO)₈] 1c. To a solution of complex **1c** (0.30 g, 0.538 mmol) in hexane (50 cm³) was added PhC≡CH (0.06 cm³, 0.54 mmol) and the resulting solution stirred at room temperature for 3 h. Work-up as for **2a** gave brown [Co₃(μ₃-Se){PhCCHC(O)Ph}(CO)₇] **2c** (0.16 g, 45%) (Found: C, 40.18; H, 1.85. C₂₂H₁₁Co₃O₈Se requires C, 40.06; H, 1.70%). FAB mass spectrum *m/z* 660 (*M*⁺) and *M*⁺ – *n*CO (*n* = 1–6).

Reactions of Bu^tNC

With [Co₃(μ-S)Ph(CO)₈] 1a. To a solution of complex **1a** (0.30 g, 0.588 mmol) in hexane (50 cm³) was added Bu^tNC (0.06 cm³, 0.58 mmol) and the resulting solution stirred at room temperature for 3 h. The solvent was then removed on a rotary evaporator, the residue taken up in the minimum volume of dichloromethane and adsorbed onto silica. The silica was pumped dry and added to the top of a silica chromatography column. Elution with hexane–dichloromethane (3:2) gave brown [Co₃(μ₃-S)(μ-Bu^tNCPH)(CO)₇] **3a** (0.22 g, 65%) (Found: C, 38.18; H, 2.42. C₁₈H₁₄Co₃NO₇S requires C, 38.23; H, 2.50%). ¹³C NMR (CDCl₃, 20 °C, ¹H composite pulse decoupled): δ 208.9 (CO), 148.3 (PhCN), 129.0–126.6 (m, Ph), 121.5 (s, Ph), 121.0 (s, Ph), 63.9 (Me₃C) and 30.95 Me. FAB mass spectrum: *m/z* 565 (*M*⁺) and *M*⁺ – *n*CO (*n* = 1–7).

With [Co₃(μ-S)Et(CO)₈] 1b. To a solution of complex **1b** (0.30 g, 0.649 mmol) in hexane (50 cm³) was added Bu^tNC (0.07 cm³, 0.64 mmol) and the resulting solution stirred at room temperature for 2 h. Work-up as for **3a** except for the use of hexane–dichloromethane (4:1) as the solvent gave brown [Co₃(μ₃-S)(μ-Bu^tNCEt)(CO)₇] **3b** (0.18 g, 55%) (Found: C, 32.30; H, 2.85. C₁₄H₁₄Co₃NO₇S requires C, 32.50; H, 2.71%). ¹³C NMR (CDCl₃, 20 °C, ¹H composite pulse decoupled): δ 208.7 (CO), 146.3 (EtCN), 60.8 (Me₃C), 40.9 (CH₂CH₃), 31.0 [(CH₃)₃C] and 11.8 (CH₃CH₂). FAB mass spectrum: *m/z* 517 (*M*⁺) and *M*⁺ – *n*CO (*n* = 1–6).

With [Co₃(μ-Se)Ph(CO)₈] 1c. To a solution of complex **1c** (0.30 g, 0.538 mmol) in hexane (50 cm³) was added Bu^tNC (0.06 cm³, 0.54 mmol) and the resulting solution stirred at room temperature for an hour. Work-up as for **3a** gave brown [Co₃(μ₃-Se)(μ-Bu^tNCPH)(CO)₇] **3c** (0.16 g, 50%) (Found: C, 35.33; H, 2.32. C₁₈H₁₄Co₃NO₇Se requires C, 35.29; H, 2.29%). ¹³C NMR (CDCl₃, 20 °C, ¹H composite pulse decoupled): δ 208.2 (CO), 146.3 (PhCN), 128.7–123.3 (m, Ph), 121.3 (s, Ph), 120.8 (s, Ph), 63.8 (Me₃C) and 30.9 [(CH₃)₃C]. FAB mass spectrum: *m/z* 612 (*M*⁺) and *M*⁺ – *n*CO (*n* = 1–7).

Crystallography

Data for [Co₃(μ₃-S){PhCCHC(O)Ph}(CO)₇] **2a** and [Co₃(μ₃-S)(μ-Bu^tNCPH)(CO)₇] **3a** were collected by the ω–2θ scan method on a Rigaku AFC7R (**2a**) or AFC5R (**3a**) four-circle diffractometer. Three standard reflections were monitored at intervals of 200 reflections. Cell parameters were obtained by least-squares refinement on diffractometer angles from 25 centred reflections (20 < 2θ < 22.5°). Semiempirical absorption corrections based on ψ-scan data were applied.^{14,15}

The structures were solved by direct methods (SIR 92¹⁶) and subsequent Fourier-difference syntheses and refined anisotropically on all non-H atoms by full-matrix least squares on *F*² (SHELXL 93¹⁷). Hydrogen atoms were placed in idealised positions and refined using a riding model or as rigid methyl groups. In the final cycles of refinement a weighting scheme was introduced which produced a flat analysis of variance. Full details are given in Table 4.

Table 4 Crystal data for $[\text{Co}_3(\mu_3\text{-S})\{\text{PhCCHC}(\text{O})\text{Ph}\}(\text{CO})_7]$ **2a** and $[\text{Co}_3(\mu_3\text{-S})(\mu\text{-Bu}^t\text{NCPH})(\text{CO})_7]$ **3a***

| | $[\text{Co}_3(\mu_3\text{-S})\{\text{PhCCHC}(\text{O})\text{Ph}\}(\text{CO})_7]$ 2a | $[\text{Co}_3(\mu_3\text{-S})(\mu\text{-Bu}^t\text{NCPH})(\text{CO})_7]$ 3a |
|------------------------------------------------------|--------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------|
| Molecular formula | $\text{C}_{22}\text{H}_{11}\text{Co}_3\text{O}_8\text{S}$ | $\text{C}_{18}\text{H}_{14}\text{Co}_3\text{NO}_7\text{S}$ |
| <i>M</i> | 612.16 | 565.15 |
| <i>a</i> /Å | 11.422(2) | 8.797(2) |
| <i>b</i> /Å | 13.324(2) | 15.793(3) |
| <i>c</i> /Å | 7.872(2) | 8.5066(11) |
| <i>a</i> /° | 91.42(2) | 104.259(13) |
| <i>β</i> /° | 98.71(2) | 101.344(13) |
| <i>γ</i> /° | 93.640(10) | 83.24(2) |
| <i>U</i> /Å ³ | 1181.1(4) | 1120.04(4) |
| <i>D_c</i> /Mg m ⁻³ | 1.721 | 1.676 |
| Crystal size/nm | 0.20 × 0.10 × 0.10 | 0.3 × 0.3 × 0.1 |
| Crystal habit | Red brown | Dark red plate |
| <i>F</i> (000) | 608 | 564 |
| <i>μ</i> /mm ⁻¹ | 2.215 | 2.326 |
| Maximum, minimum relative transmission | 0.999, 0.807 | 0.997, 0.888 |
| Data collection range/° | 5.24 < 2θ < 45.04 | 5.02 < 2θ < 45.0 |
| Index ranges | 0 ≤ <i>h</i> ≤ 12, -14 ≤ <i>k</i> ≤ 14, -8 ≤ <i>l</i> ≤ 8 | 0 ≤ <i>h</i> ≤ 9, -16 ≤ <i>k</i> ≤ 16, -9 ≤ <i>l</i> ≤ 8 |
| Reflections measured | 3295 | 3157 |
| Independent reflections (<i>R</i> _{int}) | 3102 (0.019) | 2931 (0.039) |
| Parameters, restraints | 309, 0 | 271, 0 |
| <i>wR</i> 2 (all data) | 0.086 | 0.2068 |
| <i>x</i> , <i>y</i> | 0.0327, 0.278 | 0.100, 0.00 |
| <i>R</i> 1 [<i>I</i> > 2σ(<i>I</i>)] | 0.035 | 0.053 |
| Observed reflections | 2389 | 1986 |
| Goodness of fit on <i>F</i> ² (all data) | 1.057 | 1.296 |
| Maximum shift/σ | 0.001 | 0.000 |
| Peak, hole in final difference map/e Å ⁻³ | 0.343, -0.231 | 0.837, -1.479 |
| Decay (%) | 0 | 7 |

* Data in common: graphite-monochromated Mo-Kα radiation, λ = 0.71073 Å, *T* = 293(2) K; triclinic, space group *P* $\bar{1}$; *Z* = 2; *R*1 = Σ||*F*_o| - |*F*_c||/Σ|*F*_o|, *wR*2 = [Σ*w*(*F*_o² - *F*_c²)²/Σ*wF*_o⁴]^{1/2}, *w* = 1/[σ²(*F*_o)² + (*xP*)² + (*yP*)²], *P* = (*F*_o² + 2*F*_c²)/3, where *x* and *y* are constants adjusted by the program; goodness of fit = {Σ[*w*(*F*_o² - *F*_c²)²]/(*n* - *p*)} where *n* is the number of reflections and *p* the number of parameters.

CCDC reference number 186/871.

See <http://www.rsc.org/suppdata/dt/1998/775/> for crystallographic files in .cif format.

Acknowledgements

We thank the EPSRC for funding for the X-ray diffractometer and for financial support (to J. E. D. and G. P. S.), The United Kingdom Committee of Vice-Chancellors and Principals, The Cambridge Commonwealth Trust and Downing College, Cambridge for financial support (to V. S.) and the Cambridge Crystallographic Data Centre for support (to J. E. D. and G. P. S.).

References

- G. Bor, L. Marko and E. Klumpp, *Chem. Ber.*, 1967, **100**, 1451.
- G. Bor, L. Marko and E. Klumpp, *Chem. Ber.*, 1964, **97**, 926.
- C. H. Wei and L. F. Dahl, *J. Am. Chem. Soc.*, 1968, **90**, 3960.
- C. H. Wei and L. F. Dahl, *J. Am. Chem. Soc.*, 1968, **90**, 3977.
- G. Gervasia, S. Vastag, G. Bor, G. Natile and L. Marko, *Inorg. Chim. Acta*, 1996, **251**, 35.
- A. J. M. Caffyn, M. J. Mays, G. Conole, M. McPartlin and H. R. Powell, *J. Organomet. Chem.*, 1992, **436**, 83.
- A. J. Edwards, A. Martin, M. J. Mays, D. Nazar, P. R. Raithby and G. A. Solan, *J. Chem. Soc., Dalton Trans.*, 1993, 355.
- H. Patin, G. Mignani, C. Mahe, J. Le Marouille, A. Benoit, D. Grandjean and G. Levesque, *J. Organomet. Chem.*, 1981, **208**, C39.
- C. H. Wei and L. F. Dahl, *Inorg. Chem.*, 1967, **6**, 1229.
- H. Werner, K. Leonhard, O. Kolb, E. Röttinger and H. Vahrenkamp, *Chem. Ber.*, 1980, **113**, 1654; J. Fortune, A. R. Manning and F. S. Stephens, *J. Chem. Soc., Chem. Commun.*, 1983, 1071.
- V. Divjakovic and V. Leovac, *Cryst. Struct. Commun.*, 1978, **7**, 689.
- D. P. Perrin, W. L. F. Amarego and D. R. Perrin, *Purification of Laboratory Chemicals*, 3rd edn., Pergamon, Oxford, 1988.
- A. J. M. Caffyn, M. J. Mays and P. R. Raithby, *J. Chem. Soc., Dalton Trans.*, 1991, 2349.
- TEXSAN, Version 1.7-1, Molecular Structure Corporation, The Woodlands, TX, 1985, 1992, 1995.
- A. C. T. North, D. C. Phillips and F. S. Mathews, *Acta Crystallogr., Sect. A*, 1968, **24**, 351.
- A. Altomare, G. Cascarano, C. Giacavazzo, A. Guagliardi, M. C. Burla, G. Polidori and M. Camalli, *J. Appl. Crystallogr.*, 1994, **27**, 435.
- G. M. Sheldrick, SHELXL 93, University of Göttingen, 1993.

Received 27th October 1997; Paper 7/07709B