

Synthesis and structures of new mono- and multi-nuclear benzylin carboxylates and phosphinates: a novel dibenzylin phosphinate stabilized by intramolecular hydrogen bonding


K. C. Kumara Swamy,^{*,†,a} Musa A. Said,^a S. Nagabrahmanandachari,^a Damodara M. Poojary^b and Abraham Clearfield^b

^a School of Chemistry, University of Hyderabad, Hyderabad - 500046, A. P., India






^b Department of Chemistry, Texas A & M University, College Station, Texas 77843-3255, USA

The reaction of dibenzylin dichloride **1** with 2 mol equivalents each of $\text{AgO}_2\text{P}(\text{C}_6\text{H}_{11})_2$ and $(\text{C}_6\text{H}_{11})_2\text{PO}_2\text{H}$ afforded the novel monometallic complex $\text{Sn}(\text{CH}_2\text{Ph})_2[\text{O}_2\text{P}(\text{C}_6\text{H}_{11})_2]_2[\text{HO}_2\text{P}(\text{C}_6\text{H}_{11})_2]_2$ **2**. When **1** was treated with 2 mol equivalents of $\text{AgO}_2\text{P}(\text{C}_6\text{H}_{11})_2$ only the trinuclear O-capped cluster $\{[\text{Sn}(\text{CH}_2\text{Ph})(\text{OH})[\text{O}_2\text{P}(\text{C}_6\text{H}_{11})_2]_3\text{O}][\text{O}_2\text{P}(\text{C}_6\text{H}_{11})_2]_2\}$ **4** was obtained by an Sn–C bond cleavage *via* an intermediate formulated as $\{\text{Sn}(\text{CH}_2\text{Ph})(\text{OH})[\text{O}_2\text{P}(\text{C}_6\text{H}_{11})_2]_2\}$ **3**. In the reaction with silver carboxylates the type of product isolated depends on the carboxylic acid used. Thus the ‘ladder’ $\{[(\text{PhCH}_2)\text{SnO}\cdot\text{O}_2\text{CMe}]_2(\text{PhCH}_2)\text{Sn}(\text{O}_2\text{CMe})_3\}_2$ **5**, the ‘hexagonal prismanes’ $[(\text{PhCH}_2)\text{SnO}\cdot\text{O}_2\text{CR}]_6$ (R = Me **6**, Bu[†] **7**, C₅H₄N **8**, CH=CM₂ **9** or CHPh₂ **10**) and the mononuclear compound $\text{Sn}(\text{CH}_2\text{Ph})_2(\text{O}_2\text{CCHPh}_2)_2$ **11** have been isolated. X-Ray structural analysis confirmed the identities of **2**, **4**·CHCl₃, and **5**·2CH₂Cl₂. In **2** the mononuclear structure is stabilized by internal hydrogen bonding between P–OH and P=O of the co-ordinated phosphinic acid and the phosphinate, respectively, from the same molecule. Compound **4** has a tristanoxane ring with a capping oxygen, whereas in **5** the stannoxane skeleton is that of a ‘ladder’ with three different types of tin.

Several interesting multinuclear monoorganotin carboxylate/phosphinate cages and clusters that include ‘drums’ (hexagonal prismanes) $(\text{RSnO}\cdot\text{O}_2\text{CR})_6$, ‘cubanes’ $(\text{RSnO}\cdot\text{O}_2\text{PR}_2)_4$, and ‘ladders’ $[(\text{RSnO}\cdot\text{O}_2\text{CR})_2\text{RSn}(\text{O}_2\text{CR})_3]_2$ have been reported during the last decade.^{1–3} Most of these compounds have been prepared by either (i) condensation reactions of the appropriate carboxylic/phosphorus-based acids with stannous acids, $\text{RSn}(\text{O})\text{OH}$, or (ii) treating organotin halides with silver salts of carboxylic/phosphinic acids. In one case an ester elimination reaction which gives the hexagonal prismane **I** [equation (1)] is also used.⁴



Another route which leads to mono- and multi-nuclear tin compounds involves Sn–C bond cleavage reactions of phenyltin compounds [equations (2)–(6)].^{5–8} In this connection it is



pertinent that allyltin compounds are also prone to Sn–C bond cleavage which forms the basis of their utility in allylation reactions.⁹ Since a benzyl group may be expected to behave in a manner similar to an allyl group,¹⁰ tin–carbon bond cleavage reactions must be facile for benzylin compounds also. Thus we chose to study the reactions of the readily prepared dibenzylin dichloride, $\text{Sn}(\text{CH}_2\text{Ph})_2\text{Cl}_2$ **1**,¹¹ with silver salts of carboxylic/phosphinic acids in order to see if the cages/clusters mentioned above can be obtained by this route, and also to explore whether products with structures other than the known types are formed.


Herein we report the synthesis and crystal structures of (a) the novel mononuclear compound $\text{Sn}(\text{CH}_2\text{Ph})_2[\text{O}_2\text{P}(\text{C}_6\text{H}_{11})_2]_2[\text{HO}_2\text{P}(\text{C}_6\text{H}_{11})_2]_2$ **2**, (b) the O-capped cluster $\{[\text{Sn}(\text{CH}_2\text{Ph})(\text{OH})[\text{O}_2\text{P}(\text{C}_6\text{H}_{11})_2]_3\text{O}][\text{O}_2\text{P}(\text{C}_6\text{H}_{11})_2]_2\}$ **4**·CHCl₃, which is obtained by hydrolysis of the dimer $\{\text{Sn}(\text{CH}_2\text{Ph})(\text{OH})[\text{O}_2\text{P}(\text{C}_6\text{H}_{11})_2]_2\}$ **3** and (c) the ladder compound $\{[(\text{PhCH}_2)\text{SnO}\cdot\text{O}_2\text{CMe}]_2(\text{PhCH}_2)\text{Sn}(\text{O}_2\text{CMe})_3\}_2 \cdot 2\text{CH}_2\text{Cl}_2$ **5**·2CH₂Cl₂. Compound **2** represents a unique structure stabilized by intramolecular hydrogen bonding. Whereas the formation of **2** involves no Sn–C bond cleavage, that of **4** or **5** involves such a cleavage. In addition to the above compounds the prismanes $[(\text{PhCH}_2)\text{SnO}\cdot\text{O}_2\text{CR}]_6$ (R = Me **6**, Bu[†] **7**, C₅H₄N **8**, CH=CM₂ **9** or CHPh₂ **10**) and the monometallic derivative $\text{Sn}(\text{CH}_2\text{Ph})_2(\text{O}_2\text{CCHPh}_2)_2$ **11** have been isolated from the reaction of **1** with the corresponding silver carboxylates.

† E-Mail: kckssc@uohyd.ernet.in

Results and Discussion


Synthesis and spectroscopy

Compound **2** was synthesized according to equation (7); it is




also formed as a minor product during recrystallization in the reaction of **1** with an excess (>3 mol equivalents) of silver dicyclohexylphosphinate. The ^{31}P NMR spectrum of **2** showed a broad signal at δ 54.7; in the ^1H NMR spectrum the expected integrated intensity ratio is observed with the phosphinic acid OH proton appearing at δ 7.95 [$^4J(\text{Sn}-\text{H}) \approx 25$ Hz]. An analogous reaction of dicyclooctylphosphinic acid and its silver salt with **1** gave a single product [δ (^{31}P) 64.8 (br); ^1H NMR broad with expected integrated intensities], but no crystalline product could be isolated.

In the reaction of compound **1** with 2.5 mol equivalents of $\text{AgO}_2\text{P}(\text{C}_6\text{H}_{11})_2$ alone the crystalline product **3** obtained ini-




ally showed two ^{31}P NMR signals at δ 57.2 [$^2J(\text{Sn}-\text{O}-\text{P}) = 136.4$] and 48.7 [$^2J(\text{Sn}-\text{O}-\text{P}) = 196.4$ Hz] of equal intensity. The ^1H NMR spectrum showed broad signals with a complex pattern but the integrated intensities showed that there are two phosphinates per benzyl group. The ^{119}Sn NMR exhibited a multiplet centred at δ -586.0. These features are analogous to those observed for the dimer $\{\text{SnBu}^n(\text{OH})[\text{O}_2\text{P}(\text{C}_6\text{H}_{11})_2]_2\}_2$ [^{31}P NMR: δ 56.02 [$^2J(\text{Sn}-\text{O}-\text{P}) = 124.4$], 47.60 [$^2J(\text{Sn}-\text{O}-\text{P}) = 190.8$ Hz]. ^{119}Sn NMR: δ -547.47].¹² Hence **3** is assigned an analogous structure. This product is formed most likely by cleavage of an $\text{Sn}-\text{CH}_2\text{Ph}$ bond from the intermediate $\text{Sn}(\text{CH}_2\text{Ph})_2[\text{O}_2\text{P}(\text{C}_6\text{H}_{11})_2]_2$ due to hydrolysis by adventitious moisture present.


Attempted crystallization of compound **3** in air led to the O-capped cluster **4** [equation (8)]. The ^{31}P NMR spectrum, although broad, shows two types of phosphinates [δ 41.5 (1P, free), 52.8 (3P, bridging)]; the ^{119}Sn NMR exhibits a triplet at δ -535.7 [$^2J(\text{Sn}-\text{O}-\text{P}) \approx 137$ Hz] showing that all the tin atoms are equivalent and coupled to adjacent phosphorus nuclei only. It can be noted that $\Delta\delta(^{119}\text{Sn})$ for **4** and the corresponding *n*-butyl compound [$\{\text{SnBu}^n(\text{OH})[\text{O}_2\text{P}(\text{C}_6\text{H}_{11})_2]_3\}_3^+[\text{O}_2\text{P}(\text{C}_6\text{H}_{11})_2]^-$] is 36.4 ppm; the corresponding difference between our benzyl compound **3** and the *n*-butyltin compound $\{\text{SnBu}^n(\text{OH})[\text{O}_2\text{P}(\text{C}_6\text{H}_{11})_2]_2\}_2$ is 38.5 ppm.¹² This lends credence to our assignment of the dimeric structure for **3**.



Treatment of compound **1** with silver carboxylates afforded different types of compounds as isolable products depending on the carboxylic acid used (Scheme 1). The NMR spectra (^1H ,




^{13}C and ^{119}Sn) for the ladder compound **5** show broad signals. In the ^{119}Sn NMR, the signal for the prismane $[(\text{PhCH}_2)_2\text{SnO} \cdot \text{O}_2\text{CMe}]_6$ **6** at δ -523.1 is also seen; peaks for **5** are broad [δ -530 to -580 and -630 to -640]. This solution behaviour is reminiscent of Holmes' observations for *n*-butyltin oxide carboxylate ladders.¹³ Hydrolysis of **5** to the pure prismane **6** can be readily effected [equation (9)]. Compounds **6**–**10** show charac-




teristic singlet resonances in the ^{119}Sn NMR spectra between δ -519 and -523, upfield by *ca.* 35 ppm to those of the corresponding *n*-butyltin compounds $(\text{Bu}^n\text{SnO} \cdot \text{O}_2\text{CR})_6$.^{1,2}

It is possible that the bulkiness of the substituents makes the diphenylacetate compound **11** readily isolable; in line with many reports on dialkyltin dicarboxylates,¹⁴ we assign a mono-



meric structure for **11**. This assignment is consistent with the ^{119}Sn NMR chemical shift value of $\delta -235.8$ which is close to that of $\text{SnBu}_2(\text{O}_2\text{CBu}^t)_2$ [$\delta(^{119}\text{Sn}) -232$].^{14c}

The initial product formed in the reaction of compound **1** with silver phosphinates or carboxylates is of the type $\text{Sn}(\text{CH}_2\text{Ph})_2(\text{O}-\text{O})_2$ **II** ($\text{O}-\text{O} = \text{O}_2\text{PR}_2$ or O_2CR); we isolated such a compound (**11**) in the case of diphenylacetate.† The second step presumably involves the attack of water on **II** leading to the dimer **III** after elimination of toluene, dimerization and ligand reorganization. Isolation of such a species was accomplished only for the phosphinates (e.g. **3**). It is likely that the dimer **III** is in equilibrium with the monomer **IV**. Conver-



sion of **III** to an O-capped cluster (e.g. **4**) has been discussed by Holmes *et al.*¹² For the formation of a 'ladder' (e.g. **5**), **III** can react with its monomer **IV** with the elimination of a molecule each of water and the acid, followed by ligand reorganization and dimerization. Formation of a 'prismane' (e.g. **6**) from the 'ladder' (e.g. **5**) has been studied before and hence is not elaborated here.¹³

Structural aspects

The molecular structures of compounds **2**, $4 \cdot \text{CHCl}_3$ and $5 \cdot 2\text{CH}_2\text{Cl}_2$ are depicted in Figs. 1–3 respectively. Selected bond distances and angles for **2** are provided in Table 1, important bond parameters for $4 \cdot \text{CHCl}_3$ and $5 \cdot 2\text{CH}_2\text{Cl}_2$ in Figs. 2 and 3.

Compound **2** crystallizes in the space group $P\bar{1}$ with half the molecule in the asymmetric unit. It is a mononuclear six-co-ordinated tin complex containing two Sn–C, two covalently linked Sn–O (phosphinate) and two co-ordinatively linked Sn–O (phosphinic acid) bonds; the two Sn–C bonds are *trans* to each other. The two types of Sn–O bonds are nearly of the same length [2.237(2), 2.203(2) Å] but are longer than those observed in the O-capped cluster **4** (mean: 2.097 Å). The angles at the oxygens connected to tin are also nearly the same [143.6(1) and 143.7(1)°]. Distinction between the phosphinate and the phosphinic acid is best made by comparing the P–O distances, which show that P(1) belongs to the phosphinic acid.

The most important structural feature in compound **2** is the presence of hydrogen bonding between O(2) and O(4) [relevant distances and angle: $\text{O}(2)-\text{H} \approx 1.06$, $\text{O}(4) \cdots \text{H} \approx 1.37$ Å, $\text{O}(2)-\text{H} \cdots \text{O}(4) \approx 175.9^\circ$] forming an eight-membered ring that includes the hydrogen. This probably has a stabilizing effect on the observed structure. Although reports on many other diorganotin phosphates/phosphinates are available,^{15,17} to our knowledge **2** is the only mononuclear diorganotin phosphinate to be structurally characterized. These tin phosphinates, in general, have a tendency to oligomerize.

† The compound $\text{SnMe}_2[\text{O}_2\text{P}(\text{OBu}^n)_2]_2$ has been reported; it is assigned a monomeric structure with chelating phosphates, but no crystal structure is available.¹⁵




Fig. 1 An ORTEP¹⁶ drawing of compound **2** with displacement ellipsoids at 30% probability level. Only the non-hydrogen atoms are shown






Fig. 2 An ORTEP drawing of compound $4 \cdot \text{CHCl}_3$ with displacement ellipsoids at 30% probability level. Only tin, oxygen, phosphorus and carbon atoms connected to tin or phosphorus are shown. Bond distances (Å) and angles (°): Sn–O 2.05(1)–2.16(1), Sn–C 2.15(2)–2.20(2); O–Sn–O(*cis*) 77.4(5)–92.4(5), O–Sn–O(*trans*) 161.4(5)–165.8(5), O–Sn–C(*cis*) 91.9(7)–105.0(7), O–Sn–C(*trans*) 171.6(7)–173.1(5), Sn–OH–Sn 98.2(6)–99.8(5), Sn(1)–O(10)–Sn(2) 104.2(6), Sn(1)–O(10)–Sn(3) 102.4(6), Sn(2)–O(10)–Sn(3) 103.4(6)

The cationic part in compound **4** consists of a trisnnoxane in a chair conformation with Sn(1) and O(11) on the opposite sides of the mean plane containing Sn(2), O(9), Sn(3) and O(12); the hydroxyl groups comprise the oxygen atoms of the ring. The tin atoms are capped on one side of the trisnnoxane skeleton by a three-co-ordinated oxygen atom [O(10)] at 0.87 Å above the Sn(1)–Sn(2)–Sn(3) plane. The mean Sn–O–Sn angle at the capping oxygen is $\approx 103.3^\circ$ which is close to that in $[\{\text{SnBu}^n(\text{OH})(\text{O}_2\text{PPh}_2)_3\text{O}\}^+[\text{O}_2\text{PPh}_2]^-]$ [mean $103.6(2)^\circ$]¹⁸ but smaller than that observed in analogous titanium and zirconium carboxylates¹⁹ (see **V** below). All the three tin atoms in **4** are equivalent and six-co-ordinated, each with two bridging phosphinates. As in $[\{\text{SnBu}^n(\text{OH})(\text{O}_2\text{PPh}_2)_3\text{O}\}^+[\text{O}_2\text{PPh}_2]^-]$,¹⁸ the oxygen of the free phosphinate in **4** is hydrogen bonded to the hydroxyls of the trisnnoxane skeleton in an essentially identical fashion; the relevant distances in **4** are $\text{O}(7) \cdots \text{O}(9)$ 2.71(2), $\text{O}(7) \cdots \text{O}(11)$ 2.69(2) and $\text{O}(8) \cdots \text{O}(12)$ 2.68(2) Å.

Table 1 Selected bond lengths (Å) and angles (°) for compound **2** with estimated standard deviations in parentheses

Sn–O(1)	2.237(2)	P(1)–C(7)	1.812(3)
Sn–O(3)	2.203(2)	P(2)–O(3)	1.518(2)
Sn–C(25)	2.139(3)	P(2)–O(4)	1.505(2)
P(1)–O(1)	1.509(2)	P(2)–C(13)	1.812(3)
P(1)–O(2)	1.538(2)	P(2)–C(19)	1.826(2)
P(1)–C(1)	1.809(3)		
O(1)–Sn–O(1a)	180.00	C(25)–Sn–C(25a)	180.00
O(1)–Sn–O(3)	87.40(7)	O(3)–Sn–C(25a)	89.3(1)
O(1)–Sn–O(3a)	92.60(7)	O(1)–P(1)–O(2)	114.2(1)
O(1)–Sn–C(25)	88.58(9)	O(3)–P(2)–O(4)	116.2(1)
O(1)–Sn–C(25a)	91.42(9)	Sn–O(1)–P(1)	143.6(1)
O(3)–Sn–O(3a)	180.00	Sn–O(3)–P(2)	143.7(1)
O(3)–Sn–C(25)	90.7(1)		

Hydrogen bonding appears to be a stabilizing factor for the O-capped cation. It is also found in [$\{\text{SnBu}^n(\text{OH})[\text{O}_2\text{P}(\text{C}_6\text{H}_{11})_2]\}_3\text{O}\}[\{\text{SnBu}^n\text{Cl}_2[\text{O}_2\text{P}(\text{C}_6\text{H}_{11})_2]\}_2(\text{OH})]^{20}$ as well as in similar titanium (**Va**) and zirconium (**Vb**) carboxylate clusters.¹⁹



The molecule of compound **5** (Fig. 3) is a dimer with the two halves related by a centre of symmetry; there are three crystallographically distinct tin atoms. The atoms Sn(1) and Sn(2) are six-co-ordinated, bonded to five oxygen atoms and one carbon; the geometry around tin is a distorted octahedron. The three bridging carboxylates on Sn(1) are meridional whereas the two bridging carboxylates on Sn(2) are *trans* to each other. The terminal Sn(3) is seven-co-ordinated and has a pentagonal bipyramidal geometry with C(25) and O(2) at the apical positions. The five oxygen atoms and Sn(3) which comprise the equatorial plane are coplanar to within ± 0.11 Å. The apical angle O(2)–Sn(3)–C(25) is $173.5(3)^\circ$.

Three types of oxygen atoms are present in the molecule: (a) three-co-ordinated framework, (b) two-co-ordinated bridging carboxylate and (c) two-co-ordinated chelating carboxylate. The framework Sn–O bond lengths range from 1.999(5) to 2.138(5) Å, the longest being Sn(2)–O(2). It is interesting that this has always been the longest framework Sn–O bond in all the known structures of this type.^{13,21} The shortest Sn–O bond is the one from the terminal tin to the μ_3 -oxygen [Sn(3)–O(2) in **5**]. As noted by others,^{13,21} the Sn–O bonds to the bridging carboxylates (mean 2.172 Å) are, in general, longer than the framework Sn–O bonds (mean 2.062 Å). The Sn(3)–O bonds to the chelating carboxylates are much longer (mean 2.258 Å) and their values are closer than those in the structures reported





Fig. 3 An ORTEP drawing of compound **5**·2CH₂Cl₂ with displacement ellipsoids at 30% probability level. All non-hydrogen atoms except those of the solvent and of the phenyl carbons of the SnCH₂Ph group are shown. Bond distances (Å) and angles (°): Sn–O (three-co-ordinated) 1.999(5)–2.138(5), Sn–O (carboxylate) 2.115(5)–2.262(6), Sn–C 2.136(9)–2.153(9); O(1)–Sn(1)–O(2) 77.5(2), O(1)–Sn(2)–O(1a) 76.1(2), O(1)–Sn(2)–O(2) 74.8(2), O(1a)–Sn(2)–O(2) 150.9(2), Sn(1)–O(1)–Sn(2a) 149.0(3), Sn(1)–O(1)–Sn(2) 103.2(2), Sn(2)–O(1)–Sn(2a) 103.9(2), Sn(1)–O(2)–Sn(2) 100.4(2), Sn(1)–O(2)–Sn(3) 123.7(3), Sn(2)–O(2)–Sn(3) 126.1(3)

before.^{13,21} The sum of the angles around the framework oxygen atoms is close to 360° [Σ O(1) = 356.1, Σ O(2) = 350.2]; however the geometry around O(1) is slightly non-planar and it is ca. 0.37 Å above the mean plane containing Sn(1), Sn(2) and Sn(3).

An interesting structural correlation can be made between compound **5** and the μ -S bridged compound **VI** reported by Schmid *et al.*²² Compound **VI** essentially represents one half of the molecule of **5** with O(1) replaced by a sulfur and the chelating carboxylate on Sn(3) (in **5**) replaced by a bridging carboxylate connected to another tin.



The major topological difference between the prismane (see structure **I**) and the ladder (*e.g.* **5**) is perhaps the orientation (folding) of the two peripheral Sn₂O₂ rings [Sn(1)–O(2)–Sn(2)–O(1); Sn(1a)–O(2a)–Sn(2a)–O(1a) in the ladder **5**] with respect to the central Sn₂O₂ ring [Sn(2)–O(1)–Sn(2a)–O(1a)]; whereas the folding in the ladder is *transoid*, in the hexagonal prismane it is *cisoid*. In the process of hydrolysis of the ladder, therefore, it is possible that the peripheral Sn₂O₂ rings reorient themselves.

To summarize, the present study demonstrates the utility of dibenzyltin dichloride **1** as a useful precursor to obtain monobenzyltin oxide carboxylate/phosphinate cages/clusters in addition to the novel hydrogen-bonded dibenzyltin compounds such as **2**. It remains to be seen whether a carboxylate 'cubane' (RSnO·O₂CR')₄ or an 'O-capped cluster' [$\{\text{SnR}(\text{OH})(\text{O}_2\text{CR}')\}_3\text{O}\}[\text{O}_2\text{CR}']$] can be isolated by a suitable choice of R and R'; the latter type of carboxylates have been isolated for titanium (**Va**) and zirconium (**Vb**). Since the O-capped cluster (*e.g.* **4**) can be viewed as a cubane with one tin corner missing it must be possible to prepare cubanes by further condensation reactions on the O-capped cluster.^{1,23}

Experimental

Chemicals and solvents were from Aldrich/Fluka or from local

manufacturers. Further purification was done according to standard procedures.²⁴ All operations, unless stated otherwise, were carried out under a dry nitrogen atmosphere using standard Schlenk techniques.²⁵ The ¹H, ¹³C, ³¹P (at 80.961 MHz) and ¹¹⁹Sn (at 74.544 MHz) NMR spectra were recorded on a Bruker 200 MHz spectrometer in CDCl₃ solutions with shifts referenced to SiMe₄ (¹H, ¹³C; δ 0), external 85% H₃PO₄ (δ 0) and SnMe₄ (δ 0) respectively. The IR spectra were recorded on either a Perkin-Elmer 1310 or a JASCO FT/IR 5300 spectrophotometer. Elemental analyses were carried out on a Perkin-Elmer 240C CHN analyzer. Melting points are uncorrected.

Dicyclohexylphosphinic acid²⁶ and dibenzyltin dichloride **1**¹¹ were prepared according to literature methods. Silver salts of carboxylic acids were prepared using stoichiometric amounts of the acid, sodium hydroxide and silver nitrate.

Syntheses

Sn(CH₂Ph)₂[O₂P(C₆H₁₁)₂]₂[HO₂P(C₆H₁₁)₂] **2**. A mixture of compound **1** (1.50 g, 4.04 mmol) and Ag₂O₂P(C₆H₁₁)₂ (2.72 g, 8.07 mmol) was heated under reflux in toluene (75 cm³) for 5 h and then dicyclohexylphosphinic acid (1.86 g, 8.07 mmol) added. The mixture was heated under reflux for 15 h and filtered. The solvent was removed and the residue washed with *n*-heptane to remove any unchanged acid. The residue was crystallized from dichloromethane–*n*-heptane (1:5) to afford **2**. Yield: 2.3 g, 55%; m.p. 180 °C (Found: C, 61.12; H, 8.62. Calc. for C₆₂H₁₀₄O₈P₄Sn: C, 61.03; H, 8.59%). ¹H NMR: δ 1.05–2.19 (m, 88 H, cyclohexyl), 2.70 [s, with tin satellites, ²J(Sn–H) = 180, 176, 4 Hz, CH₂Ph], 6.85–7.33 (m, 10 H, aryl H) and 7.95 [s, with tin satellites, ⁴J(Sn–H) = 25 Hz, 2 H, OH]. ³¹P NMR: δ 54.7 (br).

{Sn(CH₂Ph)(OH)[O₂P(C₆H₁₁)₂]}₂ **3** and **{[Sn(CH₂Ph)(OH)-O₂P(C₆H₁₁)₂]}₃O** **4**. A mixture of compound **1** (1.00 g, 2.69 mmol) and Ag₂O₂P(C₆H₁₁)₂ (2.27 g, 6.72 mmol) was heated in toluene (50 cm³) under reflux for 12 h and then filtered. The solvent was completely removed from the filtrate and the residue dissolved in dichloromethane (30 cm³). After filtering the insolubles, the volume of the solution was reduced to ca. 5 cm³ and *n*-heptane added to obtain **3** as a crystalline solid. Yield: 1.1 g, 60%; m.p. 198 °C (Found: C, 54.04; H, 7.25. Calc. for C₆₂H₁₀₄O₁₀P₄Sn₂: C, 54.30; H, 7.59%). ¹H NMR: δ 0.73–2.25 (m, 88 H, cyclohexyl), 2.60–2.85 (m, 4 H, CH₂Ph), 4.90 (br, 2 H, OH) and 6.90–8.00 (m, 10 H, aryl H). ³¹P NMR: δ 48.7 [s, with tin satellites, ²J(Sn–O–P) = 196.4, P(dangling)] and 57.2 [s, with tin satellites, ²J(Sn–O–P) = 134.4 Hz, P(bridging)]. ¹¹⁹Sn NMR: δ –586.0 (m).

Upon attempted crystallization of the product **3** (chloroform + heptane) in air the O-capped cluster **{[Sn(CH₂Ph)(OH)[O₂P(C₆H₁₁)₂]}₃O**⁺[O₂P(C₆H₁₁)₂][–] **4** as a solvate (CHCl₃) was obtained in >65% yield; m.p. 228–230 °C [Found: C, 51.22; H, 7.10. Calc. for C₆₉H₁₁₂O₁₂P₄Sn₃ (after removing the solvent of crystallization): C, 51.34; H, 6.99%]. ¹H NMR (after drying): δ 0.85–2.10 (m, 88 H, cyclohexyl), 2.83 [s, with tin satellites, 6 H, ²J(Sn–H) = 166, 170 Hz, CH₂Ph], 6.95–7.40 (m, 15 H, aryl H) and 8.90 (br s, 3 H, OH). ¹³C NMR: δ 25.8, 26.1, 26.6, 26.8 (cyclohexyl), 34.8 [³J(P–O–Sn–C) = 30, CH₂Ph], 37.6 [¹J(P–C) = 88, cyclohexyl], 37.8 [¹J(P–C) = 84 Hz, cyclohexyl], 124.2, 127.8, 129.5 and 139.0 (aryl). ³¹P NMR: δ 52.8 [br, 3 P, P(bridging)] and 41.5 [br, 1 P, P(free)]. ¹¹⁹Sn NMR: δ –535.7 [t, ²J(Sn–O–P) = 137 Hz].

{[(PhCH₂)SnO·O₂CMe]₂(PhCH₂)Sn(O₂CMe)₃}₂·2CH₂Cl₂ **5**·**2CH₂Cl₂** and **[(PhCH₂)SnO·O₂CMe]₆** **6**. A mixture of compound **1** (4.0 g, 10.7 mmol) and silver acetate (5.0 g, 29.95 mmol) was heated under reflux in toluene (80 cm³) for 20 h. Filtration followed by removal of solvent afforded a solid which was crystallized from dichloromethane (12 cm³) to give **5**. Yield: 0.92 g, 25%; m.p. 196–200 °C, lost crystallinity around 135 °C

(Found: C, 37.44; H, 3.49. Calc. for C₃₂H₃₈Cl₂O₁₂Sn₃: C, 36.89; H, 3.68%). IR (KBr): 3026, 1588, 1493, 1425, 1026, 762, 737, 696 and 665 cm^{–1}. ¹H NMR: δ 1.85–2.11 (m, 30 H, CH₃), 2.95 (br m, 12 H, CH₂), 5.30 (s, 2 H, CH₂Cl₂) and 7.05–7.18 (m, 30 H, aryl). ¹³C NMR: δ 23.8 (CH₃), 35.9 (CH₂), 54.0 (CH₂Cl₂), 124.5, 127.8, 129.2, 139.3 (aryl) and 179.4 (CO₂). ¹¹⁹Sn NMR: δ –523.0 (assigned to **6**), –530 to –580 and –630 to –640 (many signals).

Compound **5** (0.1 g, 0.096 mmol) was dissolved in CH₂Cl₂ (2 cm³) in air for 24 h and the solvent evaporated under reduced pressure (repeated five times) to yield **6**. Yield: 0.08 g, ≈98%; m.p. 252–256 °C (Found: C, 37.67; H, 3.63. Calc. for C₅₄H₆₀O₁₈Sn₆: C, 37.95; H, 3.54%). IR (KBr): 3025, 1589, 1542, 1493, 1447, 1209, 1028, 758, 696, 610, 546 and 482 cm^{–1}. ¹H NMR: δ 1.85 (s, 18 H, CH₃), 2.63 [s, with tin satellites, 12 H, ²J(Sn–H) = 160.0, 164.0 Hz, CH₂] and 7.08–7.13 (m, 30 H, aryl). ¹³C NMR: δ 23.7 (CH₃), 34.4 (CH₂), 124.5, 127.8, 128.2, 128.7 and 129.2 (aryl) and 179.6 (CO₂). ¹¹⁹Sn NMR: δ –523.1.

[(PhCH₂)SnO·O₂CR]₆ (**R** = Bu⁷, C₅H₉N⁸ or CH=CMe₂ **9**). These compounds were prepared in yields of 20–40% from the reaction of **1** (1–5 mmol) with 2 mol equivalents of appropriate silver carboxylates using the above method.

Compound **7**: M.p. >260 °C (Found: C, 44.59; H, 4.94. Calc. for C₇₂H₉₆O₁₈Sn₆: C, 44.12; H, 4.94%). IR (KBr) 2963, 1582, 1537, 1485, 1427, 1231, 901, 754, 693 and 640 cm^{–1}. ¹H NMR: δ 0.96 [s, with tin satellites, 54 H, C(CH₃)₃], 2.73 [s, with tin satellites, 12 H, ²J(Sn–H) = 158, 152 Hz, CH₂] and 6.96–7.32 (m, 30 H, aryl); ¹³C NMR: δ 27.1 [C(CH₃)₃], 34.4 (CH₂), 40.5 [C(CH₃)₃], 124.3, 127.9, 129.2, 139.2 (aryl) and 185.1 (CO₂); ¹¹⁹Sn NMR: δ –523.1.

Compound **8**: m.p. 198–202 °C (Found: C, 43.92; H, 3.33; N, 4.43. Calc. for C₇₈H₆₆N₆O₁₈Sn₆: C, 44.58; H, 3.17; N, 4.67%). IR (KBr) 3022, 1593, 1549, 1491, 1397, 1194, 1028, 872, 756 and 696 cm^{–1}. ¹H NMR: δ 3.13 [s, with tin satellites, 12 H, ²J(Sn–H) = 130.7, 129.8, CH₂], 6.80–7.10 (m, 30 H, aryl), 7.50 [t, 6 H, ³J(H–H) = 4, nicotinate H⁵], 8.45 [d, 6 H, ³J(H–H) = 4, nicotinate H⁶], 8.88 [d, 6 H, ³J(H–H) = 4 Hz, nicotinate H⁴] and 9.40 (s, 6 H, nicotinate H¹); ¹³C NMR: δ 35.4 (CH₂), 123.1–153.2 (11 lines, C₆H₅ + C₅H₄N) and 172.5 (CO₂); ¹¹⁹Sn NMR: δ –519.8.

Compound **9**: m.p. 264–266 °C (Found: C, 44.52; H, 4.56. Calc. for C₇₂H₈₄O₁₈Sn₆: C, 44.36; H, 4.34%). IR (KBr) 3024, 1642, 1510, 1414, 1316, 1182, 1055, 862, 760, 694, 621 and 451 cm^{–1}. ¹H NMR: δ 1.69 and 1.79 (s each, 36 H, CH₃), 2.69 [s, with tin satellites, 12 H, ²J(Sn–H) = 156.2, 149.2 Hz, CH₂], 5.36 (s, 6 H, Me₂C=CH) and 6.95–7.20 (m, 30 H, aryl); ¹³C NMR: 22.9 (CH₃), 34.9 (CH₂), 41.9 (CH₃), 124.8–130.9 (many lines, aryl), 132.7 (Me₂C=CH), 138.0 (Me₂C=CH) and 173.6 (CO₂); ¹¹⁹Sn NMR: δ –522.9 (s).

[(PhCH₂)SnO·O₂CCHPh]₆ **10** and **Sn(CH₂Ph)₂(O₂CCHPh)₂** **11**. The above procedure was again followed using compound **1** (0.58 g, 1.57 mmol) and silver diphenylacetate (1.0 g, 3.13 mmol) in toluene (50 cm³). Compound **11** was isolated from a concentrated toluene (5 cm³) solution. Yield: 0.41 g, 36%; m.p. 188–192 °C (Found: C, 70.53; H, 4.68. Calc. for C₄₂H₃₆O₄Sn: C, 70.70; H, 4.85%). IR (KBr): 3027, 1591, 1493, 1452, 1379, 1240, 1053, 1030, 720, 693, 652 and 453 cm^{–1}. ¹H NMR: δ 3.0 [s, with tin satellites, 4 H, ²J(Sn–H) ≈ 95.4 Hz, CH₂], 4.90 (s, 2H, CHPh₂) and 6.75–7.50 (m, 30 H, aryl). ¹³C NMR: δ 54.9 (CH₂), 60.1 (CHPh₂), 124.1–138.8 (six lines, aryl) and 182.1 (CO₂). ¹¹⁹Sn NMR: δ –235.8.

After isolation of compound **11** the mother-liquor was stripped of all the solvent and the residue crystallized from dichloromethane–diethyl ether–hexane (2:6:3) to give **10**. Yield: 0.21 g, 31%; m.p. 236–239 °C. IR (KBr): 3027, 1597, 1555, 1493 and 1412 cm^{–1}. ¹H NMR: δ 2.55 [s, 2 H, ²J(Sn–H) = 155, 159 Hz, CH₂], 4.55 (s, 1 H, CHPh₂) and 6.80–7.50 (m,

Table 2 Crystal data for compounds **2**, **4**·CHCl₃ and **5**·2CH₂Cl₂

	2	4 ·CHCl ₃	5 ·2CH ₂ Cl ₂
Empirical formula	C ₃₁ H ₅₂ O ₄ P ₂ Sn _{0.5}	C ₇₀ H ₁₁₃ Cl ₃ O ₁₂ P ₄ Sn ₃	C ₃₂ H ₃₈ Cl ₂ O ₁₂ Sn ₃
<i>M</i>	610.04	1732.98	1041.62
M.p. (°C), colour	180, colourless	228–230, colourless	252–256, colourless
Crystal system	Triclinic	Triclinic	Monoclinic
Space group	<i>P</i> $\bar{1}$	<i>P</i> $\bar{1}$	<i>P</i> 2 ₁ / <i>c</i>
<i>a</i> /Å	13.267(2)	16.018(3)	18.225(4)
<i>b</i> /Å	13.593(2)	19.600(4)	12.954(4)
<i>c</i> /Å	9.757(2)	15.492(2)	18.382(4)
<i>a</i> '°	96.90(2)	107.18(1)	
<i>β</i> '°	99.71(1)	113.80(1)	118.85(2)
<i>γ</i> '°	114.133(9)	80.34(2)	
<i>U</i> /Å ³	1547.3(5)	4244(1)	3801(1)
<i>Z</i>	2	2	4
<i>D</i> _c /g cm ⁻³	1.309	1.356	1.82
<i>λ</i> /Å	0.710 69	0.710 69	1.541 78
<i>T</i> /°C	23	23	-130
<i>μ</i> /cm ⁻¹	5.64	10.96	173.72
<i>F</i> (000)	650	1780	2040
Crystal size/mm	0.4 × 0.35 × 0.1	0.3 × 0.25 × 0.5	0.3 × 0.3 × 0.4
2 θ _{max} /°	55.1	45.1	120.2
Reflections collected	7459	11 633	6172
Independent reflections	7147	11 151	5963
<i>R</i> _{int}	0.032	0.026	0.135 ^a
Data [<i>I</i> > 3 σ (<i>I</i>)]	5883	6299	5665
Parameters	549	460	433
<i>R</i> ^b [<i>I</i> > 3 σ (<i>I</i>)]	0.033	0.091	0.069
<i>R</i> ' ^b	0.039	0.121	0.052
Maximum, minimum peaks in difference map/e Å ⁻³	0.38, -0.34	2.05, -1.65	1.16, -1.08

^a The higher *R*_{int} is basically due to the differences in the intensities in some very low-angle reflections. High absorption from the heavier scatterers for the Cu-K α radiation as well as the relatively significant crystal decay during the experiment probably have contributed to the *R*_{int} value. ^b $R = \sum ||F_o| - |F_c|| / \sum |F_o|$; $R' = [\sum w(|F_o| - |F_c|)^2 / \sum |F_o|^2]$.

15 H, aryl). ¹³C NMR: δ 34.4 (CH₂), 54.9 (CHPh₂), 124.2–139.3 (many lines, aryl) and 179.9 (CO₂). ¹¹⁹Sn NMR: δ -525.0.

X-Ray crystallography

A suitable crystal of compound **2** (crystallized from dichloromethane–heptane), **4** (crystallized from chloroform–heptane) or **5** (crystallized from dichloromethane–hexane) was mounted on a glass fibre. Data were collected on a Rigaku AFC5R diffractometer using graphite-monochromated Mo-K α (for **2** and **4**) or Cu-K α (for **5**) radiation. Three control reflections, collected every 150, showed no significant changes in intensities. The details pertaining to data collection and refinement are listed in Table 2. The structures were solved by conventional methods²⁷ and refined by full-matrix least squares.²⁸ An empirical absorption correction was applied and the data were corrected for Lorentz-polarization effects. Neutral atom scattering and anomalous-scattering correction were taken from ref. 29. The non-hydrogen atoms were refined anisotropically and the H atoms (for **2** and **5**) included at calculated positions. In the case of **4** some of the carbon atoms of the O₂P(C₆H₁₁)₂ groups are disordered. In particular, one C₆H₁₁ group of P(3) is highly disordered. On refinement its carbon atoms moved significantly, distorting the geometry. Therefore, these carbon atoms were fixed at positions found in Fourier-difference maps. Accordingly, no attempts were made to include H atoms for refinement. In addition, the final Fourier-difference map showed some minor residual densities (1–2 e Å⁻³), possibly corresponding to a partially occupied disordered solvent molecule. These peaks could not be modelled and therefore not included in the refinement.

CCDC reference number 186/908.

Acknowledgements

Financial support by Department of Science & Technology (New Delhi) and Council of Scientific & Industrial Research

(New Delhi) is gratefully acknowledged. Thanks are also due to University Grants Commission, India (COSIST and Special Assistance Programme) as well as the AvH Foundation (Germany) for instrumental facilities.

References

- 1 R. R. Holmes, *Acc. Chem. Res.*, 1989, **22**, 190 and refs. therein.
- 2 V. B. Mokal, V. K. Jain and E. R. T. Tiekink, *J. Organomet. Chem.*, 1991, **407**, 173.
- 3 K. C. Kumara Swamy, C. G. Schmid, R. O. Day and R. R. Holmes, *J. Am. Chem. Soc.*, 1990, **112**, 223.
- 4 J. Caruso, M. J. Hampden-Smith, A. L. Rheingold and G. Yap, *J. Chem. Soc., Chem. Commun.*, 1995, 157.
- 5 N. W. Alcock and S. M. Roe, *J. Chem. Soc., Dalton Trans.*, 1989, 1589.
- 6 V. Chandrasekhar, R. O. Day and R. R. Holmes, *Inorg. Chem.*, 1985, **24**, 1970.
- 7 A. Roy and A. K. Ghosh, *Inorg. Chim. Acta*, 1978, **29**, L275; 1977, **24**, L89.
- 8 S. Nagabrahmanandachari and K. C. Kumara Swamy, *Indian J. Chem., Sect. A*, 1995, **34**, 658.
- 9 Y. Yamamoto and N. Asao, *Chem. Rev.*, 1993, **93**, 2207; S. Kumaraswamy, S. Nagabrahmanandachari and K. C. Kumara Swamy, *Synth. Commun.*, 1996, **26**, 729.
- 10 J. March, *Advanced Organic Chemistry*, 3rd Edn, Wiley-Interscience, New York, 1986, pp. 47, 145, 153 and 164.
- 11 W. L. Jolly, *The Synthesis and Characterization of Inorganic Compounds*, Prentice-Hall, Engle-Wood Cliffs, NJ, 1970, p. 480.
- 12 R. R. Holmes, K. C. Kumara Swamy, C. G. Schmid and R. O. Day, *J. Am. Chem. Soc.*, 1988, **110**, 7060.
- 13 V. Chandrasekhar, C. G. Schmid, S. D. Burton, J. M. Holmes, R. O. Day and R. R. Holmes, *Inorg. Chem.*, 1987, **26**, 1050.
- 14 (a) T. P. Lockhart, J. C. Calabrese and F. Davidsen, *Organometallics*, 1987, **6**, 2479; (b) V. Chandrasekhar, R. O. Day, J. M. Holmes and R. R. Holmes, *Inorg. Chem.*, 1988, **27**, 958; (c) V. B. Mokal and V. K. Jain, *J. Organomet. Chem.*, 1992, **441**, 215; (d) E. R. T. Tiekink, *J. Organomet. Chem.*, 1991, **408**, 323; (e) S. W. Ng, V. G. Kumar Das, B. W. Skelton and A. H. White, *J. Organomet. Chem.*, 1989, **377**, 221.
- 15 V. K. Jain, T. Kesavadas and C. Vatsa, *Spectrochim. Acta*, 1992, **11/12**, 1583.

- 16 C. K. Johnson, ORTEP, Report ORNL-5138, Oak Ridge National Laboratory, Oak Ridge, TN, 1976.
- 17 V. K. Jain, *Coord. Chem. Rev.*, 1994, **135/136**, 809.
- 18 R. O. Day, J. M. Holmes, V. Chandrasekhar and R. R. Holmes, *J. Am. Chem. Soc.*, 1987, **109**, 940.
- 19 K. Döppert and U. Thewalt, *J. Organomet. Chem.*, 1986, **301**, 41; U. Thewalt, K. Döppert and W. Lasser, *J. Organomet. Chem.*, 1980, **308**, 303
- 20 K. C. Kumara Swamy, R. O. Day and R. R. Holmes, *Inorg. Chem.*, 1992, **31**, 4184.
- 21 R. R. Holmes, C. G. Schmid, V. Chandrasekhar, R. O. Day and J. M. Holmes, *J. Am. Chem. Soc.*, 1987, **109**, 1408.
- 22 C. G. Schmid, R. O. Day and R. R. Holmes, *Phosphorus Sulfur Silicon Relat. Elem.*, 1989, **41**, 69.
- 23 S. Nagabrahmanandachari, Ph.D. Thesis, University of Hyderabad, 1996.
- 24 D. D. Perrin, W. L. Armarego and D. R. Perrin, *Purification of Laboratory Chemicals*, Pergamon, Oxford, 1986.
- 25 D. F. Shriver and M. A. Drezdson, *The Manipulation of Air-sensitive Compounds*, Pergamon, Oxford, 1986.
- 26 D. F. Peppard, G. W. Mason and C. M. Andrejasich, *J. Inorg. Nucl. Chem.*, 1965, **27**, 697.
- 27 J. C. Calbrese, PHASE, Patterson Heavy Atom Solution Extractor, Ph.D. Thesis, University of Wisconsin, Madison, 1972; P. T. Beurskens, DIRDIF, Direct Methods for Difference Structures, An Automatic Procedure for Phase Extension and Refinement of Difference Structure Factors, Technical Report 1984/1, Crystallography Laboratory, Toernooiveld, Nijmegen, 1984.
- 28 TEXSAN, Texray Structure Analysis Package, Molecular Structure Corp, The Woodlands, TX, 1987 (revised).
- 29 *International Tables for X-Ray Crystallography*, Kynoch Press, Birmingham, 1974, vol. 4, Table 2.2A.

Received 24th November 1997; Paper 7/08460I