

Synthesis and characterisation of the first E-unidentate $[\text{Ph}_2\text{P}(\text{O})\text{NP}(\text{E})\text{Ph}_2]^-$ metal complexes $[\text{Au}\{\text{Ph}_2\text{P}(\text{O})\text{NP}(\text{E})\text{Ph}_2\}\{\text{Ph}_2\text{P}(\text{O})\text{NHPPPh}_2\}]$ and $[\text{Pd}\{\text{Ph}_2\text{P}(\text{O})\text{NP}(\text{E})\text{Ph}_2\}\{\text{H}_2\text{NCH}_2\text{CH}_2\text{NH}_2\}]$ (E = S or Se)


Alexandra M. Z. Slawin, Martin B. Smith and J. Derek Woollins*[†]

Department of Chemistry, Loughborough University, Loughborough, Leics., UK LE11 3TU


Chloride metathesis of $[\text{AuCl}(\text{HL})]$ [$\text{HL} = \text{Ph}_2\text{P}(\text{O})\text{NHPPPh}_2$] or $[\text{PdCl}_2(\text{en})]$ (en = ethane-1,2-diamine) in CH_3OH with either 1 (or 2) equivalents of $[\text{Ph}_2\text{P}(\text{O})\text{NP}(\text{E})\text{Ph}_2]^-$ (E = S or Se) gave the neutral compounds $[\text{Au}\{\text{Ph}_2\text{P}(\text{O})\text{NP}(\text{E})\text{Ph}_2\}\{\text{HL}\}]$ or $[\text{Pd}\{\text{Ph}_2\text{P}(\text{O})\text{NP}(\text{E})\text{Ph}_2\}\{\text{en}\}]$ respectively; the anionic ligands in these complexes adopt a new bonding mode and furthermore are involved in intramolecular N–H...OP hydrogen bonding with HL or en.

The co-ordination chemistry of $[\text{R}_2\text{P}(\text{E})\text{NP}(\text{E})\text{R}_2]^-$ (E = lone pair, O, S or Se) and related systems has received much interest of late.^{1,2} In particular the bis(chalcogenide) ligands $[\text{R}_2\text{P}(\text{E})\text{NP}(\text{E})\text{R}_2]^-$, closely associated with the ubiquitous acetylacetonate (acac) and its derivatives, have found relevance in diverse areas such as selective metal extractants, NMR shift reagents and more recently in catalysis.^{1,3} Asymmetric analogues of acac such as $[\text{R}_2\text{P}(\text{O})\text{NP}(\text{E})\text{R}_2]^-$ (E = S or Se) incorporating dissimilar Group 16 atoms have, until recently, been poorly studied.⁴ However the facile synthesis of $[\text{R}_2\text{P}(\text{O})\text{NP}(\text{E})\text{R}_2]^-$ now means that this type of chemistry is likely to prove much more fruitful than in the related mixed acac area.

Whereas $[\text{R}_2\text{P}(\text{E})\text{NP}(\text{E})\text{R}_2]^-$, $[\text{R}_2\text{P}(\text{O})\text{NP}(\text{E})\text{R}_2]^-$ and acac bind metal ions predominantly *via* chelation we reasoned that $[\text{R}_2\text{P}(\text{O})\text{NP}(\text{E})\text{R}_2]^-$ could also function as ambidentate ligands, ligating through either the O or E donor atom. Here we report the first examples of metal complexes with the unsymmetrical ligands $[\text{R}_2\text{P}(\text{O})\text{NP}(\text{E})\text{R}_2]^-$ (E = S **1**; E = Se **2**; R = Ph) in a novel co-ordination mode. X-Ray crystallography reveals that $[\text{Ph}_2\text{P}(\text{O})\text{NP}(\text{E})\text{Ph}_2]^-$ binds in a unidentate manner through exclusively the 'soft' sulfur (or selenium) donor centre. Moreover intramolecular PO...H–N hydrogen bonding between $[\text{Ph}_2\text{P}(\text{O})\text{NP}(\text{E})\text{Ph}_2]^-$ and the ancillary ligands results in the formation of eight- and nine-membered metallacycles.



Reaction of **1** (or **2**), as their potassium salts, with a suspension of $[\text{AuCl}(\text{HL})]$ [$\text{HL} = \text{Ph}_2\text{P}(\text{O})\text{NHPPPh}_2$]⁵ in CH_3OH gave the 1:1 complexes $[\text{Au}\{\text{Ph}_2\text{P}(\text{O})\text{NP}(\text{E})\text{Ph}_2\}\{\text{HL}\}]$ (E = S **3**; E = Se **4**) in high yields (typically >90%) [equation (1)] and



which display the expected spectroscopic properties.[‡] The structures of **3** and **4** were supported by IR [absence of $\nu(\text{AuCl})$ vibrations] and $^{31}\text{P}\{-^1\text{H}\}$ NMR spectroscopy. Notably $\delta(\text{P}_\text{O})$ for the co-ordinated ligand $[\text{Ph}_2\text{P}(\text{O})\text{NP}(\text{E})\text{Ph}_2]^-$ in **3** and **4** were similar to values previously reported for $\text{K}[\text{Ph}_2\text{P}(\text{O})\text{NP}(\text{E})\text{Ph}_2]$ [$\delta(\text{P}_\text{O})$ 13.9 **1**; 12.9 **2**] but appreciably shifted to lower frequency (typically *ca.* 15 ppm) with respect to values documented for complexes bearing these ligands in a O,E-chelating fashion.⁴ Hence the ^{31}P NMR resonances appear sensitive to the ligation mode adopted by the ligand and serve as a convenient probe. Further confirmation that the phosphoryl group of HL remains 'dangling' is reflected in the similarity of the $^{31}\text{P}\{-^1\text{H}\}$ data [$\delta(\text{P}_\text{O})$ 25.3 **3**; 25.4 **4**] with that of $[\text{AuCl}(\text{HL})]$ [$\delta(\text{P}_\text{O})$ 26.4].⁵ The molecular structure§ of **3**·0.5H₂O·0.5CH₃OH [Fig. 1(a)] shows the gold(I) centre to be co-ordinated by the P^{III} donor atom of HL and the S atom of $[\text{Ph}_2\text{P}(\text{O})\text{NP}(\text{S})\text{Ph}_2]^-$ in a slightly distorted linear geometry [$\text{P}(\text{3})-\text{Au}(\text{1})-\text{S}(\text{1})$ 174.42(12)°]. The two ligands are locked into a nine-membered AuN₂OP₃S aurocycle [Fig. 1(b)] by an intramolecular N–H...OP hydrogen bond [$\text{N}\cdots\text{O}$ 2.67, $\text{H}\cdots\text{O}$ 1.91 Å, $\text{N}-\text{H}\cdots\text{O}$ 147°] and which

‡ Compounds **1** (and **2**) were prepared as described in ref. 4. A typical synthesis is illustrated here for compound **3**. To a CH_3OH suspension (1 cm³) of $[\text{AuCl}(\text{HL})]$ (0.053 g, 0.0836 mmol) was added solid **1** (0.043 g, 0.0912 mmol) in one portion. The suspension dissolved and the product **3** precipitated almost immediately. After stirring the mixture for *ca.* 15 min the solid was collected by suction filtration and dried *in vacuo*. Yield 0.078 g, 90%. Selected spectroscopic data for complexes **3**–**6**. For **3**: NMR (CDCl_3) $^{31}\text{P}\{-^1\text{H}\}$ (referenced to 85% H₃PO₄): δ 61.9 (P) and 25.3 (P_O) [HL ligand, $J(\text{PP})$ 28.6 Hz]; 28.5 (P_S) and 14.0 (P_O) [$J(\text{PP})$ not resolved]. ^1H : δ 10.75 (NH), 8.25–7.18 (aromatic H). IR (KBr) $\nu(\text{PO})$ 1240, 1219, 1150 and $\nu(\text{PS})$ 561 cm⁻¹ (Found: C, 55.00; H, 3.75; N, 2.35. C₄₈H₄₁AuN₂O₂P₄S·0.5H₂O·0.5CH₃OH requires C, 55.15; H, 4.20; N, 2.65%). For **4**: NMR (CDCl_3) $^{31}\text{P}\{-^1\text{H}\}$: δ 63.2 (P) and 25.4 (P_O) [HL ligand, $J(\text{PP})$ 33.0 and 8.8]; 15.3 (P_{Se}) [$J(\text{PSe})$ 470 Hz] and 14.5 (P_O) [$J(\text{PP})$ not resolved]. ^1H : δ 10.63 (NH), 8.13–7.10 (aromatic H). IR (KBr) $\nu(\text{PO})$ 1235, 1217, 1151 and $\nu(\text{PSe})$ 553 cm⁻¹ (Found: C, 52.40; H, 4.00; N, 2.40. C₄₈H₄₁AuN₂O₂P₄Se·0.5H₂O·0.5CH₃OH requires C, 52.80; H, 4.05; N, 2.55%). For **5**: NMR (CDCl_3) $^{31}\text{P}\{-^1\text{H}\}$: δ 28.8 (P_S) and 14.9 (P_O) [$J(\text{PP})$ 2.2 Hz]. ^1H : δ 7.91–7.81, 7.38–7.28 (aromatic H), 4.56 (NH) and 2.25 (CH₂). IR (KBr) $\nu(\text{PS})$ 567 cm⁻¹. FAB MS: m/z 970 ($M - \text{en}$)⁺ (Found: C, 58.10; H, 4.60; N, 5.40. C₅₀H₄₈N₄O₂P₄PdS₂ requires C, 58.20; H, 4.70; N, 5.45%). For **6**: NMR (CDCl_3) $^{31}\text{P}\{-^1\text{H}\}$: δ 15.6 (P_O) and 14.2 (P_{Se}) [$J(\text{PSe})$ 515 Hz] [$J(\text{PP})$ not resolved]. ^1H : δ 7.94–7.84, 7.38–7.30 (aromatic H), 4.38 (NH) and 2.27 (CH₂). IR (KBr) $\nu(\text{PSe})$ 553 and 544 cm⁻¹. FAB MS: m/z 1065 ($M - \text{en}$)⁺ (Found: C, 52.80; H, 4.10; N, 5.10. C₅₀H₄₈N₄O₂P₄PdSe₂ requires C, 53.35; H, 4.30; N, 5.00%).

§ Crystal data for complex **3**: C₄₈H₄₁AuN₂O₂P₄S·0.5H₂O·0.5CH₃OH, $M = 1055.76$, monoclinic, space group $C2/c$, $a = 39.344(1)$, $b = 13.227(1)$, $c = 18.491(1)$ Å, $\beta = 91.47(1)^\circ$, $U = 9619.5(4)$ Å³, $Z = 8$, $\mu(\text{Mo}-\text{K}\alpha) = 3.28$ mm⁻¹, $T = 293$ K, $R1 = 0.0496$ for 6926 unique reflections.

Crystal data for complex **6**: C₅₀H₄₈N₄O₂P₄PdSe₂·0.50Et₂, $M = 1162.23$, triclinic, space group $P\bar{1}$, $a = 15.403(8)$, $b = 19.198(8)$, $c = 10.318(4)$ Å, $\alpha = 97.07(3)$, $\beta = 94.37(4)$, $\gamma = 67.02(4)^\circ$, $U = 2786(2)$ Å³, $Z = 2$, $\mu(\text{Cu}-\text{K}\alpha) = 5.59$ mm⁻¹, $T = 293$ K, $R1 = 0.044$ for 8300 unique reflections. CCDC reference number 186/970. See <http://www.rsc.org/suppdata/dt/1998/1537/> for crystallographic files in .cif format.

[†] E-Mail: j.d.woollins@lboro.ac.uk





Fig. 1 (a) The crystal structure of **3** (solvent molecules omitted for clarity). Selected bond lengths (Å) and angles (°): Au(1)–P(3) 2.262(4), Au(1)–S(1) 2.335(3), S(1)–P(1) 2.042(4), P(1)–N(1) 1.569(10), N(1)–P(2) 1.601(10), P(2)–O(2) 1.510(8), P(3)–N(3) 1.657(9), N(3)–P(4) 1.645(9), P(4)–O(4) 1.460(8); S(1)–Au(1)–P(3) 174.42(12), Au(1)–S(1)–P(1) 96.1(2), S(1)–P(1)–N(1) 120.7(4), P(1)–N(1)–P(2) 134.8(6), N(1)–P(2)–O(2) 119.2(5), Au(1)–P(3)–N(3) 106.0(4), P(3)–N(3)–P(4) 132.4(6), N(3)–P(4)–O(4) 111.8(5). (b) Core geometry in **3** (all C atoms removed for clarity)

accounts for the *anti* conformation of the NH and P=O oxygen. In addition the S(1)–Au(1)–P(3)–N(3)–O(2)–P(2) portion of the ring is planar with a maximum deviation of 0.05 Å [for O(2)]. Both P(1) and N(1) lie above [1.36 and 1.00 Å respectively] and H(3b) below [–0.34 Å] the mean plane. The Au(1)–P(3) and Au(1)–S(1) distances [2.262(4) and 2.335(3) Å respectively] are normal whilst the bond lengths and angles within the S–P–N–P–O backbone are consistent with some degree of delocalisation. The P(1)–N(1)–P(2) angle in **3** is somewhat enlarged [134.8(6)°] as would be anticipated for a terminal bound ligand. In the crystal structure of **3**·0.5H₂O·0.5CH₃OH the half weight H₂O and CH₃OH solvate molecules are H-bonded together.

Transmetalation of **1** (or **2**) with [PdCl₂(en)] (en = ethane-1,2-diamine) in a 2:1 molar ratio in CH₃OH gave [Pd{Ph₂P(O)NP(E)Ph₂-E}₂(en)] (E = S **5**; E = Se **6**) and which could be recrystallised from CH₂Cl₂–OEt₂ [or light petroleum (b.p. 60–80 °C)] [equation (2)]. Compounds **5** and **6** were char-



acterised spectroscopically[‡] and $\delta(\text{P}_\text{o})$ for the co-ordinated [Ph₂P(O)NP(E)Ph₂][–] ligand observed at δ 14.9 (for **5**) and 15.6




Fig. 2 (a) The crystal structure of **6** (solvent molecules omitted for clarity). Selected bond lengths (Å) and angles (°): Pd(1)–Se(1) 2.413(1), Pd(1)–Se(2) 2.410(2), Pd(1)–N(2) 2.043(6), Pd(1)–N(3) 2.081(6), Se(1)–P(1) 2.199(2), P(1)–N(1) 1.580(6), N(1)–P(2) 1.605(6), P(2)–O(1) 1.493(6), Se(2)–P(3) 2.198(2), P(3)–N(4) 1.560(6), N(4)–P(4) 1.603(6), P(4)–O(2) 1.507(5); Se(1)–Pd(1)–Se(2) 85.16(4), Se(1)–Pd(1)–N(2) 91.2(2), Se(1)–Pd(1)–N(3) 174.7(2), Se(2)–Pd(1)–N(2) 175.9(2), Se(2)–Pd(1)–N(3) 99.9(2), N(2)–Pd(1)–N(3) 83.7(3), Pd(1)–Se(1)–P(1) 99.64(6), Se(1)–P(1)–N(1) 117.7(3), P(1)–N(1)–P(2) 131.2(4), N(1)–P(2)–O(1) 118.8(3), Pd(1)–Se(2)–P(3) 107.06(7), Se(2)–P(3)–N(4) 120.8(3), P(3)–N(4)–P(4) 133.3(4), N(4)–P(4)–O(2) 117.7(3). (b) Core geometry in **6** (all C atoms removed for clarity)

(for **6**) are similar to those values discussed for **3** and **4**. Furthermore the absence of $\nu(\text{PdCl})$ vibrations is also consistent with the formation of **5** and **6**. A crystal structure[§] [Fig. 2(a)] of **6**·0.5OEt₂ shows the palladium centre to be bound by two terminal [Ph₂P(O)NP(Se)Ph₂-Se][–] ligands and a bidentate H₂NCH₂CH₂NH₂ ligand in a slightly distorted square planar geometry with co-ordination angles in the range 83.7(3)–99.9(2) and 174.7(2)–175.9(2)°. The Pd–Se distances in **6** [2.413(1) and 2.410(2) Å] are slightly longer than those seen in *cis*-[Pd{Ph₂P(O)NP(Se)Ph₂-O,Se₂}] [2.381(7) and 2.383(8) Å].⁴ The P–N bond lengths and angles in **6** are similar to those in **3**. Both [Ph₂P(O)NP(Se)Ph₂][–] ligands are involved in intra-

molecular N–H⋯OP hydrogen bonding with H₂NCH₂–CH₂NH₂ [N(2)⋯O(1) 2.79, H(2a)⋯O(1) 2.06 Å, N(2)–H(2a)⋯O(1) 168°; N(3)⋯O(2) 2.78, H(3b)⋯O(2) 1.85 Å, N(3)–H(3b)⋯O(2) 162°]. This leads to a spirocyclic arrangement composed of two eight-membered rings [Fig. 2(b)] about a central palladium(II) metal centre. The two PdN₂OP₂Se puckered rings are not identical [transannular separations Pd(1)⋯O(1) 4.13, Pd(1)⋯N(1) 3.71 Å; Pd(1)⋯O(2) 3.90, Pd(1)⋯N(4) 4.48 Å]. An intermolecular N–H⋯OP hydrogen bond is observed between H(2b) and O(2*) of an adjacent molecule [N(2)⋯O(2*) 2.87, H(2b)⋯O(2*) 2.05 Å, N(2)–H(2b)⋯O(2*) 128°].

Interestingly under similar conditions, reaction of the symmetrical ligands [Ph₂P(E)NP(E)Ph₂][–] (E = S or Se) gave either the three-co-ordinate complexes [Au{Ph₂P(E)NP(E)Ph₂-E,E'}-(HL)] or the known⁶ homoleptic complexes [Pd{Ph₂P(E)NP(E)Ph₂-E,E'}₂] in which substitution of all ligands has resulted. Surprisingly there are very few examples of related metal-β-diketonates (including monothio derivatives) that exhibit a unidentate mode of co-ordination akin to that observed here.^{7,8} The ambidentate nature of **1** and **2** may lead to new complexes that could be of importance in catalytic systems. Further studies are currently in progress.

Acknowledgements

We are grateful to the EPSRC for funding and to Johnson Matthey for loans of precious metals. Fast atom bombardment (FAB) mass spectra were run by the EPSRC mass spectrometry service at Swansea.

References

- 1 For a recent example, see, H. Rudler, B. Denise, J. Ribeiro Gregorio and J. Vaissermann, *Chem. Commun.*, 1997, 2299.
- 2 A. M. Z. Slawin, M. B. Smith and J. D. Woollins, *Chem. Commun.*, 1996, 2095.
- 3 O. Navrátil, E. Herrmann, G. Grossmann and J. Teplý, *Collect. Czech. Chem. Commun.*, 1990, **55**, 364.
- 4 A. M. Z. Slawin, M. B. Smith and J. D. Woollins, *J. Chem. Soc., Dalton Trans.*, 1996, 3659.
- 5 A. M. Z. Slawin, M. B. Smith and J. D. Woollins, *J. Chem. Soc., Dalton Trans.*, 1996, 1283.
- 6 P. Bhattacharyya, J. Novosad, J. Phillips, A. M. Z. Slawin, D. J. Williams and J. D. Woollins, *J. Chem. Soc., Dalton Trans.*, 1995, 1607.
- 7 S. Kawaguchi, *Coord. Chem. Rev.*, 1986, **70**, 51.
- 8 G. H. Barnett, M. K. Cooper, M. McPartlin and G. B. Robertson, *J. Chem. Soc., Dalton Trans.*, 1978, 587.

Received 6th March 1998; Communication 8/01856A