

Kinetics and mechanism of the reduction of pentacyanonitro-ferrate(III) by L-ascorbic acid in acidic aqueous solution

Alicja Wanat,^a Rudi van Eldik^b and Grażyna Stochel^{*,a,b}

^a Department of Inorganic Chemistry, Jagiellonian University, Ingardena 3, 30-060 Kraków, Poland

^b Institute for Inorganic Chemistry, University of Erlangen-Nürnberg, Egerlandstrasse 1, 91058 Erlangen, Germany

The electron transfer reaction between $[\text{Fe}(\text{CN})_5(\text{NO}_2)]^{3-}$ and ascorbic acid was subjected to a detailed kinetic and thermodynamic study as a function of pH, ascorbic acid concentration, temperature and pressure. The pH profile indicates a pH independent region in the pH 5.0–5.5 range, which is ascribed to the oxidation of the ascorbate anion HA^- present in solution under such conditions. The experimental rate and activation parameters suggest that this redox process follows an outer-sphere electron transfer mechanism. The Marcus, Fuoss and Stranks–Marcus–Hush relationships were applied to estimate the self-exchange rate constant for the $[\text{Fe}(\text{CN})_5(\text{NO}_2)]^{3-/4-}$ couple, the ion-pair formation constant, the rate constant for electron transfer and the value of $\Delta V_{\text{et}}^\ddagger$. The results are discussed in reference to related systems reported in the literature.

An increasing interest exists in the study of electron transfer processes that are important in biological systems. The oxidation of L-ascorbic acid (vitamin C) by transition metal complexes in aqueous solution is one of these. It is known that ascorbate exists in cellular systems at relatively high concentration¹ and is an important cellular antioxidant.² Reactions of ascorbic acid with various inorganic oxidants have recently been investigated in much detail.^{3–18} We have extended our studies to include the reaction between ascorbic acid and pentacyanonitroferrate(III). Our interest in the reduction of $[\text{Fe}(\text{CN})_5(\text{NO}_2)]^{3-}$ by ascorbic acid results from specific properties of the nitro complex and its potential application as a hypotensive agent. Pentacyanonitroferrate(III) together with sodium nitroprusside, molsidomine and nitroglycerin belongs to the group of drugs referred to as nitrovasodilators.¹⁸ The relaxation of smooth muscle caused by the action of these compounds is connected with their ability to produce nitric oxide, NO. From the biochemical point of view the interaction of nitrovasodilators with cellular antioxidants is particularly important as it can be one of the decisive steps in their metabolism.

It was previously found^{3,4} that pseudo-first-order rate constants for the oxidation reactions of L-ascorbic acid by metal complexes exhibit a characteristic pH dependence due to both the acid dissociations of ascorbic acid (H_2A) and the hydrolysis equilibria of the oxidant when it involves an aquated metal ion. Many of these reactions proceed according to an outer-sphere electron transfer process since the oxidant is an inert metal complex and does not possess vacant co-ordination sites, *viz.* $\text{Fe}(\text{CN})_6^{3-}$, $\text{Fe}(\text{phen})_3^{3+}$, $\text{Co}(\text{bipy})_3^{3+}$, $\text{Co}(\text{phen})_3^{3+}$, $\text{Co}(\text{C}_2\text{O}_4)_3^{3-}$, *etc.*^{3,13–16} However, in other systems where the metal complex has labile co-ordination sites, for example in aquated Mn^{III} , Co^{III} and Fe^{III} , kinetic evidence for the operation of an inner-sphere electron transfer process was presented.^{11,17}

In recent years high-pressure techniques have been applied to the study of bioinorganic reaction systems. Their application in the study of inter- and intra-molecular electron transfer reactions of co-ordination complexes in solution has added a new dimension to improve the understanding of the intimate reaction mechanism.^{19–21} Here we report our investigations on the outer-sphere electron transfer reaction between a low and high

charged anionic species, *viz.* the ascorbate anion HA^- and the $[\text{Fe}(\text{CN})_5(\text{NO}_2)]^{3-}$ complex ion, respectively. For this system a pH, temperature and pressure dependence study was performed and the activation parameters (ΔH^\ddagger , ΔS^\ddagger , ΔV^\ddagger) are reported. Theoretical calculations based on the Marcus, Fuoss and Stranks–Marcus–Hush relationships were employed to estimate the self-exchange rate constant for the $[\text{Fe}(\text{CN})_5(\text{NO}_2)]^{3-/4-}$ couple, the ion-pair formation constant, the rate constant for electron transfer, and to account for the experimentally observed pressure effects.

Experimental

Materials

Potassium pentacyanonitroferrate(III), $\text{K}_3[\text{Fe}(\text{CN})_5(\text{NO}_2)]$, was prepared as described before²² and its purity checked by elemental analyses, IR and UV/VIS spectroscopy. All other chemicals were of analytical reagent grade and used without further purification. Solutions were prepared with deionized (Millipore) water and purged with N_2 for *ca.* 15 min prior to use. An acetate buffer solution and NaOH were used to control the pH in the range 4.0–5.6, whereas a phosphate buffer and NaOH were employed for $\text{pH} > 5.6$. The ionic strength of the test solutions (0.3 M) was adjusted by the addition of NaCl.

Measurements

pH Measurements were performed on a Metrohm 623 pH meter equipped with a Sigma glass electrode. The UV/VIS spectra were recorded on a Shimadzu UV-2100 spectrophotometer equipped with a thermostatted cell compartment CPS-260. Kinetic measurements were performed on a thermostatted ($\pm 0.1^\circ\text{C}$) stopped-flow SX-17MV spectrophotometer from Applied Photophysics and on a home-made high-pressure stopped-flow unit^{23,24} at pressures up to 120 MPa. The kinetic traces were recorded on an IBM compatible computer and analysed with the OLIS KINFIT (Bogart, GA, 1989) set of programs. The redox process was followed at 384 nm, the absorbance maximum for $[\text{Fe}(\text{CN})_5(\text{NO}_2)]^{3-}$. All kinetic experiments were performed under pseudo-first-order conditions, *i.e.* at least a ten-fold excess of ascorbic acid. The studied reactions exhibit excellent pseudo-first-order behaviour for at least three half-lives. The reported rate constants are

* E-Mail: stochel@trurl.ch.uj.edu.pl




Fig. 3 Concentration dependence of k_{obs} for the reduction of $[\text{Fe}(\text{CN})_5(\text{NO}_2)]^{3-}$ by ascorbate anion. Experimental conditions: $[\text{Fe}^{\text{III}}] = 5 \times 10^{-4} \text{ M}$, 25°C , $I = 0.3 \text{ M}$, $\text{pH } 5.3$




Fig. 4 Plot of $\ln k$ versus pressure for the reduction of $[\text{Fe}(\text{CN})_5(\text{NO}_2)]^{3-}$ by ascorbate anion. Experimental conditions: $[\text{Fe}^{\text{III}}] = 5 \times 10^{-4} \text{ M}$, $[\text{H}_2\text{A}]_t = 0.015 \text{ M}$, 15°C , $I = 0.3 \text{ M}$, $\text{pH } 5.3$

$$-\frac{d[\text{Fe}^{\text{III}}]}{dt} = \frac{2kK_1}{[\text{H}^+] + K_1} [\text{H}_2\text{A}]_t [\text{Fe}^{\text{III}}] \quad (4)$$

$[\text{H}_2\text{A}]_t$ represents the total ascorbate concentration. Under these conditions, *i.e.* $[\text{H}_2\text{A}]_t \gg [\text{Fe}^{\text{III}}]$, the expression for k_{obs} is as in equation (5). On the basis of this equation, k_{obs} should

$$k_{\text{obs}} = \frac{2kK_1}{[\text{H}^+] + K_1} [\text{H}_2\text{A}]_t \quad (5)$$

increase linearly with increasing total ascorbate concentration. As shown in Table 1 and Fig. 3 the dependence of k_{obs} on $[\text{H}_2\text{A}]_t$ is linear, from which it follows that $k = 645 \pm 8 \text{ M}^{-1} \text{ s}^{-1}$ (at 25°C and $I = 0.3 \text{ M}$; $\text{p}K_1 = 3.6$).

The presence of a small intercept in the plot in Fig. 3 can be assigned to experimental error and indicates that there is no significant contribution from reverse or other parallel reaction paths. The large value of the rate constant k is characterised by a relatively low activation enthalpy, $\Delta H^\ddagger = 21 \pm 2 \text{ kJ mol}^{-1}$ (see Table 2). The activation entropy found for this process is very negative, $\Delta S^\ddagger = -119 \pm 6 \text{ J K}^{-1} \text{ mol}^{-1}$, and could result from the formation of a highly ordered transition state.

As can be seen from the results reported in Table 1, the observed rate constant increases with increasing pressure. The pressure dependence of the second-order rate constant (Fig. 4)

Table 1 Rate constants and activation parameters for the reduction of $[\text{Fe}(\text{CN})_5(\text{NO}_2)]^{3-}$ by ascorbate anion

pH	$[\text{H}_2\text{A}]/\text{M}$	$T/^\circ\text{C}$	P/MPa	$k_{\text{obs}}/\text{s}^{-1}$	$k/\text{M}^{-1} \text{ s}^{-1}$
4.00	0.015	25	0.1	11.3 ± 0.6	
4.20				11.1 ± 0.3	
4.60				19.0 ± 0.6	
4.80				19.4 ± 0.1	
5.05				20.4 ± 0.2	697 ± 7
5.15				21.0 ± 0.1	716 ± 2
5.25				21.0 ± 0.1	716 ± 4
5.30				20.6 ± 0.1	704 ± 4
5.40				21.4 ± 0.1	729 ± 4
5.50				21.3 ± 0.1	728 ± 5
5.60				22.5 ± 0.5	
5.90				23.9 ± 0.3	
6.10				25.1 ± 0.4	
6.40				26.5 ± 0.5	
6.60				29.3 ± 0.2	
6.90				37.2 ± 0.4	
7.00				48.2 ± 0.7	
5.30	0.005	25	0.1	7.0 ± 0.3	645 ± 8
	0.010			12.7 ± 0.1	
	0.015			19.0 ± 0.1	
	0.020			25.3 ± 0.4	
	0.025			32.0 ± 0.5	
5.30	0.015	20	0.1	16.4 ± 0.1	561 ± 3
		25		19.0 ± 0.1	645 ± 8
		30		23.2 ± 0.4	792 ± 12
		35		25.0 ± 0.1	858 ± 2
		40		31.2 ± 0.5	1066 ± 17
5.30	0.005	15	10	7.0 ± 0.4	755 ± 45
			30	7.4 ± 1.1	802 ± 122
			60	8.7 ± 0.4	942 ± 38
			90	9.7 ± 0.3	1050 ± 29
			120	11.0 ± 2.0	1188 ± 211

Table 2 Comparison of the activation parameters for the reduction of cyanoferrate(III) complexes by ascorbate anion at 25°C


	$[\text{Fe}(\text{CN})_6]^{3-/4- a}$	$[\text{Fe}(\text{CN})_5(\text{NO}_2)]^{3-/4- b}$
$k/\text{M}^{-1} \text{ s}^{-1}$	842.0 ± 23	645.0 ± 8.0
$\Delta G^\ddagger/\text{kJ mol}^{-1}$	56.3	56.5 ± 3.0
$\Delta H^\ddagger/\text{kJ mol}^{-1}$	20.8 ± 0.8	21.0 ± 2.0
$\Delta S^\ddagger/\text{J K}^{-1} \text{ mol}^{-1}$	-119.0 ± 3.0	-119.0 ± 6.0
$\Delta V^\ddagger/\text{cm}^3 \text{ mol}^{-1}$	-16.3 ± 4.0	-10.0 ± 0.5

^a Ref. 3. ^b This work.

resulted in a negative activation volume, $\Delta V^\ddagger = -10.0 \pm 0.5 \text{ cm}^3 \text{ mol}^{-1}$. A meaningful decrease in partial molar volume during the redox process can be ascribed to the increase in electrostriction due to significant charge concentration on going from 3- to 4-charged complex species during the electron transfer process.

It is informative to compare the results of the present system with those reported earlier for the related system $[\text{Fe}(\text{CN})_6]^{3-}$ –ascorbate anion.³ The values of the second-order rate constants and activation parameters summarized in Table 2 indicate that k , ΔG^\ddagger , ΔH^\ddagger and ΔS^\ddagger for $[\text{Fe}(\text{CN})_5(\text{NO}_2)]^{3-/4-}$ agree very well with those found for $[\text{Fe}(\text{CN})_6]^{3-/4-}$. The value of ΔV^\ddagger for the studied system is less negative.

The experimental rate constants and activation parameters clearly suggest that oxidation of ascorbate anion by $[\text{Fe}(\text{CN})_5(\text{NO}_2)]^{3-}$ follows an outer-sphere electron transfer mechanism. The electron transfer process according to the standard Marcus–Hush model consist of three steps²⁸ of which the first and the third step (precursor formation and successor dissociation to the reaction products, respectively) are diffusion controlled and the second step involving irreversible electron transfer is rate determining (see Scheme 3). Under conditions where



Scheme 3

k_{obs} depends linearly on $[\text{H}_2\text{A}]_t$ as found in this study, $k = k_{\text{et}}K_{\text{os}}$ since K_{os} is expected to be small for precursor formation between species with charges of the same sign; K_{os} and k_{et} have to be separated theoretically. The ion-pair formation constant K_{os} can be predicted by using the extended Fuoss equation (6)²⁹

$$K_{\text{os}} = \frac{4}{3}\pi N_A a^3 \exp(-w_{ij}/RT) \quad (6)$$

where a is the contact distance of the ions ($a = r_i + r_j$) and w_{ij} represents the electric work term required to bring the reactants i and j to the contact distance in the precursor complex. The latter term arises from the Debye–Hückel interionic potential to allow for ionic strength effects²⁹ and is expressed by equation (7) where z_1, z_2 are the charges on the ions, ϵ_0 the permittivity of

$$w_{12} = z_1 z_2 e_0^2 N_A / 4\pi\epsilon_0 \epsilon a (1 + \kappa a) \quad (7)$$

a vacuum, ϵ the bulk relative permittivity and $\kappa = (2e_0 N_A I / \epsilon_0 \epsilon k_B T)^{1/2}$ with ionic strength I in M. At 25 °C, $I = 0.3$ M and with ionic radii of 0.34 nm for HA^- and 0.44 nm for the $[\text{Fe}(\text{CN})_5(\text{NO}_2)]^{3-}$ complex [approximated by that of $[\text{Fe}(\text{CN})_6]^{3-}$], it follows that $w_{12} = 2.78$ kJ mol⁻¹ and $K_{\text{os}} = 0.40$ M⁻¹. The value of K_{os} is very small as expected for reactants that are both anions. The calculated value of the electron transfer rate constant $k_{\text{et}} = 1.6 \times 10^3$ s⁻¹ from $k = k_{\text{et}}K_{\text{os}}$ is in good agreement with that reported³ for the related system $[\text{Fe}(\text{CN})_6]^{3-} - \text{HA}^-$, *viz.* 1.45×10^3 s⁻¹.

The pressure dependence of the rate constant k in accordance with $k = k_{\text{et}}K_{\text{os}}$ can be expressed as in equation (8) where

$$\Delta V_{12}^\ddagger = \Delta V_{\text{et}}^\ddagger + \Delta \bar{V}_{\text{os}} \quad (8)$$

$\Delta V_{\text{et}}^\ddagger$ represents the mechanistically important term. The value of $\Delta \bar{V}_{\text{os}}$ can be evaluated by using the modified version of the Hemmes equation²⁹ (9) to allow for ionic strength effects. With

$$\Delta \bar{V}_{\text{os}} = -RT \left\{ \frac{Z_1 Z_2 e_0^2 [\delta + 0.5a\kappa(\delta + \beta)]}{4\pi\epsilon_0 \epsilon k_B T a (1 + \kappa a)^2} + \beta \right\} \quad (9)$$

$\delta = (\partial \ln \epsilon / \partial P)_T = 4.64 \times 10^{-4}$ MPa⁻¹ (ref. 30) and $\beta = (\partial \ln \rho / \partial P)_T = 4.67 \times 10^{-4}$ MPa⁻¹ (ref. 31), $\Delta \bar{V}_{\text{os}} = -3.16$ cm³ mol⁻¹.

According to the Marcus–Hush–Stranks^{32–34} theory the value of the activation volume for the electron transfer step, $\Delta V_{\text{et}}^\ddagger$, can be evaluated as a sum of calculable contributions, equation (10) where $\Delta V_{\text{COUL}}^\ddagger$ is the volume change associated

$$\Delta V_{\text{et}}^\ddagger = \Delta V_{\text{COUL}}^\ddagger + \Delta V_{\text{SR}}^\ddagger + \Delta V_{\text{DH}}^\ddagger + \Delta V_{\text{IR}}^\ddagger \quad (10)$$

with the rearrangement to bring the ionic species together, expressed by equation (11) [σ is the reactant separation distance

$$\Delta V_{\text{COUL}}^\ddagger = \frac{N_A Z_1 Z_2 e_0^2}{4\pi\epsilon_0 \epsilon \sigma} \left(\frac{\beta}{3} - \delta \right) \quad (11)$$

when the electron is transferred (set to $r_1 + r_2$) (ref. 3)], $\Delta V_{\text{SR}}^\ddagger$ is the contribution to the rearrangement of the surrounding solvent molecules, expressed by equation (12) [$\epsilon_{\text{op}} = 1.780$ is

$$\Delta V_{\text{SR}}^\ddagger = \frac{N_A e_0^2}{16\pi\epsilon_0} \left[\left(\frac{1}{2r_1} + \frac{1}{2r_2} - \frac{1}{\sigma} \right) \frac{\partial}{\partial P} \left(\frac{1}{\epsilon_{\text{op}}} - \frac{1}{\epsilon} \right)_T - \frac{\beta}{3\sigma} \left(\frac{1}{\epsilon_{\text{op}}} - \frac{1}{\epsilon} \right) \right] \quad (12)$$

the relative permittivity of optical frequencies, $\partial(\epsilon_{\text{op}}^{-1} - \epsilon^{-1})_T / \partial P = -1.17 \times 10^{-4}$ MPa⁻¹ (ref. 3)], $\Delta V_{\text{DH}}^\ddagger$ is the term due to the Debye–Hückel or other electrolyte effects, expressed by equation (13) ($a = 0.66$ nm, $B = 3.29$ M^{-0.5}, $C = 1.174$ M^{-0.5} are

$$\Delta V_{\text{DH}}^\ddagger = \frac{RT Z_1 Z_2 C \sqrt{I}}{(1 + aB\sqrt{I})^2} [\delta(3 + 2aB\sqrt{I}) - \beta] \quad (13)$$

the Debye–Hückel parameters³) and $\Delta V_{\text{IR}}^\ddagger$ is the contribution due to internal rearrangement of the two reacting species and is neglected in our calculations since it is usually very small.

From equations (11)–(13) and the quoted values for the necessary parameters it follows that $\Delta V_{\text{COUL}}^\ddagger = -2.1$, $\Delta V_{\text{SR}}^\ddagger = -8.0$, and $\Delta V_{\text{DH}}^\ddagger = +2.0$ cm³ mol⁻¹ such that $\Delta V_{\text{et}}^\ddagger = -8.1$ cm³ mol⁻¹. The theoretically evaluated activation volume for the net reaction is $\Delta V_{12}^\ddagger = -11.3$ cm³ mol⁻¹, which is in close agreement with the experimental value $\Delta V_{12(\text{exp})}^\ddagger = -10.0 \pm 0.5$ cm³ mol⁻¹.

The Marcus theory³³ is often used for the estimation of the self-exchange rate constants especially where these are impossible to measure directly. According to this theory, the free energy of activation, ΔG_{12}^\ddagger , for a cross-reaction, considering electrostatic effects is given by equations (14)–(16) and ΔG_{11}^\ddagger ,

$$\Delta G_{12}^\ddagger = 0.5 (\Delta G_{11}^\ddagger + \Delta G_{22}^\ddagger + \Delta G_{12}^{\circ} + \Delta w) \quad (14)$$

$$\Delta w = w_{12} + w_{21} - w_{11} - w_{22} \quad (15)$$

$$\Delta G_{12}^{\circ} = -nF(E_{11}^{\circ} - E_{22}^{\circ}) \quad (16)$$

ΔG_{22}^\ddagger represent the free energies of activation for the self-exchange reactions. The free energies of activation are determined from the means of the Eyring equation (17). Since the self-

$$k_{ij} = k_B T h^{-1} \exp(-\Delta G_{ij}^\ddagger / RT) \quad (17)$$

exchange rate constant k_{22} of the couple $\text{HA}^- - \text{HA}^-$ has been reported,^{3,12} we used the above equations to calculate the self-exchange rate constant k_{11} of the couple $[\text{Fe}(\text{CN})_5(\text{NO}_2)]^{3-} - [\text{Fe}(\text{CN})_5(\text{NO}_2)]^{4-}$ which was so far unknown.

The free energy change for the cross-reaction ΔG_{12}° is obtained by means of equation (16) using $E_{11}^{\circ} = 0.39$ V (ref. 35) and $E_{22}^{\circ} = 0.71$ V (ref. 3). It results in $\Delta G_{12}^{\circ} = 30.60$ kJ mol⁻¹. The self-exchange rate constant, $k_{22} = 1.6 \times 10^5$ M⁻¹ s⁻¹ for the couple $\text{HA}^- - \text{HA}^-$, is adopted that leads on the basis of equation (17) to $\Delta G_{22}^\ddagger = 43.32$ kJ mol⁻¹. The electric work terms calculated from equation (15) results in $w_{11} = 9.35$ kJ mol⁻¹, $w_{22} = 0$, $w_{12} = 2.78$ kJ mol⁻¹, $w_{21} = 0$ and therefore $\Delta w = -6.57$ kJ mol⁻¹. With the above obtained values of ΔG_{12}° , ΔG_{22}^\ddagger , Δw and the experimental value of ΔG_{12}^\ddagger , $\Delta G_{11}^\ddagger = 46.45$ kJ mol⁻¹ and thus $k_{11} = 4.5 \times 10^4$ M⁻¹ s⁻¹ for the self-exchange rate constant of the couple $[\text{Fe}(\text{CN})_5(\text{NO}_2)]^{3-} - [\text{Fe}(\text{CN})_5(\text{NO}_2)]^{4-}$ at 25 °C and $I = 0.3$ M. The large value of k_{11} can be ascribed to a low internal reorganization energy required for the self-exchange process, as also found for many other complexes.¹² The obtained value of k_{11} is of the same order of magnitude as those found for the related systems $[\text{Fe}(\text{CN})_6]^{3-/4-}$ and $[\text{Fe}(\text{C}_2\text{O}_4)_3]^{3-/4-}$.^{3,12}

The experimental rate and activation parameters, as well as the theoretical calculations, clearly suggest that the first step of the reduction of $[\text{Fe}(\text{CN})_5(\text{NO}_2)]^{3-}$ by ascorbate anion follows an outer-sphere electron transfer mechanism. The theoretical calculations are in perfect agreement with the experimental results and exhibit the same trend as those found earlier for related systems.^{3,11,12}

Acknowledgements

The authors gratefully acknowledge financial support from State Committee for Scientific Research, Republic of Poland (KBN) and TEMPUS project JEP 8169.

References

- 1 D. Hornig, *Ann. N. Y. Acad. Sci.*, 1985, **258**, 103.
- 2 B. Frei, L. England and B. N. Ames, *Proc. Natl. Acad. Sci. USA*, 1989, **86**, 6377.
- 3 B. Bänisch, P. Martinez, J. Zuluaga, D. Uribe and R. van Eldik, *Z. Phys. Chem.*, 1991, **170**, 59.
- 4 P. Martinez and D. Uribe, *An. Quim.*, 1980, **76**, 201.
- 5 P. Martinez, J. Zuluaga and D. Uribe, *Z. Phys. Chem., Neue Folge*, 1983, **137**, 43.
- 6 P. Martinez and J. Zuluaga, *An. Quim.*, 1984, **80**, 179.
- 7 P. Martinez, J. Zuluaga, D. Uribe and R. van Eldik, *Inorg. Chim. Acta*, 1987, **136**, 11.
- 8 P. Martinez, J. Zuluaga, J. Kraft and R. van Eldik, *Inorg. Chim. Acta*, 1988, **146**, 9.
- 9 P. Martinez, J. Zuluaga and A. F. Rodriguez, *Z. Phys. Chem.*, 1989, **270**, 491; 1990, **271**, 597.
- 10 P. Martinez, J. Zuluaga, P. Noheda and R. van Eldik, *Inorg. Chim. Acta*, 1992, **195**, 249.
- 11 B. Bänisch, P. Martinez, D. Uribe, J. Zuluaga and R. van Eldik, *Inorg. Chim. Acta*, 1991, **30**, 4555.
- 12 B. Bänisch, R. van Eldik and P. Martinez, *Inorg. Chim. Acta*, 1992, **201**, 75.
- 13 E. Pelizetti, E. Mentasti and E. Pramauro, *Inorg. Chem.*, 1976, **15**, 2898.
- 14 M. Kimura and M. Yamamoto, *J. Chem. Soc., Dalton Trans.*, 1982, 423.
- 15 K. Tsukahara and Y. Yamamoto, *Bull. Chem. Soc. Jpn.*, 1981, **54**, 2642.
- 16 R. A. Rickman, R. L. Sorensen, K. O. Watkins and G. Davies, *Inorg. Chem.*, 1977, **16**, 1570.
- 17 S. Gangopadhyay, S. Saha, M. Ali and P. Banerjee, *Int. J. Chem. Kinet.*, 1991, **23**, 105.
- 18 J. Oszajca, G. Stochel, E. Wasielewska, Z. Stasicka, R. J. Gryglewski, A. Jakubowski and K. Cieslik, *J. Inorg. Biochem.*, in the press.
- 19 C. D. Hubbard and R. van Eldik, in *Chemistry under extreme or non-classical conditions*, eds. R. van Eldik and C. D. Hubbard, Wiley, New York, 1997, ch. 2, pp. 53–102.
- 20 M. Meier, J. Sun, J. F. Wishart and R. van Eldik, *Inorg. Chem.*, 1996, **35**, 1564.
- 21 M. Meier and R. van Eldik, *Chem. Eur. J.*, 1997, **3**, 33.
- 22 E. Hejmo, E. Porcel-Ortega, T. Senkowski and Z. Stasicka, *Bull. Pol. Acad. Chem.*, 1988, **7-8**, 351.
- 23 R. van Eldik, D. A. Palmer, R. Schmidt and H. Kelm, *Inorg. Chim. Acta*, 1981, **50**, 131.
- 24 R. van Eldik, W. Gaede, S. Wieland, J. Kraft, M. Spitzer and D. A. Palmer, *Rev. Sci. Instrum.*, 1993, **64**, 1355.
- 25 J. H. Swinehart and P. A. Rock, *Inorg. Chem.*, 1966, **5**, 573.
- 26 G. Stochel, R. van Eldik, E. Hejmo and Z. Stasicka, *Inorg. Chem.*, 1988, **27**, 2767.
- 27 G. Stochel, J. Oszajca, A. Herdegen, E. Wasielewska and Z. Stasicka, *12th Summer School of Coordination Compounds, Book of Abstracts, Wroclaw*, 1993, p. 146.
- 28 R. D. Cannon, *Electron Transfer Reaction*, Butterworths, London, 1980, ch. 4.2.
- 29 P. Hemmes, *J. Phys. Chem.*, 1972, **76**, 895.
- 30 K. R. Srinivasan and R. L. Kay, *J. Chem. Phys.*, 1974, **60**, 3645.
- 31 R. C. West, *Handbook of Chemistry and Physics*, CRC Press, Cleveland, OH, 1976–1977.
- 32 D. R. Stranks, *Pure Appl. Chem.*, 1974, **38**, 303.
- 33 R. A. Marcus, *J. Phys. Chem.*, 1956, **24**, 966; 1957, **26**, 867; 1963, **72**, 891.
- 34 N. S. Hush, *Trans. Faraday Soc.*, 1961, **57**, 557.
- 35 H. Chein-Ho, *Soochow J. Math. Nat. Sci.*, 1975, **1**, 133.

Received 15th April 1998; Paper 8/02821D

