

Oxidation of the thiosulfate ion with a (μ -oxo)diiron(III) complex: kinetic evidence for self reaction of the thiosulfate radical ion




Beauty Chaudhuri and Rupendranath Banerjee*

Department of Chemistry, Jadavpur University, Calcutta 700032, India

Received 23rd April 1998, Accepted 13th July 1998

In aqueous Hphen⁺–phen buffer media (pH 3.9–5.2), the complex ion [Fe^{III}₂(μ -O)(phen)₄(H₂O)₂]⁴⁺ **1** (phen = 1,10-phenanthroline) has been shown to oxidise the thiosulfate ion to the tetrathionate ion. In the presence of excess phenanthroline, **1** itself was reduced to [Fe(phen)₃]²⁺ **3**²⁺, an inhibitor for the reaction. According to the proposed mechanism, an inner-sphere, one-electron reduction of **1** by S₂O₃²⁻ generates [Fe(phen)₃]³⁺ **3**³⁺, [Fe(phen)₃]²⁺ **3**²⁺ and the thiosulfate radical ion S₂O₃^{•-}; **3**³⁺ is then further reduced to **3**²⁺ by another S₂O₃²⁻ ion. The radical ion S₂O₃^{•-} either re-oxidises **3**²⁺ to **3**³⁺ and thus inhibits the reaction, or produces the tetrathionate ion through a self reaction. Kinetic activity of **1** swamps that for its hydrolysed derivatives.

Thiosulfate is an important, industrial reducing agent. Mild oxidising agents oxidise it to tetrathionate¹ *via* the transient radical ion S₂O₃^{•-}, detected by induced polymerisation of acrylonitrile in aqueous solution.² The radical ion produces tetrathionate either by self reaction [eqn. (1)], or by coupling with S₂O₃²⁻ [eqn. (2a)] followed by oxidation of the S₄O₆³⁻ radical ion [eqn. (2b)].



Many workers assumed self reaction in schemes proposed for the oxidation reactions of S₂O₃²⁻. Nevertheless, a major objection to the self reaction (1) is that such a mechanism, which involves a second-order decay of the intermediate S₂O₃^{•-} ion would have a relatively high steady-state concentration of the intermediate. Re-oxidation of product by S₂O₃^{•-} would therefore be significant and product inhibition observed. So far, no such inhibition has been reported.


While investigating the preliminary kinetics of oxidation of S₂O₃²⁻ by the (μ -oxo)diiron(III) complex ion [Fe₂(μ -O)(phen)₄(H₂O)₂]⁴⁺ **1** (phen = 1,10-phenanthroline) in slightly acidic aqueous media, we recently observed product inhibition by [Fe(phen)₃]²⁺ **3**²⁺. The results appeared interesting compared with the reactions discussed above and we conducted the more detailed studies presented here. The crystal structure of **1** [NO₃]₄·5H₂O is known³ and some time ago we investigated the kinetics and equilibria of its reaction with hydroxylamine.⁴ Besides this, kinetic studies of oxidation by multinuclear iron complexes are in general rare.⁵

Results and discussion


Stoichiometry

In the presence of [complex] \ll c_{phen} (= [phen] + [Hphen⁺]) each mol of **1** oxidises 1.85 \pm 0.05 mol of thiosulfate, and is reduced to 1.90 \pm 0.06 mol of [Fe(phen)₃]²⁺. No sulfate ion could be detected in the product solution under conditions where the lower limit of detection is 1 \times 10⁻⁵ mol dm⁻³, but the expected yield of SO₄²⁻ is 2 \times 10⁻² mol dm⁻³ provided that **1** oxidises S₂O₃²⁻ to SO₄²⁻ quantitatively. The results suggest that \leq 0.05% of the total S₂O₃²⁻ consumed changed into SO₄²⁻.

The most likely oxidation product of S₂O₃²⁻ is therefore S₄O₆²⁻ [eqn. (3)]. Formation of the tetrathionate ion was further con-




firmed by the method of Kolthoff and Belcher⁶ using reaction (4).



Solution equilibria

In slightly acidic solutions, the oxo-bridge in **1** is thermodynamically stable.⁴ Unidentate ligands such as SCN⁻ rapidly substitute the aqua ligands in **1**, thus indicating their labile nature.⁷ Under the experimental conditions used, **1** and its hydrolytic derivative **2** coexist in rapid equilibrium⁴ [eqn. (5)]



[K_{a1} = (4.03 \pm 0.05) \times 10⁻⁶ mol dm⁻³ at 25.0 $^\circ$ C, I = 1.0 mol dm⁻³] with no evidence for oxo-bridge protonation and bridge breaking. At pH < 2, however, **1** slowly cleaves to [Fe(phen)₃]²⁺ **3**²⁺ identified spectrophotometrically.

Kinetics and reaction scheme

Fig. 1 demonstrates a typical example of spectral changes observed for an aqueous solution of **1** mixed with 20–300-fold thiosulfate and excess c_{phen}. Typical absorbance *versus* time data at 510 nm are displayed in Fig. 2(b). Noticeably, the reaction lags behind a hypothetical one-step, first-order reaction [Fig. 2(a)] and demonstrates product inhibition. The highest absorbance recorded in 2(b) is also less than that expected for complete conversion of **1** into **3**²⁺. The observed inhibition increases with increasing amount of added [Fe(phen)₃]²⁺ **3**²⁺ [Fig. 2(c) and 2(d)]. Presence or absence of dissolved dioxygen had no effect on the observed absorbance–time profiles.

The S₂O₃²⁻ ion is a well-known one-electron reducing agent^{1,8,9} and most probably reduces the diiron(III) species by one-electron steps. It is however, known that Fe^{III}Fe^{II} and Fe^{II}₂ dimers are putative outside a protein environment and quickly dissociate to monomers.^{5a,10} According to available kinetic and




Fig. 1 The UV/VIS spectrum for complex **1** (1). Spectra for a mixture of **1** with 15 mmol dm⁻³ thiosulfate at different times: immediately after mixing (2) and after 1 (3), 3 (4), 5 (5), 7 (6), 15 (7), 20 (8), 25 (9) and 30 (10) min; [complex] 0.05 mmol dm⁻³; *c*_{phen} 3.0 mmol dm⁻³; pH 4.50; *T* 25.0 °C; *I* 1.0 mol dm⁻³ (NaNO₃). The isosbestic point is at 389 nm; another isosbestic point can be seen at 623 nm on an expanded scale (not shown).







Fig. 2 The absorbance versus time curves at 510 nm: [complex] 0.05 mmol dm⁻³; *c*_{phen} 3.0 mmol dm⁻³; [S₂O₃²⁻] 15.0 mmol dm⁻³; pH 4.50; *T* 25.0 °C; *I* 1.0 mol dm⁻³ (NaNO₃); [3²⁺] (mmol dm⁻³), 0 (a) and (b); 0.05 (c); 1.0 (d); [3²⁺] in aqueous blank was the same as in the experimental mixture. Simulated curve for a hypothetical, one-step reaction with rate constant 2.83 × 10⁻³ dm³ mol⁻¹ s⁻¹ (a). Real data (b)–(d). Some experimental points in the initial parts of (b)–(d) have not been shown for clarity. The solid lines in (b)–(d) are simulated curves using the reaction sequence (6–8) and rate constants (dm³ mol⁻¹ s⁻¹), *k*₁ = 2.83 × 10⁻³, *k*₂ = 1.6 × 10⁶, *k*₋₂ = 3.5 × 10⁹, *k*₃ = 6.7 × 10⁶, *k*₋₃ = (3.3 ± 0.2) × 10⁻² s⁻¹; *ε* (dm³ mol⁻¹ cm⁻¹) values: **1** 2400; **3**²⁺ 1.1 × 10⁴; **3**³⁺ 0. The residual plots for (b), (c) and (d) are shown in the inset.

thermodynamic data^{11,12} all such iron(III) and iron(II) monomers must change completely to their respective tris chelates, viz., **3**³⁺ or **3**²⁺ in the presence of excess phenanthroline. The time-scale for such changes is very much smaller than in the present kinetic experiments. As discussed, the observed product inhibition suggests the self reaction of S₂O₃²⁻. Eqns. (6)–(8)



therefore form a simple and logical sequence for the reaction between **1** and S₂O₃²⁻; the product inhibition arising from re-oxidation of **3**²⁺ by S₂O₃²⁻.

This sequence permits excellent simulation (Fig. 2) of the observed kinetics using the package KFSIM40,^{13a} which expresses this reaction sequence as a series of differential equations and performs numerical integration to yield the optimised rate constants and the concentrations of the participating species at desired time intervals. The KFSIM40 package consists of two programs, KINSIM40 and FITSIM40. The program KINSIM40 compiles a reaction sequence written with the DOS editor, and simulates concentration (or any linear function of it) at desired time intervals corresponding to a given mechanism and a given set of rate constants. Incorporation of known absorption coefficients (*ε*) yields calculated values for the absorbance at different time. The simulated absorbance–time curve can be visually compared with the real curve. One can change the mechanism, any rate constant and *ε* value to produce a better fit.

The program FITSIM40 involves regression analysis utilising the Marquardt technique by default. It yields optimised rate constants, associated errors and concentrations (absorbance) for a real data set and a given mechanism. The absorbance–time data simulated by the optimised rate constants may be read from the DOS editor and hence residual plots constructed.

We fitted the observed absorbance–time data in the proposed sequence using eight input parameters: three known *ε* values (see caption to Fig. 2) and five rate constants, out of which *k*₂ and *k*₋₂ were kept fixed. Reasonably good estimates for these two constants are possible. Available redox potentials for the couples [Fe(phen)₃]^{3+/2+} (1.06 V)¹⁴ and S₂O₃^{-1/-2} (1.30 V)^{1a} are sufficiently precise and may be used to yield a reliable value for the ratio *k*₂/*k*₋₂ = 4.5 × 10⁻⁴. Again the *k*₋₂ step should be diffusion controlled since the analogous reaction between [Fe(bipy)₃]³⁺ and S₂O₃²⁻ is known to be so. A good estimate for *k*₋₂ is therefore 3.5 × 10⁹ dm³ mol⁻¹ s⁻¹.¹⁵ This along with the *k*₂/*k*₋₂ ratio, gives *k*₂ = 1.6 × 10⁶ dm³ mol⁻¹ s⁻¹. We therefore optimised only the three unknown rate constants *k*₁, *k*₃ and *k*₋₃ with FITSIM40 and thus obtained unique solutions. The observed fit is quite sensitive to the individual rate constants. For example, decreasing *k*₂ by more than 5%, keeping other constants unchanged, appreciably decreases the simulated absorbances and evidently worsens the fit, which cannot be improved by altering *k*₁, *k*₃ or *k*₋₃ in either direction. One may increase the simulated *A_t* by increasing *k*₁ and thus partially balance the adverse effect of decreasing *k*₂ on the fit, but then the *A_t* values at longer times far exceed the real value. The fit is affected less dramatically, though quite appreciably, when *k*₂ is kept constant whilst *k*₋₂ is decreased up to 95% of the input value. Further decrease in *k*₂ has no effect on the fit.

The observed kinetics and its simulation demonstrate that the proposed steps are necessary and sufficient to explain the observed inhibition of the reaction by **3**²⁺. We did test other different sequences. In one of them, for example, steps (6) and (7) were followed by eqn. (2). In another, a sequence of consecutive irreversible steps were considered: **1** → **3**²⁺ + **3**³⁺; **3**³⁺ → **3**²⁺. None reproduced the observed inhibition.

Variations in rate constants

The optimised second-order rate constant *k*₁ (Table 1) decreases with increasing [S₂O₃²⁻]. The calculated first-order rate constant *k*₁⁰ (= *k*₁[S₂O₃²⁻]) increases with increasing [S₂O₃²⁻], gradually saturating at high [S₂O₃²⁻] (Fig. 3). A similar variation is

observed for k_1^0 with $[H^+]$ (Fig. 3). However, k_1^0 does not change when c_{phen} is increased. The average values of optimised $k_3 = (6.7 \pm 0.3) \times 10^6 \text{ dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$ and $k_{-3} = (3.3 \pm 0.2) \times 10^{-2} \text{ s}^{-1}$ are independent of pH, c_{phen} , $[S_2O_3^{2-}]$ and $[3^{2+}]$ as expected.

Rate and equilibrium constants of some reactions analogous to eqn. (8) have been listed in Table 2. The equilibrium constants are moderate to high and reflect the strong nucleophilicity of the S-atom in its different oxoanions and oxoanion radicals. The equilibrium constant ($k_3/k_{-3} = 2 \times 10^8 \text{ dm}^3 \text{ mol}^{-1}$) for the formation of $S_4O_6^{2-}$ [eqn. (8)] is much larger than that ($800 \text{ dm}^3 \text{ mol}^{-1}$) for the formation of $S_4O_6^{3-}$ [eqn. (2a)] and reflects the higher stability of the $S_4O_6^{2-}$ ion.

The tendency of k_1^0 to saturate at higher $[H^+]$ suggests that a protonated species dominates in kinetics over its conjugate base. In the experimental pH range, the near-exclusive form of the reductant is $S_2O_3^{2-}$ since at 25 °C pK_{a1} and pK_{a2} for $H_2S_2O_3$ are low, 0.35 and 1.01 respectively.¹⁸ Some workers proposed that reduction of $[IrCl_6]^{2-}$ and I_2^9 with $S_2O_3^{2-}$ involved an asymmetrical form of $S_2O_3^{2-}$. Its conjugate acid (HS)SO₃⁻ is believed to be a weak acid ($pK_{SH} \approx 10$) and Mehrotra and co-workers¹⁹ assumed the protic equilibria, $(HS)SO_3^- \rightleftharpoons H^+ + S_2O_3^{2-}$ to explain their observed increase in rate with increasing pH in the reduction of $[IrCl_6]^{2-}$ with $S_2O_3^{2-}$. In the present system, rate increased with decreasing pH, and kinetically one cannot accommodate any of the possible protic equilibria involving the $S_2O_3^{2-}$ ion. Hence, the hydrolysis equilibrium of **1** [eqn. (5)] is the only plausible explanation for the observed dependence of k_1^0 on $[H^+]$, provided **1** dominates over **2** kinetically.

Table 1 Optimum k_1 values^a for the reduction of the $[Fe^{III}_2(\mu-O)(phen)_4(H_2O)_2]^{4+}$ ion with thiosulfate in aqueous solutions

$[S_2O_3^{2-}]/$ mmol dm ⁻³	$c_{\text{phen}}/$ mmol dm ⁻³	pH ^b	$10^3 k_1/\text{dm}^3$ mol ⁻¹ s ⁻¹
1.0	3.0	4.50	2.83
2.5			2.73
5.0			2.40
8.0			2.06
10.0			1.88
12.5			1.61
15.0 ^c			1.36
		5.16	1.22
		4.76	1.34
		4.50	1.37
		4.22	1.49
		4.10	1.53
		3.99	1.54
		3.93	1.55
	6.0		1.37
	9.0		1.37
	12.0		1.36


^a Temperature 25.0 °C; [complex] 0.05 mmol dm⁻³; $I = 1.0 \text{ mol dm}^{-3}$ (NaNO₃). Reported k_1 are the average of 2–3 determinations, precision of an individual set lies within 2–5%; standard deviation 4–8%. ^b Average of initial and final experimental pH. ^c A parallel experiment under conditions of this set was carried out after passing N₂ gas; rate constant changed only 2.5%. Experiment also repeated in the presence of 0.05 and 1.0 mmol dm⁻³ 3^{2+} added externally; rate constants did not change significantly.

Table 2 Rate constants for the self reaction of the $S_2O_3^-$ radical ion and some analogous reactions in aqueous solution at 25 °C

Reaction	Forward rate constant ^a	Reverse rate constant	Equilibrium constant	Ref.
$S_2O_3^{2-} + S_2O_3^- \rightleftharpoons S_4O_6^{3-}$	2×10^9	2.5×10^{6b}	800 ^c	2, 16
$SO_3^- + SO_3^- \rightleftharpoons S_2O_6^{2-}$	1.8×10^8	—	—	17
$S_2O_3^- + S_2O_3^- \rightleftharpoons S_4O_6^{2-}$	6.7×10^6	3.3×10^{-2b}	2×10^{8c}	This work
$I_2 + S_2O_3^{2-} \rightleftharpoons I_2S_2O_3^{2-}$	7.8×10^9	2.5×10^{2b}	3.2×10^{7c}	9
$I_3^- + S_2O_3^{2-} \rightleftharpoons I_2S_2O_3^{2-} + I^-$	4.2×10^8	9.5×10^{3a}	4.42×10^4	9

^a In $\text{dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$. ^b In s^{-1} . ^c In $\text{dm}^3 \text{ mol}^{-1}$.

The tendency of k_1^0 to saturate at higher $[S_2O_3^{2-}]$ suggests adduct formation between the oxidant and reductant. This is expected from the labile nature^{4,7} of the terminal H₂O in **1** and other oxo-bridged iron(III) complexes.^{5,20} Adduct formation is indicated also from the observed increase in the absorbance at the isosbestic point (389 nm) with increase in $[S_2O_3^{2-}]$. Measured absorbances at 389 nm are 0.155, 0.226, 0.240 and 0.248 for $[S_2O_3^{2-}] = 0, 5.0, 10.0$ and $15.0 \text{ mmol dm}^{-3}$ respectively. The standard deviation in three independent measurements for A_{389} at a fixed $[S_2O_3^{2-}]$ is ≤ 0.003 , which shows that the changes in A_{389} with $[S_2O_3^{2-}]$ are reproducible and systematic, albeit small, and do not arise out of experimental uncertainties. It is noticeable in this instance that the spectrum of an aqueous solution of **1** without $S_2O_3^{2-}$ does not pass through the isosbestic point (Fig. 1). One can conclude that **1** and $S_2O_3^{2-}$ together in solution rapidly form some new species, (**I**, say) which slowly changes to $[Fe(\text{phen})_3]^{2+}$ maintaining the isosbestic point [eqn. (9)]. This is a somewhat elaborated form



of eqn. (6) and leads to eqn. (10) provided that the hydrolysed Fe^{III}_2 species **2** does not react appreciably with $S_2O_3^{2-}$.

$$k_1^0 = (k_i K_1 [H^+] [S_2O_3^{2-}]) / \{ [H^+] (1 + K_1 [S_2O_3^{2-}]) + K_{a1} \} \quad (10)$$

As expected from eqn. (10), plots of $1/k_1^0$ vs. $1/[S_2O_3^{2-}]$ at fixed pH and plots of $1/k_1^0$ vs. $1/[H^+]$ at fixed $[S_2O_3^{2-}]$ are excellent straight lines. Slopes and intercepts of these two straight lines yield $k_i K_1 = (0.41 \pm 0.02) \text{ dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$ [$k_i = (3.7 \pm 0.2) \times 10^{-3} \text{ s}^{-1}$], $K_1 = 111 \pm 5 \text{ dm}^3 \text{ mol}^{-1}$ and $K_{a1} = 5.1 \times 10^{-6} \text{ mol}$




Fig. 3 Variation of the first-order rate constant $k_1^0 (= k_1 [S_2O_3^{2-}])$ with change in $[S_2O_3^{2-}]$ at fixed pH (A) and with $[H^+]$ at fixed $[S_2O_3^{2-}]$ (B); [complex] 0.05 mmol dm⁻³; T 25.0 °C; I 1.0 mol dm⁻³ (NaNO₃). The solid lines represent the best visual fits.

dm⁻³ in good agreement with its spectrophotometrically measured value⁴, 4.0×10^{-6} mol dm⁻³. This agreement lends support to eqn. (10) and the proposed inactivity of **2**. It is also indicated that the intermediate adduct **I** contains only one S₂O₃²⁻. The Fuoss equation²¹ yields $K_1 = 5.7$ dm³ mol⁻¹ for a purely electrostricted ion pair formed between **1** and S₂O₃²⁻ ions, whose respective radii are 71 nm (ref. 3) and 33 nm (ref. 8). The kinetically measured value for K_1 is much larger than this value and lies close to the first formation constant (126 dm³ mol⁻¹ at 25 °C, $I = 0.47$ mol dm⁻³) for the Fe³⁺-S₂O₃²⁻ system.²² Moreover, S₂O₃²⁻ is a strong nucleophile ($n = 6.36$ on the Swain-Scott scale),^{18,23} its co-ordination precedes the redox reactions^{1,9} with [Ag(OH)₄]⁻, [Fe(H₂O)₆]³⁺, Cr(IV), [Au(NH₃)₄]³⁺, [AuCl₄]⁻, [OsO₄(OH)₂]²⁻, [Co(NTA)(H₂O)₂], [Cu(NH₃)₄]²⁺, I₂ and I₃⁻. Deaquation of an iron(III) site in **1** and its inner-sphere interaction with thiosulfate is therefore anticipated. However, oxidation of S₂O₃²⁻ by [Fe(bipy)₃]³⁺, is known to be outer-sphere.¹⁴ The analogous reaction of [Fe(phen)₃]³⁺ with S₂O₃²⁻ should also follow a similar mechanism since ligand substitution in both complexes are slow compared to redox. The outer-sphere mechanism for [Fe(phen)₃]³⁺ is further supported by the Marcus relation, which along with the self exchange rate for the [Fe(phen)₃]^{3+/2+} couple²⁴ yields $k_2 = 1.8 \times 10^5$ dm³ mol⁻¹ s⁻¹. The value agrees reasonably well with our previous estimate.

The assumption regarding non-reactivity of **2** seems justified. Inner-sphere adduct formation with S₂O₃²⁻ by **2** and further by **I** should both be less favourable than **1** due to their lower charges. The lesser number of replaceable water molecules in **I** and **2** than in **1** should also disfavour adduct formation on statistical grounds. Not only should **1** form a better adduct but it should also dominate kinetically over **2** because **1** is more electron deficient and hence can be reduced more easily. It is true for most redox systems involving cationic complexes and anionic reductants that protonation of the oxidant increases its reactivity whereas the opposite is true for protonation of the reductant.²⁵

We chose a somewhat restricted set of conditions of pH and c_{phen} for the present experiments to avoid some potential complications. The large excess of c_{phen} over the iron complexes suppresses the loss of phen⁴ by **1** and thus minimises the number of kinetically active iron(III) species. Excess c_{phen} lightly buffers the solution to within ± 0.08 (but mostly within ± 0.04) pH units and also sequesters any trace metal ion, known to strongly catalyse^{14,26} the dissociation of S₂O₃²⁻ (a potential complication in the study of redox reactions of S₂O₃²⁻). Use of excess c_{phen} was essential also to ensure quantitative formation of the tris(phenanthroline) complexes **3**³⁺ and **3**²⁺ from any transient bis intermediates. Otherwise, interpretation of kinetic data becomes difficult and oxidation of [Fe(phen)₃]²⁺ to **1** may be an additional complication since Cyfert²⁷ showed that hydrogen peroxide oxidises [Fe(phen)₃]²⁺ to a {Fe₂O}⁴⁺ species, possibly **1** in the absence of excess phenanthroline.

Substitution or redox

Spectral observations and tendency of k_1^0 to saturate at high [S₂O₃²⁻] suggested a role for the intermediate **I** in kinetics. Which of the two processes is the slower one, formation of **I** or its subsequent decomposition to products, is an interesting question.

Chelation labilises the co-ordinated water molecules. For example, the water exchange rate constant²⁸ for [Fe(H₂O)₆]³⁺ is 280 s⁻¹ at 25 °C, but the value for²⁹ [Fe(EDTA)(H₂O)]⁻ is $\approx 10^5$ s⁻¹. Candle and Crumbliss³⁰ observed a similar labilising influence of hydroxamic acids on water molecules co-ordinated to iron(III) centres while investigating the dissociation reactions of some iron(III) complexes. Labilisation by the ligand Tiron (4,5-dihydroxy-1,3-benzene disulfonic acid, disodium salt monohydrate) was noted also by Zhang and Jordan³¹ while studying the kinetics of dissociation reactions of mono-, bis- and tris-

Table 3 Rate constants ($k/\text{dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$) for the reduction of some iron(III) complexes in aqueous media, 25.0 °C

Complex	Reducing agent	k	Ref.
[Fe(phen) ₃] ³⁺	S ₂ O ₃ ²⁻	1.8×10^5 ^a	This work
	Ascorbate(1-)	1.5×10^9 ^b	32
	ClO ₂ ⁻	2.3×10^7	33
[Fe ₂ O(phen) ₄ (H ₂ O) ₄] ⁴⁺ ^c	S ₂ O ₃ ²⁻	0.41	This work
	NH ₂ OH	2.31	4
[Fe ₃ O(O ₂ CMe) ₆] ^d	NH ₂ OH	4.6×10^{-2}	5(b)
[Fe ₂ O(L)(CH ₃ CN) ₂] ^e	Hydroquinone	4.3	5(a)
	Catechol	1.7	5(a)
	Picric acid	0.4	5(a)
	Triphenylphosphine	0.02	5(a)

^a Estimated value, see text. ^b At 20.0 °C. ^c Phenanthroline buffer. ^d Acetate buffer, 35.0 °C. ^e L is 2dmg·2BF₂, CH₃CN media.

complexes of this ligand with iron(III) in acidic aqueous solution. Based on these observations, one expects that the water exchange rate constant for **1** should be much higher than 280 s⁻¹ and the observed, small value for k_i [$(3.7 \pm 0.2) \times 10^{-3}$ s⁻¹] better fits the intramolecular electron-transfer step in **I**. Accordingly, we conclude that in the present system rapid adduct formation precedes the slow intramolecular electron transfer step.

The k_i step itself must represent a multistep reaction. Its most likely components are: (a) slow intramolecular electron transfer and formation of a Fe^{III}Fe^{II} dimer (k_i), (b) fast cleavage of the oxo-bridge in the Fe^{III}Fe^{II} dimer, and generation of Fe^{III} and Fe^{II} monomers, (c) rapid and quantitative conversion of the monomers to **3**²⁺ and **3**³⁺ in the presence of excess phenanthroline.

Rate constants for oxidation by some iron(III) complexes have been gathered in Table 3. Note that the oxo-bridged species react at a slower rate than the comparable monomers.

To our knowledge, the present kinetic investigation is the first example of product inhibition in the oxidation of S₂O₃²⁻ by a metal ion complex and constitutes strong evidence in favour of the self reaction of S₂O₃²⁻ leading to the tetrathionate ion.

Experimental

Materials

Crystals of [Fe₂(μ-O)(phen)₄(H₂O)₂][NO₃]₄·5H₂O were prepared by a known³ method. The prepared samples gave satisfactory electronic spectra^{3,4} and elemental analyses (Found: C, 47.2; H, 3.7; Fe, 9.1; N, 13.8. Calc. for C₄₈H₃₂Fe₂N₁₂O₁₃·7H₂O: C, 47.15; H, 3.8; Fe, 9.1; N, 13.75%). Stock solutions of sodium thiosulfate (AR, BDH) were renewed daily and standardised iodometrically against potassium dichromate.³⁴ Nitrate salts of **3**³⁺ and **3**²⁺ were prepared according to the literature.³⁵ 1,10-Phenanthroline monohydrate (Sigma) was used without further purification. Solutions of recrystallised NaNO₃ (AR, BDH) were standardised by the ion exchange technique.³⁶ All other reagents used were of analytical grade. Chromium(II)-scrubbed dinitrogen gas and triply distilled water were used throughout. Reported kinetic data are for 25.0 °C and $I = 1.0$ mol dm⁻³. The molar absorption coefficient (ϵ) of an aqueous solution of **1** in the presence of S₂O₃²⁻ was measured immediately after mixing the reagents under the experimental conditions; ϵ for **3**²⁺ was measured under experimental conditions with a fresh solution of the prepared sample (see Table 1, footnote a); **3**³⁺ does not absorb at 510 nm, the experimental wavelength.

Kinetics

Kinetics was monitored at 510 nm, the visible maximum of the reaction product [Fe(phen)₃]²⁺, *in situ* in the electrically controlled thermostatted cell housing (CPS-240A) of a Shimadzu spectrophotometer (UV-1601 PC). A large excess of thiosulfate

and c_{phen} over the complex ($0.05 \text{ mmol dm}^{-3}$) were used. For some faster reactions, the reductant was rapidly and directly injected into the cell containing other materials in solution and thermally equilibrated in the cell housing. Absorbance data were collected in the 'Kinetic mode' of the spectrophotometer and data collection started almost simultaneously with mixing of the reagents. The pH of experimental solutions was measured with an Orion (710 A) pH-meter using a calibrated³⁶ combined electrode.

Stoichiometry

A 100 ml portion of $0.10 \text{ mmol dm}^{-3}$ complex solution containing three different concentrations ($0.40, 0.60, 1.0 \text{ mmol dm}^{-3}$) of $\text{S}_2\text{O}_3^{2-}$ but a fixed c_{phen} (3.0 mmol dm^{-3}) at pH 4.50 was allowed to equilibrate up to a time after which no absorbance change could be noticed over a period of 3–4 h. The amount of 3^{2+} generated was determined from the solution absorbance measured at 510 nm after appropriate dilution. The whole solution was then eluted through a column of cation exchange resin in the Na^+ -form. The amount of $\text{S}_2\text{O}_3^{2-}$ in the eluted solution was quantified iodometrically using starch indicator.³⁴ In another sample (P) 100 ml, 4.0 mmol dm^{-3} complex were taken with $20 \text{ mmol dm}^{-3} \text{ S}_2\text{O}_3^{2-}$. The solution was eluted similarly after equilibrium was attained and the volume was reduced to $\approx 5 \text{ ml}$ by evaporation at 50°C . A concentrated solution of $\text{Sr}(\text{NO}_3)_2$ was added to test for the presence of sulfate ion in the product.³⁷ Under experimental conditions similar to sample P, we tested for the presence of tetrathionate in the eluted solution by the cyanolysis method of Kolthoff and Belcher.⁶

Acknowledgements

The work has been carried out with financial assistance from the Department of Science and Technology (New Delhi). This and the award of a Senior Research Fellowship to B. C. by the University Grants Commission (New Delhi) are gratefully acknowledged. We are grateful to Professor C. Frieden and to Dr. Joydeep Mitra for their kind gift of the Program KFSIM40. Some recent applications of this Program may be found in ref. 13(b).

References

- (a) R. Sarala and D. M. Stanbury, *Inorg. Chem.*, 1992, **31**, 2771; (b) J. J. Byerley, S. A. Fouda and G. L. Rampel, *J. Chem. Soc., Dalton Trans.*, 1973, 889; (c) J. D. Rush and L. J. Kirschenbaum, *Inorg. Chem.*, 1985, **24**, 744.
- R. Mehnert and O. Brede, *Radiat. Phys. Chem.*, 1984, **23**, 463.
- J. E. Plowman, T. M. Loehr, C. K. Schauerand and O. P. Anderson, *Inorg. Chem.*, 1984, **23**, 3553.
- B. Chaudhuri and R. Banerjee, *Can. J. Chem.*, 1998, in the press.
- (a) H. Noglik, D. W. Thompson and D. V. Stynes, *Inorg. Chem.*, 1991, **30**, 4571; (b) K. Arora, P. Bhatnagar, A. P. Bhargava and Y. K. Gupta, *J. Chem. Soc., Dalton Trans.*, 1991, 1081.
- I. M. Kolthoff and R. Belcher, *Volumetric Analysis*, Interscience, New York, 3rd edn., 1957.
- A. F. M. Nazer and C. F. Wells, *J. Chem. Soc., Dalton Trans.*, 1980, 1532.
- R. Sarala, S. B. Rabin and D. M. Stanbury, *Inorg. Chem.*, 1991, **30**, 3999.
- W. M. Scheper and D. W. Matgerum, *Inorg. Chem.*, 1992, **31**, 5466.
- D. M. Kurtz, jun., *Chem. Rev.*, 1990, **90**, 585; S. Menage, B. A. Brennan, C. Juarez-Garcia, E. Munck and L. Que, jun., *J. Am. Chem. Soc.*, 1990, **112**, 6423; L. A. Bottomley, C. Ercolani, J. N. Gorce, G. Pennesi and G. Rassi, *Inorg. Chem.*, 1986, **25**, 2339; A. L. Feig, A. Masschelein, A. Bakac and S. J. Lippard, *J. Am. Chem. Soc.*, 1997, **119**, 334.

- C. V. Banks and R. I. Bystroff, *J. Am. Chem. Soc.*, 1959, **81**, 6153; M. J. Blandamer, J. Burgess, H. J. Cowles, I. M. Horn, J. B. F. N. Engberts, S. A. Galema and C. D. Hubbard, *J. Chem. Soc., Faraday Trans.*, 1989, **85**, 3733; J. C. Thompson and H. A. Mottola, *Anal. Chem.*, 1984, **56**, 755; T. S. Lee, I. M. Kolthoff and D. L. Leussing, *J. Am. Chem. Soc.*, 1948, **70**, 2348, 3596.
- I. A. Brikun, I. A. Storozhev and E. A. Mambetkaziev, *Izv. Akad. Nauk Kaz. SSR, Ser. Khim.*, 1984, **3**, 66 (referred to in *Coord. Chem. Rev.*, 1992, **113**, 131); A. S. P. Vittal, P. V. Rao and K. J. M. Rao, *React. Catal. Lett.*, 1983, **23**, 175.
- (a) B. A. Barshop, R. F. Wrenn and C. Frieden, *Anal. Biochem.*, 1983, **130**, 134; C. Frieden, *Trends Biochem. Sci.*, 1994, **240**, 311; (b) K. Wang and J. M. Mayer, *J. Am. Chem. Soc.*, 1997, **119**, 1470; M. C. Ghosh, J. W. Reed, R. N. Bose and E. S. Gould, *Inorg. Chem.*, 1994, **33**, 73.
- D. M. Stanbury, W. K. Wilmarth, S. Khalof, H. N. Po and J. E. Byrd, *Inorg. Chem.*, 1980, **19**, 2715; W. K. Wilmarth, D. M. Stanbury, J. E. Byrd, H. N. Po and C. P. Chua, *Coord. Chem. Rev.*, 1983, **51**, 155.
- B. S. Brunshwig and N. Sutin, *Inorg. Chem.*, 1979, **18**, 1731.
- R. Mehnert, O. Brede and I. Janovsky, *Radiochem. Radioanal. Lett.*, 1982, **53**, 299.
- S. J. Waygood and W. J. McEtroy, *J. Chem. Soc., Faraday Trans.*, 1992, **88**, 1525.
- J. Hine, *Physical Organic Chemistry*, McGraw-Hill, New York, 1962, p. 161.
- B. Goyal, S. Solanki, S. Arora, A. Prakash and R. N. Mehrotra, *J. Chem. Soc., Dalton Trans.*, 1995, 3109.
- S. Menage, J. M. Vincent, C. Lambeaux, G. Chottard, A. Grand and M. Fantecane, *Inorg. Chem.*, 1993, **32**, 4766; R. H. Beer, W. B. Tolman, S. G. Bott and S. J. Lippard, *Inorg. Chem.*, 1991, **30**, 2082; S. Ito, T. Okuno, H. Matsushima, T. Tokii and Y. Nishida, *J. Chem. Soc., Dalton Trans.*, 1996, 4037; R. M. Buchanan, S. Chen, J. F. Richardson, M. Bressan, L. Forti, A. Morvillo and R. H. Fish, *Inorg. Chem.*, 1994, **33**, 3208.
- W. R. Gilkerson, *J. Chem. Phys.*, 1956, **25**, 1199; R. M. Fuoss, *J. Am. Chem. Soc.*, 1957, **79**, 3301; 1958, **80**, 5059.
- F. M. Page, *Trans. Faraday Soc.*, 1954, **50**, 120.
- C. G. Swain and C. B. Scott, *J. Am. Chem. Soc.*, 1953, **75**, 141.
- M. Chan and A. C. Wahl, *Abstract from 167th National Meeting of the American Chemical Society*, Los Angeles, CA, 1974, paper INOR 97.
- S. Mukhopadhyay and R. Banerjee, *J. Chem. Soc., Dalton Trans.*, 1994, 1349; S. K. Ghosh, R. N. Bose and E. S. Gould, *Inorg. Chem.*, 1987, **26**, 2688; K. D. Asmus, M. Banifacie, P. Toffel, P. O'Neill, D. S. Frohlinde and S. Steenken, *J. Chem. Soc., Faraday Trans. 1*, 1978, 1820.
- M. Kimura and M. Ishibashi, *Inorg. Chim. Acta*, 1987, **129**, 69; K. K. Sengupta, S. Das and S. SenGupta, *Transition Met. Chem. (London)*, 1988, **13**, 155.
- M. Cyfert, *Inorg. Chim. Acta*, 1985, **98**, 25.
- F. Basolo and R. G. Pearson, *Mechanisms of Inorganic Reactions*, Wiley, New York, 2nd edn., 1967, 198.
- M. D. Ritler and R. Van Eldik, *J. Chem. Soc., Dalton Trans.*, 1992, 1037 and references therein.
- M. T. Candle and A. L. Crumbliss, *Inorg. Chem.*, 1994, **33**, 4077.
- Z. Zhang and R. B. Jordan, *Inorg. Chem.*, 1996, **35**, 1571.
- E. Pellizzetti, E. Mentasti and E. Pramauro, *Inorg. Chem.*, 1976, **15**, 2898.
- L. A. Lednicky and D. M. Stanbury, *J. Am. Chem. Soc.*, 1983, **105**, 3098.
- A. I. Vogel, *Quantitative Inorganic Analysis*, ELBS, London, 3rd edn., 1961, p. 297.
- N. Sutin and B. M. Gordon, *J. Am. Chem. Soc.*, 1961, **83**, 70; T. S. Lee, I. M. Kolthoff and D. L. Leussing, *J. Am. Chem. Soc.*, 1948, **70**, 2348.
- S. Kundu, A. Bhattacharya and R. Banerjee, *J. Chem. Soc., Dalton Trans.*, 1996, 3951.
- A. I. Vogel, *Qualitative Chemical Analysis*, Longmans, Green and Co., London, 3rd edn., 1942, p. 329.