

The first alkyl bismuthates: tris(trifluoromethyl)fluoro- and tetrakis(trifluoromethyl)-bismuthate

Wieland Tyrra,^a Dieter Naumann,^{*a} Natalya V. Kirij,^b Alexander A. Kolomeitsev^b and Yurii L. Yagupolskii^b

^a Institut für Anorganische Chemie, Universität zu Köln, D-50939 Köln, Germany.

E-mail: d.naumann@uni-koeln.de

^b Institute of Organic Chemistry, National Academy of Sciences of Ukraine, 253660 Kiev, Ukraine

Received 11th January 1999, Accepted 25th January 1999


Tris(trifluoromethyl)bismuth reacts with fluoride ions and intermediately generated trifluoromethyl anions to form the first representatives of hitherto unknown fluoroalkyl-bismuthates: tris(trifluoromethyl)fluoro- and tetrakis(trifluoromethyl)-bismuthates.

Hypervalent bismuthate(III) anions with five electron pairs in the bismuth valence shell are important models for the transition states in nucleophilic substitution reactions at bismuth(III) centers. Therefore, such compounds are of considerable interest. To date some stable bismuthates have been isolated,¹⁻⁴ but to our knowledge there is no indication in the literature on the synthesis of acyclic hypervalent alkylbismuthate(III) species bearing at least one alkyl group on bismuth. Meanwhile such species have not only theoretical interest, but could be potentially used as precursors to prepare unknown tetra- and pentaalkyl derivatives of Bi(V), which are not available *via* reactions of trialkylbismuth dihalides with organometallic compounds. The only example of a tetramethylbismuth compound is [Bi(CH₃)₄]OSO₂CF₃ as the product of the reaction of Bi(CH₃)₃ and CH₃OSO₂CF₃.⁵ Some of us have developed a convenient procedure to synthesize perfluorinated trialkyl bismuthanes.^{6a,b}

We suppose that introduction of trifluoromethyl groups to bismuth(III) increases the Lewis acidity of the bismuth compounds which as a result should enhance the interaction of such fluorinated bismuthanes with nucleophiles, stabilizing hypervalent anionic four-coordinated bismuthates and increasing their ability for CF₃ group transfer. Successful fluoride induced C₆F₅ group transfer reactions presumably involving tris(pentafluorophenyl)fluorobismuthates, [Bi(C₆F₅)₃F]⁻, have been established in the literature for several years.^{7a,b}

The stabilising influence of fluorine and trifluoromethyl groups on phosphoranes is well documented. There are also two reports on stabilisation by fluorine atoms and CF₃ groups of phosphites, which can be regarded as the conjugate bases of phosphoranes.^{8,9}


The reaction of Bi(CF₃)₃ with the fluoride ion proceeds in THF, glyme (H₃COCH₂CH₂OCH₃) or diethyl ether. As sources of F⁻ we used [{N(C₂H₅)₂]₃S]⁺F⁻, [N(PPh₃)₂]⁺F⁻, [N(CH₃)₄]⁺F⁻ or [N(C₂H₅)₂]₃PF₂, as shown in eqn. (1), where




$\text{A}^+ = \{[\text{N}(\text{C}_2\text{H}_5)_2]_3\text{S}\}^+, [\text{N}(\text{PPh}_3)_2]^+, [\text{N}(\text{CH}_3)_4]^+$ or $\{[\text{N}(\text{C}_2\text{H}_5)_2]_3\text{PF}_2\}^+$.

The stability of hypervalent bismuthates formed depends on the counter ions and the nature of the fluoride ion source. The most stable solid bismuthate, [N(CH₃)₄]⁺[Bi(CF₃)₃F]⁻ **1**,[†] was prepared using [N(CH₃)₄]⁺F⁻ as a source of the fluoride ion. Surprisingly, tetramethylammonium fluoride was a better fluoride ion source than [{N(C₂H₅)₂]₃S}F. Covalent tris(diethylamino)difluorophosphorane, a mild donor of fluoride ion, reacts reversibly with Bi(CF₃)₃. The tris(diethylamino)fluoro-

phosphonium cation formed in the course of this reaction is electrophilic enough to react reversibly with [Bi(CF₃)₃F]⁻ to form [N(C₂H₅)₂]₃PF₂. This behaviour can be interpreted as a first hint of the fluorinating properties of [Bi(CF₃)₃F]⁻, eqn. (2).



Tris(trifluoromethyl)bismuth reacts with the intermediately formed CF₃⁻ anions (generated from (CH₃)₃SiCF₃ and [N(CH₃)₄]⁺F⁻) in diethyl ether, THF or glyme to form tetrakis(trifluoromethyl)bismuthate [N(CH₃)₄]⁺[Bi(CF₃)₄]⁻ **2**,[‡] eqn. (3). The reaction proceeds at -70 to -50 °C. An increase



in temperature causes the formation of [N(CH₃)₄]⁺[Bi(CF₃)₃F]⁻ **1** as a by-product. The formation of **1** in reactions above -50 °C in the presence of (CH₃)₃SiCF₃ indicates that at lower temperatures primarily [(CH₃)₃Si(CF₃)F]⁻ is formed which undergoes a replacement reaction with Bi(CF₃)₃ below -50 °C. At elevated temperatures (>-50 °C) Bi(CF₃)₃ appears to be a harder Lewis acid than (CH₃)₃SiCF₃ and reacts preferably with the fluoride ion. Therefore, temperature constancy (-70 to -50 °C) is necessary for the preparation of [Bi(CF₃)₄]⁻.

Hypervalent anionic bismuth species **1,2** are solids, which are stable in a dry nitrogen atmosphere at ambient temperature for some days and can be stored without decomposition in a refrigerator. CF₃-containing bismuthates are extremely reactive towards all common aprotic and especially protic solvents. Trifluoromethane formation in THF is observed even at -70 °C.

The reactivity of both bismuthate anions was studied in reactions with some organic and inorganic electrophilic reagents. To date, only reactions accompanied by a cleavage of the Bi-F or one Bi-CF₃ bond have been found.

Compound **1** reacts as a fluorinating agent. The interaction of this compound with trimethylchlorosilane and trichlorofluoromethane leads to chlorine exchange affording (CH₃)₃SiF and CCl₂F₂, respectively, and Bi(CF₃)₃.

We have found, despite the high reactivity with aprotic solvents, that **2** is an excellent source of the trifluoromethyl anion even at -78 °C. Successful trifluoromethylations with this hypervalent bismuthate were carried out with pentafluoropyridine, (CH₃)₃SnCl and SO₂Cl₂ to give the trifluoromethylated compounds 4-(trifluoromethyl)tetrafluoropyridine, trimethyl-(trifluoromethyl)tin and trifluoromethylsulfurylchloride, respectively.

The bismuthates **1,2** are unambiguously identified by a combination of NMR techniques supported by a satisfactory elemental analysis for all elements. In the ¹⁹F NMR spectrum (-58 °C, THF-d₈) of the [Bi(CF₃)₃F]⁻ anion the CF₃ group

resonance is detected as a broad singlet at $\delta -39.84$, remarkably shifted to higher field in comparison with the parent molecule $\text{Bi}(\text{CF}_3)_3$, $\delta -33.40$, and the fluorine atom as a singlet at $\delta -106.33$. The ^{13}C NMR resonance of the CF_3 groups is detected as a quartet at $\delta 183.16$ ($^1J_{\text{CF}} = 395.0$ Hz). The ^{19}F NMR resonance of the trifluoromethyl groups in $[\text{Bi}(\text{CF}_3)_4]^-$ is located at higher field ($\delta -41.21$). The quartet of the trifluoromethyl groups in the ^{13}C NMR spectrum is detected at $\delta 167.37$ ($^1J_{\text{CF}} = 405$ Hz).

According to VSEPR theory, both salts should show resonances of two different types of CF_3 groups in the ^{19}F NMR spectrum. The fact that only one resonance is observed may be explained by a fast exchange of axial and equatorial CF_3 groups in solution (even at low temperature), which is not detectable on the NMR time scale.

Acknowledgements

We gratefully acknowledge the support and financial assistance of this work by the Deutsche Forschungsgemeinschaft and the Fonds der Chemischen Industrie.

Notes and references

† Preparation of **1**: tetramethylammonium fluoride (1 mmol) was gradually added under argon to a solution of tris(trifluoromethyl)bismuth (1 mmol) in 5 ml diethyl ether at -50°C and the reaction mixture was stirred for 2 h at -40°C . The solvent was removed *in vacuo* below -20°C and the remaining product was washed twice with diethyl ether at -30°C and finally dried *in vacuo*. The salt was obtained in nearly quantitative yield and was stable at -30°C under argon for two weeks. ^{19}F NMR (282.35 MHz, THF- d_8 , -58°C , ppm) $\delta -39.84$ (CF_3 , 9F), -106.33 (F, 1F); ^{13}C NMR (75.48 MHz, THF- d_8 , -58°C , ppm) $\delta 183.16$ (CF_3 , q, $^1J_{\text{CF}} 395$), 53.28 (CH_3 , q, $^1J_{\text{CH}} 146$ Hz) (Found: C,

16.88; H, 2.64; Bi, 40.68; F, 37.20; N, 2.41. $\text{C}_7\text{H}_{12}\text{BiF}_{10}\text{N}$ requires C, 16.51; H, 2.38; Bi, 41.05; F, 37.31; N, 2.75%).

‡ Preparation of **2**: to a solution of tris(trifluoromethyl)bismuth (1 mmol) in 5 ml diethyl ether at -70°C trifluoromethyltrimethylsilane (3 mmol) and tetramethylammonium fluoride (1 mmol) were added. The mixture was stirred at -60 to -65°C for 1 h. All volatile products were removed *in vacuo* at -30°C , the remaining product was washed twice with diethyl ether at -30°C and dried *in vacuo*. The salt was obtained in nearly quantitative yield and was stable at -30°C under argon for 2–3 days. ^{19}F NMR (282.35 MHz, THF- d_8 , -58°C , ppm) $\delta -41.21$ (CF_3); ^{13}C NMR (75.48 MHz, THF- d_8 , -58°C , ppm) $\delta 167.37$ (CF_3 , q, $^1J_{\text{CF}} 405$), 55.10 (CH_3 , q, $^1J_{\text{CH}} 146$ Hz) (Found: C, 16.81; H, 2.45; Bi, 37.72; F, 39.43; N, 2.09. $\text{C}_8\text{H}_{12}\text{BiF}_{12}\text{N}$ requires C, 17.18; H, 2.16; Bi, 37.37; F, 40.77; N, 2.51%).

- 1 J.-P. Finet, *Chem. Rev.*, 1989, **89**, 1487.
- 2 G. Faraglia, *J. Organomet. Chem.*, 1969, **20**, 99.
- 3 T. Allman, R. Foel and H. S. Prasad, *J. Organomet. Chem.*, 1979, **166**, 365.
- 4 X. Chen, Y. Yamamoto, K. Akiba, S. Yoshida, M. Yasui and F. Iwasaki, *Tetrahedron Lett.*, 1992, **33**, 6656.
- 5 S. Wallenhauer and K. Seppelt, *Angew. Chem.*, 1994, **106**, 1044.
- 6 (a) D. Naumann and W. Tyrre, *J. Organomet. Chem.*, 1987, **334**, 323; (b) D. Naumann, R. Schlengermann and W. Tyrre, *J. Fluorine Chem.*, 1994, **66**, 79.
- 7 (a) H. J. Frohn and H. Maurer, *J. Fluorine Chem.*, 1986, **34**, 129; (b) A. O. Miller and G. G. Furin, *Russ. Chem. Bull.*, 1994, **43**, 168; *Izv. Akad. Nauk, Ser. Khim.*, 1994, 171.
- 8 A. A. Kolomeitsev, M. Goerg, U. Dieckbreder, E. Lork and G.-V. Roeschenthaler, *Phosphorus, Sulfur, Silicon Relat. Elem.*, 1996, **109–111**, 597.
- 9 K. O. Christe, D. A. Dixon, H. P. A. Mercier, J. C. P. Sanders, G. J. Schrobilgen and W. W. Wilson, *J. Am. Chem. Soc.*, 1994, **116**, 2850.

Communication 9/00308H