

Easy ring expansion and contraction in Pt–Sn bonded metallacycles

Michael C. Janzen, Hilary A. Jenkins, Michael C. Jennings, Louis M. Rendina and Richard J. Puddephatt*


Department of Chemistry, University of Western Ontario, London, Canada N6A 5B7.
E-mail: pudd@uwo.ca

Received 14th April 1999, Accepted 22nd April 1999

The alkyne RCCR, R = CO₂Me, reacts with 5-membered metallacycles [PtMe₂{SnMe₂ESnMe₂E}(bu₂bpy)], E = S, Se or Te, bu₂bpy = 4,4'-di-*tert*-butyl-2,2'-bipyridine, yielding the corresponding 7-membered metallacycles [PtMe₂{SnMe₂ESnMe₂CR=CRE}(bu₂bpy)] which slowly eliminate "Me₂SnE" to give new 5-membered metallacycles [PtMe₂{SnMe₂CR=CRE}(bu₂bpy)]; further reaction of which with excess RCCR gives [PtMe₂(CR=CRH)(CCR)(bu₂bpy)], a complex which contains alkyl, alkenyl and alkynyl functionalities in the same molecule.

A key property of metallacyclic compounds is their ability to undergo easy ring expansion/contraction reactions and these reactions are central to such useful catalytic reactions as alkene or alkyne metathesis, dimerization or trimerization.¹ This article reports that easy ring expansion and contraction can occur in reactions of an alkyne with the metallacycles [PtMe₂(SnMe₂ESnMe₂E)(bu₂bpy)]; bu₂bpy = 2,2'-di-*tert*-butyl-4,4'-bipyridine, E = S, 2; Se, 3; or Te, 4, which are easily prepared by reaction of (Me₂SnE)₃ with [PtMe₂(bu₂bpy)] 1, as shown in Scheme 1.² These appear to be unique examples of such reactions in heteronuclear metallacycles; the closest analogy appears to be the single chalcogen atom abstraction, which converts the 6-membered Pt^{II}ENC(Ph)NE (E = S or Se) ring to a 5-membered ring.³

The electrophilic alkyne dimethyl acetylenedicarboxylate inserts regioselectively into a Sn–E bond of the 5-membered metallacycle 2, 3, or 4 to form the corresponding 7-membered metallacycle 5, 6, or 7. These reactions are complete in about 1 h at room temperature and the products are yellow and air-stable when E = S or Se, but dark brown and air sensitive when E = Te. They were characterized by their NMR spectra⁴ and, for complex 6 by an X-ray structure determination.⁵ The ¹H NMR spectra of 5–7 contain four MeSn, two MePt and two MeO resonances, each corresponding to three protons, and so demonstrate that one equivalent of alkyne has been added. The ¹¹⁹Sn NMR spectrum of 5 contained two resonances, one of which displayed a coupling ¹J(PtSn) = 11 860 Hz, thus showing that the Pt–Sn bond was still present, and both resonances exhibited a coupling ²J(Sn¹–Sn²) = 151 Hz, thus showing that the PtSnMe₂SSnMe₂ unit was still present. Final proof that insertion occurred into the remaining PtE–Sn bond of 2–4 was obtained from the structure determination for complex 6 (Fig. 1).⁵ The conformation of the 7-membered ring leads to relatively short transannular distances Sn2...Se1 = 3.42 Å, Sn1...Se1 = 3.48 Å, perhaps indicating weak secondary bonding between these atoms. The stereochemistry at the C=C bond is *cis* and one CO₂Me group stacks below the bipyridyl ligand. It is interesting that the Sn–E bonds in the precursor molecules (Me₂SnE)₃ are unreactive towards this alkyne, and so the Sn–E bond is activated within the platinum complex. We suggest that the reaction is initiated by nucleophilic attack from a lone pair of electrons of the PtE group on the electrophilic alkyne, and that the nucleophilicity of E is increased by donation of electron density from the *trans* MePt group. The complexes 2–4 are unreactive towards less electrophilic alkynes such as PhCCPh.

Scheme 1 R = CO₂Me.

The complexes 5–7 decompose, over a period of about 8 hours at room temperature in solution in CH₂Cl₂ by elimination of (Me₂SnE)₃ (identified by its NMR spectrum)^{2,3} to form the 5-membered metallacycles 8–10, Scheme 1. The ¹H NMR of complexes 8–10 each contained two MeSn, two MePt and two MeO resonances, and the ¹¹⁹Sn NMR spectra each contain only one resonance with a large coupling due to ¹J(PtSn).⁴ The structure of 9 is shown in Fig. 2.⁵ The 5-membered PtSnC=CSe ring is only slightly distorted from planarity (torsional angle Pt–Se–C=C = –8.6°; Pt–Sn–C=C = 9.3°), in contrast to the twisted conformation adopted by the 7-membered ring in 6 and the envelope conformations of 2–4.^{2,3} In both 6 and 9, the Pt–N distance *trans* to tin is longer than that *trans* to methyl, as a result of the very high *trans* influence of tin.

Complex 9 reacts catalytically with excess (Me₂SnSe)₃ and RCCR to give a mixture of products. When reactions were monitored by ¹H NMR, 9 was shown to remain as the catalyst




Fig. 1 The molecular structure of **6**. Selected bond distances (Å): Pt–N(2) 2.156(3), Pt–N(1) 2.228(4), Pt–C(1) 2.085(5), Pt–C(2) 2.065(5), Pt–Sn(2) 2.5625(4), Sn(2)–Se(2) 2.5671(6), Se(2)–Sn(1) 2.5137(6), Sn(1)–C(3) 2.161(6), C(3)–C(6) 1.323(7), C(6)–Se(1) 1.919(4), Se(1)–Pt 2.5380(5). Bond angles (°): Pt–Sn(2)–Se(2) 114.52(2), Sn(2)–Se(2)–Sn(1) 103.19(2), Se(2)–Sn(1)–C(3) 112.4(1), Sn(1)–C(3)–C(6) 122.2(4), C(3)–C(6)–Se(1) 121.4(4), C(6)–Se(1)–Pt 106.4(1), Se(1)–Pt–Sn(2) 84.12(1).




Fig. 2 The molecular structure of **9**. Selected bond distances (Å): Pt–C(1) 2.062(9), Pt–C(2) 2.091(8), Pt–Sn(1) 2.5578(7), Pt–Se(1) 2.5303(9), Pt–N(1) 2.144(6), Pt–N(2) 2.244(7). Bond angles (°): Se(1)–Pt–Sn(1) 88.60(3), Pt–Sn(1)–C(7) 99.0(2), Sn(1)–C(7)–C(8) 120.6(6), C(7)–C(8)–Se(1) 109.4(5), C(8)–Se(1)–Pt 104.5(2).

“resting state”. The organoselenium complexes were separated chromatographically and identified as a mixture of the known⁶ selenole **11** and the bis(*Z*-alkenyl)selenium compound **12**,^{4,7} but an organotin product, **13**, shown to be present in the reaction mixture by its NMR spectra,⁴ was decomposed on the column and so was not isolated in pure form or structurally characterized. The same organoselenium and organotin compounds were formed stoichiometrically by reaction of **9** with excess alkyne but, in this case, a new organoplatinum complex **14** was also formed as shown in Scheme 2. Complex **14** is stable and fails to react with either (Me₂SnSe)₃ or excess alkyne; it has been characterized by its ¹H and ¹³C NMR spectra and by a structure determination (Fig. 3).^{4,5} One methyl group and the alkenyl group are *trans* to nitrogen donors while the other methyl group and the alkynyl group are mutually *trans*. The formation of the alkenyl groups present in compounds **12** and **14** requires that an H-atom abstraction step must occur and, since the alkenyl proton is still observed in the ¹H NMR when the reaction is carried out in deuterated solvents such as CD₂Cl₂ or C₆D₆, the source of the H-atom in the CR=CRH group must be one of the reagents used. The formation of **14** also requires cleavage of a C–C bond of the alkyne and the fate of the CO₂Me fragment





Fig. 3 The molecular structure of **14**. Selected bond distances (Å): Pt–C(35) 2.01(1), Pt–C(28) 2.07(1), Pt–C(21) 2.09(1), Pt–C(27) 2.10(1), Pt–N(1) 2.134(8), Pt–N(2) 2.114(9).



that is eliminated is unknown. It had been envisioned that the platinum complex **9** might catalyze the reaction of (Me₂SnSe)₃ with RCCR to give organotin metallacycles of the form {(Me₂SnSe)_n(RCCR)_m}, but the actual catalytic reactions are clearly more complex. The nature of the organoselenium products suggests that reaction of **9** with alkyne may be initiated by nucleophilic attack by selenium on the electrophilic alkyne, but the mechanisms of subsequent steps are still to be determined. This work is significant in showing that Sn–E bonds are strongly activated within organoplatinum metallacycles, that easy ring expansion and contraction can occur in reactions with an electrophilic alkyne, and that catalytic reactions may be developed.

Acknowledgements

We thank the NSERC (Canada) for financial support.

Notes and references

- See for example: P. W. Jennings and L. L. Johnson, *Chem. Rev.*, 1994, **94**, 2241.
- L. M. Rendina, J. J. Vittal and R. J. Puddephatt, *Organometallics*, 1996, **15**, 1749; M. C. Janzen, H. A. Jenkins, L. M. Rendina, J. J. Vittal and R. J. Puddephatt, *Inorg. Chem.*, in the press.

- 3 N. Feeder, R. J. Less, J. M. Rawson and J. N. B. Smith, *J. Chem. Soc., Dalton Trans.*, 1998, 4091.
- 4 Selected spectroscopic data: NMR in CD₂Cl₂ (refs. SiMe₄, Me₄Sn, K₂PtCl₄). **5**: $\delta(^1\text{H})$ 3.41 [s, 3H, β -CO₂Me]; 2.80 [s, 3H, α -CO₂Me]; 0.86 [s, 3H, $^2J(\text{PtH}) = 59$ Hz, Pt–Me]; 0.76 [s, 3H, $^2J(\text{SnH}) = 46$ Hz, Pt–Sn–Me^a]; 0.59 [s, 3H, $^2J(\text{SnH}) = 52$ Hz, Sn–Me^a]; 0.39 [s, 3H, $^3J(\text{PtH}) = 4$ Hz, $^2J(\text{SnH}) = 46$ Hz, Pt–Sn–Me^b]; 0.31 [s, 3H, $^2J(\text{SnH}) = 55$ Hz, Sn–Me^b]; 0.11 [s, 3H, $^2J(\text{PtH}) = 62$ Hz, Pt–Me]; $\delta(^{119}\text{Sn})$ 6.81 [$^2J(\text{SnSn}) = 151$ Hz, Pt–S–C=C–Sn]; –81.45 [$^1J(\text{PtSn}) = 11$ 860 Hz, Pt–Sn]; $\delta(^{195}\text{Pt})$ –1770 [$^1J(\text{SnPt}) = 11$ 860 Hz]. **6**: $\delta(^1\text{H})$ 3.43 [s, 3H, β -CO₂Me]; 2.83 [s, 3H, α -CO₂Me]; 0.87 [s, 3H, Pt–Sn–Me]; 0.81 [s, 3H, $^2J(\text{PtH}) = 59$ Hz, Pt–Me]; 0.70 [s, 3H, $^2J(\text{SnH}) = 51$ Hz, Sn–Me^a]; 0.46 [s, 3H, $^2J(\text{SnH}) = 44$ Hz, Pt–Sn–Me^b]; 0.34 [s, 3H, $^2J(\text{SnH}) = 53$ Hz, Sn–Me^b]; 0.20 [s, 3H, $^2J(\text{PtH}) = 62$ Hz, Pt–Me]. **7**: $\delta(^1\text{H})$ 3.46 [s, 3H, β -CO₂Me]; 2.88 [s, 3H, α -CO₂Me]; 0.98 [s, 3H, $^2J(\text{SnH}) = 56$ Hz, Sn–Me]; 0.92 [s, 3H, Sn–Me]; 0.74 [s, 3H, $^2J(\text{PtH}) = 60$ Hz, Pt–Me]; 0.55 [s, 3H, $^2J(\text{SnH}) = 45$ Hz, Sn–Me]; 0.41 [s, 3H, $^2J(\text{SnH}) = 52$ Hz, Sn–Me]; 0.30 [s, 3H, $^2J(\text{PtH}) = 61$ Hz, Pt–Me]. **8**: $\delta(^1\text{H})$ 3.65 [s, 3H, CO₂Me]; 3.61 [s, 3H, CO₂Me]; 0.94 [s, 3H, $^2J(\text{PtH}) = 61$ Hz, $^3J(\text{SnH}) = 6$ Hz, Pt–Me]; 0.60 [s, 3H, $^2J(\text{SnH}) = 49$ Hz, $^3J(\text{PtH}) = 6$ Hz, Sn–Me]; 0.34 [s, 3H, $^2J(\text{SnH}) = 55$ Hz, $^3J(\text{PtH}) = 4$ Hz, Sn–Me]; 0.29 [s, 3H, $^2J(\text{PtH}) = 57$ Hz, Pt–Me]; $\delta(^{119}\text{Sn})$ –2.1 [$^1J(\text{SnPt}) = 9904$ Hz]. **9**: $\delta(^1\text{H})$ 3.66 [s, 3H, CO₂Me]; 3.62 [s, 3H, CO₂Me]; 1.03 [s, 3H, $^2J(\text{PtH}) = 60.3$ Hz, $^3J(\text{SnH}) = 5$ Hz, Pt–Me]; 0.58 [s, 3H, $^2J(\text{SnH}) = 48$ Hz, $^3J(\text{PtH}) = 5$ Hz, Sn–Me]; 0.34 [s, 3H, $^2J(\text{SnH}) = 54$ Hz, $^3J(\text{PtH}) = 4$ Hz, Sn–Me]; 0.32 [s, 3H, $^2J(\text{PtH}) = 57$ Hz, Pt–Me]; $\delta(^{119}\text{Sn})$ –5.8 [$^1J(\text{SnPt}) = 10$ 031 Hz]. **10**: $\delta(^1\text{H})$ 3.65 [s, 3H, CO₂Me]; 3.62 [s, 3H, CO₂Me]; 1.03 [s, 3H, $^2J(\text{PtH}) = 61$ Hz, Pt–Me]; 0.55 [s, 3H, $^2J(\text{SnH}) = 48$ Hz, $^3J(\text{PtH}) = 5$ Hz, Sn–Me]; 0.34 [s, 3H, $^2J(\text{PtH}) = 55$ Hz, Pt–Me]; 0.32 [s, 3H, $^2J(\text{SnH}) = 53$ Hz, $^3J(\text{PtH}) = 4$ Hz, SnMe]. **12**: $\delta(^1\text{H})$ 6.46 [s, 2H, $^3J(\text{SeH}) = 5$ Hz, Se–C=C–H]; 3.83 [s, 6H, CO₂Me]; 3.75 [s, 6H, CO₂Me]; MS: $m/z = 366$. **13**: $\delta(^1\text{H})$ 0.54 [s, $^2J(\text{SnH}) = 66$ Hz, MeSn]; no other proton resonances. **14**: $\delta(^1\text{H})$ 6.54 [s, 1H, $^3J(\text{PtH}) = 80$ Hz, Pt–C=C–H]; 3.79 [s, 3H, CO₂Me]; 3.68 [s, 3H, CO₂Me]; 3.51 [s, 3H, CO₂Me]; 1.30 [s, 3H, $^3J(\text{PtH}) = 69$ Hz, Pt–Me]; –0.04 [s, 3H, $^3J(\text{PtH}) = 51$ Hz, Pt–Me]; $\delta(^{13}\text{C})$ –5.38 [$^1J(\text{PtC}) = 575$ Hz, PtMe *trans* to N]; 3.88 [$^1J(\text{PtC}) = 477$ Hz, PtMe *trans* to C]; 119.26 [$^1J(\text{PtC}) = 780$ Hz, PtC(alkenyl)]; 142.17 [$^1J(\text{PtC}) = 901$ Hz, PtC(alkynyl)].
- 5 Crystal data: **6**: C₃₀H₄₈N₂O₄PtSe₂Sn₂, $M = 1091.09$, triclinic, space group $P\bar{1}$, $a = 10.8110(3)$, $b = 12.5192(2)$, $c = 18.975(3)$ Å, $\alpha = 78.895(1)$, $\beta = 70.862(1)$, $\gamma = 86.721(1)^\circ$, $V = 1889.60(7)$ Å³, $D_c = 1.918$ g cm^{–3}, $Z = 2$, $T = 294$ K, $R = 0.0267$, $R_w = 0.0621$, $\mu = 6.96$ mm^{–1}, 6616 independent reflections. **9**: C₂₈H₄₂N₂O₄PtSeSn, $M = 863.38$, orthorhombic, $Pca2_1$, $a = 30.899(6)$, $b = 12.156(2)$, $c = 17.311(4)$ Å, $V = 6502(2)$ Å³, $D_c = 1.764$ g cm^{–3}, $Z = 8$, $T = 150(2)$ K, $R = 0.0293$, $R_w = 0.0630$, $\mu = 6.22$ mm^{–1}, 6852 independent reflections. **14**: C₃₀H₄₀N₂O₆Pt, $M = 719.73$, monoclinic, $P2(1)/n$, $a = 10.2375(6)$, $b = 16.035(1)$, $c = 18.889(1)$ Å, $\beta = 102.624(3)^\circ$, $V = 3025.9(3)$ Å³, $D_c = 1.580$ g cm^{–3}, $Z = 4$, $T = 150(2)$ K; $R = 0.039$, $R_w = 0.0959$, $\mu = 4.68$ mm^{–1}, 3057 independent reflections. CCDC reference number 186/1434. See <http://www.rsc.org/suppdata/dt/1999/1713/> for crystallographic files in .cif format.
- 6 M. R. J. Dorrity, J. F. Malone, C. P. Morley and R. R. Vaughan, *Phosphorus, Sulfur Silicon Relat. Elem.*, 1992, **68**, 37.
- 7 The Z stereochemistry for **12** is defined from the magnitude of $^3J(\text{SeH}) = 5$ Hz; the coupling would be *ca.* 20 Hz in the *E*-isomer. C. Paulmier, *Selenium Reagents and Intermediates in Organic Synthesis*, Pergamon, Toronto, 1986.

Communication 9/02991E