

Reaction of thiones with dihalogens; comparison of the solid state structures of 4,5-bis(methylsulfanyl)-1,3-dithiole-2-thione–diiodine, –dibromine and –iodine monobromide

Neil Bricklebank,^{*a} Peter J. Skabara,^{*a} David E. Hibbs,^b Michael B. Hursthouse^{†b} and K. M. Abdul Malik^b

^a Division of Chemistry and Materials Research Institute, Sheffield Hallam University, Sheffield, UK S1 1WB. E-mail: N.Bricklebank@shu.ac.uk; P.J.Skabara@shu.ac.uk

^b Department of Chemistry, Cardiff University, P.O. Box 912, Park Place, Cardiff, UK CF1 3TB

Received 19th April 1999, Accepted 30th June 1999

Reaction of 4,5-bis(methylsulfanyl)-1,3-dithiole-2-thione **1** with diiodine or iodine monobromide in CH₂Cl₂ resulted in the formation of molecular charge-transfer complexes **1**·I₂ and **1**·IBr respectively. Both complexes have been characterised crystallographically and contain a linear S–I–X (X = I or Br) moiety with the sulfur adopting a tetrahedral geometry taking into account the stereochemically active lone pairs. The S–I [2.716(3)] and I–I [2.808(3) Å] bond lengths in **1**·I₂ are similar to those reported for diiodine complexes of related thione donors. The adduct **1**·IBr is the first crystallographically characterised thione–iodine monobromide charge-transfer complex. The S–I distance [2.589(2) Å] is shorter than that in **1**·I₂, consistent with IBr being a stronger acceptor than I₂. The I–Br distance [2.7138(11) Å] is lengthened with respect to that in unco-ordinated IBr, but within bonding distance when compared to the sum of the van der Waals radii for iodine and bromine (3.75 Å). Treatment of **1** with dibromine under identical conditions resulted in the formation of the adduct **1**·Br₂ and the dithioliium salt [C₅H₆S₄Br][Br₃][−]· $\frac{1}{2}$ Br₂ **2**. Treatment of **1** with Br₂ in toluene led to the isolation of **1**·Br₂ only. The crystal structure of **1**·Br₂ shows the compound to contain a linear Br–S–Br moiety with the sulfur in a T-shaped or Ψ-trigonal bipyramidal environment (taking into account the stereochemically active lone pairs). The structure of **2** reveals a three component system consisting of the [C₅H₆S₄Br]⁺ cation, the [Br₃][−] anion and a molecule of Br₂ in a 2:2:1 ratio. These components are held in the lattice by a series of weak intermolecular interactions which link the tribromide ions and dibromine molecules into zigzag chains.

Introduction

The adducts formed through the interaction of dihalogens with triorgano Group 15^{1,2} and diorgano Group 16^{3–19} molecules play pivotal roles in a variety of chemical scenarios, ranging from organic and inorganic synthesis to biological processes and materials chemistry. Therefore, detailed knowledge of the structural nature of these adducts is essential if we are fully to understand their efficacy and the mechanisms by which they react.


The donor–dihalogen interaction can proceed *via* a number of pathways, resulting in the formation of a variety of products, for example: a simple molecular ‘donor–acceptor’ or ‘charge-transfer’ complex;^{1–14} homolytic cleavage of the dihalogen bond with the concomitant formation of two donor–halide bonds;^{1,15,16} the dihalogen may undergo heterolytic cleavage with the formation of halogen(I) salts;¹⁷ in the presence of an excess of dihalogen, polyhalide salts may result;^{3e,9,10} the adduct formed *in situ* may experience further spontaneous chemical reaction resulting in the formation of new products which are not simple adducts. The past decade has witnessed a massive increase in the quantity of available structural data on all the above types of product (led by seminal contributions from the group of McAuliffe¹ and others), which show that the pathway followed, and hence the product formed, is influenced not only by the donor and the dihalogen acceptor, but also by external factors such as the molar ratio of the reactants, the solvent and the reaction conditions.

Thiones (thiocarbonyls) present a large class of particularly useful donors, the dihalogen adducts of which have received attention in areas as diverse as the treatment of hyperthyroidism¹⁸ and molecular electronics.¹⁹ The structural nature of the diiodine adducts is, in general, well established.^{3–11} However, there have been fewer studies into the interaction of thiones with the lighter dihalogens or interhalogens and only limited structural data are available.¹⁵ Furthermore, few comparative studies have been carried out, in contrast to the situation for other classes of donor. Here we describe the structural nature of the adducts formed from the interaction of diiodine, iodine monobromide and dibromine with 4,5-bis(methylsulfanyl)-1,3-dithiole-2-thione **1**,²⁰ a heterocyclic sulfur rich thione which has been used as a building-block in tetrathiafulvalene-type materials chemistry.²¹ This work represents the first comparative study of the interaction of a thione with these acceptors.

Results and discussion

The reactions of compound **1** with X₂ (X₂ = I₂, IBr or Br₂) are summarised in Scheme 1. Addition of I₂ or IBr to an equimolar quantity of **1** dissolved in refluxing dichloromethane resulted in the formation of an orange-red solution which deposited large brown crystals of **1**·I₂ (85% yield) or **1**·IBr, in quantitative yield, when cooled to −5 °C overnight. However, treatment of a solution of **1** in refluxing dichloromethane with Br₂ produced, after cooling, a brown microcrystalline powder, **1**·Br₂, in 50% yield. The brown solid was removed by filtration and the dichloromethane of the filtrate allowed to evaporate, yielding a purple oil, from which a small crop of purple crystals was

[†] Present address: Department of Chemistry, University of Southampton, Highfield, Southampton, UK SO17 1BJ.



Scheme 1

deposited (10% yield). These purple crystals, which emitted pungent fumes of dibromine, were identified as dithiolium salt **2** by X-ray crystallography. The salt **2** was found to be hygroscopic and decomposed slowly upon standing. The reaction of **1** and Br_2 was repeated using toluene as the solvent; on cooling, $\mathbf{1}\cdot\text{Br}_2$ was obtained as an orange-brown crystalline solid in 54% yield. It is worth noting that the presence of **2** was not detected in this reaction.

We have investigated the three dihalogen adducts of compound **1** using a combination of low frequency FT-Raman and FTIR spectroscopy. In particular, Raman spectroscopy provides a sensitive probe for the elucidation of charge-transfer interactions between donors and dihalogens, by means of a shift in the $\nu(\text{X-X})$ vibrations in the complex, with respect to those in the free acceptors.¹¹ The Raman spectrum of $\mathbf{1}\cdot\text{I}_2$ contains a strong band at 135 cm^{-1} due to $\nu(\text{I-I})$ in the charge-transfer complex. As expected, the value is shifted to a lower frequency from a value of 180 cm^{-1} seen for solid I_2 , consistent with the decrease in the I-I bond order when diiodine interacts with a donor. The position of the $\nu(\text{I-I})$ band of $\mathbf{1}\cdot\text{I}_2$, and the magnitude of the shift of $\nu(\text{I-I})$ from free I_2 (45 cm^{-1}), concur with those for I_2 complexes of other thione donors.^{6,7,11} Similarly, $\mathbf{1}\cdot\text{IBr}$ exhibits a $\nu(\text{IBr})$ band at 144 cm^{-1} , which is shifted considerably from the value of 262 cm^{-1} of solid IBr . The magnitude of the shift for $\mathbf{1}\cdot\text{IBr}$ (118 cm^{-1}) is much greater than that for $\mathbf{1}\cdot\text{I}_2$ (45 cm^{-1}), since IBr is a better acceptor than I_2 . The Raman spectrum of $\mathbf{1}\cdot\text{Br}_2$ contains a strong band at 162 cm^{-1} , attributable to symmetric Br-S-Br stretching.

The FTIR spectra of adducts $\mathbf{1}\cdot\text{I}_2$, $\mathbf{1}\cdot\text{IBr}$ and $\mathbf{1}\cdot\text{Br}_2$ are very similar to that of the unco-ordinated donor **1**. The principal peak in the IR spectrum of **1** is a band at 1067 cm^{-1} , which can be attributed to an out-of-phase combination of the $\nu(\text{C-S})$ and $\nu(\text{SCS})$ stretching vibrations, coupled with the $\delta(\text{CSS})$ deformation.^{6,7} As expected, the co-ordination of a dihalogen to **1** results in a shift to a lower frequency for this band; thus, strong peaks are observed at 1017 , 1030 and 991 cm^{-1} for $\mathbf{1}\cdot\text{I}_2$, $\mathbf{1}\cdot\text{IBr}$ and $\mathbf{1}\cdot\text{Br}_2$, respectively.

The solid state structures of compounds $\mathbf{1}\cdot\text{I}_2$, $\mathbf{1}\cdot\text{IBr}$ and $\mathbf{1}\cdot\text{Br}_2$ are shown in Figs. 1, 2 and 3 respectively. Selected geometry parameters for the three compounds are given in Table 1. The $\text{S}\cdots\text{X}$ and X-X distances for some related dihalogen complexes are collated in Table 2 for comparison. Compounds $\mathbf{1}\cdot\text{I}_2$ and $\mathbf{1}\cdot\text{IBr}$ are molecular "charge-transfer" adducts containing linear $\text{S}\cdots\text{I-X}$ linkages. The $\text{S}\cdots\text{I-I}$ and $\text{S}\cdots\text{I-Br}$ angles are $177.46(7)$ and $175.63(5)^\circ$ respectively. The molecular structure of "donor-acceptor" or "charge-transfer" complexes is relatively well understood; transfer of electron density from a lone pair of the donor to the σ^* antibonding orbital of the





Fig. 1 Solid state structure of adduct $\mathbf{1}\cdot\text{I}_2$ showing the atom numbering scheme. The thermal ellipsoids are drawn at 50% probability level.

dihalogen leads to the formation of a donor-halogen bond with the concomitant elongation of the X-X bond. This is illustrated for thiones in Scheme 2. The sulfur atom of the thione (thio-carbonyl) donor is sp^2 hybridised with a double bond to the adjacent carbon atom and should be a reasonably good donor of electrons towards dihalogens. The ability of thiones to delocalise the positive charge onto the carbon atom of the C-S bond provides an additional factor which helps to stabilise the S-X-X interaction (Scheme 2). Thus, thiones should form



Scheme 2

stronger S-X bonds than, for example, the sp^3 hybridised sulfur in a thioether, which does not have the ability to delocalise the positive charge. Moreover, the complexes derived from the donor **1** are further stabilised by the resonance contribution of the 1,3-dithiole sulfur atoms (Scheme 2), such that the positive charge is delocalised over the $\text{S}(1)\text{-C}(2)\text{-S}(3)$ fragment of the ring, forming a 6π pseudo-aromatic system. All three dihalogen adducts reported here provide good evidence for this. Ligand **1** contains sulfur atoms in three distinct environments; the thione sulfur (sp^2 hybridised), the two dithiole sulfurs and the two thioether sulfurs (all of which are sp^3 hybridised). In $\mathbf{1}\cdot\text{I}_2$, $\mathbf{1}\cdot\text{IBr}$ and $\mathbf{1}\cdot\text{Br}_2$ the halogen atoms only interact with the thione sulfur. No interaction is observed between halogen atoms and the thioether or dithiole sulfurs. The thione C-S bonds in $\mathbf{1}\cdot\text{I}_2$ and $\mathbf{1}\cdot\text{IBr}$ are longer than a typical carbon-sulfur double bond (1.60 \AA)²² but shorter than a carbon-sulfur single bond (1.82 \AA)²² consistent with the delocalisation of some positive charge on the C-S bond but retaining significant double bond character. The $\text{S}\cdots\text{I}$ distances in $\mathbf{1}\cdot\text{I}_2$ and $\mathbf{1}\cdot\text{IBr}$ are $2.716(3)$ and $2.589(2)\text{ \AA}$ respectively; these values are longer than those expected on the basis of the sum of the covalent radii of sulfur




Fig. 2 Solid state structure of adduct **1·IBr** showing the atom numbering scheme. The thermal ellipsoids are drawn at 50% probability level.




Fig. 3 Solid state structure of adduct **1·Br₂** showing the atom numbering scheme. The thermal ellipsoids are drawn at 50% probability level.

and iodine ($1.02 + 1.33 = 2.35 \text{ \AA}$),²² but are also much shorter than the corresponding sum of the van der Waals radii ($1.80 + 1.95 = 3.75 \text{ \AA}$),²² and are therefore indicative of definite S–I bonds. The S–I distance in **1·IBr** is shorter than that in **1·I₂** reflecting the greater acceptor strength of IBr over I₂. No other structural data have been reported for a thione–IBr donor–acceptor complex, though data are available for IBr complexes of several other sulfur donors (Table 2). The S–I distance in **1·IBr** is shorter and the I–Br distance longer than the equivalent distances in IBr complexes of other sulfur donors.^{12–14} This is not unexpected, since these complexes all contain sp³ hybridised sulfurs or, in the case of Ph₃PS·IBr, an sp² hybridised sulfur atom bonded to a triphenylphosphine group, which is less effective at stabilising the accepted positive charge than is the heterocycle **1**. Consistent with this, the $\nu(\text{I–Br})$ band in the Raman spectrum of **1·IBr** (144 cm^{-1}) occurs at a much lower frequency than the equivalent bands of [14]aneS₄·2IBr (184 cm^{-1}),¹² [16]aneS₄·4IBr ($184, 164 \text{ cm}^{-1}$),¹² [18]aneS₆·2IBr (182 cm^{-1})¹² and Ph₃PS·IBr ($206, 164 \text{ cm}^{-1}$).¹³ The I–I and I–Br bonds in **1·I₂** and **1·IBr**, respectively, are lengthened compared to the bonds in unco-ordinated I₂ [$d_{\text{I–I}} = 2.715(6) \text{ \AA}$


Table 1 Selected bond lengths (Å) and angles (°) in adducts **1·I₂**, **1·IBr** and **1·Br₂**

	1·I₂	1·IBr	1·Br₂
I(1)–I(2)/Br(1)	2.8083(14)	2.7138(11)	
S(5)–I(1)	2.716(3)	2.589(2)	
S(5)–Br(1)			2.496(4)
S(5)–Br(2)			2.437(4)
S(5)–C(1)	1.666(11)	1.688(8)	1.75(2)
S(1)–C(1)	1.663(13)	1.693(7)	1.71(2)
S(2)–C(1)	1.701(12)	1.712(8)	1.680(12)
S(1)–C(2)	1.703(11)	1.728(8)	1.739(14)
S(2)–C(3)	1.742(11)	1.730(8)	1.75(2)
C(2)–C(3)	1.31(2)	1.348(11)	1.34(2)
S(5)–I(1)–I(2)/Br(1)	177.46(7)	175.63(5)	
Br(1)–S(5)–Br(2)			171.6(2)
C(1)–S(5)–I(1)	101.9(4)	105.5(3)	
C(1)–S(5)–Br(1)			92.7(5)
C(1)–S(5)–Br(2)			95.7(5)
S(1)–C(1)–S(5)	125.0(7)	126.1(5)	120.7(7)
S(2)–C(1)–S(5)	119.9(7)	119.6(4)	123.9(9)
S(1)–C(1)–S(2)	115.1(7)	114.3(5)	115.4(8)

in the solid state, $2.667(2) \text{ \AA}$ ²⁴ in the gas phase] and IBr [$d_{\text{I–Br}} = 2.521(4) \text{ \AA}$ ²⁵ in the solid state, $2.470(5) \text{ \AA}$ ²⁶ in the gas phase], but are well within bonding distance when compared with the van der Waals radii of I₂ (3.40 \AA)²² and IBr (3.75 \AA).²² As can be seen from Figs. 1 and 2, the orientation of the S···I–X units in these two compounds is similar, whereas those of the S–CH₃ substituents are quite different.

The S–I distance in thione–diiodine complexes (Table 2) lies in the range 2.61 to 2.92 Å [shorter distances are observed in complexes where the diiodine is interacting with a second molecule of diiodine, e.g. ethylenethiourea–diiodine(1/2),¹⁰ $d_{\text{S–I}} = 2.487(3) \text{ \AA}$]. The S–I distance in **1·I₂** falls within this range. Not surprisingly, the S–I distance in **1·I₂** is virtually identical to those determined for the diiodine complexes of other sulfur-rich thiones, i.e. **3·I₂**, **4·I₂** and **5·I₂**. Since there is usually a reciprocal relationship between the D–I and I–I bond lengths in D–I–I complexes one might expect the I–I distance in **1·I₂** to be similar to those in **3·I₂**, **4·I₂** and **5·I₂**. However, it is significantly shorter, indicating the absence of any secondary intermolecular interactions involving the S–I–I unit in **1·I₂**. In **1·I₂** the molecules exist as discrete units within the crystal and pack parallel to (110) as shown in Fig. 4. The shortest contacts involving the neighbouring molecules are: 3.549 \AA [S(3)···C(1) ($0.5 - x, 0.5 + y, -0.5 - z$)], 3.653 \AA [S(3)···S(5) ($0.5 - x, 0.5 + y, -0.5 - z$)] and 3.660 \AA [S(5)···I(2) ($0.5 + x, 0.5 - y, -0.5 - z$)], which indicate the absence of any significant stacking interactions. In **1·IBr** the molecules are packed parallel to (010) as shown in Fig. 5, but in this case several intermolecular contacts are identified which are significant and indicative of stacking interactions: 3.357 \AA [S(3)···S(3) ($1 - x, 1 - y, 2 - z$)], 3.379 \AA [S(2)···Br(1) ($x, -1 + y, z$)] and 3.445 \AA [S(5)···Br(1) ($x, -1 + y, z$)].

The structure of adduct **1·Br₂** is very different from those of **1·I₂** and **1·IBr**, since the Br–Br bond of dibromine is completely broken and the compound has a three-co-ordinated sulfur atom and two terminal S–Br bonds. The geometry about the thione-sulfur atom is T-shaped or Ψ-trigonal bipyramidal taking into account the stereochemically active lone pairs (VSEPR model).²² This contrasts neatly with **1·I₂** and **1·IBr** in which the thione-sulfur has a bent or approximately tetrahedral geometry. The Br(1)–S(5)–Br(2) angle at $171.6(2)^\circ$ is almost linear. The S(5)–Br distances, $2.437(4)$ and $2.496(4) \text{ \AA}$, are of unequal length and considerably longer than the appropriate covalent radii sum ($1.02 + 1.14 = 2.16 \text{ \AA}$),²² but well within bonding distance when compared with the sum of the van der Waals radii of bromine and sulfur ($1.8 + 1.8 = 3.6 \text{ \AA}$).²² Similar asymmetry has been observed in the related compound **6·Br₂** (Table 2) and

Table 2 Geometrical details for S–I–X and X–S–X moieties in dihalogen adducts of sulfur donor ligands

Thione-diiodine complexes	S–I	I–I	S–I–I	Ref.
6 ·I ₂	2.616(6)	2.967(12)	175.71(1)	9
1,1'-Ethylene-bis(3-methyl-4-imidazoline-2-thione)·2I ₂	2.607(6)	2.984(12)	177.11(4)	
1,5-Diphenylthiocarbazonone·I ₂	2.642(3)	2.903(2)	177.0(1)	5(a)
1,1'-Methylene-bis(3-methyl-4-imidazoline-2-thione)·2I ₂	2.664(3)	2.918(1)	178.4(1)	10
<i>N</i> -Methylthiocaprolactam·I ₂	2.683(2)	2.897(1)	175.71(3)	5(a)
4 ·I ₂	2.688(2)	2.880(1)	176.21(4)	5(d)
3 ·I ₂	2.715(4)	2.823(2)	176.5(1)	6
1 ·I ₂	2.715(5)	2.843(2)	176.7(1)	7
5 ·I ₂	2.716(3)	2.8083(14)		<i>a</i>
5,5-Dimethylimidazolidine-2,4-dithione·2I ₂	2.719(6)	2.841(2)		8
	2.737(1)	2.849(1)	177.94(4)	11
	2.844(1)	2.767(1)	173.74(4)	
5,5-Dimethylimidazolidine-2,4-dithione·I ₂	2.748(1)	2.817(1)	176.89(2)	11
5,5-Dimethyl-2-thioxoimidazolidine-4-one·I ₂	2.737(1)	2.802(1)	176.14(2)	11
4,5,6,7-Tetrathiocino[1,2,- <i>b</i> :3,4- <i>b'</i>]-diimidazolyl-1,3,8,10-tetraethyl-2,9-dithiole·2I ₂	2.775(4)	2.822(2)	174.8(1)	5(b)
2(Morpholinothiocarbonyl)·2I ₂	2.920(6)	2.751(2)	175.5(1)	5(c)
	2.789(6)	2.805(3)	174.7(1)	
	2.892(6)	2.738(2)	173.9(2)	
	2.801(6)	2.787(3)	176.5(2)	
Sulfur donor–iodine monobromide complexes	S–I	I–Br	S–I–Br	
1 ·IBr	2.589(2)	2.7138(11)		<i>a</i>
[14]aneS ₄ ·2IBr	2.678(1)	2.654(2)	175.53(4)	12
[16]aneS ₄ ·4IBr	2.618(2)	2.7049(11)	177.65(5)	12
	2.687(2)	2.6445(12)	177.57(5)	
[18]aneS ₆ ·2IBr	2.619(3)	2.695(2)	175.00(6)	12
Ph ₃ PS·IBr	2.665(1)	2.668(1)	175.08(2)	13
Dithiacyclohexane·2IBr	2.687(2)	2.646(1)	178.2(4)	14
Thione-dibromine adducts	S–Br		Br–S–Br	–
1 ·Br ₂	2.496(4)		171.6(2)	<i>a</i>
	2.437(4)			
6 ·Br ₂	2.520			15
	2.477			

^a This work.

in the dibromine complexes of several selenium donors.^{16,27,28} In the latter cases the differences in the Se–Br bond lengths have been attributed to the existence of weak Se···Br²⁷ or Br···Br²⁸ interactions within the crystal lattice. The packing of molecules in **1**·Br₂ (Fig. 6) is similar to that of **1**·IBr, with interlayer distances of 3.675 [S(1)···Br(1) (–*x*, 1 – *y*, –*z*)], 3.685 [S(1)···Br(1) (1 – *x*, 1 – *y*, –*z*)], 3.683 [S(2)···Br(2) (–*x*, 1 – *y*, –*z*)], 3.514 [S(5)···C(3) (1 – *x*, 1 – *y*, –*z*)] and 3.357 Å [S(1)···C(5) (–*x*, 1 – *y*, –*z*)]. The thiocarbonyl bond in **1**·Br₂ is significantly longer than those in **1**·I₂ or **1**·IBr and is closer to that expected for a single C–S bond. We have calculated the C–S bond orders in the three complexes (using the method of Husebye *et al.*);⁴ the values are 1.515, 1.366 and 1.05 for **1**·I₂, **1**·IBr and **1**·Br₂ respectively. Thus, both **1**·I₂ and **1**·IBr retain significant multiple bond character whereas **1**·Br₂ shows very little.

The C₃S₂ heterocycles in all three complexes are planar, the maximum deviations of any atom from the respective mean

planes being 0.019(5), 0.005(5) and 0.019(5) Å for **1**·I₂, **1**·IBr and **1**·Br₂, respectively. The S(5) atom is coplanar with the ring plane in **1**·I₂ [deviation = 0.001(10) Å] but is significantly out of the respective planes in **1**·IBr [0.093(15) Å] and **1**·Br₂ [0.051(15) Å]. The S(5)–I(1)–X (X = I or Br) moieties in **1**·I₂ and **1**·IBr are coplanar with their respective C₃S₂ heterocycles as shown by the I(1)–S(5)–C(1)–S(1) torsion angles of –1.5(9)° in **1**·I₂ and 6.5(6)° in **1**·IBr. The Br(2)–S(5)–C(1)–S(1) torsion angle in **1**·Br₂ is 1.5(7)°, which shows that the SBr₂ moiety in this molecule is also coplanar with the associated C₃S₂ ring.

The formation of dithiolylium salt **2**, together with **1**·Br₂, on addition of Br₂ to compound **1** in dichloromethane was totally unexpected, although the analogous bromine salt **7** has been prepared by the reaction of dibromine with *N*-methyl-1,3-thiazolidine-2(3*H*)-selone.^{16,27} Additionally, previous workers²⁹ have reported the isolation of the 2-chloro-1,3-dithiolylium salt **8**, which was obtained *via* the reaction of **1** and phosphorus pentachloride. The formation of **2** proceeds in dichloromethane




Fig. 4 Unit-cell contents of adduct $1 \cdot I_2$ viewed along the b axis.




Fig. 6 Unit-cell contents of adduct $1 \cdot Br_2$ viewed along the b axis.




Fig. 5 Unit-cell contents of adduct $1 \cdot IBr$ viewed along the b axis.

but not in toluene, suggesting that the polarity of the solvent and hence the polarisation of the intermediates/reagents is an important factor in the reaction mechanism.





Fig. 7 Unit-cell contents of compound **2** viewed along the a axis. The thermal ellipsoids are drawn at 50% probability level. The atom numbering is shown for the asymmetric unit.

The structure of compound **2** is shown in Fig. 7. Selected bond lengths and angles are listed in Table 3. The compound consists of a dithiolylium cation and a tribromide anion together with a molecule of dibromine occluded in the lattice, akin to a solvent of crystallisation. The ratio of the above species in this compound is 2:2:1. The tribromide anion is virtually linear [$178.14(5)^\circ$], as expected for a trihalide anion,³⁰ and lies sandwiched between two dithiolylium cations. The Br–Br bond lengths are unsymmetrical, Br(3)–Br(4) [$2.555(2) \text{ \AA}$], Br(2)–Br(3) [$2.512(2) \text{ \AA}$], but are similar to those reported in other salts containing this ion.³⁰ The Br–Br bond of the

Table 3 Selected bond lengths (Å) and angles (°) in compound **2**

Br(2)–Br(3)	2.512(2)	S(2)–C(3)	1.711(10)
Br(3)–Br(4)	2.555(2)	S(3)–C(2)	1.743(10)
Br(5)–Br(5) ^a	2.329(2)	S(3)–C(4)	1.776(10)
Br(1)–C(1)	1.865(9)	S(4)–C(3)	1.739(10)
S(1)–C(1)	1.648(11)	S(4)–C(5)	1.773(10)
S(1)–C(2)	1.719(10)	C(2)–C(3)	1.36(2)
S(2)–C(1)	1.668(11)		
Br(4)⋯Br(5)	3.214	Br(2)⋯Br(2) ^b	3.439
Br(2)–Br(3)–Br(4)	178.14(5)	C(3)–S(4)–C(5)	100.8(5)
C(1)–S(1)–C(2)	95.9(5)	S(1)–C(1)–S(2)	117.7(6)
C(1)–S(2)–C(3)	95.4(5)	S(1)–C(1)–Br(1)	121.5(6)
C(2)–S(3)–C(4)	102.0(5)	S(2)–C(1)–Br(1)	120.7(6)
Br(3)–Br(4)–Br(5)	117.69	Br(4)–Br(5)–Br(5a)	172.37

Symmetry operator: ^a 1 – x, 1 – y, 1 – z. ^b –x, 2 – y, –z.



occluded Br₂ molecule [2.329(2) Å] is considerably shorter than the Br–Br distances in the [Br₃][–] ion, but somewhat longer than that in crystalline Br₂ [2.27 Å].³¹ There are several short contacts involving the [Br₃][–] ion and of the Br₂ molecule with the dithiolylium cation, the most important of which are: 3.525 [Br(3)⋯C(1)], 3.196 [Br(1)⋯Br(3) (–1 + x, y, z)], 3.643 [Br(2)⋯S(1) (x, 1 + y, z)], 3.506 Å [Br(2)⋯S(3) (–1 + x, 1 + y, z)], 3.627 [Br(3)⋯C(2) (x, 1 + y, z)], 3.557 [Br(3)⋯C(3) (x, 1 + y, z)], 3.668 [Br(4)⋯S(4) (–1 + x, 1 + y, z)] and 3.501 Å [Br(4)⋯C(5) (–x, 1 – y, 1 – z)]. All these weak interactions hold the three different chemical species in the crystal. However, the most important are weak interactions between the [Br₃][–] ions and the Br₂ molecule [Br(4)⋯Br(5) 3.214 Å] and between two adjacent [Br₃][–] anions [Br(2)⋯Br(2) 3.439 Å], which link the tribromide ions and the dibromine molecules into ‘zigzag’ chains. Polybromide ions, other than the tribromide ion, are rare^{30,32} and can generally be considered as an assemblage of tribromide ions and dibromine molecules linked by weak interactions, as is observed here for **2**; for example, the [Br₁₀]^{2–} ion consists of two tribromide ions and two dibromine molecules.³² Similarly, the inclusion of a molecule of dibromine within the lattice of **2** is also rare, but not unknown: the salts [Me₄N][Sb₂Br₉]·Br₂³³ and [TeBr₃][AuBr₄]^{1/2}Br₂³⁴ have been characterised crystallographically. The dibromine molecule in [Me₄N][Sb₂Br₉]·Br₂ acts as a bridge between two [Sb₂Br₉][–] anions. The Br–Br bond distance [2.31(3) Å] is similar to that in **2** [2.329(2) Å], and both are significantly longer than the Br–Br distance in dibromine [2.27 Å]; the elongation in the Br–Br distance is probably due to intermolecular interactions. The dibromine molecule in [TeBr₃][AuBr₄]^{1/2}Br₂ has a Br–Br distance of 2.294(8) Å, much shorter than those in **2** or [Me₄N][Sb₂Br₉]·Br₂, indicative of the absence of any significant intermolecular interactions.

In the structure of compound **2** the cationic heterocycle is planar within 0.015(4) Å and the Br(1) atom shows a displacement of 0.170(13) Å from this plane with a concomitant C(2)–S(1)–C(1)–Br(1) torsion angle of 174.0(6)°. Replacement of the thione-sulfur atom with a bromine atom and the associated positive charge do not have any appreciable effect on the bond lengths and angles of the ring, other than a slight increase in the C(1)–S(1)/(2) bond distances and an increase in the S(1)–C(1)–S(2) bond angle. As observed for **1**·I₂, **1**·IBr and **1**·Br₂, the two S–CH₃ groups on the ring are oriented differently as shown by the torsion angles C(4)–S(3)–C(2)–C(3) 154.5(8) and C(5)–S(4)–C(3)–C(2) 172.9(8)°.

Conclusion

We have demonstrated that the solid-state structures adopted by thione–dihalogen adducts, R₂C=S·X₂, are dependent upon the nature of the halogen X₂. When X₂ = I₂ or IBr the molecular ‘charge-transfer’ adducts are obtained, containing linear S⋯I–X (X = I or Br) moieties in which the sulfur atom is in a Ψ-tetrahedral environment (taking into account the stereochemically active lone pairs). However, when X₂ = Br₂ homolytic cleavage of dibromine occurs, with concomitant formation of two S–Br bonds. The Br–S–Br moiety is linear and the sulfur atom adopts a T-shaped (or Ψ-trigonal bipyramidal) geometry. The geometric dependence of the thionyl sulfur atom in R₂C=S·X₂ upon X (X = I or Br) is clearly established.

The FT-Raman spectra of the three adducts confirm the observed structural features. Although thione–diiodine complexes have been widely investigated, there is a paucity of comparative data on adducts of the lighter halogens and interhalogens which is readdressed to some extent in this study. The solvent dependency of the reactions of Group 15 or 16 donors with dihalogens or interhalogens is illustrated by the reaction of **1** with Br₂: when the reaction is carried out in toluene, the adduct of **1**·Br₂ is the sole product; however, in dichloromethane **1**·Br₂ is obtained, together with the dithiolylium salt **2** which contains an unusual dibromine of crystallisation.

Experimental

4,5-Bis(methylsulfanyl)-1,3-dithiole-2-thione **1** was prepared as described.²⁰ Iodine, iodine monobromide and bromine were purchased from Aldrich and used as received.

Melting points were taken using an Electrothermal Melting Point apparatus and are uncorrected. Infrared spectra were recorded on a Mattson Genesis Series FTIR spectrometer. Elemental analyses were carried out by MEDAC Ltd. Raman spectra were obtained from a Renishaw Ramascope 1000 Raman Microscope. The laser used was a 780 nm near IR laser. All spectra were recorded using a 200 exposure ‘extended’ scan between 3400 and 300 cm^{–1} and a 0.5 s static scan at lower wavenumbers.

Preparations

4,5-Bis(methylsulfanyl)-1,3-dithiole-2-thione–diiodine 1·I₂. To a solution of compound **1** (260 mg, 1.15 mmol) in dichloromethane (50 ml) was added iodine (290 mg, 1.14 mmol) in one portion. The mixture was stirred at room temperature for 2 h, followed by cooling at –5 °C over 16 h. The reaction contents were filtered and the solids washed with diethyl ether (2 × 25 ml) to give **1**·I₂ as a brown crystalline material (470 mg, 85% yield); mp 62–64 °C (Found: C, 12.77; H, 1.32; S, 33.77. C₅H₆I₂S₅ requires C, 12.51; H, 1.26; S, 33.38%); $\tilde{\nu}_{\max}/\text{cm}^{-1}$ 1443, 1308, 1017 (C–S), 532 and 458.

4,5-Bis(methylsulfanyl)-1,3-dithiole-2-thione–iodine monobromide 1·IBr. To a solution of compound **1** (400 mg, 1.77 mmol) in dichloromethane (100 ml) was added iodine monobromide (400 mg, 1.93 mmol) in one portion. The reaction mixture was refluxed for 5 min, followed by cooling at –5 °C over 16 h. The product was filtered off and washed with diethyl ether (2 × 25 ml) to give **1**·IBr as an orange-brown crystalline solid (760 mg, 99% yield); mp 98–100 °C (Found: C, 13.74; H, 1.46; S, 37.74. C₅H₆BrIS₅ requires C, 13.86; H, 1.40; S, 37.00%); $\tilde{\nu}_{\max}/\text{cm}^{-1}$ 1435, 1302, 1030 (C–S), 973 and 463.

4,5-Bis(methylsulfanyl)-1,3-dithiole-2-thione–dibromine 1·Br₂ (Method A) and 2-bromo-4,5-bis(methylsulfanyl)-1,3-dithiolylium tribromide–bromine [2/1] 2. To a solution of compound **1** (400 mg, 1.77 mmol) in dichloromethane (100 ml) was added bromine (0.1 ml, 1.94 mmol) dropwise. The reaction mixture was refluxed for 5 min, followed by cooling at –5 °C

Table 4 Crystal data and details of data collection and structure refinement for adducts **1**·I₂, **1**·IBr, **1**·Br₂ and **2**

	1 ·I ₂	1 ·IBr	1 ·Br ₂	2
Chemical formula	C ₅ H ₆ I ₂ S ₅	C ₅ H ₆ BrIS ₅	C ₅ H ₆ Br ₂ S ₅	C ₅ H ₆ Br ₄ S ₄ ·½Br ₂
Formula weight	480.21	433.21	386.22	593.89
Crystal system	Monoclinic	Triclinic	Triclinic	Triclinic
Space group	<i>P</i> 2 ₁ / <i>n</i> (no. 14)	<i>P</i> 1̄ (no. 2)	<i>P</i> 1̄ (no. 2)	<i>P</i> 1̄ (no. 2)
<i>a</i> /Å	9.037(3)	7.5933(12)	6.9403(9)	6.4872(6)
<i>b</i> /Å	9.606(2)	8.601(2)	8.3128(14)	7.0403(9)
<i>c</i> /Å	13.906(5)	10.8848(14)	10.554(4)	17.166(4)
<i>α</i> /°	—	110.710(8)	69.042(13)	84.060(10)
<i>β</i> /°	98.57(2)	91.227(9)	87.146(11)	80.958(9)
<i>γ</i> /°	—	111.989(13)	87.880(8)	69.198(9)
<i>V</i> /Å ³	1193.7(6)	606.8(2)	567.8(2)	722.8(2)
<i>Z</i>	4	2	2	2
<i>μ</i> (Mo-Kα)/cm ⁻¹	60.92	67.41	80.03	144.40
Reflections collected	3811	2681	1981	3011
Unique reflections	1729	1694	1181	1973
<i>R</i> _{int}	0.0646	0.0672	0.0778	0.0615
Final <i>R</i> 1, <i>wR</i> 2	0.0675 (0.0514) ^a	0.0418 (0.0382),	0.0590 (0.0492),	0.0534 (0.0450),
	0.1258 (0.1187) ^a	0.1042 (0.1031)	0.1346 (0.1325)	0.1136 (0.1116)

^a Values for all unique data; those calculated for data with $I > 2\sigma(I)$ are given in parentheses.

over 16 h. The solution was filtered and the solid washed with diethyl ether (2 × 25 ml) to give **1**·Br₂ as a brown powder (340 mg, 50% yield). After slow evaporation of the filtrate a purple oil was obtained, from which dark red crystals of **2** were obtained after 16 h under ambient conditions (120 mg, 10% yield). **1**·Br₂: mp 72 °C (decomp.) (Found: C, 16.00; H, 1.70; S, 41.09. C₅H₆Br₂S₅ requires C, 15.55; H, 1.57; S, 41.51%); $\tilde{\nu}_{\max}/\text{cm}^{-1}$ 1434s, 1312, 1058, 1043, 991 (C–S), 964 and 465. **2**: $\tilde{\nu}_{\max}/\text{cm}^{-1}$ 1262, 1096, 1024 and 803.

Method B. To a solution of compound **1** (400 mg, 1.77 mmol) in toluene (50 ml) was added bromine (0.1 ml, 1.94 mmol) dropwise. The reaction mixture was refluxed for 5 min, followed by cooling at –5 °C for 2 h. After filtering and washing with diethyl ether (2 × 25 ml) the product was isolated as an orange-brown crystalline solid (370 mg, 54% yield); all spectroscopic data were identical to those of **1**·Br₂ obtained *via* Method A.

X-Ray crystallography

All crystallographic measurements for complexes **1**·I₂, **1**·IBr, **1**·Br₂ and **2** were made at 150 K on a Delft Instruments FAST TV area detector diffractometer positioned at the window of a rotating anode generator, using Mo-Kα radiation ($\lambda = 0.71069$ Å) in a manner described previously.³⁵ All data sets were corrected for absorption using DIFABS.³⁶ The structures were solved by direct methods (SHELXS 86),³⁷ developed *via* difference syntheses, and refined on F^2 by full-matrix least squares (SHELXL 93)³⁸ using all unique data with intensities greater than 0. In all cases the non-hydrogen atoms were anisotropic, and the hydrogen atoms included in calculated positions (riding model). The crystallographic data and refinement details are presented in Table 4.

CCDC reference number 186/1543.

See <http://www.rsc.org/suppdata/dt/1999/3007/> for crystallographic files in .cif format.

Acknowledgements

We thank Professor Jack Yarwood and Dr. Chris Sammon for their help with Raman measurements.

References

- S. M. Godfrey, C. A. McAuliffe, R. G. Pritchard and J. M. Sheffield, *Chem. Commun.*, 1998, 921; S. M. Godfrey, C. A. McAuliffe, I. Mushtaq, R. G. Pritchard and J. M. Sheffield, *J. Chem. Soc., Dalton Trans.*, 1998, 3815; N. Bricklebank, S. M. Godfrey, H. P. Lane, C. A. McAuliffe, R. G. Pritchard and J. M. Moreno, *J. Chem. Soc., Dalton Trans.*, 1995, 3873; N. Bricklebank, S. M. Godfrey, H. P. Lane, C. A. McAuliffe and R. G. Pritchard, *J. Chem. Soc., Dalton Trans.*, 1995, 1759; N. Bricklebank, S. M. Godfrey, H. P. Lane, C. A. McAuliffe, R. G. Pritchard and J. M. Moreno, *J. Chem. Soc., Dalton Trans.*, 1994, 2421.
- F. Ruthe, W. W. du Mont and P. G. Jones, *Chem. Commun.*, 1997, 1947; R. D. Bailey, G. W. Drake, M. Grabarczyk, T. W. Hanks, L. L. Hook and W. T. Pennington, *J. Chem. Soc., Perkin Trans. 2*, 1997, 2773.
- (a) D. C. Apperley, N. Bricklebank, S. L. Burns, D. E. Hibbs, M. B. Hursthouse and K. M. A. Malik, *J. Chem. Soc., Dalton Trans.*, 1998, 1289; (b) S. M. Godfrey, S. L. Jackson, C. A. McAuliffe and R. G. Pritchard, *J. Chem. Soc., Dalton Trans.*, 1997, 4499; (c) A. J. Blake, F. A. Devillanova, A. Garau, L. M. Gilby, R. O. Gould, F. Isaia, V. Lippolis, S. Parsons, C. Radek and M. Schroder, *J. Chem. Soc., Dalton Trans.*, 1998, 2037; (d) A. J. Blake, F. Cristiani, F. A. Devillanova, A. Garau, L. M. Gilby, R. O. Gould, F. Isaia, V. Lippolis, S. Parsons, C. Radek and M. Schroder, *J. Chem. Soc., Dalton Trans.*, 1997, 1337; (e) F. Cristiani, F. Delmartin, F. A. Devillanova, F. Isaia, V. Lippolis and G. Verani, *Inorg. Chem.*, 1994, **33**, 6315.
- M. D. Rudd, S. V. Lindeman and S. Husebye, *Acta Chem. Scand.*, 1997, **51**, 689.
- (a) F. Bigoli, P. Deplano, M. L. Mercuri, M. A. Pellinghelli, A. Sabatini, E. F. Trogu and A. Vacca, *J. Chem. Soc., Dalton Trans.*, 1996, 3583; (b) D. Atzei, P. Deplano, E. F. Trogu, F. Bigoli, M. A. Pellinghelli, A. Sabatini and A. Vacca, *Can. J. Chem.*, 1989, **67**, 1416; (c) D. Atzei, P. Deplano, E. F. Trogu, F. Bigoli, M. A. Pellinghelli and A. Vacca, *Can. J. Chem.*, 1988, **66**, 1483; (d) E. L. Ahlson and K. O. Stromme, *Acta Chem. Scand., Ser. A*, 1974, **28**, 175.
- F. Bigoli, P. Deplano, M. L. Mercuri, M. A. Pellinghelli and E. F. Trogu, *Phosphorus Sulfur Silicon Relat. Elem.*, 1992, **70**, 175.
- F. Bigoli, P. Deplano, M. L. Mercuri, M. A. Pellinghelli and E. F. Trogu, *Phosphorus Sulfur Silicon Relat. Elem.*, 1992, **72**, 65.
- F. L. Lu, M. Keshavarz, G. Srdanov, R. H. Jacobson and F. Wudl, *J. Org. Chem.*, 1989, **54**, 2165.
- F. Freeman, J. W. Ziller, H. N. Po and M. C. Keindl, *J. Am. Chem. Soc.*, 1988, **110**, 2568.
- F. H. Herbstein and W. Schwotzer, *J. Am. Chem. Soc.*, 1984, **106**, 2367.
- F. Cristiani, F. Delmartin, F. A. Devillanova, F. Isaia, G. Saba and G. Verani, *J. Chem. Soc., Dalton Trans.*, 1992, 3553.
- A. J. Blake, F. A. Devillanova, A. Garau, F. Isaia, V. Lippolis, S. Parsons and M. Schröder, *J. Chem. Soc., Dalton Trans.*, 1999, 525.
- M. Arca, F. A. Devillanova, F. A. Garau, F. Isaia, V. Lippolis, G. Verani and F. Delmartin, *Z. Anorg. Allg. Chem.*, 1998, **624**, 745.
- C. Knobler, C. Baker, H. Hope and J. D. McCullough, *Inorg. Chem.*, 1971, **10**, 697.
- A. J. Arduengo and E. M. Burgess, *J. Am. Chem. Soc.*, 1977, **99**, 2376.
- F. A. Devillanova, P. Deplano, F. Isaia, V. Lippolis, M. L. Mercuri, S. Piludu, G. Verani and F. Delmartin, *Polyhedron*, 1998, **17**, 305.

- 17 F. Delmartin, P. Deplano, F. A. Devillanova, F. Isaia, V. Lippolis and G. Verani, *Inorg. Chem.*, 1993, **32**, 3694; D. L. Nosco, M. J. Hegg, M. D. Glick, R. C. Elder and E. Deutsch, *J. Am. Chem. Soc.*, 1980, **102**, 7784; G. H. Y. Lin and H. Hope, *Chem. Commun.*, 1970, 169; *Acta Crystallogr., Sect. B*, 1972, **28**, 643; O. Hassel and H. Hope, *Acta Chem. Scand.*, 1961, **15**, 407.
- 18 J. Buxeraud, A. C. Absil, J. Claude, C. Raby, G. Catanzano and C. Beck, *Eur. J. Med. Chem. Chim. Ther.*, 1985, **20**, 43.
- 19 J. R. Ferraro and J. M. Williams, *Introduction to Synthetic Electric Conductors*, Academic Press, New York, 1987.
- 20 G. Steimecke, R. Kirmse and E. Hoyer, *Z. Chem.*, 1975, **15**, 28; G. Steimecke, H.-J. Sieler, R. Kirmse and E. Hoyer, *Phosphorus Sulfur Relat. Elem.*, 1979, **7**, 49.
- 21 N. Svenstrup and J. Becher, *Synthesis*, 1995, 215.
- 22 J. E. Huheey, *Inorganic Chemistry: Principles of Structure and Reactivity*, 3rd edn., Harper and Row, New York, 1983.
- 23 F. van Bolhuis, P. B. Koster and T. Mighelsen, *Acta Crystallogr.*, 1967, **23**, 940.
- 24 I. L. Karle, *J. Chem. Phys.*, 1955, **23**, 1739.
- 25 L. N. Swink and G. B. Carpenter, *Acta Crystallogr., Sect. B*, 1968, **24**, 429.
- 26 L. E. Selin, *Naturwissenschaften*, 1960, **47**, 104.
- 27 F. Bigoli, P. Deplano, F. A. Devillanova, V. Lippolis, M. L. Mercuri, M. A. Pellinghelli and E. F. Trogu, *Eur. J. Inorg. Chem.*, 1998, 137; M. Arca, F. Demartin, F. A. Devillanova, A. Garau, F. Isaia, V. Lippolis, S. Piludu and G. Verani, *Polyhedron*, 1998, **17**, 3111.
- 28 S. M. Godfrey, S. L. Jackson, C. A. McAuliffe and R. G. Pritchard, *J. Chem. Soc., Dalton Trans.*, 1998, 4201.
- 29 E. Fanghänel, K. H. Kühnemund and A. M. Richter, *Synth. Commun.*, 1984, 319.
- 30 K. N. Robertson, P. K. Bakshi, T. S. Cameron and O. Kemp, *Z. Anorg. Allg. Chem.*, 1997, **623**, 101.
- 31 B. Vonnegut and B. E. Warren, *J. Am. Chem. Soc.*, 1936, **58**, 2459; J. Donohue and S. H. Goodman, *Acta Crystallogr.*, 1965, **18**, 568.
- 32 C. W. Cunningham, G. R. Burns and V. McKee, *Inorg. Chim. Acta*, 1990, **167**, 135.
- 33 C. R. Hubbard and R. A. Jacobson, *Inorg. Chem.*, 1972, **11**, 2247.
- 34 C. Freireddbrugger, D. Jentsch, P. G. Jones and E. Schwarzmann, *Z. Naturforsch., Teil B*, 1987, **42**, 1553.
- 35 J. A. Darr, S. R. Drake, M. B. Hursthouse and K. M. A. Malik, *Inorg. Chem.*, 1993, **32**, 5704.
- 36 N. P. C. Walker and D. Stuart, *Acta Crystallogr., Sect. A*, 1983, **39**, 158.
- 37 G. M. Sheldrick, *Acta Crystallogr., Sect. A*, 1990, **46**, 467.
- 38 G. M. Sheldrick, SHELXL 93, program for crystal structure refinement, University of Göttingen, 1993.

Paper 9/03068I