

A new assembly of diiodine molecules at the triphenylphosphine sulfide template†

Massimiliano Arca,^a Francesco Demartin,^b Francesco A. Devillanova,^{*a} Alessandra Garau,^a Francesco Isaia,^a Vito Lippolis^{*a} and Gaetano Verani^a

^a Dipartimento di Chimica e Tecnologie Inorganiche e Metallorganiche, Università di Cagliari, Via Ospedale 72, I-09124, Cagliari, Italy. E-mail: lippolis@vaxca1.unica.it

^b Dipartimento di Chimica Strutturale e Stereochimica Inorganica e Centro CNR, Università di Milano, Via G. Venezian 21, I-20133 Milano, Italy

Received 27th May 1999, Accepted 5th July 1999

The new I₂ adduct TPPS·3I₂ (TPPS = triphenylphosphine sulfide) has been synthesized and characterised by X-ray diffraction on a single crystal. It consists of a TPPS·I₂ 1 : 1 molecular adduct interacting with two further diiodine molecules in a “head to tail” sequence to give an overall TPPS·I₂ ··· I₂ ··· I₂ arrangement in the asymmetric unit. Considering the I ··· I interactions below 3.6 Å, five of the six iodine atoms of the asymmetric unit are arranged in infinite polycyclic ribbons. The iodine atoms, to which the TPPS molecules are connected, are almost perpendicular appendages on opposite sides of each ribbon. Weaker interactions between the ribbons give rise to unprecedented continuous stepped layers of diiodine molecules anchored to TPPS donor units. The structural features of TPPS·3I₂ are discussed in comparison with those of the previously reported 2TPPS·3I₂, TPPS·I₂, TPPS·IBr and TPPS·ICl, and its FT-Raman spectrum compared with those of 2TPPS·3I₂ and TPPS·I₂. A re-examination of the equilibrium reaction between TPPS and I₂ by ³¹P NMR confirms the formation of the 1 : 1 adduct in CH₂Cl₂ solution and gives a formation constant value of 137 ± 11 dm³ mol⁻¹.

Introduction

Dihalogen charge-transfer complexes of tertiary phosphine chalcogenides R₃PE (R = alkyl or aryl; E = S or Se) were first reported by Zingaro and co-workers^{1–5} who mainly used UV-visible and infrared spectroscopies to obtain information on the nature of R₃PE·X₂ 1 : 1 adducts (X₂ = I₂, IBr or ICl) both in solution (CCl₄) and in the solid state. The first reported crystallographic study concerned the compound of formula 2Ph₃PS·3I₂ that was unexpectedly obtained from the stoichiometric reaction of triphenylphosphine sulfide (TPPS) with diiodine in CCl₄.⁶ In this compound one diiodine molecule bridges two TPPS·I₂ 1 : 1 adduct units *via* soft–soft I ··· I interactions of 3.570(1) Å to give a “Z”-shaped D·I₂ ··· I₂ ··· I₂·D arrangement (D = TPPS). Initially, it seemed that the power of TPPS as donor was not enough to stabilise a discrete 1 : 1 adduct. However, very recently Bricklebank and co-workers⁷ reported the crystal structure of the 1 : 1 adduct TPPS·I₂, obtained by treating TPPS and I₂ in a 1 : 1 molar ratio in CH₂Cl₂.⁸ At the same time, the 1 : 1 adducts TPPS·IBr and TPPS·ICl were structurally characterised by us,⁹ clearly illustrating that triphenylphosphine sulfide can form 1 : 1 charge-transfer complexes with both I₂ and stronger acceptors such as ICl and IBr.

Diiodine is the halogen with the highest catenating ability, and the polyiodide anions which stem from the donor/acceptor interaction of I₃⁻ and/or I⁻ anions with I₂ molecules can assume a wide range of structural motifs.¹⁰ Polyiodide cations such as I₄²⁺, I₅⁺ and I₅³⁺ have also been reported;¹¹ nevertheless, neutral extended poly(I₂) arrangements held up by soft–soft I₂ ··· I₂ interactions are extremely rare, and the best example so far reported is the structure of solid I₂ itself.¹² This

consists of an orthorhombic layer lattice: the I–I bond distance [2.715(6) Å] is longer than that in the gas phase [2.667(2) Å];¹³ the inter-atomic I ··· I distances between two I₂ molecules are 3.497(6) and 3.972(7) Å within a layer (still less than the sum of the van der Waals radii for I₂ [4.2 Å]¹⁴) and 4.269(6), 4.332(7) and 4.412(7) Å between layers; the angles between I₂ molecules within each layer have values very close to 90 and 180°. The (I₂)₇ chain observed in the compound bis(1,5-diphenylthiocarbazono) heptakis(diiodine) was described¹⁵ on the basis of structural data, as a D–I⁺ ··· [I₅]⁻ ··· I₂ ··· [I₅]⁻ ··· [I–D] system, and not properly as a chain of neutral diiodine molecules. Several neutral I₂ adducts with a diiodine content higher than the 1 : 1 stoichiometry have been reported; normally, their crystal structures do not feature extended poly(I₂) networks, but only discrete D·I₂ ··· I₂ or D·I₂ ··· I₂ ··· I₂·D arrangements (see for example the 1 : 2 adduct between *N*-methylbenzoxazole-2(3*H*)-selone¹⁶ and I₂ or the above described 2TPPS·3I₂); rarely, as in *N*-methylbenzothiazole-2(3*H*)-selone–diiodine(1/2),¹⁷ extended neutral networks of diiodine molecules can be observed.

Considering the ability of TPPS to give I₂ adducts with different stoichiometry on changing the reaction solvent, we argued that the same ligand could be used as a donor to stabilise extended neutral poly(I₂) networks when using a TPPS : I₂ molar ratio higher than 1 : 1. We report herein the first example of a genuine infinite three dimensional poly(I₂) assembly obtained by treating TPPS with an excess of I₂ in CH₂Cl₂ solution. The new adduct corresponding to the formulation TPPS·3I₂ has been characterised by single crystal X-ray diffraction, FT-IR and FT-Raman spectroscopic techniques.

Results and discussion

The reaction of triphenylphosphine sulfide with an excess of I₂ (1 : 3 molar ratio) in CH₂Cl₂ solution affords dark red crystals

† Supplementary data available: FT-IR and FT Raman spectra. Available from BLDSC (No. SUP 57603, 3 pp.). See Instructions for Authors, 1999, Issue 1 (<http://www.rsc.org/dalton>).

after slow evaporation of the solvent. An X-ray diffraction study shows that the crystals consist of a $\text{TPPS}\cdot\text{I}_2$ 1:1 molecular adduct interacting with two further diiodine molecules in a "head to tail" sequence (Fig. 1) to give an overall $\text{TPPS}\cdot 3\text{I}_2$ stoichiometry. The S-donor atom of the ligand binds the I(1)–I(2) diiodine molecule almost linearly [S–I(1) 2.591(1), I(1)–I(2) 2.982(1) Å, S–I(1)–I(2) 178.11(3)°] (Table 1), with a coordination geometry that is very similar to that of the related compounds, $\text{TPPS}\cdot\text{I}_2$,⁷ $2\text{TPPS}\cdot 3\text{I}_2$,⁶ $\text{TPPS}\cdot\text{IBr}$ ⁹ and $\text{TPPS}\cdot\text{ICl}$.⁹ Comparison of the structural features of the P–S–I–X (X = I, Br or Cl) moiety for all the structurally characterised charge-transfer adducts of TPPS with halogens and interhalogens (Table 2) shows that in all cases the P–S bond is lengthened with respect to the value found in pure TPPS [1.950(3)Å],¹⁸ and that




Fig. 1 Partial view of the ribbon of iodine atoms with the TPPS appendages.

Table 1 Selected interatomic distances (Å) and angles (°) in $\text{TPPS}\cdot 3\text{I}_2$

I(1)–I(2)	2.982(1)	I(3)–I(4)	2.744(1)
I(2)–I(3)	3.373(1)	I(4)–I(5)	3.582(1)
I(2)–I(4 ⁱ)	3.561(1)	I(5)–I(6)	2.710(1)
I(2)–I(6 ⁱⁱ)	3.503(1)	I(2)⋯I(2 ⁱⁱⁱ)	4.104(1)
I(3)⋯I(4 ^{iv})	4.169(1)	I(4)⋯I(4 ^{iv})	4.158(1)
I(1)–S	2.591(1)	S–P	2.030(2)
P–C(11)	1.797(5)	P–C(21)	1.805(5)
P–C(31)	1.795(4)		
I(2)–I(1)–S	178.11(3)	I(1)–S–P	105.47(6)
S–P–C(11)	112.9(2)	S–P–C(21)	104.0(2)
S–P–C(31)	113.1(2)	C(11)–P–C(21)	110.6(2)
C(11)–P–C(31)	107.4(2)	C(21)–P–C(31)	108.7(2)
I(1)–I(2)–I(3)	93.02(2)	I(2)–I(3)–I(4)	175.35(2)
I(3)–I(4)–I(5)	86.06(2)	I(4)–I(5)–I(6)	171.24(3)
I(1)–I(2)–I(6 ⁱⁱ)	82.75(2)	I(3)–I(2)–I(4)	175.78(2)
I(2)–I(6 ⁱⁱ)–I(5 ⁱⁱ)	177.27(3)		

Symmetry codes: i $x, 1 + y, z$; ii $-x, \frac{1}{2} + y, \frac{3}{2} - z$; iii $-x, -y, 2 - z$; iv $-x, -1 - y, 2 - z$.

Table 2 Structural features of the P–S–I–X (X = I, Br or Cl) framework in all the structurally characterised adducts of triphenylphosphine sulfide with I–X

	X = I			X = Br TPPS·IBr ⁹	X = Cl TPPS·ICl ⁹
	TPPS·I ₂ ⁷	2TPPS·3I ₂ ⁶	TPPS·3I ₂		
P–S ^a	1.998(2)	2.007(3)	2.030(2)	2.014(1)	2.014(1)
S–I(1)	2.753(2)	2.729(2)	2.591(1)	2.665(1)	2.641(1)
I(1)–X	2.823(1)	2.838(1)	2.9818(5)	2.668(1)	2.586(1)
S–I(1)–X	175.51(3)	175.23(5)	178.11(3)	175.08(2)	174.86(3)
P–S–I(1)	108.78(7)	107.0(1)	105.47(6)	107.98(3)	108.29(5)

^a The P–S bond distance found in pure TPPS is 1.950(3) Å.¹⁸

the new adduct $\text{TPPS}\cdot 3\text{I}_2$ has the shortest S–I bond. Even shorter S–I bonds have been observed only for some adducts of ethylenethiourea and 1,5-diphenylthiocarbazono with I₂.^{15a} However, the S–I and I–I bond distances found in $\text{TPPS}\cdot 3\text{I}_2$ fall in the $d(\text{S–I})$ versus $d(\text{I–I})$ correlation reported for the adducts of sulfur containing donors and diiodine.¹⁹

It is interesting to point out the decrease and increase of the S–I and I–I bond distances respectively on passing from $\text{TPPS}\cdot\text{I}_2$ to $2\text{TPPS}\cdot 3\text{I}_2$ and $\text{TPPS}\cdot 3\text{I}_2$. Since the donor molecule is the same in the three compounds, the changes in such lengths in $2\text{TPPS}\cdot 3\text{I}_2$ and $\text{TPPS}\cdot 3\text{I}_2$ should originate from the additional short contacts of the terminal iodine atom of the TPPS co-ordinated diiodine molecule with further iodine atoms. While in $\text{TPPS}\cdot\text{I}_2$ the adduct units are discrete and are held together by normal van der Waals contacts, in $2\text{TPPS}\cdot 3\text{I}_2$ two $\text{TPPS}\cdot\text{I}_2$ units act as donors towards a bridging diiodine molecule that acts as an acceptor to give a $\text{TPPS}\cdot\text{I}_2\cdots\text{I}_2\cdots\text{I}_2\cdots\text{TPPS}$ "Z"-shaped system [I₂⋯I₂ 3.57(2) Å]. In a different way, in $\text{TPPS}\cdot 3\text{I}_2$, the $\text{TPPS}\cdot\text{I}_2$ unit interacts with a second diiodine molecule [I(3)–I(4), I₂⋯I₂ 3.373(1) Å], which in turn behaves as a donor towards a third diiodine molecule through the I(4) iodine atom [I(5)–I(6), I₂⋯I₂ 3.582(1) Å] to give an overall $\text{TPPS}\cdot\text{I}_2\cdots\text{I}_2\cdots\text{I}_2$ system (Fig. 1). Therefore, a similar (I₂)₃ chain is present in $2\text{TPPS}\cdot 3\text{I}_2$ and $\text{TPPS}\cdot 3\text{I}_2$; however, in $2\text{TPPS}\cdot 3\text{I}_2$ both terminal diiodine molecules of the I₂⋯I₂⋯I₂ arrangement act as donors towards the central I₂, whereas in $\text{TPPS}\cdot 3\text{I}_2$ the terminal diiodine molecule I(5)–I(6) behaves as an acceptor. In other words, we can schematically represent the (I₂)₃ chain in $2\text{TPPS}\cdot 3\text{I}_2$ and $\text{TPPS}\cdot 3\text{I}_2$ as $\text{TPPS}\cdot\text{I}_2\cdots\text{I}_2\cdots\text{D}'$ and $\text{TPPS}\cdot\text{I}_2\cdots\text{I}_2\cdots\text{A}$ respectively (D' = $\text{TPPS}\cdot\text{I}_2$ as donor, A = I₂ as acceptor). Clearly, in $\text{TPPS}\cdot 3\text{I}_2$ the partial negative charge induced on I(2) upon interaction of the I(1)–I(2) molecule with TPPS will be more effectively delocalised by the additional rather strong I(2)⋯I₂ bonding interactions, with consequent lengthening and shortening of the I(1)–I(2) and S–I(1) bonds respectively. These interactions are at the basis of the crystal packing in $\text{TPPS}\cdot 3\text{I}_2$.

In this new compound, if I⋯I interactions below 3.6 Å are considered, the iodine atoms I(2), I(3), I(4), I(5) and I(6) and their equivalents are arranged to form infinite polycyclic ribbons running along [010] and made up of 9-membered rings sharing eight iodine atoms. Iodine atoms I(1) are almost perpendicular appendages on opposite sides of each ribbon, to which the TPPS molecules are connected (Fig. 1). Weaker interactions between the ribbons (still below the value of the sum of the van der Waals radii for I₂), in the 4.104(1)–4.169(1) Å range (Table 1), give rise to unprecedented continuous stepped layers of diiodine molecules anchored to TPPS donor units (Fig. 2). Each of the I₂ molecules in the asymmetric unit displays a different environment. The I(1)–I(2) molecule, which interacts strongly with the TPPS ligand, is the most perturbed, and accordingly displays the largest I–I distance. Three other I₂ molecules within the ribbon interact at almost 90° with the I(1)–I(2) molecule via the I(2) iodine atom [I(2)–I(3) 3.373(1),

I(2)–I(4ⁱ) 3.561(1), I(2)–I(6ⁱⁱ) 3.503(1) Å, i $x, 1 + y, z$; ii $-x, \frac{1}{2} + y, \frac{3}{2} - z$]. This further contributes to the lengthening of the I(1)–I(2) bond. The I(2)–I(3) molecule is also engaged in a weaker inter-ribbon interaction with its symmetry equivalent [I(2)⋯I(2ⁱⁱⁱ) 4.104(1) Å, iii $-x, -y, 2 - z$]. The I(3)–I(4)




Fig. 2 Perspective view of a stepped layer of iodine atoms. Darker I–I bonds correspond to distances shorter than 3.0 Å. Thin lines correspond to I–I bonds within the 3.0–3.6 Å range. The remaining I⋯I interactions, shown as dashed lines, are shorter than 4.2 Å.

molecule has six interactions with the surrounding iodine atoms in the 3.373(1)–4.169(1) Å range (Table 1) and lengthens to 2.744(1) Å. Finally the I(5)–I(6) molecule, which only has two short interactions [3.503(1) and 3.582(1) Å (Table 1)], is almost unperturbed and displays an I–I bond length [2.710(1) Å] very close to that observed in the pure solid I₂ [2.715 Å].¹² The TPPS molecules, located between the poly(I₂) layers, are arranged as centrosymmetric pairs (Fig. 3); each has a phenyl ring parallel to the corresponding one of the symmetry-related molecule with a distance between the phenyl planes of about 3.83 Å, typical of a “graphite-like” stacking.

In the past, we classified 1:1 I₂ adducts in three classes¹⁹ according to the value of the I–I bond order (n)²⁰ defined by the equation $d(\text{I–I}) = d_0 - c \log n$ [d_0 (2.67 Å) is the I–I bond distance for I₂ in the gas phase, c an empirical constant with a value of 0.85]: (1) weak or medium-weak [I–I bond order $n > 0.6$]; (2) strong [I–I bond order $0.4 > n > 0.6$]; (3) very strong [I–I bond order $n < 0.4$]. Weak or medium-weak I₂ adducts are characterised by the presence of only one Raman peak in the $\nu(\text{I–I})$ stretching region, whose frequency depends on the I–I bond elongation; this is clearly pointed out by the linear correlation observed between the measured $\nu(\text{I–I})$ Raman frequencies and $d(\text{I–I})$ bond distances, with $\nu(\text{I–I})$ moving towards lower values compared to the stretching frequency reported for pure I₂ (180 cm⁻¹).¹⁹ For strong adducts, two symmetric (ν_1) and antisymmetric (ν_3) stretching vibrations should be expected in the FT-Raman spectra for the E–I–I (E = donor atom) three-body system, whereas the spectra of very strong adducts should be characterised by the stretching vibration of the [(D)E–I]⁺ system.

For TPPS·3I₂ only two peaks at 173 and 140 cm⁻¹ have been observed in the FT-Raman spectrum (SUP 57603) at exactly the same frequencies as the peaks found for 2TPPS·3I₂ [172.5 and 140 cm⁻¹],⁹ despite their different structural features. For 2TPPS·3I₂ the two peaks were respectively assigned to the bridging I₂ molecule [I–I 2.757(2) Å; $n = 0.79$] and to the S-bonded diiodine molecules [I–I 2.838(1) Å; $n = 0.63$].⁹




Fig. 3 Crystal packing of TPPS·3I₂, seen along [010].




Fig. 4 Linear correlation between $\Delta\delta_{\text{exp}}[\text{TPPS}]^\circ$ vs. $[\text{TPPS}\cdot\text{I}_2]_{\text{calc}}$. Correlation coefficient = 0.997; $[\text{TPPS}]^\circ$ = analytical concentration of TPPS. The ^{31}P NMR chemical shift of the adduct $\delta_{\text{TPPS}\cdot\text{I}_2}$ calculated from the slope is 40.68.

In $\text{TPPS}\cdot 3\text{I}_2$ the structural features show that the S-bonded diiodine molecule is very elongated [I–I 2.982(1) Å; $n = 0.43$]; consequently, the $\text{TPPS}\cdot\text{I}_2$ moiety can be considered as an adduct falling on the borderline between the two classes of strong and very strong adducts. The band at 173 cm^{-1} falls fairly well in the above mentioned correlation of $d(\text{I}–\text{I})$ versus $\nu(\text{I}–\text{I})$ considering both the bond distances found in $\text{I}(3)–\text{I}(4)$ and $\text{I}(5)–\text{I}(6)$ molecules. The other peak at 140 cm^{-1} should be tentatively assigned to the P–S–I system. The FT-IR spectrum confirms these assignments, since no bands are present around 173 cm^{-1} , while the most intense absorption falls at 141 cm^{-1} . We have synthesized the 1:1 adduct $\text{TPPS}\cdot\text{I}_2$, recently characterised by X-ray diffraction by Bricklebank and co-workers;⁷ its Raman spectrum shows a very strong peak, which dominates the whole spectrum, whose frequency at 145 cm^{-1} correlates very well with the elongation of the co-ordinated diiodine [I–I 2.823(1) Å].

Since contradictory results^{3,21,22} have been reported on the $\text{TPPS}\cdot\text{I}_2$ 1:1 adduct formation constant in solution, we re-examined the equilibrium reaction using the ^{31}P NMR chemical shifts of several CH_2Cl_2 solutions containing the same concentration of TPPS and increasing amounts of diiodine (see Experimental section). The linearity of the plot of $\Delta\delta[\text{TPPS}]^\circ$ vs. $[\text{TPPS}\cdot\text{I}_2]_{\text{calc}}$ (see Fig. 4) confirms formation of only the 1:1 molecular adduct in solution.^{23,24} The formation constant has been calculated by the non-linear least-squares program KEPS 11, as previously described.^{23,24} The calculated K value of $137 \pm 11\text{ dm}^3\text{ mol}^{-1}$ (21 °C, CH_2Cl_2) is very different from that reported in ref. 22 but it is close to the values reported by Tefeller and Zingaro³ ($106\text{ dm}^3\text{ mol}^{-1}$; 25 °C, CHCl_3) and Sobczyk and co-workers²¹ ($160\text{ dm}^3\text{ mol}^{-1}$; 25 °C, CCl_4).

Conclusion

An iodine-rich novel compound $\text{TPPS}\cdot 3\text{I}_2$, obtained by treating triphenylphosphine sulfide with diiodine, has been synthesized and characterised by X-ray single crystal diffraction. In $\text{TPPS}\cdot 3\text{I}_2$ the donor molecule is able to template a unique three-dimensional arrangement of diiodine molecules. This structure represents a very good example of the high catenating ability of diiodine in a very wide range of structural motifs.

Another very important point is related to FT-Raman spectroscopy. The compounds $2\text{TPPS}\cdot 3\text{I}_2$ and $\text{TPPS}\cdot 3\text{I}_2$ represent a case in which characterisation by FT-Raman spectroscopy alone could lead to erroneous conclusions on their structures, since both compounds have identical spectra but different structural features. This shows that crystal structure determination is indispensable to elucidate the nature of this kind of compound.

Experimental

Triphenylphosphine sulfide and CH_2Cl_2 were used as purchased from Aldrich; $\text{TPPS}\cdot\text{I}_2$ was synthesized according to the literature.⁷

Synthesis of $\text{TPPS}\cdot 3\text{I}_2$

A solution of triphenylphosphine sulfide (100 mg, 0.34 mmol) and diiodine (258.8 mg, 1.02 mmol) in CH_2Cl_2 (50 cm^3) was stirred at room temperature for 2 h. Subsequently *n*-hexane (50 cm^3) was added and the resulting mixture was stored at 5 °C. Brown-red crystals were formed in a few days. Found (Calc. for $\text{C}_{18}\text{H}_{15}\text{I}_6\text{PS}$): C, 20.48 (20.33); H, 1.43(1.50); S, 3.04(2.98)%. FT-IR spectrum (300–50 cm^{-1}): 270m, 257m, 249m, 201ms, 193ms, 141s(br), 91w. FT-Raman spectrum (300–50 cm^{-1} ; intensity of peaks related to the strongest taken equal to 10, in parentheses): 173 (10) and 140 (2.6).

Spectroscopic measurements

The FT-IR spectra (300–50 cm^{-1} , resolution 2 cm^{-1}) as polyethylene pellets were recorded on a Bruker IFS 66 spectrometer, FT-Raman spectra on pure powders (300–50 cm^{-1} , resolution 2 cm^{-1} ; excitation frequency 1064 nm; Nd:YAG laser; power 100 mW) on a Bruker RFS 100 FTR spectrometer and ^{31}P NMR spectra in solution on a Varian VXR 300 FT-NMR spectrometer, operating at 120 MHz and using 85% H_3PO_4 (aq) as an external reference.

Solution studies

The redetermination of the formation constant of the 1:1 adduct between TPPS and diiodine was carried out in CH_2Cl_2 by measuring the ^{31}P NMR signals for ten different solutions of TPPS (0.015 mol dm^{-3}) with increasing amounts of diiodine ($[\text{I}_2]/[\text{TPPS}]$ ranging between 0 and 5). The procedures for testing the presence of the 1:1 adduct in solution and for determining its formation constant have been described previously.²³

Crystal structure determination

Crystal data for $\text{TPPS}\cdot 3\text{I}_2$. $\text{C}_{18}\text{H}_{15}\text{I}_6\text{PS}$, $M = 1055.78$, monoclinic, space group $P2_1/c$ (no. 14), $a = 13.335(6)$, $b = 9.672(3)$, $c = 21.022(5)$ Å, $\beta = 102.87(3)^\circ$, $U = 2643(2)$ Å³, $T = 293$ K, $Z = 4$, $\mu(\text{Mo-K}\alpha) = 71.2\text{ cm}^{-1}$, 4745 reflections measured, 4632 unique ($R_{\text{int}} = 0.013$).

Intensity data were collected at room temperature with an Enraf-Nonius CAD4 diffractometer using graphite monochromated Mo-K α radiation. Lorentz-polarisation and an empirical absorption²⁵ correction were applied to the data. The structure was solved by direct methods²⁶ and refined by full-matrix least squares with anisotropic parameters for all non-hydrogen atoms. The final R and R' indices were 0.036 and 0.049. All H atoms were seen in Fourier difference maps but placed at calculated positions and not refined. Sources of neutral atomic scattering factors for all atoms are given in ref. 27; the values of δf and $\delta f'$ were those of ref. 28. All calculations were performed using the Personal SDP software.²⁹

CCDC reference number 186/1552.

See <http://www.rsc.org/suppdata/dt/1999/3069/> for crystallographic files in .cif format.

Acknowledgements

We thank the Regione Autonoma della Sardegna and Ministero dell'Università e della Ricerca Scientifica e Tecnologica (60%) for support in this research.

References

- 1 R. A. Zingaro and E. A. Meyers, *Inorg. Chem.*, 1962, **1**, 771.

- 2 R. A. Zingaro, R. E. McGlothlin and E. A. Meyers, *J. Phys. Chem.*, 1962, **66**, 2579.
- 3 W. Tefteller, Jr. and R. A. Zingaro, *Inorg. Chem.*, 1966, **5**, 2151.
- 4 R. A. Zingaro and R. M. Hedges, *J. Phys. Chem.*, 1961, **65**, 1132.
- 5 R. A. Zingaro, *Inorg. Chem.*, 1963, **2**, 192.
- 6 W. W. Schweikert and E. A. Meyers, *J. Phys. Chem.*, 1968, **72**, 1561; J. W. Bransford and E. A. Meyers, *Cryst. Struct. Commun.*, 1978, **7**, 697.
- 7 D. C. Apperley, N. Bricklebank, S. L. Burns and D. E. Hibbs, *J. Chem. Soc., Dalton Trans.*, 1998, 1289.
- 8 The results obtained by Bricklebank and co-workers⁷ are a clear example of how important the solvent can be in determining the nature of the final products in this type of reaction. Some other examples have been reported by us: F. Demartin, F. A. Devillanova, F. Isaia, V. Lippolis and G. Verani, *Inorg. Chim. Acta*, 1997, **255**, 203 and refs. therein; F. Bigoli, P. Deplano, M. L. Mercuri, M. A. Pellinghelli and E. F. Trogu, *J. Chem. Res.*, 1993, (S) 90.
- 9 M. Arca, F. A. Devillanova, A. Garau, F. Isaia, V. Lippolis and G. Verani, *Z. Anorg. Allg. Chem.*, 1998, **624**, 745.
- 10 A. J. Blake, F. A. Devillanova, R. O. Gould, W.-S. Li, V. Lippolis, S. Parsons, C. Radek and M. Schröder, *Chem. Soc. Rev.*, 1998, **27**, 195.
- 11 R. Faggiani, R. J. Gillespie, R. Kapoor, C. J. L. Lock and J. E. Vekris, *Inorg. Chem.*, 1988, **27**, 4350.
- 12 F. van Bolhuis, P. B. Koster and T. Migeheisen, *Acta Crystallogr.*, 1967, **23**, 90.
- 13 I. L. Karle, *J. Chem. Phys.*, 1955, **23**, 1739.
- 14 S. C. Nyburg and C. H. Faerman, *Acta Crystallogr., Sect. B*, 1985, **41**, 274.
- 15 F. H. Herbstein and W. Schwotzer, (a) *J. Am. Chem. Soc.*, 1984, **106**, 2367; (b) *Angew. Chem., Int. Ed. Engl.*, 1982, **21**, 219.
- 16 F. Cristiani, F. A. Devillanova, F. Isaia, V. Lippolis, G. Verani and F. Demartin, *Polyhedron*, 1995, **14**, 2937.
- 17 F. Cristiani, F. Demartin, F. A. Devillanova, F. Isaia, V. Lippolis and G. Verani, *Inorg. Chem.*, 1994, **33**, 6315.
- 18 P. W. Coddling and K. A. Kerr, *Acta Crystallogr., Sect. B*, 1978, **34**, 3785.
- 19 M. C. Aragoni, M. Arca, F. Demartin, F. A. Devillanova, A. Garau, F. Isaia, V. Lippolis and G. Verani, *Research Trends*, 1999, in the press and refs. therein.
- 20 L. Pauling, *The Nature of the Chemical Bond*, 3rd edn., Cornell University Press, Ithaca, NY, 1960.
- 21 F. Lux, R. Paetzold, J. Danel and L. Sobczyk, *J. Chem. Soc., Faraday Trans. 2*, 1975, 1610.
- 22 S. Kaur and T. S. Lobana, *J. Indian Chem. Soc.*, 1983, **60**, 126.
- 23 M. Arca, F. Cristiani, F. A. Devillanova, A. Garau, F. Isaia, V. Lippolis, G. Verani and F. Demartin, *Polyhedron*, 1997, **16**, 1983.
- 24 G. Carta, G. Crisponi and A. Lai, *J. Magn. Reson.*, 1982, **48**, 341.
- 25 A. C. North, D. C. Phillips and F. S. Mathews, *Acta Crystallogr., Sect. A*, 1968, **24**, 351.
- 26 P. Main, S. J. Fiske, S. E. Hull, L. Lessinger, G. Germain, J. P. Declercq and M. N. Wolfson, MULTAN 80, Universities of York and Louvain, 1980.
- 27 D. T. Cromer and J. T. Waber, *International Tables for X-Ray Crystallography*, Kynoch Press, Birmingham, 1974, vol. IV, Table 2.2B.
- 28 D. T. Cromer, *International Tables for X-Ray Crystallography*, Kynoch Press, Birmingham, 1974, vol. IV, Table 2.3.1.
- 29 B. Frenz, *Comput. Phys.*, 1988, **2**, No. 3 (May/June), 42–8; B. Frenz, *Crystallographic Computing 5*, Oxford University Press, 1991, ch. 11, pp. 126–135.

Paper 9/04271G