

# A kinetic study of the formation of a model high-valent diiron non-heme complex, $[\text{Fe}^{\text{III}}\text{Fe}^{\text{IV}}(\mu\text{-O})_2(\text{tpa})_2]^{3+}$ (tpa = tris(2-pyridylmethyl)amine), by cryogenic stopped-flow techniques†

Sergey V. Kryatov and Elena V. Rybak-Akimova\*

Department of Chemistry, Tufts University, Medford, Massachusetts 02155, USA

Received 9th June 1999, Accepted 18th August 1999

$[\text{Fe}_2(\mu\text{-O})(\text{tpa})_2(\text{OH})(\text{H}_2\text{O})]^{3+}$  reacts with  $\text{H}_2\text{O}_2$  in acetonitrile at  $-40^\circ\text{C}$  forming a transient diferric peroxo intermediate ( $\lambda_{\text{max}} = 700\text{ nm}$ ,  $\epsilon = 1800\text{ dm}^3\text{ mol}^{-1}\text{ cm}^{-1}$ ), which then reacts with a second  $\text{H}_2\text{O}_2$  molecule to give  $[\text{Fe}^{\text{III}}\text{Fe}^{\text{IV}}(\mu\text{-O})_2(\text{tpa})_2]^{3+}$ .

Non-heme diiron redox enzymes have been one of the ‘‘hot topics’’ of bioinorganic chemistry for the past decade. Their common mechanistic feature is the formation of a diiron(III) peroxo complex, which then transforms into a high-valent diiron species responsible for the oxidative action of an enzyme.<sup>1–4</sup> The actual need for modeling such a process on simple diiron complexes has been recently stated.<sup>1,2</sup> The only well characterized structural and functional models for the high-valent diiron species are  $\text{Fe}^{\text{III}}\text{Fe}^{\text{IV}}(\mu\text{-O})_2$  diamond core complexes with tpa and its derivatives, formed from corresponding  $\text{Fe}^{\text{III}}_2$ -complexes and  $\text{H}_2\text{O}_2$  at low temperature.<sup>2,5–7</sup> A relatively long-lived peroxo complex has been observed as an intermediate in the formation of  $[\text{Fe}^{\text{III}}\text{Fe}^{\text{IV}}(\mu\text{-O})_2(6\text{-Me}_3\text{-tpa})_2]^{3+}$  (6-Me<sub>3</sub>-tpa = tris(6-methyl-2-pyridylmethyl)amine),<sup>8</sup> with the general scheme similar to the peroxide shunt of diiron non-heme enzymes.<sup>1–3</sup> In this paper, we communicate the direct observation of a transient model diferric peroxo complex quickly converting into a high-valent species.

Mixing of  $[\text{Fe}^{\text{III}}_2(\mu\text{-O})(\text{tpa})_2(\text{OH})(\text{H}_2\text{O})]^{3+}$  and  $\text{H}_2\text{O}_2$  solutions in acetonitrile at  $-40.0 \pm 0.1^\circ\text{C}$  under pseudo-first order conditions (10-fold or larger excess of  $\text{H}_2\text{O}_2$ ) produces kinetic traces (Fig. 1), which are well fit by two exponential functions yielding wavelength-independent sequential rate constants  $k'_{\text{obs}}$  and  $k''_{\text{obs}}$ .† Treatment of the data by IS-2 Rapid Kinetics Software (Hi-Tech) using the  $\text{A} \rightarrow \text{B} \rightarrow \text{C}$  model gave electronic spectra of the three species involved (Fig. 1). The initial and final spectra agree well with those reported for  $[\text{Fe}^{\text{III}}_2(\mu\text{-O})(\text{tpa})_2(\text{OH})(\text{H}_2\text{O})]^{3+}$  and  $[\text{Fe}^{\text{III}}\text{Fe}^{\text{IV}}(\mu\text{-O})_2(\text{tpa})_2]^{3+}$ , correspondingly.<sup>5,7</sup> The spectrum of the intermediate  $\text{B}_{\text{peroxo}}$  has a broad maximum at about 700 nm ( $\epsilon = 1800\text{ dm}^3\text{ mol}^{-1}\text{ cm}^{-1}$ ), which is very similar to the  $\text{H}_{\text{peroxo}}$  intermediate of methane monooxygenase ( $\lambda_{\text{max}} = 700\text{ nm}$ ,  $\epsilon = 1800\text{ dm}^3\text{ mol}^{-1}\text{ cm}^{-1}$ )<sup>4</sup> and the peroxo intermediate of a mutant ribonucleotide reductase ( $\lambda_{\text{max}} = 700\text{ nm}$ ,  $\epsilon = 1500\text{ dm}^3\text{ mol}^{-1}\text{ cm}^{-1}$ ).<sup>9</sup> Such a band is characteristic of  $\text{Fe}^{\text{III}}_2(\mu\text{-peroxo})$  complexes<sup>10,11</sup> and is referred to LMCT.<sup>12</sup> A similar  $\text{Fe}^{\text{III}}_2(\mu\text{-O})(\mu\text{-O}_2)(6\text{-Me}_3\text{-tpa})_2^{2+}$  complex has been reported, which converts into  $[\text{Fe}^{\text{III}}\text{Fe}^{\text{IV}}(\mu\text{-O})_2(6\text{-Me}_3\text{-tpa})_2]^{3+}$  upon the addition of one equiv. of acid.<sup>8</sup> Thus, the species  $\text{B}_{\text{peroxo}}$  is most probably  $[\text{Fe}^{\text{III}}_2(\mu\text{-OH})(\mu\text{-O}_2)(\text{tpa})_2]^{3+}$  or  $[\text{Fe}^{\text{III}}_2(\mu\text{-O}_2)(\text{tpa})_2(\text{OH})(\text{H}_2\text{O})]^{3+}$ . It can be also a mixture of these and other diferric peroxo complexes.<sup>10</sup> The rate constant  $k'_{\text{obs}}$  corresponds to the formation of  $\text{B}_{\text{peroxo}}$ , and  $k''_{\text{obs}}$  to the transformation of  $\text{B}_{\text{peroxo}}$  into the  $\text{Fe}^{\text{III}}\text{Fe}^{\text{IV}}$ -species. Concentration dependences for the constants  $k'_{\text{obs}}$  and  $k''_{\text{obs}}$  allow us to propose a mechanistic scheme for these two consecutive steps (Scheme 1).

† Supplementary data available: plots of kinetic data. For direct electronic access see <http://www.rsc.org/suppdata/dt/1999/3335/>, otherwise available from BLDSC (No. SUP 57624, 4 pp.) or the RSC Library. See Instructions for Authors, 1999, Issue 1 (<http://www.rsc.org/dalton>).





Fig. 1 Spectra of the initial complex  $[\text{Fe}_2(\mu\text{-O})(\text{tpa})_2(\text{OH})(\text{H}_2\text{O})]^{3+}$  ( $2.5 \times 10^{-4}\text{ mol dm}^{-3}$ ) ( $\blacklozenge$ ), transient species  $\text{B}_{\text{peroxo}}$  ( $\blacksquare$ ) and  $[\text{Fe}^{\text{III}}\text{Fe}^{\text{IV}}(\mu\text{-O})_2(\text{tpa})_2]^{3+}$  ( $\blacktriangle$ ) calculated from kinetic data.  $[\text{H}_2\text{O}_2]_0 = 2.5 \times 10^{-3}\text{ mol dm}^{-3}$  (after mixing).



Scheme 1

The formation of the transient intermediate  $\text{B}_{\text{peroxo}}$  is decelerated by water and accelerated by hydrogen peroxide. Plots of kinetic data  $k'_{\text{obs}}^{-1}$  vs.  $[\text{H}_2\text{O}]$  at constant  $[\text{H}_2\text{O}_2]$  and  $k'_{\text{obs}}^{-1}$  vs.

$[\text{H}_2\text{O}_2]^{-1}$  at constant  $[\text{H}_2\text{O}]$  are straight lines with practically the same intercept. (SUP 57624) Such behavior suggests the formation of another intermediate ( $A_1$ ), which exists in equilibrium with the starting complex  $[\text{Fe}^{\text{III}}_2(\mu\text{-O})(\text{tpa})_2(\text{OH})(\text{H}_2\text{O})]^{3+}$  and converts irreversibly into  $B_{\text{peroxo}}$ . Spectral kinetic data (Fig. 1) show that the intermediate  $A_1$  does not accumulate in the reaction mixture to an appreciable amount, and that the reaction  $[\text{Fe}^{\text{III}}_2(\mu\text{-O})(\text{tpa})_2(\text{OH})(\text{H}_2\text{O})]^{3+} \rightarrow A_1$  is the rate-limiting step.

The simplest model to account for the observations is shown in Scheme 1. §


$$k'_{\text{obs}} = \frac{k_1(k_2/k_{-1})[\text{H}_2\text{O}_2]}{(k_2/k_{-1})[\text{H}_2\text{O}_2] + [\text{H}_2\text{O}]}$$

The constant  $k_1 = 0.5 \pm 0.1 \text{ s}^{-1}$  corresponds to the rate of the  $\text{H}_2\text{O}$  molecule dissociation in  $[\text{Fe}^{\text{III}}_2(\mu\text{-O})(\text{tpa})_2(\text{OH})(\text{H}_2\text{O})]^{3+}$ . The ratio  $k_2/k_{-1} = 6 \pm 2$  reflects the larger nucleophilicity of  $\text{H}_2\text{O}_2$  compared to  $\text{H}_2\text{O}$  ( $\alpha$ -effect).<sup>13</sup> The dissociative mechanism is common for  $\text{Fe}^{\text{III}}$  octahedral aqua species due to the labilizing effect of the  $\text{OH}^-$  ligand.<sup>14</sup> A similar action can be expected from the  $\text{O}^{2-}$  ligand. Thus, the intermediate  $A_1$  probably is  $[\text{Fe}^{\text{III}}_2(\mu\text{-O})(\text{tpa})_2(\eta^1\text{-OH})]^{3+}$  or  $[\text{Fe}^{\text{III}}_2(\mu\text{-O})(\mu\text{-OH})(\text{tpa})_2]^{3+}$ .<sup>15</sup> It can react with  $\text{H}_2\text{O}_2$  to form  $[\text{Fe}^{\text{III}}_2(\mu\text{-O})(\text{tpa})_2(\text{OH})(\text{H}_2\text{O}_2)]^{3+}$ , which then converts into  $[\text{Fe}^{\text{III}}_2(\mu\text{-O}_2)(\mu\text{-OH})(\text{tpa})_2]^{3+}$  or  $[\text{Fe}^{\text{III}}_2(\mu\text{-O}_2)(\text{tpa})_2(\text{OH})(\text{H}_2\text{O})]^{3+}$  by proton transfers. The apparent  $\varepsilon$  of  $B_{\text{peroxo}}$  (calculated from kinetic data) does not significantly depend on water concentration up to  $[\text{H}_2\text{O}]/[\text{H}_2\text{O}_2] = 40$ . It confirms the presence of a practically irreversible stage in the formation of the peroxo intermediate. Peroxide is apparently a much better ligand for  $\text{Fe}^{\text{III}}$  than oxide or hydroxide, as stable  $\text{Fe}^{\text{III}}(\text{O}_2^{2-})$ -complexes can form even in aqueous solution.<sup>16</sup>

The observed pseudo-first order rate constant ( $k''_{\text{obs}}$ ) of the following transformation  $B_{\text{peroxo}} \rightarrow [\text{Fe}^{\text{III}}\text{Fe}^{\text{IV}}(\mu\text{-O})_2(\text{tpa})_2]^{3+}$  does not depend on  $[\text{H}_2\text{O}]$ , but is proportional to  $[\text{H}_2\text{O}_2]$ .

$$k''_{\text{obs}} = k_3[\text{H}_2\text{O}_2]$$

The graph  $k''_{\text{obs}}$  vs.  $[\text{H}_2\text{O}_2]$  is a straight line with a practically zero intercept and a slope of  $k_3 = 10 \pm 2 \text{ dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$ , corresponding to a second-order rate constant for the reaction between  $B_{\text{peroxo}}$  and  $\text{H}_2\text{O}_2$  (SUP 57624). The reaction is apparently a redox process with tentative overall stoichiometry:



It should be noted that the presence of a reductant is essential for the action of native ribonucleotide reductase, presumably in order to reduce the  $\text{Fe}^{\text{III}}_2$ -peroxo intermediate to the high-valent  $\text{Fe}^{\text{III}}\text{Fe}^{\text{IV}}$  intermediate  $X$ .<sup>3</sup> In the model system studied here,  $\text{H}_2\text{O}_2$  is the most probable reductant.

The yields of  $[\text{Fe}^{\text{III}}\text{Fe}^{\text{IV}}(\mu\text{-O})_2(\text{tpa})_2]^{3+}$  have never been quantitative, reaching the maximum of ca. 70% (based on the reported value  $\varepsilon = 5500 \text{ dm}^3 \text{ mol}^{-1} \text{ cm}^{-1}$ )<sup>6</sup> at 40-fold or more excess of  $\text{H}_2\text{O}_2$ , which corresponds to  $[\text{H}_2\text{O}_2]_0 \geq 10 \text{ mM}$  under our conditions. The same maximum yield has been reported by Que's group with only 1.5-fold excess of  $\text{H}_2\text{O}_2$ , which corresponds also to ca. 10 mM  $\text{H}_2\text{O}_2$ .<sup>7</sup> These data can be rationalized by invoking an independent pathway of  $B_{\text{peroxo}}$  decomposition, which plays a larger role at insufficient absolute (rather than relative)  $\text{H}_2\text{O}_2$  concentrations.

At room temperature the  $[\text{Fe}_2(\mu\text{-O})(\text{tpa})_2(\text{OH})(\text{H}_2\text{O})]^{3+}$  complex causes intense effervescence of the  $\text{H}_2\text{O}_2$  acetonitrile solution. Significant catalytic activity has been reported for  $[\text{Fe}_2(\mu\text{-O})(\mu\text{-O}_2\text{CCH}_3)(\text{tpa})_2]^{2+}$  in such conditions.<sup>17</sup> To determine, if the excess of  $\text{H}_2\text{O}_2$  survives at low temperature, additional double-mixing experiments were carried out. The 5 mM  $\text{H}_2\text{O}_2$  solution was first mixed with 0.5 mM  $[\text{Fe}_2(\mu\text{-O})(\text{tpa})_2(\text{OH})(\text{H}_2\text{O})]^{3+}$  solution and then, upon the completion of apparent spectral changes, with a fresh portion of 0.25 mM  $[\text{Fe}_2(\mu\text{-O})(\text{tpa})_2(\text{OH})(\text{H}_2\text{O})]^{3+}$  solution. Kinetic quantitation showed that no more than 2 moles of  $\text{H}_2\text{O}_2$  per mole of the  $\text{Fe}_2$ -complex had been consumed in the first mixing. Thus, the

catalytic process in this system is negligible and suppressed at  $-40^\circ\text{C}$ .

In the other series of double-mixing experiments the  $[\text{Fe}_2(\mu\text{-O})(\text{tpa})_2(\text{OH})(\text{H}_2\text{O})]^{3+}$  solution was first mixed with  $\text{H}_2\text{O}_2$  solution, and upon the completion of apparent spectral changes, with a fresh portion of more concentrated  $\text{H}_2\text{O}_2$  solution. It was found that the ratio  $\text{H}_2\text{O}_2:\text{Fe}_2 \geq 1.5 \pm 0.1$  was enough to consume all of the initial diiron complex and prevent the formation of new  $B_{\text{peroxo}}$  in the second mixing. This confirms the presence of an irreversible stage in the formation of  $B_{\text{peroxo}}$  and suggests that the starting complex  $[\text{Fe}_2(\mu\text{-O})(\text{tpa})_2(\text{OH})(\text{H}_2\text{O})]^{3+}$  is not regenerated in the system. The stoichiometry of the initial interaction between  $[\text{Fe}_2(\mu\text{-O})(\text{tpa})_2(\text{OH})(\text{H}_2\text{O})]^{3+}$  and  $\text{H}_2\text{O}_2$  to give  $B_{\text{peroxo}}$  is most probably 1:1, with some  $\text{H}_2\text{O}_2$  consumed in the concomitant formation of the  $\text{Fe}^{\text{III}}\text{Fe}^{\text{IV}}$  complex and in a minor catalytic process.

Other kinetic experiments regarding the formation and reactivity of  $\text{Fe}_2\text{O}_2(\text{H})$  diamond cores are currently under way.

## Acknowledgements

This work was supported by Tufts University.

## Notes and references

‡ The initial complex,  $[\text{Fe}_2(\mu\text{-O})(\text{tpa})_2(\text{OH})(\text{H}_2\text{O})](\text{ClO}_4)_3$  was prepared by a published procedure.<sup>5</sup> Hydrogen peroxide, 30% aqueous solution (ACS certified grade) and acetonitrile (HPLS grade) were purchased from Fisher. Stopped-flow experiments were carried out by using a Hi-Tech Scientific (Salisbury, UK) SF-43 cryogenic stopped-flow apparatus with stainless steel plumbing. Measurements were made at wavelengths from 360 to 800 nm with 20 nm intervals to get spectral information.

§ More complicated molecular interpretations can also describe the observations. However, another simple model, with the fast preequilibrium  $[\text{Fe}_2(\mu\text{-O})(\text{tpa})_2(\text{OH})(\text{H}_2\text{O})]^{3+} + \text{H}_2\text{O}_2 \rightleftharpoons A_1 + \text{H}_2\text{O}$  and the rate-limiting step  $A_1 \rightarrow B_{\text{peroxo}}$ , can be discounted. It implies that  $A_1$  is an  $\text{Fe}^{\text{III}}$ -peroxo complex, which is formed in substantial amount in the reaction mixture (the preequilibrium constant  $K_{\text{eq}} = 6 \pm 2$ ). Such an assumption contradicts the spectral changes calculated from kinetic data.

- 1 A. M. Valentine and S. J. Lippard, *J. Chem. Soc., Dalton Trans.*, 1997, 3925.
- 2 L. Que, Jr., *J. Chem. Soc., Dalton Trans.*, 1997, 3933.
- 3 B. J. Wallar and J. D. Lipscomb, *Chem. Rev.*, 1996, **96**, 2625.
- 4 A. M. Valentine, S. S. Stahl and S. J. Lippard, *J. Am. Chem. Soc.*, 1999, **121**, 3876.
- 5 Y. Dong, H. Fujii, M. P. Hendrich, R. A. Leising, G. Pan, C. R. Randall, E. C. Wilkinson, Y. Zang, L. Que, Jr., B. G. Fox, K. Kauffmann and E. Münck, *J. Am. Chem. Soc.*, 1995, **117**, 2778; H.-F. Hsu, Y. Dong, L. Shu, V. G. Young, Jr. and L. Que, Jr., *J. Am. Chem. Soc.*, 1999, **121**, 5230.
- 6 C. Kim, Y. Dong and L. Que, Jr., *J. Am. Chem. Soc.*, 1997, **119**, 3635.
- 7 R. A. Leising, B. A. Brennan, L. Que, Jr., B. G. Fox and E. Münck, *J. Am. Chem. Soc.*, 1991, **113**, 3988.
- 8 Y. Dong, Y. Zang, L. Shu, E. C. Wilkinson, L. Que, Jr., K. Kauffmann and E. Münck, *J. Am. Chem. Soc.*, 1997, **119**, 12683.
- 9 J. M. Bollinger, Jr., C. Krebs, A. Vicol, S. Chen, B. A. Ley, D. E. Edmondson and B. H. Huynh, *J. Am. Chem. Soc.*, 1998, **120**, 1094; P. Moënne-Loccoz, J. Baldwin, B. A. Ley, T. M. Loehr and J. M. Bollinger, Jr., *Biochemistry*, 1998, **37**, 14659.
- 10 C. Duboc-Toia, S. Ménage, R. Y. N. Ho, L. Que, Jr., C. Lambeaux and M. Fontecave, *Inorg. Chem.*, 1999, **38**, 1261.
- 11 H. Sugimoto, T. Nagayama, S. Maruyama, S. Fujinami, Y. Yasuda, M. Suzuki and A. Uehara, *Bull. Chem. Soc. Jpn.*, 1998, **71**, 2267.
- 12 T. C. Brunold, N. Tamura, N. Kitajima, Y. Moro-oka and E. I. Solomon, *J. Am. Chem. Soc.*, 1998, **120**, 5674.
- 13 S. Hoz and E. Bunzel, *Isr. J. Chem.*, 1985, **26**, 313.
- 14 J. O. Edwards, F. Monacelli and G. Ortaggi, *Inorg. Chim. Acta*, 1974, **11**, 47; A. E. Merbach, *Pure Appl. Chem.* 1987, **59**, 161.
- 15 H. Zheng, Y. Zang, Y. Dong, V. G. Young, Jr. and L. Que, Jr., *J. Am. Chem. Soc.*, 1999, **121**, 2226.
- 16 F. Neese and E. I. Solomon, *J. Am. Chem. Soc.*, 1998, **120**, 12829.
- 17 T. Okuno, S. Ito, S. Ohba and Y. Nishida, *J. Chem. Soc., Dalton Trans.*, 1997, 3547.

Communication 9/04626G