

Preparation of new diazene complexes of ruthenium and osmium †

Gabriele Albertin,^{*a} Stefano Antoniutti,^a Alessia Bacchi,^b Mara Boato^a and Giancarlo Pelizzi^b

^a Dipartimento di Chimica, Università Ca' Foscari di Venezia, Dorsoduro 2137, 30123 Venezia, Italy

^b Dipartimento di Chimica Generale ed Inorganica, Chimica Analitica, Chimica Fisica, Università di Parma, Parco Area delle Scienze 17/a, 43100 Parma, Italy

Received 21st March 2002, Accepted 25th June 2002

First published as an Advance Article on the web 9th August 2002

The hydrazine complexes $[\text{Ru}(\text{NH}_2\text{NH}_2)\text{L}_5](\text{BPh}_4)_2$ **1**, **8** or $[\text{Ru}(\text{NH}_2\text{NH}_2\text{L}'\text{L}_4)(\text{BPh}_4)_2$ **2** [$\text{L} = \text{P}(\text{OEt})_3$, $\text{L}' = \text{PPh}(\text{OEt})_2$] were prepared by allowing the corresponding hydride species $[\text{RuHL}_5]\text{BPh}_4$ or $[\text{RuHL}'\text{L}_4]\text{BPh}_4$ to react first with HBF_4 and then with hydrazine. Oxidation of these hydrazine complexes or $[\text{Os}(\text{NH}_2\text{NH}_2)\text{L}_5](\text{BPh}_4)_2$ with $\text{Pb}(\text{OAc})_4$ at -30°C led to the corresponding stable and isolable 1,2-diazene complexes $[\text{M}(\text{NH}=\text{NH})\text{L}_5](\text{BPh}_4)_2$ **3**, **5** ($\text{M} = \text{Ru}$, Os) or $[\text{Ru}(\text{NH}=\text{NH})\text{L}'\text{L}_4](\text{BPh}_4)_2$ **4**. The phenyldiazene derivatives $[\text{M}(\text{PhN}=\text{NH})\text{L}_5](\text{BPh}_4)_2$ **6**, **7** ($\text{M} = \text{Ru}$, Os) were also prepared by treating the hydride $[\text{MHL}_5]\text{BPh}_4$ species with phenyldiazonium tetrafluoroborate. The aquo-complex, $[\text{Ru}(\text{H}_2\text{O})\{\text{P}(\text{OEt})_3\}_5](\text{BPh}_4)_2$ **8**, was obtained by substitution of the $\text{NH}=\text{NH}$ ligand in $[\text{Ru}(\text{NH}=\text{NH})\text{L}_5](\text{BPh}_4)_2$ and was characterised by X-ray crystal structure determination. Oxidation reactions of the bis(hydrazine) complexes $[\text{Ru}(\text{NH}_2\text{NH}_2)_2\text{L}_4](\text{BPh}_4)_2$ or $[\text{Ru}(\text{CH}_3\text{NHNH}_2)_2\text{L}_4](\text{BPh}_4)_2$ [$\text{L} = \text{P}(\text{OEt})_3$] with $\text{Pb}(\text{OAc})_4$ were reinvestigated and were found to give the diazene complexes $[\text{Ru}(\text{NH}=\text{NH})_2\text{L}_4](\text{BPh}_4)_2$ **9**, $[\text{Ru}(\text{CH}_3\text{N}=\text{NH})_2\text{L}_4](\text{BPh}_4)_2$ **11** or $[\text{Ru}(\text{CH}_3\text{N}=\text{NH})(\text{CH}_3\text{NHNH}_2)\text{L}_4](\text{BPh}_4)_2$ **10**.

Introduction

Diazene $\text{NH}=\text{NH}$ is a very reactive molecule in the free state, and disproportionates¹ at -150°C into N_2 and NH_2NH_2 . Nevertheless, it is an important molecule of possible relevance as a metal-bound intermediate in inorganic and bioinorganic N_2 fixation processes,² and useful as a reagent in stereoselective *cis*-hydrogenation of unsaturated organic compounds.³ Stabilisation of $\text{NH}=\text{NH}$ in standard conditions has only been achieved in a few cases by coordination to transition metals, mainly in bimetallic complexes containing a μ - $\text{NH}=\text{NH}$ ligand.⁴ Stable derivatives containing monodentate diazene have also been reported.⁵

We are interested in the chemistry of partially reduced dinitrogen ligands, and have previously reported⁶ the synthesis of both methylidiazene and arylidiazene complexes of ruthenium and osmium, obtained by oxidation of the related hydrazine derivatives. However, no evidence of formation of stable complexes containing coordinated 1,2-diazene was obtained. We have therefore extended those studies, searching for an appropriate metal fragment stabilising the diazene $\text{NH}=\text{NH}$ molecule by coordination, and results are reported here. Reinvestigation of our previous reports^{6a,c} on oxidation of bis(hydrazine) $[\text{M}(\text{NH}_2\text{NH}_2)_2\text{L}_4]^{2+}$ ($\text{M} = \text{Ru}$, Os) derivatives, which allows bis(1,2-diazene) complexes to be prepared, is also described here.

Experimental

All synthetic work was carried out in an inert atmosphere using standard Schlenk techniques or a Vacuum Atmosphere dry-box. Once isolated, the complexes were found to be relatively stable in air, but were stored in an inert atmosphere at -25°C .

† Electronic supplementary information (ESI) available: observed and calculated $^{31}\text{P}\{^1\text{H}\}$ NMR spectra of compounds $[\text{Ru}(\text{NH}_2\text{NH}_2)\{\text{PPh}(\text{OEt})_2\}\{\text{P}(\text{OEt})_3\}_4](\text{BPh}_4)_2$ **2** (Fig. S1) and $[\text{Ru}(\text{NH}=\text{NH})\{\text{PPh}(\text{OEt})_2\}\{\text{P}(\text{OEt})_3\}_4](\text{BPh}_4)_2$ **4** (Fig. S2). See <http://www.rsc.org/suppdata/dt/b2/b202888n/>

All solvents were dried over appropriate drying agents, degassed on a vacuum line and distilled into vacuum-tight storage flasks. The phosphite $\text{P}(\text{OEt})_3$ (Aldrich) was purified by distillation under nitrogen; $\text{PPh}(\text{OEt})_2$ was prepared by the method of Rabinowitz and Pellon.⁷ Diazonium salts were obtained in the usual way.⁸ Hydrazine NH_2NH_2 was prepared by decomposition of hydrazine cyanurate (Fluka) following the reported method.⁹ $\text{RuCl}_3 \cdot 3\text{H}_2\text{O}$ salt was a ChemPur product and $(\text{NH}_4)_2\text{OsCl}_6$ was purchased from Johnson Matthey; both salts were used as received. High-grade (99.99%) lead(IV) acetate was purchased from Aldrich. Other reagents were purchased from commercial sources in the highest available purity and used as received. Infrared spectra were recorded on a Nicolet Magna 750 FT-IR spectrophotometer. NMR spectra (^1H , ^{31}P) were obtained on a Bruker AC200 spectrometer at temperatures varying between -90 and $+30^\circ\text{C}$, unless otherwise noted. ^1H spectra are referred to internal tetramethylsilane. $^{31}\text{P}\{^1\text{H}\}$ chemical shifts are reported with respect to 85% H_3PO_4 , with downfield shifts considered positive. The conductivity of 10^{-3} mol dm^{-3} solutions of the complexes in CH_3NO_2 at 25°C was measured with a Radiometer CDM 83 instrument.

Preparation of complexes

Hydrides RuH_2L_4 [$\text{L} = \text{P}(\text{OEt})_3$, $\text{PPh}(\text{OEt})_2$] and $[\text{OsH}\{\text{P}(\text{OEt})_3\}_5]\text{BPh}_4$ were prepared following previous methods.^{6a,10}

$[\text{RuHL}_5]\text{BPh}_4$ [$\text{L} = \text{P}(\text{OEt})_3$ and $\text{PPh}(\text{OEt})_2$]. Hydrides $[\text{RuHL}_5]\text{BPh}_4$ were prepared by a modification of the method previously reported.¹¹ An equimolar amount of the appropriate phosphite (0.56 mmol) was added to a solution of RuH_2L_4 (0.56 mmol) in 5 cm^3 of ethanol and the mixture cooled to -196°C . A slight excess of $\text{HBF}_4 \cdot \text{Et}_2\text{O}$ (0.57 mmol, 82 μL) was then added and the reaction mixture, brought to room temperature, stirred for about 2 h. The addition of an excess of NaBPh_4 (1.2 mmol, 0.41 g) caused the separation of a white solid, which was filtered and crystallised from CH_2Cl_2 and ethanol; yield $\geq 80\%$. (For $[\text{RuH}\{\text{P}(\text{OEt})_3\}_5]\text{BPh}_4$ Found: C, 51.64; H, 7.85. $\text{C}_{54}\text{H}_{96}\text{BO}_{15}\text{P}_5\text{Ru}$ requires C, 51.80; H, 7.73%. $M_{\text{M}} = 56.2$ S cm^2

mol⁻¹. IR (KBr) ν/cm^{-1} : 1915 (m) $\nu(\text{Ru-H})$. δ_{H} [(CD₃)₂CO, 25 °C]: 7.40–6.78 (m, 20 H, Ph), 4.11 (m, 30 H, CH₂), 1.31, 1.30 (t, 45 H, CH₃), -9.25 to -10.03 (m, 1 H, H⁻). δ_{P} [(CD₃)₂CO, 25 °C]: A₄B spin system, δ_{A} 142.0, δ_{B} 140.4, J_{AB} = 46.6 Hz).

[RuH{PPh(OEt)₂}₃{P(OEt)₃}₄](BPh₄)₂. A slight excess of HBF₄·Et₂O (0.42 mmol, 60 μL) was added to a solution of RuH₂{P(OEt)₃}₄ (0.4 mmol, 0.31 g) in 10 cm³ of C₂H₅OH and the reaction mixture was brought to about -30 °C and stirred for 30 min. A slight excess of PPh(OEt)₂ (0.42 mmol, 83 μL) was added and the resulting solution brought to room temperature. After 1 h of stirring, an excess of NaBPh₄ (0.8 mmol, 0.27 g) was added and the white solid obtained filtered and crystallised from CH₂Cl₂ and ethanol; yield \geq 80%. (Found: C, 54.34; H, 7.45. C₅₈H₉₆BO₁₄P₅Ru requires C, 54.25; H, 7.53%. A_{M} = 57.5 S cm² mol⁻¹. IR (KBr) ν/cm^{-1} : 1936 (w) $\nu(\text{Ru-H})$. δ_{H} (CD₂Cl₂, 25 °C): 8.01–6.70 (m, 25 H, Ph), 4.20–3.50 (m, 28 H, CH₂), 1.26 (m, 42 H, CH₃), -8.81 to -9.70 (m, 1 H, H⁻). δ_{P} (CD₂Cl₂, 25 °C): ABC₂M spin system, δ_{M} 166.4, δ_{A} 141.0, δ_{B} 140.1, δ_{C} 137.6, J_{AM} = -435.7, J_{BM} = 56.0, J_{CM} = 31.6, J_{AB} = 67.7, J_{AC} = 38.3, J_{BC} = 55.4 Hz).

[Ru(NH₂NH₂)₂{P(OEt)₃}₅](BPh₄)₂ **1**. An excess of HBF₄·Et₂O (0.4 mmol, 57 μL) was added to a solution of hydride [RuHL₅](BPh₄) (0.125 g, 0.1 mmol) in 10 cm³ of CH₂Cl₂ cooled to -196 °C and placed under a hydrogen atmosphere. The reaction mixture was brought to room temperature, stirred for 2 h and then, after replacing the H₂ atmosphere with argon, an excess of NH₂NH₂ (1 mmol, 31 μL) was added. The resulting solution was stirred for 20 h and then the solvent was removed under reduced pressure. The oil obtained was triturated with ethanol containing an excess of NaBPh₄ (0.2 mmol, 68 mg) giving a white solid, which was filtered and crystallised from CH₂Cl₂ and ethanol; yield \geq 60%. (Found: C, 58.35; H, 7.58; N, 1.72. C₇₈H₁₁₉B₂N₂O₁₅P₅Ru requires C, 58.47; H, 7.49; N, 1.75%. A_{M} = 114.9 S cm² mol⁻¹).

[Ru(NH₂NH₂)₂{PPh(OEt)₂}₃{P(OEt)₃}₄](BPh₄)₂ **2**. This compound was prepared exactly like the related pentakis(phosphite) complex **1** by reacting the hydride [RuH{PPh(OEt)₂}₂{P(OEt)₃}₄](BPh₄) first with HBF₄·Et₂O and then with NH₂NH₂; yield \geq 65%. (Found: C, 60.39; H, 7.45; N, 1.76. C₈₂H₁₁₉B₂N₂O₁₄P₅Ru requires C, 60.26; H, 7.34; N, 1.71%. A_{M} = 115.1 S cm² mol⁻¹).

[Ru(NH₂NH₂)₂{PPh(OEt)₂}₄](BPh₄)₂. This complex was obtained in an attempt to prepare the pentakis(phosphite) [Ru(NH₂NH₂)₂{PPh(OEt)₂}₅](BPh₄)₂ following the method reported for the related compounds **1** and **2**. In this case, the known^{6a} bis(hydrazine) derivative was obtained as the only isolated product, with a yield of about 75%. (Found: C, 66.12; H, 6.95; N, 3.45. C₈₈H₁₀₈B₂N₄O₈P₄Ru requires C, 66.21; H, 6.82; N, 3.51%. A_{M} = 114.7 S cm² mol⁻¹).

[Os(NH₂NH₂)₂L₅](BPh₄)₂ [L = P(OEt)₃]. This complex was prepared following the method previously reported.^{6c}

[Ru(NH=NH){P(OEt)₃}₅](BPh₄)₂ **3** and **[Os(NH=NH){P(OEt)₃}₅](BPh₄)₂** **5**. A sample of the appropriate hydrazine complex [M(NH₂NH₂)₂L₅](BPh₄)₂ (0.16 mmol) was placed in a three-necked 25 cm³ round-bottomed flask fitted with a solid-addition sidearm containing an equimolar amount of Pb(OAc)₄ (0.16 mmol, 71 mg). Dichloromethane (10 cm³) was added, the solution cooled to -30 °C and Pb(OAc)₄ added portionwise over 20–30 min to the cold stirring solution. The solution was then brought to 0 °C, stirred for 10 min and the solvent removed under reduced pressure. The oil obtained was treated at 0 °C with ethanol (2 cm³) containing an excess of NaBPh₄ (0.2 mmol, 68 mg). A white solid slowly separated out,

which was filtered and crystallised by dissolving in CH₂Cl₂ and, after filtration and concentration, by fast precipitation with ethanol; yield \geq 60%. (Found: C, 58.70; H, 7.47; N, 1.70. C₇₈H₁₁₇B₂N₂O₁₅P₅Ru **3** requires C, 58.54; H, 7.37; N, 1.75%. A_{M} = 118.4 S cm² mol⁻¹. Found: C, 55.38; H, 7.04; N, 1.75. C₇₈H₁₁₇B₂N₂O₁₅OsP₅ **5** requires C, 55.45; H, 6.98; N, 1.66%. A_{M} = 120.7 S cm² mol⁻¹).

[Ru(NH=NH){PPh(OEt)₂}₃{P(OEt)₃}₄](BPh₄)₂ **4**. This complex was prepared exactly like the related species **3** and **5** by oxidation with Pb(OAc)₄ of the starting hydrazine compound **2**; yield \geq 45%. (Found: C, 60.16; H, 7.30; N, 1.61. C₈₂H₁₁₇B₂N₂O₁₄P₅Ru requires C, 60.33; H, 7.22; N, 1.72%. A_{M} = 116.9 S cm² mol⁻¹).

[Ru(PhN=NH){P(OEt)₃}₅](BPh₄)₂ **6** and **[Os(PhN=NH){P(OEt)₃}₅](BPh₄)₂** **7**. In a three-necked 25 cm³ round-bottomed flask were placed solid samples of the appropriate hydrides [MHL₅](BPh₄) (0.1 mmol) and an excess of phenyldiazonium tetrafluoroborate [PhN₂]⁺BF₄⁻ (0.5 mmol, 96 mg) and the mixture was cooled to -196 °C. Acetone (10 cm³) was slowly added and the reaction mixture, brought to room temperature, stirred for 4 h. The solvent was removed under reduced pressure giving a brown oil which was treated with ethanol (5 cm³) containing an excess of NaBPh₄ (0.2 mmol, 68 mg). A yellow solid slowly separated out, which was filtered and crystallised from CH₂Cl₂ and ethanol; yield \geq 75%. (Found: C, 60.02; H, 7.35; N, 1.60. C₈₄H₁₂₁B₂N₂O₁₅P₅Ru **6** requires C, 60.18; H, 7.27; N, 1.67%. A_{M} = 120.3 S cm² mol⁻¹. Found: C, 57.01; H, 6.99; N, 1.67. C₈₄H₁₂₁B₂N₂O₁₅OsP₅ **7** requires C, 57.14; H, 6.91; N, 1.59%. A_{M} = 117.8 S cm² mol⁻¹).

[Ru(H₂O){P(OEt)₃}₅](BPh₄)₂ **8**. This complex was obtained in an attempt at crystallisation of the diazene derivative [Ru(NH=NH){P(OEt)₃}₅](BPh₄)₂ **3**. In fact, by slow cooling to -25 °C of a saturated solution of **3**, prepared by treating the solid sample (150 mg) with ethanol (8 cm³) and enough CH₂Cl₂ to obtain a saturated solution at room temperature, white microcrystals of the aquo-complex **8** were obtained. These crystals were also suitable for X-ray analysis; yield \geq 80%. (Found: C, 59.11; H, 7.49. C₇₈H₁₁₇B₂O₁₆P₅Ru requires C, 58.98; H, 7.42%. A_{M} = 118.2 S cm² mol⁻¹).

[Ru(CH₃N=NH)(CH₃NHNH₂)₂{P(OEt)₃}₄](BPh₄)₂ **10** and **[Ru(CH₃N=NH)₂{P(OEt)₃}₄](BPh₄)₂** **11**. These complexes were obtained by oxidation of the [Ru(CH₃NHNH₂)₂{P(OEt)₃}₄](BPh₄)₂ complex^{6a} with Pb(OAc)₄ in CH₂Cl₂ at -30 °C following the method used for **3** and **5**. In this case, by reacting 0.2 mmol (0.300 g) of [Ru(CH₃NHNH₂)₂{P(OEt)₃}₄](BPh₄)₂ with 0.4 mmol (0.177 g) of Pb(OAc)₄, a mixture of both compounds **10** and **11** was obtained. Their separation involves the slow cooling to -25 °C of a saturated solution of the mixture prepared at room temperature using as solvent ethanol and enough CH₂Cl₂ to dissolve the solid. The first solid obtained was the mixed-ligand compound **10** in about 25% yield, while from the mother liquor, after repeated crystallisation, the bis(methyldiazene) **11** was separated in about 10% yield. (Found: C, 59.31; H, 7.48; N, 3.66. C₇₄H₁₁₀B₂N₄O₁₂P₄Ru **10** requires C, 59.48; H, 7.42; N, 3.75%. A_{M} = 118.5 S cm² mol⁻¹. Found: C, 59.42; H, 7.41; N, 3.63. C₇₄H₁₀₈B₂N₄O₁₂P₄Ru **11** requires C, 59.56; H, 7.29; N, 3.75%. A_{M} = 108.9 S cm² mol⁻¹).

Oxidation of [Ru(NH₂NH₂)₂L₄](BPh₄)₂ [L = P(OEt)₃ and PPh(OEt)₂] and [Os(NH₂NH₂)₂{P(OEt)₃}₄](BPh₄)₂

Also the oxidation of the bis(hydrazine) complexes^{6a,c} was carried out with Pb(OAc)₄ at -30 °C, following the method used for the other hydrazine complexes **1**, **2**, [Os(NH₂NH₂)₂L₅](BPh₄)₂ and [Ru(CH₃NHNH₂)₂L₄](BPh₄)₂. A typical experiment involves the addition of solid Pb(OAc)₄ (0.4 mmol,

Table 1 Crystal data and structure refinement for [Ru(H₂O){P(OEt)₃}₅](BPh₄)₂ **8**

Empirical formula	C ₇₈ H ₁₁₇ B ₂ O ₁₆ P ₅ Ru
Formula weight	1588.26
Temperature/K	293(2)
Wavelength/Å	0.71069
Crystal system	Triclinic
Space group	<i>P</i> $\bar{1}$
<i>a</i> /Å	14.763(5)
<i>b</i> /Å	16.613(5)
<i>c</i> /Å	18.646(5)
β ^o	82.05(5)
Volume/Å ³	4529(2)
<i>Z</i>	2
Absorption coefficient/mm ⁻¹	0.318
<i>F</i> (000)	1684
Reflections collected	26755
Independent reflections	18872 [<i>R</i> (int) = 0.0555]
Data/restraints/parameters	18872/6/845
Final <i>R</i> indices [<i>I</i> > 2σ(<i>I</i>)]	<i>R</i> 1 = 0.0948, <i>wR</i> 2 = 0.2383
<i>R</i> indices (all data)	<i>R</i> 1 = 0.1649, <i>wR</i> 2 = 0.2968

0.177 g) to a cooled solution (−30 °C) of the appropriate bis(hydrazine) complex (0.2 mmol) in CH₂Cl₂. Removal of the solvent at the end of the reaction gave an oil, which was triturated at 0 °C with ethanol (2 cm³) containing an excess of NaBPh₄ (0.4 mmol, 136 mg). The white solid that slowly separated out was filtered and dried under vacuum. The samples contained a mixture of bis(diazene) [M(NH=NH)₂-L₄](BPh₄)₂ and acetate [M(κ²-O₂CCH₃)₂L₄]BPh₄ complexes, which cannot be separated in pure form. However, in the case of [Ru(NH=NH)₂{P(OEt)₃}₄](BPh₄)₂ **9**, the sample contains the bis(diazene) as the major product (about 60–70%), while in the other cases only little amounts (5–10%) of bis(diazene) are present in the reaction product. For the mixture containing [Ru(NH=NH)₂{PPh(OEt)₂}₄](BPh₄)₂ δ_H (CD₂Cl₂, 20 °C): 16.23, 15.62 (dm, 2 H, NH=NH; ³J_{HH} = 32.0 Hz). For the mixture containing [Os(NH=NH)₂{P(OEt)₃}₄](BPh₄)₂ δ_H (CD₂Cl₂, 20 °C): 16.63, 16.31 (dm, 2 H, NH=NH).

Crystallography


Crystals were air sensitive, a white irregular prism single crystal was sealed in a glass capillary along with some drops of solution. X-Ray diffraction data were collected on a Bruker-Siemens SMART AXS 1000 equipped with a CCD detector, using graphite monochromated Mo-Kα radiation (λ = 0.71069 Å). Data collection details are: crystal to detector distance = 5.0 cm, hemisphere mode, time per frame = 30 s, oscillation Δω = 0.300°. Data reduction was performed up to θ = 28° by the SAINT package¹² and data were corrected for absorption effects by the SADABS¹³ procedure. Data collection and refinement results are summarised in Table 1. The crystal suffered a severe merohedral twinning expressed by a mirror plane perpendicular to *b*, which simulated a 2/*m* Laue symmetry, whilst the actual structure is described in *P* $\bar{1}$, with *a* and *β* close to 90°. The phase problem was solved by direct methods¹⁴ and the structure was refined by full matrix least squares on all *F*² by taking into account the twinning as implemented in Shelxl 97,¹⁵ using the WinGX package.¹⁶ The refinement without considering the twinning was unsuccessful. Anisotropic displacement parameters were refined for all non-hydrogen atoms, while hydrogen atoms were introduced in calculated positions, except for those belonging to the water, located on the Fourier maps. Phenyls were treated as rigid bodies. Use of the Cambridge Crystallographic Database¹⁷ facilities was made for structure discussion. Meaningless electron density residues were left around the metal in the final map.

CCDC reference number 182442.

See <http://www.rsc.org/suppdata/dt/b2/b202888n/> for crystallographic data in CIF or other electronic format.


Results and discussion

Hydrazine complexes of ruthenium [Ru(NH₂NH₂){P(OEt)₃}₅](BPh₄)₂ **1** and [Ru(NH₂NH₂){PPh(OEt)₂}{P(OEt)₃}₄](BPh₄)₂ **2** were prepared by reacting the hydride [RuHL'L₄]⁺ first with an excess of fluoroboric acid and then with hydrazine, as shown in Scheme 1.



Scheme 1 L' = L = P(OEt)₃ **1**; L' = PPh(OEt)₂, L = P(OEt)₃ **2**.

Reactions of the pentakis(phosphite)hydride [RuHL'L₄]⁺ with an excess of HBF₄ gives the η²-H₂ dihydrogen intermediate [M(η²-H₂)L'L₄]²⁺ (by ¹H NMR), which reacts with NH₂NH₂ to give final hydrazine complexes **1** or **2**. In contrast, the reaction of the hydride [RuH{PPh(OEt)₂}₅]BPh₄, first with HBF₄ and then with hydrazine, in all conditions afforded the known^{6a} bis(hydrazine) complex [Ru(NH₂NH₂)₂{PPh(OEt)₂}₄]²⁺, as shown in Scheme 2.



Scheme 2 L = PPh(OEt)₂.

In the pentakis(phosphite) [Ru(η²-H₂)L₅]²⁺ intermediate, hydrazine replaces not only the labile η²-H₂ ligand, but also one phosphite, giving the bis(hydrazine) species. Attempts to prepare other pentakis(phosphite) hydrazine complexes different from **1** and **2** failed, because the reaction of [RuHL'L₄]⁺ first with a Brønsted acid and then with an excess of CH₃NHNH₂ or C₆H₅NHNH₂ only gave mixtures of solid products not containing the hydrazine ligand. It seems that only the NH₂NH₂ species can bind to the metal centre in the ML'L₄ fragment to give stable and isolable derivatives.

Hydrazine complexes **1** and **2** were isolated as white [BPh₄]⁻ salts, stable in air and in solutions of polar organic solvents, in which they behave as 2 : 1 electrolytes. Analytical and spectroscopic data (Table 2) support the proposed formulation. The presence of the hydrazine ligand was supported by infrared spectroscopy, which showed the characteristic νNH medium-intensity bands of NH₂NH₂ between 3387 and 3266 cm⁻¹. Further support for the presence of NH₂NH₂ came from ¹H NMR spectra, which showed two broad signals at 4.37 and 3.36 ppm (**1**) or at 4.41 and 3.46 ppm (**2**), due to the two NH₂ proton resonances. Integration measurements and homodecoupling experiments confirmed the proposed assignment. In the temperature range between +30 and −80 °C, the ³¹P{¹H} NMR spectrum of [M(NH₂NH₂)L₅]²⁺ **1** appeared as an AB₄ multiplet, fitting the proposed formulation (geometry **I**, Fig. 1). In contrast, the spectra of [Ru(NH₂NH₂){PPh(OEt)₂}{P(OEt)₃}₄](BPh₄)₂ **2** showed a rather complicated pattern, simulated as a set of two systems of AB₄ and ABC₂M, partly overlapping, and with the parameters listed in Table 2. This result may be interpreted as due to two geometries in solution of the type shown in Fig. 1, with the hydrazine and PPh(OEt)₂ ligands in mutually *trans* (**II**) or *cis* (**III**) positions, respectively.

Pentakis(phosphite)hydrazine complexes of both ruthenium **1**, **2** and osmium^{6c} [Os(NH₂NH₂)L₅](BPh₄)₂ reacted with Pb(OAc)₄ at low temperature (−30 °C) with the selective oxidation of the NH₂NH₂ ligand to give 1,2-diazene complexes [M(NH=NH)L₅]²⁺ (M = Ru **3**, Os **5**) and [Ru(NH=NH){PPh-



Table 2 IR and NMR data for ruthenium and osmium complexes

Compound	IR ^a		¹ H NMR ^{b,c}		Spin system	³¹ P- ¹ H NMR ^{b,d}				
	ν/cm^{-1}	Assignment	$\delta(\text{J/Hz})$	Assignment		$\delta(\text{J/Hz})$				
1 [Ru(NH ₂ NH ₂)- {P(OEt) ₃ } ₃](BPh ₄) ₂	3368 m	$\nu(\text{NH})$	4.37 s, br	RuNH ₂	AB ₄	δ_{A} 131.9 δ_{B} 120.9 $J_{\text{AB}} = 58.3$				
	3328 m		4.07 m	CH ₂						
	3275 w		3.97 qnt							
	3267 w		3.36 s, br	NH ₂						
2 [Ru(NH ₂ NH ₂){PPh(OEt) ₂ }- {P(OEt) ₃ } ₄](BPh ₄) ₂	3387 w	$\nu(\text{NH})$	4.41 s, br	RuNH ₂	AB ₄	δ_{A} 160.7 δ_{B} 120.4 $J_{\text{AB}} = 50.5$				
	3340 w		4.15–3.70 m	CH ₂						
	3327 w		3.46 q, br	NH ₂	ABC ₂ M	δ_{A} 131.5 δ_{B} 122.0 δ_{C} 120.5 δ_{M} 150.4 $J_{\text{AB}} = 55.5$ $J_{\text{AC}} = 61.7$ $J_{\text{AM}} = 48.6$ $J_{\text{BC}} = 63.2$ $J_{\text{BM}} = -475.5$ $J_{\text{CM}} = 54.0$				
	3266 w		1.36 t	CH ₃						
			1.35 t							
			1.33 t							
			1.32 t							
			1.29 t							
	3 [Ru(NH=NH){P(OEt) ₃ } ₃]- (BPh ₄) ₂						AB ₄ XY spin syst.	=NH	AB ₄	δ_{A} 129.1 δ_{B} 118.3 $J_{\text{AB}} = 59.3$
							δ_{X} 16.76			
							δ_{Y} 15.36			
			$J_{\text{AX}} = 12.4$							
			$J_{\text{AY}} = 9.2$							
			$J_{\text{BX}} = 1.5$							
			$J_{\text{BY}} = 2.6$							
			$J_{\text{XY}} = 32.5$							
			4.06 m	CH ₂						
			1.37 t	CH ₃						
4 [Ru(NH=NH){PPh(OEt) ₂ }- {P(OEt) ₃ } ₄](BPh ₄) ₂			16.74 m	=NH	AB ₄	δ_{A} 158.8 δ_{B} 120.5 $J_{\text{AB}} = 51.1$ δ_{A} 128.8 δ_{B} 119.0 δ_{C} 118.2 δ_{M} 147.3 $J_{\text{AB}} = 53.8$ $J_{\text{AC}} = 65.2$ $J_{\text{AM}} = 45.8$ $J_{\text{BC}} = 61.2$ $J_{\text{BM}} = -460.5$ $J_{\text{CM}} = 52.7$				
			15.38 m							
			3.93 m	CH ₂						
			1.34 t	CH ₃						
			1.32 t							
			1.31 t							
			1.27 t							
			1.25 t							
	5 [Os(NH=NH){P(OEt) ₃ } ₃]- (BPh ₄) ₂			A ₄ BXY spin syst.			=NH	A ₄ B ^e	δ_{A} 84.2 δ_{B} 63.3 $J_{\text{AB}} = 40.3$	
				δ_{X} 16.82						
			δ_{Y} 15.73							
			$J_{\text{AX}} = 1.65$							
			$J_{\text{AY}} = 3.85$							
			$J_{\text{BX}} = 11.0$							
			$J_{\text{BY}} = 7.65$							
			$J_{\text{XY}} = 32.2$							
			4.11 m	CH ₂						
			4.00 qnt							
6 [Ru(C ₆ H ₅ N=NH){P(OEt) ₃ } ₃]- (BPh ₄) ₂			13.72 m ^e	=NH	AB ₄ ^e	δ_{A} 133.0 δ_{B} 121.3 $J_{\text{AB}} = 60.4$				
			4.30 m	CH ₂						
			4.19 qnt							
			1.40 t	CH ₃						
			1.32 t							
			1.35 t							
7 [Os(C ₆ H ₅ N=NH){P(OEt) ₃ } ₃]- (BPh ₄) ₂			14.15 m ^e	=NH	A ₄ B ^e	δ_{A} 86.6 δ_{B} 69.7 $J_{\text{AB}} = 41.0$				
			4.35 m	CH ₂						
			4.11 qnt							
			1.40 t	CH ₃						
			1.34 t							
			4.07 m	CH ₂						
8 [Ru(H ₂ O){P(OEt) ₃ } ₃](BPh ₄) ₂			2.65 s, br	H ₂ O	AB ₄	δ_{A} 131.7 δ_{B} 119.1 $J_{\text{AB}} = 59.5$				
			1.36 t	CH ₃						
			1.32 t							
			1.32 t							
9 [Ru(NH=NH) ₂ {P(OEt) ₃ } ₄]- (BPh ₄) ₂ ^f			16.71 dm	=NH	A ₂ B ₂	δ_{A} 128.2 δ_{B} 118.1 $J_{\text{AB}} = 60.0$				
			15.83 dm							
			$J_{\text{HH}} = 32$							
			4.07 m	CH ₂						
			1.37 t	CH ₃						
			1.32 t							

Table 2 IR and NMR data for ruthenium and osmium complexes

Compound	IR ^a		¹ H NMR ^{b,c}			³¹ P- ¹ H NMR ^{b,d}
	ν/cm^{-1}	Assignment	$\delta(\text{J/Hz})$	Assignment	Spin system	$\delta(\text{J/Hz})$
10 [Ru(CH ₃ N=NH)(CH ₃ NHNH ₂)-{P(OEt) ₃ } ₄](BPh ₄) ₂	3347 w	$\nu(\text{NH})$	13.74 m, br	=NH	ABC ₂ ^e	δ_{A} 128.9
	3304 m		4.64 s, br	RuNH ₂		δ_{B} 127.4
	3295 w		4.10 m	CH ₂		δ_{C} 118.5
			3.76 s	CH ₃ N=		J_{AB} = 68.7
			3.45 m, br	NH ₂ hydrazine		J_{AC} = 61.2
			2.65 d	CH ₃ hydrazine		J_{BC} = 60.1
			1.37 t	CH ₃		
11 [Ru(CH ₃ N=NH) ₂ {P(OEt) ₃ } ₄](BPh ₄) ₂			1.34 t		A ₂ B ₂	δ_{A} 129.8
			1.32 t			δ_{B} 119.8
			13.58 m, br	=NH		J_{AB} = 61.6
			4.10 m	CH ₂		
			3.81 s	CH ₃ N=		
			1.39 t	CH ₃		

^a In KBr pellets. ^b In CD₂Cl₂ at 25 °C, unless otherwise noted. ^c Phenyl proton resonances are omitted. ^d Positive shift downfield from 85% H₃PO₄. ^e In (CD₃)₂CO. ^f The sample contains the acetate complex [Ru(κ^2 -O₂CCH₃){P(OEt)₃}₄](BPh₄).

**Fig. 1** L = P(OEt)₃.**Scheme 3** M = Ru **3**, Os **5**; L = P(OEt)₃.


(OEt)₂]₄]²⁺ **4**, which were isolated as white solids and characterised (Scheme 3).

It was essential for the successful synthesis of diazene derivatives **3–5** to carry out reactions at low temperature (–30 °C) and to use high-purity Pb(OAc)₄ (99.99%) in equimolar amounts. Otherwise, mixtures of products not containing the 1,2-diazene complex were obtained. In particular, the use of common Pb(OAc)₄ (95%) did not yield pure samples of **3–5**, but only mixtures containing small amounts (10–15%) of the diazene complex.

All NH=NH derivatives **3–5** are stable as solids, but slowly decompose in solution, losing the diazene ligand. An attempt to crystallise **3** in a mixture of ethanol and dichloromethane as solvent gave aquo-complex [Ru(H₂O){P(OEt)₃]₅](BPh₄)₂ **8**,

which was isolated and characterised. The H₂O molecule probably comes from the traces of water always present in the common “anhydrous” solvents. In the solution the presence of hydrazine was also detected and confirms the decomposition of free NH=NH.

Diazene complexes **3–5** are 2 : 1 electrolytes, and their analytical and spectroscopic data (Table 2) support the proposed formulation. In the low-field region of the ¹H NMR spectrum of complex [Ru(NH=NH)L₅]²⁺ **3**, two multiplets (Fig. 2) at

**Fig. 2** Observed (bottom) and calculated (top) ¹H NMR spectra, in the diazene region, of [Ru(NH=NH){P(OEt)₃]₅](BPh₄)₂ **3**, in CD₂Cl₂ at 25 °C. The simulated spectrum was obtained using the parameters reported in Table 2.

16.76 and 15.36 ppm appeared, which were attributed to the two NH protons of a monohapto NH=NH ligand. This pattern is due to coupling with ³¹P nuclei (AB₄ multiplet), as confirmed by simulation of the spectrum using an AB₄XY model (X = HA, Y = HB), with the parameters listed in Table 2. The ³J_{HH} value of 32.5 Hz (J_{XY}) also suggested^{4d,5} a probable *trans*-NH=NH geometry. The related osmium complex [Os(NH=NH)L₅]²⁺ **5**, in the low-field region of the proton spectra, also showed the characteristic A₄BXY multiplet of the diazene ligands. In the temperature range between –80 and +30 °C, the ³¹P{¹H} NMR spectra of both complexes **3** and **5** appeared as AB₄ or A₄B multiplets, fitting geometry **IV** (Fig. 3).

The presence of the diazene ligand was also confirmed in the mixed-phosphite derivative [Ru(NH=NH){PPh(OEt)₂]₂{P(OEt)₃]₄](BPh₄)₂ **4**, the ¹H NMR spectrum of which showed the characteristic =NH multiplet between 17 and 15 ppm (Table 2). In the temperature range between –80 and +30 °C, the ³¹P{¹H} NMR spectrum appeared as a complicated pattern, simulated




Fig. 3 L = P(OEt)₃.

as a set of one AB₄ and one ABC₂M multiplets, partially overlapping. On this basis, and taking into account the ³¹P spectra of hydrazine precursor **2**, we propose the existence of two isomers of the type shown in Fig. 4 for our complex **4**, containing the NH=NH and PPh(OEt)₂ ligands in mutually *cis* (V) or *trans* (VI) positions, respectively.





Fig. 4 L = P(OEt)₃, L' = PPh(OEt)₂.

Complexes containing the monodentate 1,2-diazene ligand are very rare,⁵ and only one example^{5c} of isolable species has been reported for ruthenium and osmium metal centres. Using the pentakis(phosphite) ML₅ (or ML'₄) fragment stabilises the NH=NH molecule through coordination, affording stable and isolable species for both Ru(II) and Os(II).

Unfortunately, no other [M(RN=NH)L₅]²⁺ diazene complexes besides **3** and **5** could be prepared by oxidation of coordinated hydrazine, owing to the absence of the corresponding precursors. In our hands, hydrazines RNHNH₂ different from NH₂NH₂ could not be coordinated to the ML₅ fragment to give stable species. However, taking into account that aryldiazene can also be prepared by insertion of aryldiazonium cations into a M–H bond, we treated the [MHL₅]⁺ (M = Ru, Os) hydrides with the phenyldiazonium cation PhN₂⁺BF₄[−] and observed the formation of aryldiazene derivatives [M(PhN=NH)L₅]²⁺ (M = Ru **6**, Os **7**) which were isolated as BPh₄[−] salts and characterised (Scheme 4).




Scheme 4 M = Ru **6**, Os **7**; L = P(OEt)₃.

The formation of complexes **6** and **7** may indicate that the steric requirement of aryldiazene PhN=NH is lower than that of arylhydrazine PhNHNH₂, whose coordination on the ML₅ fragment did not take place. Aryldiazene [M(PhN=NH)L₅](BPh₄)₂ derivatives **6** and **7** are stable as solids and in solutions of polar organic solvents, in which they behave as 2 : 1 electrolytes. The presence of the diazene ligand was confirmed by the ¹H NMR spectra, which displayed the characteristic high-frequency NH proton resonances at 13.72 (**6**) and 14.15 (**7**) ppm. In the temperature range between −90 and +30 °C, the ³¹P{¹H} NMR spectrum appeared as an AB₄ multiplet, fitting the proposed formulation.

Table 3 Selected bond lengths (Å) and angles (°) for [Ru(H₂O){P(OEt)₃]₅(BPh₄)₂ **8**, with s.u.s in parentheses

Ru–O16	2.219(7)	Ru–P3	2.352(3)
Ru–P4	2.238(3)	Ru–P2	2.356(3)
Ru–P1	2.348(3)	Ru–P5	2.372(3)
O16–Ru–P4	178.3(2)	P1–Ru–P2	88.2(1)
O16–Ru–P1	89.9(2)	P3–Ru–P2	167.7(1)
P4–Ru–P1	91.1(1)	O16–Ru–P5	86.8(2)
O16–Ru–P3	80.7(2)	P4–Ru–P5	92.1(1)
P4–Ru–P3	98.0(1)	P1–Ru–P5	176.6(1)
P1–Ru–P3	91.81(9)	P3–Ru–P5	88.7(1)
O16–Ru–P2	87.0(2)	P2–Ru–P5	90.6(1)
P4–Ru–P2	94.3(1)		



Scheme 5 L = P(OEt)₃, M = Ru.

Although aryldiazenes **6** and **7** were stable in solution, related 1,2-diazene complexes **3–5** slowly decomposed giving aquo-complex [M(H₂O)L₅]²⁺, probably by substitution of the labile NH=NH ligand with traces of water present in the solvent. In the case of ruthenium, compound [Ru(H₂O)L₅](BPh₄)₂ **8** (Scheme 5) was isolated as white microcrystals and characterised spectroscopically (Table 2) and crystallographically (Table 1). The presence of H₂O as a ligand in the complex was indicated by the proton NMR signal at 2.65 ppm, and the ³¹P{¹H} spectra confirmed the ML₅ fragment.

The structure and atom labelling of [Ru(H₂O){P(OEt)₃]₅²⁺ is shown in Fig. 5, and the coordination geometry is summarised in Table 3. The metal coordination is octahedral, with ligand





Fig. 5 Perspective view of the solid-state molecular structure of cation [Ru(H₂O){P(OEt)₃]₅²⁺ **8**²⁺. Anisotropic displacement parameters are at the 50% probability level. Ethoxy groups are omitted for clarity.

at P4 located *trans* to the water molecule, the remaining phosphites being sterically equivalent to each other. The steric crowding of the bulky phosphite ligands is partly relieved by pushing them towards the water position, thus reducing the width of the *cis* O–Ru–P bond angles, which range between 80.7(2) and 89.9(2)°. The presence of the water ligand also affects the Ru–P4 bond *trans* to it [2.238(3) Å], which is far shorter than the remaining *cis* ones [2.348(3)–2.372(3) Å]. The

large *trans* influence of the unique P(OEt)₃ ligand is evidenced by the fact that the Ru–O bond is among the longest found in Ru²⁺ water complexes (*e.g.* Ru–O = 2.139 Å in the hexa-aquo cation¹⁸). There are no intermolecular interactions among non-hydrogen atoms below 3.55 Å in the structure.

Results from oxidation reactions of hydrazine complexes [M(NH₂NH₂)L₅]²⁺ using the new high-purity Pb(OAc)₄ (99.99%) prompted us to reinvestigate the oxidation of the bis(hydrazine) complexes [M(NH₂NH₂)₂L₄]²⁺, that were previously studied by us,^{6a,c} using low-purity Pb(OAc)₄ (95%) as oxidising agent. These earlier reactions did not give diazene complexes, but only acetate derivative [M(κ²-O₂CCH₃)L₄]⁺. The reaction was studied in CH₂Cl₂ at –30 °C using high-purity Pb(OAc)₄ in a Ru : Pb ratio 1 : 2, and showed that, in the case of [Ru(NH₂NH₂)₂{P(OEt)₃}₄]²⁺ it does give a white solid containing a mixture of bis(diazene) [Ru(NH=NH)₂L₄](BPh₄)₂ **9** and acetate [Ru(κ²-O₂CCH₃)L₄]BPh₄ derivatives, as shown in Scheme 6.



Scheme 6 L = P(OEt)₃.


From this mixture, we attempted to separate the diazene complex by fractional crystallisation at low temperature, but the instability in solution of [Ru(NH=NH)₂L₄]²⁺ on the one hand, and the comparable solubility of the two complexes on the other, always gave samples containing more products. However, the ¹H NMR spectra of the mixture strongly supports the formation of bis(diazene) **9**, showing two doublets of multiplets at 16.71 and 15.83 ppm, attributed to the two NH protons of the two diazene ligands. The ³J_{HH} value of 32 Hz also suggests^{4d,5} a *trans*-NH=NH geometry. The ³¹P{¹H} NMR spectra of the oxidation product also shows two A₂B₂ multiplets, due to acetate compound^{6a} [Ru(κ²-O₂CCH₃)L₄]⁺ and bis(diazene) complex **9**, fitting the proposed formulation. The A₂B₂ pattern also suggests the mutually *cis* positions of the two diazene ligands, as in geometry VIII, Scheme 6.

Oxidation of the related [Ru(NH₂NH₂)₂{PPh(OEt)₂}₄](BPh₄)₂ and [Os(NH₂NH₂)₂{P(OEt)₃}₄](BPh₄)₂ derivatives afforded white solids, with ¹H NMR spectra showing diazene signals, probably attributable to bis(diazene) [M(NH=NH)₂L₄](BPh₄)₂ (M = Ru, Os) complexes. In these cases, however, the predominant species is acetate compound [M(κ²-O₂CCH₃)L₄]⁺, with the diazene complex present only in a small amount (from 5 to 10%).

These studies on the oxidation of bis(hydrazine) complexes with high-purity Pb(OAc)₄ (99.99%) show that 1,2-diazene complexes can also be formed in these cases, but their stabilisation on the ML₄ fragment is restricted, as compared with related [M(NH=NH)L₅]²⁺ derivatives, and does not allow separation of the samples in pure form. However, although substitution of the NH=NH ligand by the acetate ion also takes place using high-purity Pb(OAc)₄, a solid sample containing the coordinated 1,2-diazene (**9**) as the predominant species was isolated. In addition, these results confirm our previous hypothesis^{6a,c} on the formation of a diazene complex in the oxidation of [M(NH₂NH₂)₂L₄]²⁺ cations, and highlight the importance of the purity of Pb(OAc)₄ in determining the nature of the final oxidation products.

On this basis, we extended the oxidation reaction to methylhydrazine complexes [Ru(CH₃NHNH₂)₂L₄]²⁺ and found that, using high-purity Pb(OAc)₄ (99.99%), the reaction gives a mixture of bis(methyldiazene) [Ru(CH₃N=NH)₂L₄]²⁺ **11** and methyldiazene–methylhydrazine [Ru(CH₃N=NH)(CH₃NH-NH₂)L₄]²⁺ **10** complexes, which were isolated as BPh₄[–] salts and characterised (Scheme 7).

Complexes **10** and **11** were separated by fractional crystallisation into two pale yellow solids, stable in air and in solutions of polar organic solvents, in which they behave as 2 : 1 electrolytes. Analytical and spectroscopic data (Table 2) support the proposed formulation. The ¹H NMR spectrum of bis(methyl-



Scheme 7 L = P(OEt)₃.

diazene) complex **11** shows a slightly broad multiplet at 13.58 ppm, characteristic of the diazene ligand. Beside the signals of phosphite and BPh₄ ion, the spectrum also contains, a singlet at 3.81 ppm, due to the methyl–proton resonance of the methyldiazene group. In the temperature range between –80 and +30 °C, the ³¹P{¹H} NMR spectra appear as an A₂B₂ multiplet, fitting the presence of two diazene ligands in a mutual *cis* position (**IX**).

In mixed-ligand complex [Ru(CH₃N=NH)(CH₃NHNH₂)L₄](BPh₄)₂ **10**, the ¹H NMR spectra show the presence of both nitrogenous ligands. The broad multiplet at 13.74 ppm of the NH proton and the singlet at 3.76 ppm of the methyl resonances confirm the presence of the CH₃N=NH ligand, whereas the broad signals at 4.64 and 3.45 ppm, due to NH₂ and NH resonances respectively, and the doublet at 2.65 ppm are diagnostic of the hydrazine group. The ABC₂ multiplet present in the ³¹P{¹H} spectra also suggests mutual *cis* position of the hydrazine and diazene ligands in the complex (**X**).

The RuL₄ fragment is therefore able to stabilise both 1,2-diazene and methyldiazene by coordination, affording the related bis(diazene) species, which are very stable, in the case of CH₃N=NH. Partial substitution was observed with the NH=NH ligand, affording a mixture of diazene (**9**) and acetate [Ru(κ²-O₂CCH₃)L₄]⁺ derivatives. However, isolation of solid samples containing diazene complexes of ruthenium was only achieved by using high-purity Pb(OAc)₄ (99.99%), and the absence in previous studies⁶ of any evidence of these diazene species is probably due to the fact that standard-grade Pb(OAc)₄ (95%) was used, which leads to decomposition of diazene derivatives (see ESI).

Acknowledgements

The financial support of MURST (Rome)—Programmi di Ricerca Scientifica di Rilevante Interesse Nazionale, Cofinanziamento 2000–2001—is gratefully acknowledged. We thank Daniela Baldan for technical assistance.

References

- 1 N. Wiberg, H. Bachhuber and G. Fischer, *Angew. Chem., Int. Ed. Engl.*, 1972, **11**, 829.

- 2 (a) J. Chatt and R. L. Richards, in *The Chemistry and Biochemistry of Nitrogen Fixation*, J. R. Postgate, ed., Plenum, London, 1971; (b) R. N. F. Thorneley, R. R. Eady and D. J. Lowe, *Nature*, 1978, **272**, 5578; (c) R. A. Henderson, G. J. Leigh and C. J. Pickett, *Adv. Inorg. Chem. Radiochem.*, 1983, **27**, 198; (d) B. F. G. Johnson, B. L. Haymore and J. R. Dilworth, in *Comprehensive Coordination Chemistry*, G. Wilkinson, R. D. Gillard and J. A. McCleverty, eds., Pergamon Press, Oxford, UK, 1987, vol. 2, p. 130; (e) D. Coucouvanis, *Acc. Chem. Res.*, 1991, **24**, 1.
- 3 E. J. Corey, D. J. Pasto and W. L. Mock, *J. Am. Chem. Soc.*, 1961, **83**, 2895; E. E. van Tamelen and R. J. Timmons, *J. Am. Chem. Soc.*, 1962, **84**, 1067; D. J. Pasto and D. M. Chipman, *J. Am. Chem. Soc.*, 1979, **101**, 2290; H. S. Rzepa and D. K. Agrafiotis, *J. Chem. Soc., Perkin Trans. 2*, 1989, 475.
- 4 (a) D. Sellmann, A. Brandl and R. Endell, *Angew. Chem., Int. Ed. Engl.*, 1973, **12**, 1019; (b) G. Huttner, W. Gartzke and K. Allinger, *Angew. Chem., Int. Ed. Engl.*, 1974, **13**, 822; (c) D. Sellmann, A. Brandl and R. Endell, *J. Organomet. Chem.*, 1973, **49**, C22; (d) D. Sellmann and K. Jödden, *Angew. Chem., Int. Ed. Engl.*, 1977, **16**, 464; (e) D. Sellmann, E. Böhlen, M. Waeber, G. Huttner and L. Zsolnai, *Angew. Chem., Int. Ed. Engl.*, 1985, **24**, 981; (f) D. Sellmann, W. Soglowek, F. Knoch and M. Moll, *Angew. Chem., Int. Ed. Engl.*, 1989, **28**, 1271; (g) J. P. Collman, J. E. Hutchinson, M. A. Lopez, R. Guilard and R. A. Reed, *J. Am. Chem. Soc.*, 1991, **113**, 2794; (h) D. Sellmann, J. Käßler, M. Moll and F. Knoch, *Inorg. Chem.*, 1993, **32**, 960; (i) D. Sellmann and A. Hennige, *Angew. Chem., Int. Ed. Engl.*, 1997, **36**, 276.
- 5 (a) M. R. Smith III, T.-Y. Cheng and G. L. Hillhouse, *J. Am. Chem. Soc.*, 1993, **115**, 8638; (b) T.-Y. Cheng, J. C. Peters and G. L. Hillhouse, *J. Am. Chem. Soc.*, 1994, **116**, 204; (c) T.-Y. Cheng, A. Ponce, A. L. Rheingold and G. L. Hillhouse, *Angew. Chem., Int. Ed. Engl.*, 1994, **33**, 657.
- 6 (a) G. Albertin, S. Antoniutti, A. Bacchi, E. Bordignon, P. M. Dolcetti and G. Pelizzi, *J. Chem. Soc., Dalton Trans.*, 1997, 4435; (b) G. Albertin, S. Antoniutti, E. Bordignon and S. Pattaro, *J. Chem. Soc., Dalton Trans.*, 1997, 4445; (c) G. Albertin, S. Antoniutti, A. Bacchi, M. Bergamo, E. Bordignon and G. Pelizzi, *Inorg. Chem.*, 1998, **37**, 479.
- 7 R. Rabinowitz and J. Pellon, *J. Org. Chem.*, 1961, **26**, 4623.
- 8 A. I. Vogel, *Practical Organic Chemistry*, 3rd edn., Longmans, Green and Co., New York, 1956.
- 9 E. Nachbaur and G. Leiseder, *Monatsh. Chem.*, 1971, **102**, 1718.
- 10 W. G. Peet and D. H. Gerlach, *Inorg. Synth.*, 1974, **15**, 38; D. H. Gerlach, W. G. Peet and E. L. Muettterties, *J. Am. Chem. Soc.*, 1972, **94**, 4545.
- 11 P. Amendola, S. Antoniutti, G. Albertin and E. Bordignon, *Inorg. Chem.*, 1990, **29**, 318.
- 12 SAINT: SAX, Area Detector Integration, Siemens Analytical Instruments Inc., Madison, Wisconsin, USA.
- 13 SADABS: Siemens Area Detector Absorption Correction Software, G. Sheldrick, University of Goettingen, Germany, 1996.
- 14 Sir 97: A New Program for Solving and Refining Crystal Structures, A. Altomare, M. C. Burla, M. Camalli, G. Cascarano, C. Giacovazzo, A. Guagliardi, A. G. Moliterni, G. Polidori and R. Spagna, Istituto di Ricerca per lo Sviluppo di Metodologie Cristallografiche C N R, Bari, 1997.
- 15 Shelxl 97. Program for Structure Refinement, G. Sheldrick, University of Goettingen, Germany, 1997.
- 16 L. J. Farrugia, *J. Appl. Crystallogr.*, 1999, **32**, 837.
- 17 F. H. Allen, O. Kennard and R. Taylor, *Acc. Chem. Res.*, 1983, **16**, 146.
- 18 P. Bernhard, H.-B. Burgi, J. Hauser, H. Lehmann and A. Ludi, *Inorg. Chem.*, 1982, **21**, 3936.