

Iridium-assisted C≡C bond cleavage of 1-alkyne by water: preparation of new alkyl derivatives

Gabriele Albertin,^{*a} Stefano Antoniutti,^a Alessia Bacchi,^b Giancarlo Pelizzi^b and Francesca Piasente^a

^a Dipartimento di Chimica, Università Ca' Foscari di Venezia, Dorsoduro 2137, 30123 Venezia, Italy

^b Dipartimento di Chimica Generale ed Inorganica, Chimica Analitica, Chimica Fisica, Università di Parma, Parco Area delle Scienze 17/A, 43100 Parma, Italy

Received 1st April 2003, Accepted 23rd May 2003

First published as an Advance Article on the web 18th June 2003

Alkyl complexes $\text{IrCl}_2(\eta^1\text{-CH}_2\text{Ar})(\text{CO})(\text{PPh}_3)_2$ (**1**) [Ar = Ph (**a**), *p*-tolyl (**b**)] were prepared by allowing the hydride *mer*- or *fac*- $\text{IrHCl}_2(\text{PPh}_3)_3$ to react with terminal alkynes $\text{ArC}\equiv\text{CH}$ in the presence of water. The complexes were characterized spectroscopically (IR and ^1H , ^{31}P , ^{13}C NMR) and by the X-ray crystal structure determination of **1b**. The acyl complex $\text{IrCl}_2\{\text{C}(\text{O})\text{CH}_2\text{C}(\text{CH}_3)_3\}(\text{PPh}_3)_2$ (**2**) was also prepared by reacting *mer*- $\text{IrHCl}_2(\text{PPh}_3)_3$ with *tert*-butylacetylene $\text{HC}\equiv\text{CC}(\text{CH}_3)_3$ in the presence of H_2O . A reaction path for the hydration of terminal alkyne in the presence of the Ir(III) complex leading to the cleavage of the C≡C bond, with the formation of the complexes **1** or **2** is also proposed. Acetylide complexes $\text{IrHCl}(\text{C}\equiv\text{CAr})(\text{PPh}_3)_3$ (**3**), $\text{IrHCl}(\text{C}\equiv\text{CAr})(\text{AsPh}_3)_3$ (**4**) [Ar = Ph (**a**), *p*-tolyl (**b**)] and $\text{IrHCl}(\text{C}\equiv\text{CPh})\{\text{PPh}(\text{OEt})_2\}(\text{PPh}_3)_2$ (**5**) were prepared by reacting IrHCl_2L_3 (L = PPh_3 , AsPh_3) or $\text{IrHCl}_2\{\text{PPh}(\text{OEt})_2\}(\text{PPh}_3)_2$ with lithium acetylide. Protonation reaction with Brønsted acids of the acetylide **3** was also studied and led to unstable vinyl derivatives. The stable vinyl $[\text{IrCl}\{\eta^2\text{-CH}=\text{C}(\text{H})\text{COOMe}\}\text{L}_2]\text{BPh}_4$ (**6,7**) [L = PPh_3 (**6**), AsPh_3 (**7**)] complexes, instead, were prepared by allowing IrHCl_2L_3 to react first with AgCF_3SO_3 and then with methyl propiolate.

Introduction

Transition metal complexes are known to promote hydration of unactivated alkyne affording species such as aldehydes, ketones, or vinyl alcohol complexes.¹ In some cases the metal-assisted reaction of 1-alkynes with water has been reported to give a C–C triple bond cleavage with the formation of CO and a saturated hydrocarbon with one fewer carbon atom. They are generally present as a carbonyl and η^1 -alkyl ligand.^{2–4} Among the metal centres, a prominent role is played by those of the iron-group. A mechanistic study on the stoichiometric Ru(II)-assisted C–C fission of phenylacetylene by water involving the participation of a Ru(II)-vinylidene intermediate has also been recently reported.^{3a} Less is known for other metal centres and, for the iridium, the formation of an acyl $[\text{Ir}(\text{CO})\text{CH}_2\text{Ph}]$ derivative by hydrolysis of phenylacetylene was reported in one case with a binuclear complex,^{4b} while a carbonyl complex was obtained in the reaction of $\text{PhC}\equiv\text{CH}$ with both a metallacyclo complex^{4a} and, very recently, with a water-soluble iridium derivative.^{4c} However, what is known of the hydrolysis of alkyne is still rather limited and further studies on the influence of the metal centre, the ancillary ligands and the nature of the alkyne on the reaction course and the nature of the final product should therefore be desirable.

We have previously reported some studies on the reactivity of metal complexes of the iron triad and of cobalt with 1-alkynes, which allows the synthesis of the acetylide, vinyl and vinylidene derivatives.⁵ We have now extended these investigations to include iridium as a metal centre and found a new example of hydration of terminal alkyne promoted by a transition metal complex which yields both alkyl-carbonyl and acyl derivatives. The results of these studies, which also include the synthesis of both alkynyl and vinyl complexes of Ir(III) and a detailed study on the hydrolysis of the alkyne, are reported here.

Experimental

General

All synthetic work was carried out under an inert atmosphere (argon) using standard Schlenk techniques or a Vacuum

Atmosphere dry-box. Once isolated, the complexes turned out to be quite air-stable and were stored at $-25\text{ }^\circ\text{C}$. All solvents used were dried over appropriate drying agents, degassed on a vacuum line, and distilled into vacuum-tight storage flasks. Phosphine $\text{PPh}(\text{OEt})_2$ was prepared by the method of Rabino-witz and Pellon.⁶ Alkynes $\text{RC}\equiv\text{CH}$ (R = Ph, *p*-tolyl, Bu' or Pr', Si(CH₃)₃, COOMe) were Aldrich products, used without further purification. Lithium acetylide $\text{Li}^+[\text{ArC}\equiv\text{C}]^-$ (Ar = Ph, *p*-tolyl) was prepared by reacting a slight excess of 1-alkyne (22 mmol) with lithium (20 mmol, 0.14 g) in 10 cm³ of tetrahydrofuran (THF). Oxygen-18-labelled water (95 atom% ^{18}O) was an Aldrich product. Other reagents were purchased from commercial sources in the highest available purity and used as received. Infrared spectra were recorded on Digilab Bio-Rad FTS-40 or Nicolet Magna 750 FT-IR spectrophotometers. NMR spectra (^1H , ^{13}C , ^{31}P) were obtained on Bruker AC200 or AVANCE 300 spectrometer at temperatures between -90 and $+30\text{ }^\circ\text{C}$, unless otherwise stated. ^1H and $^{13}\text{C}\{^1\text{H}\}$ (Table 2) spectra are referred to internal tetramethylsilane; $^{31}\text{P}\{^1\text{H}\}$ chemical shifts are reported with respect to 85% H_3PO_4 , with downfield shifts considered positive. The COSY and HMQC NMR experiments were obtained on the Bruker AVANCE 300 instrument using its standard programs. The SwaN-MR software package⁷ was used to treat NMR data. The conductivity of $10^{-3}\text{ mol dm}^{-3}$ solutions of the complexes in MeNO_2 at $25\text{ }^\circ\text{C}$ was measured with a Radiometer CDM 83 instrument.

Synthesis of the complexes

The hydrides *mer*- and *fac*- $\text{IrHCl}_2(\text{PPh}_3)_3$, *mer*- $\text{IrHCl}_2(\text{AsPh}_3)_3$ and $\text{IrHCl}_2\{\text{PPh}(\text{OEt})_2\}(\text{PPh}_3)_2$ were prepared following the reported methods.⁸

$\text{IrCl}_2(\eta^1\text{-CH}_2\text{Ar})(\text{CO})(\text{PPh}_3)_2$ (**1**) [Ar = Ph (**a**), *p*-tolyl (**b**)]

An excess of the appropriate alkyne (0.2 mmol) was added to a solution of *mer*- $\text{IrHCl}_2(\text{PPh}_3)_3$ (0.100 g, 0.095 mmol) in 10 cm³ of THF and the reaction mixture was refluxed for 1 h. The solvent was removed under reduced pressure giving an oil which was triturated with ethanol (5 cm³). An orange–yellow solid slowly separated out from the resulting solution, which

was filtered and recrystallized by cooling to $-25\text{ }^{\circ}\text{C}$ a saturated solution prepared at room temperature by treating the solid with 10 cm^3 of ethanol and enough CH_2Cl_2 to dissolve it. It can be noted that the addition of an excess of H_2O (0.2 mmol, 36 μL) to the reaction mixture does not change the yield in the products; yield $\geq 80\%$ (Found: C, 58.45; H, 4.06; Cl, 7.70%. $\text{C}_{44}\text{H}_{37}\text{Cl}_2\text{IrOP}_2$ (**1a**) requires C, 58.28; H, 4.11; Cl, 7.82%); $\bar{\nu}_{\text{max}}/\text{cm}^{-1}$ (CO) 2048s (KBr); δ_{H} (CD_2Cl_2 , 293 K) 7.95–6.06 (35H, m, Ph) and 3.55 (2H, t, CH_2 , $J_{\text{PH}} = 6\text{ Hz}$); δ_{P} (CD_2Cl_2 , 293 K) -15.41 (s). (Found: C, 58.85; H, 4.19; Cl, 7.88%. $\text{C}_{45}\text{H}_{39}\text{Cl}_2\text{IrOP}_2$ (**1b**) requires C, 58.69; H, 4.27; Cl, 7.70%); $\bar{\nu}_{\text{max}}/\text{cm}^{-1}$ (CO) 2048s (KBr); δ_{H} (CD_2Cl_2 , 293 K) 7.95–5.80 (m), 6.69, 5.95 (d) (34H, Ph), 3.52 (2H, t, CH_2 , $J_{\text{PH}} = 6\text{ Hz}$) and 2.23 (3H, s, CH_3); δ_{P} (CD_2Cl_2 , 293 K) -15.68 (s).

$\text{IrCl}_2(\eta^1\text{-CD}_2\text{Ph})(\text{CO})(\text{PPh}_3)_2$ (**1a-d**)

This complex was prepared in the same manner as the related **1a** by adding first an excess of $\text{PhC}\equiv\text{CH}$ (0.2 mmol, 22 μL) and then an excess of D_2O (0.4 mmol, 7.2 μL) to a solution of $\text{mer-IrHCl}_2(\text{PPh}_3)_3$ (0.100 g, 0.095 mmol) in 10 cm^3 of THF. After 1 h of reflux of the resulting solution and the evaporation of the solvent at reduced pressure, the addition of EtOD allowed to separate the solid. Proton and ^{13}C NMR spectra confirm the selective replacement of the methylenic hydrogen atoms of the benzylic group by deuterium.

$\text{IrCl}_2(\eta^1\text{-CH}_2\text{Ph})(\text{C}^{18}\text{O})(\text{PPh}_3)_2$ (**1a- ^{18}O**)

This complex was prepared in the same manner as the related **1a-d**, using H_2^{18}O (95 atom% ^{18}O) instead of D_2O . $\bar{\nu}_{\text{max}}/\text{cm}^{-1}$ (C^{18}O) 2005 (KBr).

$\text{IrCl}_2\{\eta^1\text{-C}(\text{O})\text{CH}_2\text{C}(\text{CH}_3)_3\}(\text{PPh}_3)_2$ (**2**)

An excess of $\text{Bu}^t\text{C}\equiv\text{CH}$ (0.3 mmol, 38 μL) was added to a solution of $\text{mer-IrHCl}_2(\text{PPh}_3)_3$ (0.100 g, 0.095 mmol) in 10 cm^3 of CH_2Cl_2 and the reaction mixture was stirred at room temperature for 40 h. The solvent was removed under reduced pressure to give a red-brown oil which was triturated with ethanol (5 cm^3). A yellow solid separated out from the resulting solution, which was filtered and recrystallized from CH_2Cl_2 -ethanol. It can be noted that the addition of an excess of H_2O (0.3 mmol, 5.4 μL) to the reaction mixture does not change either the reaction course, or the yield in the final product; yield $\geq 70\%$ (Found: C, 56.61; H, 4.78; Cl, 8.18%. $\text{C}_{42}\text{H}_{41}\text{Cl}_2\text{IrOP}_2$ requires C, 56.88; H, 4.66; Cl, 8.00%); $\bar{\nu}_{\text{max}}/\text{cm}^{-1}$ ($\text{C}=\text{O}$) 1681s (KBr); δ_{H} (CD_2Cl_2 , 293 K) 7.70–6.90 (30H, m, Ph), 2.76 (2H, s, CH_2) and 0.69 (9H, s, CH_3); δ_{P} (CD_2Cl_2 , 293 K) 8.25 (s).

$\text{IrCl}_2\{\eta^1\text{-C}(\text{O})\text{CD}_2\text{C}(\text{CH}_3)_3\}(\text{PPh}_3)_2$ (**2-d**)

This complex was prepared in the same manner as the related **2**, by adding first an excess of $\text{Bu}^t\text{C}\equiv\text{CH}$ (0.3 mmol, 38 μL) and then an excess of D_2O (0.4 mmol, 7.2 μL) to a solution of $\text{mer-IrHCl}_2(\text{PPh}_3)_3$ (0.100 g, 0.095 mmol) in 10 cm^3 of THF. The reaction mixture was stirred for 40 h and, after evaporation of the solvent, the solid complex was recovered by stirring the oil obtained with EtOD. The ^1H NMR spectra confirm the selective replacement of the methylenic hydrogen atom of the neopentyl group by deuterium.

$\text{IrCl}_2\{\eta^1\text{-C}(\text{O})\text{CH}_2\text{C}(\text{CH}_3)_3\}(\text{PPh}_3)_2$ (**2- ^{18}O**)

This complex was prepared in the same manner as **2** by adding first an excess of $\text{Bu}^t\text{C}\equiv\text{CH}$ (0.3 mmol, 38 μL) and then an excess of H_2^{18}O (95 atom% ^{18}O) (0.3 mmol, 5.4 μL) to a solution of $\text{mer-IrHCl}_2(\text{PPh}_3)_3$ in 10 cm^3 of CH_2Cl_2 . After 40 h of stirring of the reaction mixture and work-up, a yellow solid was obtained. $\bar{\nu}_{\text{max}}/\text{cm}^{-1}$ (C^{18}O) 1649 (KBr).

Reaction of *mer*- or *fac*- $\text{IrHCl}_2(\text{PPh}_3)_3$ with phenylacetylene in NMR tube

In a vacuum-atmosphere dry box a sample of *mer*- or *fac*- $\text{IrHCl}_2(\text{PPh}_3)_3$ (20 mg, 0.019 mmol) was placed in a screw-cap 5-mm NMR tube and 1 cm^3 of the appropriate deuterated solvent (THF-d_8 , CD_2Cl_2 or C_6D_6) was added. An excess of phenylacetylene (0.04 mmol, 4.5 μL) was added by syringe to the resulting suspension and the NMR tube was transferred into the probe of the spectrometer. Owing to the low solubility of $\text{IrHCl}_2(\text{PPh}_3)_3$ complex in THF-d_8 and C_6D_6 , the reaction was carried out at $40\text{ }^{\circ}\text{C}$ preheating the probe to this temperature. To the reaction mixture in the NMR tube was also added in some cases an excess of H_2O , D_2O or H_2^{18}O and the reaction was monitored by recording successive spectra.

$\text{IrHCl}(\text{C}\equiv\text{CAr})(\text{PPh}_3)_3$ (**3**) [Ar = Ph (**a**), *p*-tolyl (**b**)]

To a solution of *mer*- $\text{IrHCl}_2(\text{PPh}_3)_3$ (0.100 g, 0.095 mmol) in 10 cm^3 of THF was added an excess of the appropriate $\text{Li}^+[\text{ArC}\equiv\text{C}]^-$ (0.38 mmol, 0.19 cm^3 of a 2.0 M solution in THF) and the reaction mixture refluxed for about 1 h. The solvent was removed by evaporation under reduced pressure to give a red-brown oil which was treated with ethanol (5 cm^3). By vigorous stirring of the resulting solution, a red-brown solid slowly separated out, which was filtered and recrystallized from toluene-ethanol; yield $\geq 75\%$ (Found: C, 66.52; H, 4.68; Cl, 3.35%. $\text{C}_{62}\text{H}_{51}\text{ClIrP}_3$ (**3a**) requires C, 66.69; H, 4.60; Cl, 3.17%); $\bar{\nu}_{\text{max}}/\text{cm}^{-1}$ (IrH) 2168w and ($\text{C}\equiv\text{C}$) 2098m (KBr); δ_{H} (C_6D_6 , 293 K) 7.70–6.85 (50H, m, Ph) and -11.62 (1H, dt, H^- , $^2J_{\text{PH}^{\text{cis}}} = 18\text{ Hz}$, $^2J_{\text{PH}^{\text{trans}}} = 140\text{ Hz}$); δ_{P} (C_6D_6 , 293 K) spin syst A_2B , $\delta_{\text{A}} -4.92$, $\delta_{\text{B}} -22.36$, $J_{\text{AB}} = 15\text{ Hz}$. (Found: C, 66.78; H, 4.68; Cl, 3.01%. $\text{C}_{63}\text{H}_{53}\text{ClIrP}_3$ (**3b**) requires C, 66.92; H, 4.72; Cl, 3.14%); $\bar{\nu}_{\text{max}}/\text{cm}^{-1}$ (IrH) 2162w and ($\text{C}\equiv\text{C}$) 2099m (KBr); δ_{H} (C_6D_6 , 293 K) 7.70–6.80 (49H, m, Ph), 2.18 (3H, s, CH_3) and -11.61 (1H, dt, H^- , $^2J_{\text{PH}^{\text{cis}}} = 18\text{ Hz}$, $^2J_{\text{PH}^{\text{trans}}} = 140\text{ Hz}$); δ_{P} (C_6D_6 , 293 K) spin syst A_2B , $\delta_{\text{A}} -5.02$, $\delta_{\text{B}} -22.41$, $J_{\text{AB}} = 15\text{ Hz}$.

$\text{IrHCl}(\text{C}\equiv\text{CAr})(\text{AsPh}_3)_3$ (**4**) [Ar = Ph (**a**), *p*-tolyl (**b**)]

These complexes were prepared exactly in the same manner as the related complexes **3** by reacting *mer*- $\text{IrHCl}_2(\text{AsPh}_3)_3$ with an excess of $\text{Li}^+[\text{ArC}\equiv\text{C}]^-$ in THF at reflux for 3 h; yield $\geq 70\%$ (Found: C, 59.81; H, 4.05; Cl, 2.65%. $\text{C}_{62}\text{H}_{51}\text{As}_3\text{ClIr}$ (**4a**) requires C, 59.65; H, 4.12; Cl, 2.84%); $\bar{\nu}_{\text{max}}/\text{cm}^{-1}$ (IrH) 2166w and ($\text{C}\equiv\text{C}$) 2095m (KBr); δ_{H} (C_6D_6 , 293 K) 7.74–6.80 (50H, m, Ph) and -14.18 (1H, s, H^-). (Found: C, 59.70; H, 4.32; Cl, 3.02%. $\text{C}_{63}\text{H}_{53}\text{As}_3\text{ClIr}$ (**4b**) requires C, 59.93; H, 4.23; Cl, 2.81%); $\bar{\nu}_{\text{max}}/\text{cm}^{-1}$ (IrH) 2156w and ($\text{C}\equiv\text{C}$) 2098m (KBr); δ_{H} (CD_2Cl_2 , 293 K) 7.60–6.50 (49H, m, Ph), 2.27 (3H, s, CH_3) and -14.73 (1H, s, H^-).

$\text{IrHCl}(\text{C}\equiv\text{CPh})\{\text{PPh}(\text{OEt})_2\}(\text{PPh}_3)_2$ (**5**)

This complex was prepared in the same manner as the related compounds **3** and **4** by refluxing a 10 cm^3 THF solution of $\text{IrHCl}_2\{\text{PPh}(\text{OEt})_2\}(\text{PPh}_3)_2$ (0.100 g, 0.101 mmol) containing an excess of $\text{Li}^+[\text{PhC}\equiv\text{C}]^-$ (0.4 mmol, 0.2 cm^3 of a 2.0 M solution in THF) for about 18 h; yield $\geq 30\%$ (Found: C, 61.85; H, 4.81; Cl, 3.21%. $\text{C}_{54}\text{H}_{51}\text{ClIrO}_2\text{P}_3$ requires C, 61.62; H, 4.88; Cl, 3.37%); $\bar{\nu}_{\text{max}}/\text{cm}^{-1}$ (IrH) 2166w and ($\text{C}\equiv\text{C}$) 2099s (KBr); δ_{H} (CD_2Cl_2 , 293 K) 7.60–6.67 (40H, m, Ph), 3.80, 3.61 (4H, m, CH_2), 1.09 (6H, t, CH_3) and -11.78 (1H, dt, H^- , $^2J_{\text{PH}} = 4\text{ Hz}$, $^2J_{\text{PH}} = 18\text{ Hz}$); δ_{P} (CD_2Cl_2 , 293 K) spin syst AB_2 , $\delta_{\text{A}} 100.23$, $\delta_{\text{B}} -1.97$, $J_{\text{AB}} = 18\text{ Hz}$.

$[\text{IrCl}\{\eta^2\text{-CH}=\text{C}(\text{H})\text{COOMe}\}(\text{PPh}_3)_2]\text{BPh}_4$ (**6**)

In a 25-cm^3 three-necked round-bottomed flask were placed a solid sample of *mer*- $\text{IrHCl}_2(\text{PPh}_3)_3$ (0.150 g, 0.143 mmol) and an equimolar amount of AgCF_3SO_3 (0.143 mmol, 0.037 g). Dichloromethane (10 cm^3) was added and the reaction mixture

stirred for about 24 h in the dark. After filtration to remove the solid AgCl, an equimolar amount of methyl propiolate HC≡CCOOMe was added (0.143 mmol, 13 μL) and the reaction mixture stirred for 24 h. The solvent was removed by evaporation under reduced pressure to give an oil which was treated with ethanol (3 cm³). The addition of an excess of NaBPh₄ (0.3 mmol, 0.103 g) in 2 cm³ of ethanol to the resulting solution caused the separation of a pale-yellow solid which was filtered and crystallized by slow cooling to −25 °C of its saturated solution prepared at room temperature using a mixture of ethanol and CH₂Cl₂ as solvent; yield ≥55% (Found: C, 66.22; H, 4.69; Cl, 3.24%. C₆₄H₅₅BClIrO₂P₂ requires C, 66.46; H, 4.79; Cl, 3.07%); $\nu_{\max}/\text{cm}^{-1}$ (CO) 1570m (KBr); δ_{H} (CD₂Cl₂, 293 K) 7.80–6.80 (50H, m, Ph), 9.47 (1H, dt, C_αH, $J_{\text{HH}} = 15$ Hz, $J_{\text{PH}} = 9$ Hz), 6.11 (1H, d, C_βH, $J_{\text{HH}} = 15$ Hz) and 3.71 (3H, s, CH₃); δ_{P} (CD₂Cl₂, 293 K) spin syst AB, $\delta_{\text{A}} -15.2$, $\delta_{\text{B}} -17.8$, $J_{\text{AB}} = 13.8$ Hz.

[IrCl₂{η²-CH=C(H)COOMe}(AsPh₃)₂]BPh₄ (7)

This complex was prepared exactly in the same manner as the related complex **6** starting from *mer*-IrHCl₂(AsPh₃)₃; yield ≥45% (Found: C, 61.60; H, 4.38; Cl, 3.02%. C₆₄H₅₅As₂BClIrO₂ requires C, 61.77; H, 4.45; Cl, 2.85%); $\nu_{\max}/\text{cm}^{-1}$ (CO) 1580m (KBr); (CD₂Cl₂, 293 K) 8.00–7.80 (50H, m, Ph), 9.66, 6.02 (2H, d, C_αH and C_βH, $J_{\text{HH}} = 16$ Hz) and 3.64 (3H, s, CH₃).

X-Ray crystal structure determination of IrCl₂(η¹-CH₂(*p*-tolyl))(CO)(PPh₃)₂·3H₂O (**1b**·3H₂O).

Suitable crystals were obtained by slow cooling to −20 °C of a saturated solution of the complex in an ethanol–dichloromethane mixture (10 : 1 ratio). The crystals were analysed as obtained and contained three water molecules per iridium atom. A yellow irregular prism single crystal was mounted on a glass fiber and X-ray diffraction data were collected on a Philips PW1100 diffractometer, using graphite monochromated Mo-Kα radiation ($\lambda = 0.71069$ Å). Crystal decay resulted negligible. The phase problem was solved by direct methods⁹ and refined by full-matrix least squares on all F^2 ,¹⁰ with the support of the WinGX package.¹¹ Extinction effects were accounted for by refining an extinction parameter as implemented in SHELXL97.¹⁰ Anisotropic displacement parameters were refined for all non carbon atoms, while carbon atoms and carbonyl oxygen were located from Fourier maps and refined isotropically. Hydrogens were introduced in calculated positions. All aromatic rings were constrained to behave as rigid bodies. Three water molecules were found in the asymmetric unit, but the relative hydrogens were not located in the final map. Use of the Cambridge Crystallographic Database¹² facilities was made for structure discussion. The final map was featureless, except for large residues close to the iridium. Data collection and refinement results are summarized in Table 1.

CCDC reference number 207370.

See <http://www.rsc.org/suppdata/dt/b3/b303635a/> for crystallographic data in CIF or other electronic format.

Results and discussion

Both *mer*- and *fac*-iridium(III) complexes IrHCl₂(PPh₃)₃ react with terminal alkynes HC≡CAr in THF or CH₂Cl₂ to give the alkyl-carbonyl derivatives IrCl₂(η¹-CH₂Ar)(CO)(PPh₃)₂ (**1**), which were then isolated and characterized (Scheme 1).




Table 1 Crystal data and structure refinement for IrCl₂(η¹-CH₂(*p*-tolyl))(CO)(PPh₃)₂·3H₂O (**1b**·3H₂O)

Empirical formula	C ₄₅ H ₄₅ Cl ₂ IrO ₄ P ₂
Formula weight	974.85
Temperature/K	293(2)
Wavelength/Å	0.71069
Crystal system	Orthorhombic
Space group	<i>P2₁ab</i>
Unit cell dimensions	
<i>a</i> /Å	17.814(3)
<i>b</i> /Å	25.489(5)
<i>c</i> /Å	10.209(2)
<i>V</i> /Å ³	4635(1)
<i>Z</i>	4
<i>D_c</i> /Mg m ^{−3}	1.397
Absorption coefficient/mm ^{−1}	3.102
<i>F</i> (000)	1952
Crystal size/mm	0.4 × 0.4 × 0.6
Theta range for data collection/°	3–22
Index ranges	0 ≤ <i>h</i> ≤ 18, 0 ≤ <i>k</i> ≤ 26, 0 ≤ <i>l</i> ≤ 10
Reflections collected	2958
Independent reflections	2958
Observed independent reflections [<i>I</i> > 2σ(<i>I</i>)]	1872
Refinement method	Full-matrix least-squares on <i>F</i> ²
Data/restraints/parameters	2958/1/175
Goodness-of-fit on <i>F</i> ²	1.041
Final <i>R</i> indices [<i>I</i> > 2σ(<i>I</i>)]	<i>R</i> 1 = 0.0842, <i>wR</i> 2 = 0.2056
<i>R</i> indices (all data)	<i>R</i> 1 = 0.1416, <i>wR</i> 2 = 0.2466
Absolute structure parameter	−0.01(4)
Extinction coefficient	0.0009(5)
Largest Δ <i>F</i> max., min./e Å ^{−3}	2.181, −2.998

The reaction entails the hydrolysis of 1-alkyne with a C≡C bond cleavage and the formation of **1**, in agreement with the reaction shown in Scheme 1. The stoichiometry of this reaction is confirmed by the separation of both complex **1** and free PPh₃ from the reaction mixture carried out in “anhydrous” THF or CH₂Cl₂ as a solvent in about quantitative yield (≥90%). The addition of H₂O, in fact, is not necessary for the progress of the reaction, because the traces of H₂O present even in highly anhydrous solvents is shown to be enough to afford the final complex **1**. However, that H₂O is involved in the metal-mediated hydrolysis of the alkyne, yielding the alkyl-carbonyl derivative **1**, is unambiguously confirmed by the addition of D₂O or H₂¹⁸O to the reaction mixture, which yield either the labelled IrCl₂(η¹-CD₂Ph)(CO)(PPh₃)₂ (**1a-d₂**) or the IrCl₂(η¹-CH₂Ph)(C¹⁸O)(PPh₃)₂ (**1a-¹⁸O**) complexes, respectively. In the reaction mixture, the presence of free H₂ (singlet at 4.6 ppm in CD₂Cl₂)¹³ was also detected by ¹H NMR so confirming the proposed stoichiometry for the reaction (Scheme 1).

The alkyl-carbonyl complexes **1** are orange–yellow solids which are stable in the air, diamagnetic, soluble in the more common organic solvents and non-electrolytes. Their formulation is confirmed by analytical and spectroscopic data (Table 2) and by the X-ray crystal structure determination of the IrCl₂(η¹-CH₂(*p*-tolyl))(CO)(PPh₃)₂·3H₂O (**1b**·3H₂O) complex.

Its molecular structure is shown in Fig. 1 along with the labeling scheme. The most relevant geometric parameters are listed in Table 3. The metal coordination is octahedral, with the two triphenylphosphine ligands *trans* to each other, while the CH₂(*p*-tolyl) and CO molecules are *cis* to each other. The coordination is completed by two chlorine atoms, and the Ir–Cl bond length is significantly affected by the different *trans* influence of the carbonyl and CH₂(*p*-tolyl) groups, the latter inducing a certain weakening of the Ir–Cl₂ bond (2.506(9) Å) relative to the Ir–Cl₁ one (2.42(1) Å). The geometry of the CO ligand (Ir–C 1.75(6), C–O 1.24(5) Å) apparently deviates from the average values found in literature for hexa-coordinated Ir–CO systems (1.88 and 1.14 Å, respectively), but the observed bond lengths are probably affected by a thermal motion bias. The bonds relating to the coordination of the CH₂(*p*-tolyl) group (Ir–C 2.24(5), CH₂–C(Ar) 1.62(5) Å) are apparently

Table 2 $^{13}\text{C}\{^1\text{H}\}$ NMR Data for selected iridium complexes

Complex	$^{13}\text{C}\{^1\text{H}\}$ NMR ^{a,b} δ/ppm J/Hz	Assignment
$\text{IrCl}_2(\eta^1\text{-CH}_2\text{Ph})(\text{CO})(\text{PPh}_3)_2$ 1a	160.8 t	C=O
	$J(^{13}\text{C}^{31}\text{P}) = 7.5$	
	12.8 t	CH ₂
$\text{IrCl}_2(\eta^1\text{-CH}_2(p\text{-tolyl}))(\text{CO})(\text{PPh}_3)_2$ 1b	160.9 t	C=O
	$J(^{13}\text{C}^{31}\text{P}) = 7.5$	
	21.0 s	CH ₃
	12.7 t	CH ₂
$\text{IrCl}_2\{\text{C}(\text{O})\text{CH}_2\text{C}(\text{CH}_3)_3\}(\text{PPh}_3)_2$ 2	176.7 t	C=O
	$J(^{13}\text{C}^{31}\text{P}) = 3$	
	65.9 s	CH ₂
	33.7 s	C(CH ₃) ₃
	28.6 s	CH ₃
$\text{IrHCl}(\text{C}\equiv\text{CPh})(\text{PPh}_3)_3$ 3a	110.8 br	C _β
	96.1 dt	C _α
	$J(^{13}\text{C}^{31}\text{P})_{\text{cis}} = 11$	
	$J(^{13}\text{C}^{31}\text{P})_{\text{trans}} = 14$	
	113.2 s	C _β
$\text{IrHCl}(\text{C}\equiv\text{CPh})(\text{AsPh}_3)_3$ 4a	88.2 s	C _α
	114.6 s	C _β
	92.4 m, br	C _α
	66.2 d	CH ₂
	16.4 s	CH ₃
$[\text{IrCl}\{\eta^2\text{-CH}=\text{C}(\text{H})\text{COOMe}\}(\text{PPh}_3)_2]\text{BPh}_4$ 6	164.5 s	C=O
	135.9 s, br	C _α
	128.5 s	C _β
	51.3 s	CH ₃
	163.9 s	C=O
$[\text{IrCl}\{\eta^2\text{-CH}=\text{C}(\text{H})\text{COOMe}\}(\text{AsPh}_3)_2]\text{BPh}_4$ 7	136.1 s	C _α
	128.1 s	C _β
	51.0 s	CH ₃

^a In CD_2Cl_2 at 25 °C, unless otherwise noted. ^b Phenyl carbon resonances are omitted. ^c In C_6D_6 .




Fig. 1 Perspective view and labeling of the molecular structure of $\text{IrCl}_2(\eta^1\text{-CH}_2(p\text{-tolyl}))(\text{CO})(\text{PPh}_3)_2$ (**1b**). Phenyl groups of triphenylphosphine are omitted for clarity. Thermal ellipsoids and spheres are represented at the 50% probability level.

weaker with respect to those observed for a related $\text{Ir}(\text{III})\text{-CH}_2\text{Ar}$ complex: Ir–C 2.102(3)–2.125(3) Å, $\text{CH}_2\text{-C}(\text{Ar})$ 1.496(5) Å, in a pentacoordinated dibenzyl(bis(diphenylphosphinomethyl(dimethyl)silyl)amino-*P,P',N*)iridium derivative¹⁴ (structure determined at –40 °C). The intramolecular geometry for $\text{IrCl}_2(\eta^1\text{-CH}_2(p\text{-tolyl}))(\text{CO})(\text{PPh}_3)_2$ is characterized by a *syn* orientation of the $\text{CH}_2(p\text{-tolyl})$ group with respect to the carbonyl ligand (C44–Ir–C43–C25 21°, C25 ⋯ C44 3.21 Å), with the C43–C25 bond almost eclipsed with respect to the Ir–C44–O1 vector. The $\text{CH}_2(p\text{-tolyl})$ ligand is in fact sterically confined between the bulky triphenylphosphine groups and the equatorial orientation of the aryl ring on the carbonyl side is the least hindered by intramolecular non-bonded interactions. Accordingly, the carbonyl is displaced from the $\text{CH}_2(p\text{-tolyl})$ group, causing the widening of the C43–Ir–C44 angles (102(2)°) and the compression of the C44–Ir–Cl2 (77(2)°) angles. The

Table 3 Selected bond lengths (Å) and angles (°) for $\text{IrCl}_2(\eta^1\text{-CH}_2(p\text{-tolyl}))(\text{CO})(\text{PPh}_3)_2$ (**1b**)


Ir–C44	1.75(6)	C44–Ir–C43	102(2)
Ir–C43	2.24(5)	C44–Ir–P1	83(2)
Ir–P1	2.37(1)	C43–Ir–P1	94(1)
Ir–P2	2.42(1)	C44–Ir–P2	99(2)
Ir–Cl1	2.42(1)	C43–Ir–P2	88(1)
Ir–Cl2	2.506(9)	P1–Ir–P2	177.0(4)
C43–C25	1.62(5)	C44–Ir–Cl1	168(2)
O1–C44	1.24(5)	C43–Ir–Cl1	87(1)
C28–C46	1.58(6)	P1–Ir–Cl1	88.5(4)
		P2–Ir–Cl1	89.7(4)
		C44–Ir–Cl2	77(2)
		C43–Ir–Cl2	176(1)
		P1–Ir–Cl2	89.4(4)
		P2–Ir–Cl2	88.4(4)
		Cl1–Ir–Cl2	94.2(4)
		C25–C43–Ir	113(3)
		O1–C44–Ir	169(5)

two triphenylphosphine ligands eclipse each other (torsion C31–P2–P1–C7 6.6°).

The IR spectra of compounds **1** show a strong band at 2048 cm^{-1} , attributed to the ν_{CO} of the carbonyl ligand. In the labelled compound $\text{IrCl}_2(\eta^1\text{-CH}_2\text{Ph})(\text{C}\equiv^{18}\text{O})(\text{PPh}_3)_2$ (**1a- ^{18}O**) the $\nu_{\text{C}^{18}\text{O}}$ falls at 2006 cm^{-1} with $\Delta\nu$ of 42 cm^{-1} , which is in good agreement with the value calculated¹⁵ ($\Delta\nu = 47 \text{ cm}^{-1}$) from the change in reduced mass.


The NMR data of $\text{IrCl}_2(\eta^1\text{-CH}_2\text{Ar})(\text{CO})(\text{PPh}_3)_2$ **1** derivatives agree with their formulation and with the presence in solution of a geometry **I** like those of the solid state. In fact, the ^1H NMR spectra show, beside the signals of the phenyl protons of the PPh_3 ligands, a triplet at 3.55 ppm for **1a** and at 3.52 ppm for **1b** attributed to the methylene protons of the alkyl CH_2Ar ligand. In the spectra of **1b** a singlet at 2.18 ppm of the *p*-tolyl

substituent was also observed. The ^{13}C spectra of complexes **1** (Table 2) confirm the presence of CO and CH_2Ar ligands, showing a triplet at 160.8 (**1a**) and 160.9 (**1b**) ppm of the carbonyl carbon atom and a triplet at 12.8 (**1a**) and 12.7 (**1b**) ppm of the methylene CH_2Ar group. A ^1H , ^{13}C HMQC experiment clearly correlates this latter signal with the proton triplet at 3.55 ppm of the $\text{Ir}-\text{CH}_2-\text{Ar}$ group, in agreement with the proposed assignment. In the temperature range between +20 and -80°C , the $^{31}\text{P}\{^1\text{H}\}$ NMR spectra of **1** appear as a sharp singlet suggesting the magnetic equivalence of the two phosphine ligands. On the basis of these data a *cis-trans* structure **1** (Fig. 2) can be proposed.



1
cis-trans
Fig. 2

The iridium complex *mer*- or *fac*- $\text{IrHCl}_2(\text{PPh}_3)_3$ also reacts with *tert*-butylacetylene, $\text{HC}\equiv\text{C}(\text{CH}_3)_3$ to give a yellow solid which does not result to be the expected carbonyl-alkyl derivative like **1**, but an acyl complex of the $\text{IrCl}_2\{\eta^1-\text{C}(\text{O})-\text{CH}_2\text{C}(\text{CH}_3)_3\}(\text{PPh}_3)_2$ type (**2**), as shown in Scheme 2.




Scheme 2

The reaction with *tert*-butylacetylene involves also in this case the hydrolysis of the alkyne promoted by the traces of water present in the solvent to give the acyl complex **2**. This is confirmed by the red shift of the ν_{CO} observed in the labelled $\text{IrCl}_2\{\eta^1-\text{C}(^{18}\text{O})\text{CH}_2\text{C}(\text{CH}_3)_3\}(\text{PPh}_3)_2$ (**2- ^{18}O**) complex, prepared by adding H_2^{18}O to the reaction mixture, and by the preparation of $\text{IrCl}_2\{\eta^1-\text{C}(\text{O})\text{CD}_2\text{C}(\text{CH}_3)_3\}(\text{PPh}_3)_2$ (**2-d $_2$**) in the reaction with D_2O . The formulation of **2** as an acyl complex is supported by IR and NMR spectroscopic data. The IR spectrum shows a medium intensity band at 1681 cm^{-1} attributed to the ν_{CO} of the $\text{C}(\text{O})\text{CH}_2\text{C}(\text{CH}_3)_3$ group. This band shifts to 1649 cm^{-1} in the labelled **2- ^{18}O** complex ($\Delta\nu = 32\text{ cm}^{-1}$), according to the proposed attribution. The ^1H NMR spectra of **2** exhibits a singlet at 2.76 ppm, attributed to the CH_2 protons and a singlet at 0.69 of the CH_3 protons of the neopentyl $\text{CH}_2\text{C}(\text{CH}_3)_3$ moiety. However, strong support for the presence of the acyl ligand comes from the ^{13}C spectrum, which shows a triplet at 176.7 ppm due to the carbonyl carbon atom and three singlets at 65.9, 33.7 and 28.6 ppm attributed to the CH_2 , $\text{C}(\text{CH}_3)_3$ and CH_3 carbon atoms of the neopentyl group, respectively. APT NMR studies confirmed the proposed attribution, whereas a ^1H , ^{13}C HMQC NMR experiment clearly correlates the proton singlets at 2.76 and at 0.69 ppm with the $^{13}\text{C}\{^1\text{H}\}$ signals at 65.9 and at 28.6 ppm of the CH_2 and CH_3 groups of the neopentyl, in agreement with the proposed formulation.

In the temperature range between +20 and -80°C the ^{31}P spectrum appears as a sharp singlet suggesting the magnetic equivalence of the two phosphine ligands. The spectroscopic data, however, do not give conclusive information on the geometry of the complexes, which contain five ligands, but may reach the octahedral structure through an agostic interaction between the iridium centre and a C-H bond of a methyl substituent of the acyl ligand, or of the phenyl groups of the PPh_3 ligands.

A complex containing an acyl $\text{C}(\text{O})\text{R}$ ligand may be considered as a precursor of the derivative containing both the carbonyl and the alkyl ligand as in the $\text{IrCl}_2(\text{CO})(\eta^1-\text{CH}_2\text{Ar})-(\text{PPh}_3)_2$ (**1**) complexes obtained in the reaction of $\text{IrHCl}_2(\text{PPh}_3)_3$ with arylalkynes. We have therefore studied the behaviour of complex **2** in solution by NMR spectra and observed that while at room temperature it is very stable, at 60°C it slowly reacts to give a new species whose spectra indicate it to be the alkyl-carbonyl $\text{IrCl}_2\{\eta^1-\text{CH}_2\text{C}(\text{CH}_3)_3\}(\text{CO})(\text{PPh}_3)_2$ (**1c**) derivative formed by CO deinsertion of the acyl ligand (Scheme 3).



Scheme 3


The proton NMR spectra of a solution of **2** in THF-d_8 show the disappearance of the two singlets at 2.76 and at 0.69 ppm of the $\text{C}(\text{O})\text{CH}_2\text{C}(\text{CH}_3)_3$ group and the appearance of one triplet at 2.49 ppm and one singlet at 0.46 ppm attributable to the $\eta^1-\text{CH}_2\text{C}(\text{CH}_3)_3$ ligand of the alkyl compound **1c**. Also in the $^{31}\text{P}\{^1\text{H}\}$ NMR spectra a new singlet at -13.00 ppm appears, concurrent with the disappearance of the signal of **2**.

Unfortunately, this reaction is too slow as compared with the decomposition that takes place at 60°C to allow the carbonyl $\text{IrCl}_2\{\eta^1-\text{CH}_2\text{C}(\text{CH}_3)_3\}(\text{CO})(\text{PPh}_3)_2$ (**1c**) to be prepared in pure form. However, a comparison of its ^1H and ^{31}P NMR data with those of the related complexes **1a** and **1b** strongly supports the proposed formulation for this species.

We have also tested other acetylenes such as $\text{CH}_3\text{CH}_2-\text{CH}_2\text{C}\equiv\text{CH}$, $(\text{CH}_3)_3\text{SiC}\equiv\text{CH}$ and $\text{HC}\equiv\text{CCO}_2\text{R}$ in the reaction with $\text{IrHCl}_2(\text{PPh}_3)_3$, but neither alkyl- nor acyl-derivatives were obtained. The reaction seems to proceed with color change of the reaction mixture, but the NMR spectra do not reveal the presence of the hypothesized species and only intractable oils were separated from the reaction.

Although the number of tested acetylenes is limited, these results seem to indicate that the hydration of alkynes containing aryl substituents proceeds with a complete cleavage of the $\text{C}\equiv\text{C}$ bond affording both CO and the alkyl CH_2Ar groups, each behaving as a ligand. With alkyl- or silyl-substituents, instead, the hydration of $\text{RC}\equiv\text{CH}$ acetylenes does not give isolable products, except with the *tert*-butyl substituent which has the property to stabilize an acyl-iridium derivative of the type $\text{IrCl}_2\{\text{C}(\text{O})\text{CH}_2\text{C}(\text{CH}_3)_3\}(\text{PPh}_3)_2$.

The hydrolysis of terminal alkynes yielding **1** and **2** was studied extensively in order to obtain information useful to propose a mechanism for the reaction. Treating both *mer*- and *fac*- $\text{IrHCl}_2(\text{PPh}_3)_3$ in CD_2Cl_2 or THF-d_8 solution with phenylacetylene in a NMR tube one can observe the slow formation of **1a** and free PPh_3 . The addition of H_2O increases the reaction rate but does not change the profile of the spectra of the reaction mixture which shows the appearance of the signal of free PPh_3 immediately after the addition of $\text{PhC}\equiv\text{CH}$ whose signal slowly disappears. The lability of the PPh_3 ligand in $\text{IrHCl}_2(\text{PPh}_3)_3$, which was observed also in other reactions,^{8f,16} may suggest that the first step of the reaction is the substitution of PPh_3 with the alkyne giving an $\eta^1-\text{PhC}\equiv\text{CH}$ intermediate **[A]** (Scheme 4). A tautomerization of the π -coordinate alkyne on the iridium centre could then take place giving a vinylidene derivative **[B]**. This tautomerization is expected on the basis




Scheme 4

of the large number of precedents^{17,18} involving d⁶ metal complexes and should occur *via* a 1,2-hydrogen shift mechanism. We therefore attempted to detect this intermediate by monitoring the reaction course by NMR spectra (¹H, ¹³C, ³¹P), but did not observe any signal attributable to a vinylidene ligand. In fact, the absence of the characteristic carbene carbon signal¹⁷ between 350 and 280 ppm in the ¹³C NMR spectrum seems to exclude the presence of any vinylidene species, at least in detectable amounts. Therefore, the only new signals that appear in the spectrum of the reaction mixture, *i.e.* a singlet at 3.64 ppm for ¹H and a singlet at 114.1 ppm for ¹³C, may be tentatively attributed to the CH resonances (APT NMR experiment) of an η²-PhC≡CH ligand formed in the initial step of the reaction. However, the failure to detect the vinylidene intermediate does not mean that [Ir]=C(H)Ph species can be ruled out, because its further reaction with H₂O to give the final product can be faster than its formation, making its concentration undetectable.


Stable and isolable vinylidene complexes of iridium are well known^{17a,19} but are generally tetracoordinate complexes of Ir(I), with stable species of Ir(III) still unknown. This agrees with our difficulty to detect a Ir(III) vinylidene intermediate. Nevertheless, our results with D₂O and H₂¹⁸O which give **2-d₂** and **2-¹⁸O**, respectively, and the isolation, with HC≡CC(CH₃)₃, of acyl-complex **2** prompt us to propose a pathway similar to those recently proposed by Bianchini *et al.*^{3a} for the hydrolysis of terminal alkyne giving **1**, which involves a [Ir]-vinylidene complex [B] as the key intermediate.²⁰ This mechanism involves the attack of H₂O on the vinylidene [B] C_α carbon atom to give an hydroxycarbene^{21,22} IrHCl₂P₂[C(OH)CH₂Ar] intermediate [C], which can eliminate H₂ yielding the unsaturated acyl derivative IrCl₂P₂[C(O)CH₂Ar] [D]. From this acyl intermediate, the formation of the final alkyl-carbonyl complex **1** can be readily explained by CO deinsertion of the acyl ligand.

Vinyl and acetylide complexes

In parallel to the studies with terminal alkynes, we have also tested the behaviour of *mer*-IrHCl₂(PPh₃)₃ towards Li⁺[ArC≡C]⁻ and found that the reaction proceeds to give the alkynyl IrHCl(ArC≡C)(PPh₃)₃ (**3**) derivatives (Scheme 5). The related mixed-ligand IrHCl₂{PPh(OEt)₂}(PPh₃)₂ compound, as well as the arsino IrHCl₂(AsPh₃)₃ complex also react with Li⁺[ArC≡C]⁻ to give the alkynyl IrHCl(ArC≡C){PPh(OEt)₂}(PPh₃)₂ (**5**) and IrHCl(ArC≡C)(AsPh₃)₃ (**4**) derivatives which were isolated and characterized.



L = PPh₃ (**3**), AsPh₃ (**4**); Ar = Ph (**a**), *p*-tolyl (**b**)




L = PPh₃

Scheme 5

The complexes **3–5** are red–brown solids which are stable in the air, diamagnetic, moderately soluble in the more common organic solvents and non-electrolytes. The analytical and spectroscopic data support the proposed formulation. The IR spectra show a medium-intensity band at 2095–2099 cm⁻¹ attributed to the ν_{C≡C} of the acetylide ligand.²³ In the spectra a weak absorption at 2168–2156 cm⁻¹ also appears due to the ν_{IrH} of the hydride ligand. Support for the presence of the hydride

comes from the proton spectra which show, in the case of the phosphine complexes **3** and **5**, one doublet of triplets in the characteristic low-frequency region at –11.62 (**3a**), at –11.61 (**3b**) and at –11.78 (**5**) ppm. Taking into account that the ³¹P spectra are A₂B or AB₂ multiplets, the hydride pattern can be easily simulated using the appropriate A₂BX or AB₂X (X = H) model, with the parameters reported in the Experimental section. The high value of 140 Hz found for one of the two J_{PH} in complexes **3a** and **3b** also suggest that the hydride ligand is in a mutually *trans* position with one phosphine and in *cis* with the other two, as in a type-II geometry (Fig. 3). The presence of the acetylide ligand in the complexes is also confirmed by the ¹³C spectra (Table 2) which show the C_α and C_β carbon atoms of the C≡CPh group of **3a** at 96.1 and at 110.6 ppm, respectively. The C_α appears as one doublet of triplets due to the coupling with the phosphorus nuclei of the phosphine and the comparable values observed for the two J(¹³C³¹P) of 11 and 14 Hz, respectively, agrees with a type-II geometry for the complex.



II
Fig. 3



In the related mixed-ligand compound **5** the values of the two J_{PH} suggest a mutually *cis* position of the hydride with all the phosphines. Unfortunately, the broad multiplet that appears at 92.4 ppm in the ¹³C NMR spectrum for the C_α carbon atom of the C≡CPh ligand does not allow to determine the ²J(¹³CP) value and this prevents the unambiguous assignment of a geometry for the complex.

The ¹H NMR spectra of the arsino-complexes IrHCl(ArC≡C)(AsPh₃)₃ (**4**) show the hydride signal as a singlet at –14.18 (**4a**) and at –14.73 (**4b**) ppm, whereas in the ¹³C spectrum of **4a** the C_α and C_β carbon resonances of the acetylide ligand appear at 82.2 and at 113.2 ppm, respectively. These data support the proposed formulation but, also in this case, do not allow structural information to be obtained in solution.

Studies on the properties of the acetylide complexes **3–5** indicate that they are robust complexes, whose substitution reactions with several ligands (phosphine, CO, Cl⁻, *etc.*) are slow even in reflux conditions. Protonation reactions with Brønsted acids were also studied in order to test whether the proton attack takes place on the acetylide ligand to yield a [Ir]=C=C(H)Ar vinylidene complex, or on the hydride to give a dihydrogen [Ir]–(η²-H₂) derivative. The reaction was studied at different temperatures in a NMR tube by adding progressive amounts of HBF₄·Et₂O to **3a** in CD₂Cl₂ and the spectra show that the disappearance of the hydride ligand was already observed at –80 °C. However, neither signals attributable to an η²-H₂ complex formed by the protonation of the hydride, nor due to free H₂ were detected in the spectra of the protonated solution. Furthermore, even after the addition of an excess of HBF₄·Et₂O, the formation of a vinylidene complex was not detected by ¹H and ¹³C spectra. Instead, two doublets of multiplets at 8.38 and 5.53 ppm, respectively, appear in the proton spectra, which strongly suggest the formation of a vinyl complex.²⁴ The protonation reaction seems therefore to take place on the acetylide ligand, giving a coordinated terminal alkyne which can react with the hydride to yield the final vinyl complex (Scheme 6).


This vinyl complex is, unfortunately, thermally unstable and decomposed already at –10 °C, preventing its separation to the solid state.

These results prompted us to attempt a different method to prepare stable vinyl complexes and the reactions sequence reported in Scheme 7 does allow this synthesis to be achieved.



Because the reaction of *mer*-IrHCl₂(PPh₃)₃ with terminal alkyne does not give the insertion into the Ir–H bond yielding the vinyl species, but the hydrolysis of the ArC≡CH group giving **1**, we prepared the new hydride [**E**] by reacting IrHCl₂(PPh₃)₃ with AgCF₃SO₃. This complex (tentatively formulated as [IrHCl(PPh₃)₃]⁺CF₃SO₃[−]) was not isolated, but was treated in solution with terminal alkyne and, while with phenylacetylene only intractable products were obtained, with methyl propiolate a yellow solid was isolated after precipitation with NaBPh₄ and work-up, whose analytical and spectroscopic data indicated to be the chelate vinyl complex **6**. The related arsino derivative [IrCl{η²-CH=C(H)COOMe}(AsPh₃)₂]BPh₄ (**7**) was also obtained by reacting IrHCl₂(AsPh₃)₃ first with AgCF₃SO₃ and then with HC≡CCOOMe. These vinyl complexes **6**, **7** were formed through the insertion of methyl propiolate into the Ir–H bond of the hydride [**E**] and probably involved an initial coordination of the alkyne on the metal. The O-coordination of the carbonyl group to give the chelate vinyl ligand is followed by the dissociation of one PPh₃ ligand yielding a final pentacoordinate species.

Complexes **6** and **7** are yellow solids which are stable in the air and in solution of polar organic solvents, where they behave as 1 : 1 electrolytes.²⁵ Support for the presence of a chelate O-bonded vinyl ligand comes from the IR spectra which show the ν_{CO} as a medium-intensity band at a relatively low frequency²⁶ of 1570–1580 cm^{−1}. The ¹H NMR spectrum of the phosphine complex **6** shows, beside the signals of the phenyl protons, one doublet of triplets at 9.47 ppm attributed to the H_α proton and a doublet at 6.11 ppm due to the H_β proton of the C_αH_α=C_β(H_β)COOMe vinyl ligand. A singlet at 3.71 ppm of the methyl protons was also observed. Homodecoupling experiments and a ¹H COSY spectrum confirmed the proposed attribution. A value of 15 Hz for ³J_{HH} was also observed and, although it is somewhat high for a *cis* arrangement of the two vinyl protons,^{26,27} the IR spectra support the presence of the chelate (**III**) vinyl ligand (Fig. 4). The ¹³C spectra of **6** confirm the presence of the CH=C(H)COOCH₃ ligand showing a singlet at 164.5 ppm of the carbonyl carbon atom, a singlet at



III
Fig. 4

51.3 ppm of the methyl group, while the C_α and C_β vinyl carbons appear at 135.9 and 128.5 ppm, respectively. A ¹H, ¹³C HMQC experiment allowed us to correlate the vinyl C_α and C_β carbon signals to the proton multiplets at 9.47 and 6.11 ppm, in agreement with the proposed formulation. In the temperature range between +30 and −80 °C the ³¹P{¹H} NMR spectrum of **6** appears as an AB multiplet indicating the magnetic inequivalence of the two phosphine ligands. On the basis of the spectroscopic data alone we cannot unambiguously assign a geometry to our derivative **6**, although type **III** is plausible and can be proposed.

The proton NMR spectra of the arsino complex [IrCl{η²-CH=C(H)COOMe}(AsPh₃)₂]BPh₄ (**7**) show two doublets at 9.66 and 6.02 ppm with ³J_{HH} of 16 Hz attributed to the two vinyl protons H_α and H_β, respectively, of the vinyl group. The ¹³C spectra confirm the presence of the η²-ligand showing the characteristic signals of the C_α and C_β vinyl carbon atoms at 136.1 and at 128.1 ppm and those of the carbonyl and of the methyl group at 163.9 and 51.0 ppm, respectively. These data, therefore, support the proposed formulation for the arsino-vinyl derivative.

Conclusions

In this contribution we have reported a new example of hydrolysis of terminal alkynes promoted by Ir(III) hydride complexes which gives, depending on the nature of the alkyne, an alkyl-carbonyl IrCl₂(η¹-CH₂Ar)(CO)(PPh₃)₂ or an acyl IrCl₂{η¹-C(O)CH₂C(CH₃)₃}(PPh₃)₂ derivative. The structural characterization of an alkyl-carbonyl iridium complex is also described. Furthermore, the synthesis of new alkynyl IrHCl(ArC≡C)L₃ and chelate [IrCl{η²-CH=C(H)COOMe}-L₂]BPh₄ (L = PPh₃, AsPh₃) vinyl derivatives was achieved using *mer*-IrHCl₂L₃ as a precursor.

References and notes

- (a) J. Halpern, B. R. James and A. L. W. Kemp, *J. Am. Chem. Soc.*, 1966, **88**, 5142; (b) N. X. Hu, Y. Aso, T. Otsubo and F. Ogura, *Tetrahedron Lett.*, 1986, **27**, 6099; (c) W. D. Harman, J. C. Dobson and H. Taube, *J. Am. Chem. Soc.*, 1989, **111**, 3061; (d) T. M. M. Khan, S. B. Halligudi and S. Shukla, *J. Mol. Catal.*, 1990, **58**, 299; (e) W. Hiscox and P. W. Jennings, *Organometallics*, 1990, **9**, 1997; (f) Y. Fukuda and K. Utimoto, *J. Org. Chem.*, 1991, **56**, 3729; (g) J. Blum, H. Hummer and H. Alper, *J. Mol. Catal.*, 1992, **75**, 153; (h) I. K. Meier and J. A. Marsella, *J. Mol. Catal.*, 1993, **78**, 31; (i) M. Tokunaga, T. Suzuki, N. Koga, T. Fukushima, A. Horiuchi and Y. Wakatsuki, *J. Am. Chem. Soc.*, 2001, **123**, 11917.
- (a) O. M. Abu Salah and M. I. Bruce, *J. Chem. Soc., Dalton Trans.*, 1974, 2302; (b) M. I. Bruce and A. G. Swincer, *Aust. J. Chem.*, 1980, **33**, 1471; (c) B. P. Sullivan, R. S. Smythe, E. M. Kober and T. J. Meyer, *J. Am. Chem. Soc.*, 1982, **104**, 4701; (d) C. Mountassir, T. Ben Hadda and H. Le Bozec, *J. Organomet. Chem.*, 1990, **388**, C13; (e) S. Davies, J. P. McNally and A. J. Smallbridge, *Adv. Organomet. Chem.*, 1990, **30**, 30; (f) W. Knaup and H. Werner, *J. Organomet. Chem.*, 1991, **411**, 471.
- (a) C. Bianchini, J. A. Casares, M. Peruzzini, A. Romerosa and F. Zanobini, *J. Am. Chem. Soc.*, 1996, **118**, 4585; (b) M. L. Buil, M. A. Esteruelas, A. M. López and E. Oñate, *Organometallics*, 1997, **16**, 3169.
- (a) J. M. O'Connor and L. Pu, *J. Am. Chem. Soc.*, 1990, **112**, 9013; (b) M. V. Jiménez, E. Sola, A. P. Martínez, F. J. Lahoz and L. A. Oro, *Organometallics*, 1999, **18**, 1125; (c) C. S. Chin, D. Chong, B. Maeng, J. Ryu, H. Kim, L. Kim and H. Lee, *Organometallics*, 2002, **21**, 1739.
- (a) G. Albertin, P. Amendola, S. Antoniutti, S. Iannelli, G. Pelizzi and E. Bordignon, *Organometallics*, 1991, **10**, 2876; (b) G. Albertin, S. Antoniutti, E. Del Ministro and E. Bordignon, *J. Chem. Soc., Dalton Trans.*, 1992, 3203; (c) G. Albertin, S. Antoniutti and E. Bordignon, *J. Chem. Soc., Dalton Trans.*, 1995, 719; (d) G. Albertin, S. Antoniutti, E. Bordignon, F. Cazzaro, S. Iannelli and G. Pelizzi, *Organometallics*, 1995, **4**, 4114; (e) G. Albertin, S. Antoniutti, A. Bacchi, E. Bordignon and G. Pelizzi, *Organometallics*, 1995, **14**, 4126; (f) G. Albertin, S. Antoniutti, M. Bettiol, E. Bordignon and F. Busatto, *Organometallics*, 1997, **16**, 4959; (g) G. Albertin, S. Antoniutti, E. Bordignon and D. Bresolin, *J. Organomet. Chem.*, 2000, **609**, 10.

- 6 R. Rabinowitz and J. Pellon, *J. Org. Chem.*, 1961, **26**, 4623.
- 7 G. Balacco, *J. Chem. Inf. Comput. Sci.*, 1994, **34**, 1235.
- 8 (a) S. Park, A. J. Lough and R. H. Morris, *Inorg. Chem.*, 1996, **35**, 3001; (b) L. Vaska, *J. Am. Chem. Soc.*, 1961, **83**, 751; (c) R. G. Hayter, *J. Am. Chem. Soc.*, 1961, **83**, 1259; (d) N. Ahmad, S. D. Robinson and M. F. Uttley, *J. Chem. Soc., Dalton Trans.*, 1972, 843; (e) G. L. Geoffroy and R. Pierantozzi, *J. Am. Chem. Soc.*, 1976, **98**, 8054; (f) G. Albertin, S. Antoniutti, E. Bordignon and F. Menegazzo, *J. Chem. Soc., Dalton Trans.*, 2000, 1181.
- 9 SIR97: A New Program for Solving and Refining Crystal Structures: A. Altomare, M. C. Burla, M. Camalli, G. Cascarano, C. Giacovazzo, A. Guagliardi, A. G. Moliterni, G. Polidori and R. Spagna, *J. Appl. Crystallogr.*, 1999, **32**, 115.
- 10 G. M. Sheldrick, SHELXL97: Program for Structure Refinement; University of Goettingen, Germany, 1997.
- 11 L. J. Farrugia, *J. Appl. Crystallogr.*, 1999, **32**, 837.
- 12 F. H. Allen, O. Kennard and R. Taylor, *Acc. Chem. Res.*, 1983, **16**, 146.
- 13 R. H. Crabtree, M. Lavin and L. Bonneviot, *J. Am. Chem. Soc.*, 1986, **108**, 4032.
- 14 M. D. Fryzuk, P. A. MacNeil, R. L. Massey and R. G. Ball, *J. Organomet. Chem.*, 1989, **368**, 231.
- 15 D. M. P. Mingos, in *Comprehensive Organometallic Chemistry*, ed. E. W. Abel, F. G. A. Stone and G. Wilkinson, Pergamon Press, Oxford, UK, 1982, vol. 3, p. 1.
- 16 G. Albertin *et al.* manuscript in preparation.
- 17 (a) M. I. Bruce, *Chem. Rev.*, 1991, **91**, 197; (b) C. Bianchini, M. Peruzzini, A. Vacca and F. Zanobini, *Organometallics*, 1991, **10**, 3697; (c) M. D. Fryzuk, L. Huang, N. T. McManus, P. Paglia, S. J. Rettig and G. S. White, *Organometallics*, 1992, **11**, 2979; (d) J. R. Lumprey and J. P. Selegue, *J. Am. Chem. Soc.*, 1992, **114**, 5518; (e) I. de los Rios, M. Jimenez-Tenorio, M. C. Puerta and P. Valerga, *J. Am. Chem. Soc.*, 1997, **119**, 6529; (f) M. I. Bruce, B. C. Hall, N. N. Zaitseva, B. W. Skeldon and A. H. White, *J. Chem. Soc., Dalton Trans.*, 1998, 1793; (g) C. Bruneau and P. H. Dixneuf, *Acc. Chem. Res.*, 1999, **32**, 311.
- 18 (a) J. Silvestre and R. Hoffmann, *Helv. Chim. Acta*, 1985, **68**, 1461; (b) Y. Wakatsuki, N. Koga, H. Yamazaki and K. Morokuma, *J. Am. Chem. Soc.*, 1994, **116**, 8105.
- 19 (a) F. J. Garcia Alonso, A. Höhn, J. Wolf, H. Otto and H. Werner, *Angew. Chem., Int. Ed. Engl.*, 1985, **24**, 406; (b) A. Höhn, H. Otto, M. Dziallas and H. Werner, *J. Chem. Soc., Chem. Commun.*, 1987, 852; (c) H. Werner, A. Höhn and M. Schulz, *J. Chem. Soc., Dalton Trans.*, 1991, 777; (d) H. Werner, R. W. Lass, O. Gevert and J. Wolf, *Organometallics*, 1997, **16**, 4077; (e) K. Ilg and H. Werner, *Organometallics*, 2001, **20**, 3782.
- 20 A cationic vinylidene intermediate may also be formed by protonation of the coordinate alkyne in $\text{IrHCl}_2(\eta^2\text{-PhC}\equiv\text{CH})(\text{PPh}_3)_2$ [A] to give a formal Ir(v) cationic vinyl species $[\text{IrHCl}_2\{\text{CH}=\text{C}(\text{H})\text{Ph}\}(\text{PPh}_3)_2]^+$ which can undergo the Caulton-type^{20a} reaction to give the vinylidene $[\text{IrCl}_2\{\text{C}=\text{C}(\text{H})\text{Ph}\}(\text{PPh}_3)_2]^+$ and H_2 . Reaction of this intermediate with OH^- or with H_2O followed by deprotonation can then give the acyl complex [D]: (a) M. Olivan, E. Clot, O. Eisenstein and K. G. Caulton, *Organometallics*, 1998, **17**, 3091.
- 21 Hydroxycarbene metal complexes are generally unstable and thermally degrade to give acyl derivatives via HCl-elimination when chloride ligands are present in the complex framework.^{21a} Deprotonation of hydroxycarbene complexes with weak bases, giving formyl derivatives, was also observed.^{21b} (a) K. R. Grundy and J. Jenkins, *J. Organomet. Chem.*, 1984, **265**, 77; (b) M. A. Lilga and J. A. Ibers, *Organometallics*, 1985, **4**, 590.
- 22 For hydroxycarbene metal complexes, see: (a) R. A. M. Robertson, A. D. Poole, M. J. Payne and D. J. Cole-Hamilton, *J. Chem. Soc., Dalton Trans.*, 2000, 1817; (b) C. P. Casey, C. J. Czerwinski and R. K. Hayashi, *J. Am. Chem. Soc.*, 1995, **117**, 4189; (c) D. S. Barrett, C. Glidewell and D. J. Cole-Hamilton, *J. Chem. Soc., Chem. Commun.*, 1988, 1079; (d) V. Guerchais and C. Lapinte, *J. Chem. Soc., Chem. Commun.*, 1986, 663; (e) D. S. Barrett and D. J. Cole-Hamilton, *J. Organomet. Chem.*, 1986, **306**, C41; (f) W. E. Buhro, A. Wong, J. H. Merrifield, G.-Y. Lin, A. C. Constable and J. A. Gladysz, *Organometallics*, 1983, **2**, 1852; (g) E. O. Fischer and A. Riedel, *Chem. Ber.*, 1968, **101**, 156.
- 23 J. Manna, K. D. John and M. D. Hopkins, *Adv. Organomet. Chem.*, 1995, **38**, 79.
- 24 For vinyl complexes see: refs. 5a,c and (a) C. Bianchini, P. Barbaro, A. Meli, M. Peruzzini, A. Vacca and F. Vizza, *Organometallics*, 1993, **12**, 2505; (b) C. Bohanna, M. A. Esteruelas, F. J. Lahoz, E. Oñate and L. A. Oro, *Organometallics*, 1995, **14**, 4685.
- 25 W. J. Geary, *Coord. Chem. Rev.*, 1971, **7**, 81.
- 26 G. Albertin, S. Antoniutti and E. Bordignon, *J. Chem. Soc., Dalton Trans.*, 1995, 719 and references therein.
- 27 (a) B. Chamberlain and R. J. Mawby, *J. Chem. Soc., Dalton Trans.*, 1991, 2067; (b) J. M. Bray and R. J. Mawby, *J. Chem. Soc., Dalton Trans.*, 1989, 589.