



pounds, CNP was the most efficient; its use improved yields of cyanation products and led to a shortening of irradiation times. Photocyanation of *m*-dimethoxybenzene and mesitylene occurred efficiently when CNN



was used. On the other hand, 9,10-dicyanoanthracene, a typical electron acceptor, was not useful because of its rapid consumption during the irradiation and its low solubility in 9 : 1 acetonitrile-water.

Photocyanation under N_2 .—The photocyanation of phenanthrene under N_2 in the presence of DCNB gave 9,10-dihydrophenanthrene-9-carbonitrile (2a) and 9-cyanoanthracene (3a) in 55 and 16% yields, respectively, with 72% of DCNB being recovered. In the absence of DCNB, formation of the cyanation products was much slower. In the case of anthracene, (2b) and (3b) were obtained in 67% combined yield. Although (2b) could not be purified because of rapid aromatisation

naphthalene (1c) and 2,3-dimethylnaphthalene (1g) gave the corresponding 1-cyanonaphthalene compounds (3c and g) and their dihydro-isomers (2c and g). Although analytically pure (2g) could not be obtained, the structure was inferred from the spectroscopic properties; the n.m.r. spectrum showed a singlet for 2- CH_3 at δ 2.2 and a doublet for 3- CH_3 at δ 1.0, but none for olefinic protons. On the other hand, the other arenes gave complex



mixtures upon irradiation under N_2 . The results are summarised in Table 1.

Photocyanation under O_2 .—When the irradiation was carried out for oxygen-saturated solutions under O_2 , various arenes were efficiently cyanated to give the corresponding aromatic nitriles in fairly good yields, even in cases where irradiation under N_2 gave complex mix-

TABLE I
Photocyanation of arenes under nitrogen ^a

Arene (mmol)	A ^b (mmol)	NaCN (mmol)	Irradn. time/h	Products, yield (%) ^c	Conversion (%)	Recovery of A (%)
Phenanthrene (1a) (6.0)	DCNB (3.0)	(120)	10	(2a) 55; (3a) 16	89	72
	DCNB ^d		8	(2a) [52] ^d ; (3a) [11] ^d	(79)	
	^{d,e}		8	(2a) [4] ^d ; (3a) [3] ^d	(11)	
Anthracene (1b) (3.6)	DCNB (3.0)	(36)	12	(2b) + (3b) 25	79	0
	CNP (0.9)	(36)	4	(2b) 19 ^f ; (3b) 48 ^f	81	0
	CNP (1.25)	(100)	41	(2c) 44; (3c) 18	94	57
Naphthalene (1c) (10)						
2,3-Dimethylnaphthalene (1g) (10)	CNP (1.25)	(100)	6	(2g) [49]; (3g) [35]	100	71

^a For solutions in acetonitrile-water (100 ml; 9 : 1 v/v). ^b Electron acceptor: DCNB = *p*-dicyanobenzene, CNP = 9-cyanoanthracene. ^c Based on arenes unrecovered. Values in parentheses are g.l.c. yields. ^d Control runs for solutions in acetonitrile-water (3 ml; 9 : 1 v/v) of phenanthrene (0.2 mmol) and NaCN (4 mmol) [and DCNB (0.1 mmol)]. Yields are based on phenanthrene used. ^e In the absence of DCNB. ^f The product ratio was determined by n.m.r.

to (3b) during recrystallisation, its formation was indicated by the n.m.r. spectrum of a mixture of (2b) and (3b), which showed two singlets at δ 4.3 and 4.7 in a 2 : 1 ratio. The mixture of (2b) and (3b) was quantitatively converted into (3b) by refluxing a *p*-cymene solution in the presence of 5% Pd-C. Similarly, photocyanation of

tures. The results are shown in Table 2. In the case of anthracene, however, the irradiation of an oxygen-saturated solution gave no definite cyanation products but mainly intractable brownish material. Probably, photo-oxidation of anthracene (and also the products) would prevail.

It was again confirmed that the use of electron acceptors is remarkably advantageous in most cases, because of the occurrence of clean and rapid cyanation reactions. For example, the Figure shows that the


portant in determining whether or not the cyanation proceeds efficiently (Scheme 2). Under N_2 , the cyanated radicals are reduced by the anion radical of electron acceptors or undergo disproportionation, dimerisation,

TABLE 2
Photocyanation of arenes under oxygen ^a

Arene [mmol]	A (mmol) ^b	Irradn. time/h	Products, yield (%) ^c	Recovery of A (%)
Phenanthrene (1a) [6.0]	CNP (0.92)	10	(3a) 78 ^d	
Naphthalene (1c) [10]	CNP (1.25)	15	(3c) 65	59
1-Methylnaphthalene (1d) [10]	CNP (1.25)	15	(3d) 64	70
2-Methylnaphthalene (1e) [10]	CNP (1.25)	14	(3e) 63	72
2-Acetonaphthone ethylene glycol acetal (1f) [10]	CNP (1.25)	23	(3f) 51; (3f') 9	68
2,3-Dimethylnaphthalene (1g) [10]	CNP (1.25)	7	(3g) 63	51
2,6-Dimethylnaphthalene (1h) [10]	CNP (1.25)	10	(3h) 60	23
1-Methoxynaphthalene (1i) [10]	CNP (1.25)	15	(3i) 21; (3c) 35	79
2-Methoxynaphthalene (1j) [10]	CNP (1.25)	10	(3j) 51	53
<i>m</i> -Dimethoxybenzene (1k)	CNN (2.5)	24	(3k) 45	59
Mesitylene (1l) [10]	CNN (2.5)	36	(3l) 14	50

^a For solutions in acetonitrile-water (100 ml; 9 : 1 v/v) of NaCN (100 mmol). ^b Electron acceptor: DCNB = *p*-dicyanobenzene, CNP = 9-cyanophenanthrene, CNN = 1-cyanonaphthalene. ^c Yields based on arene used. ^d Yield based on (1a) and CNP used.

photocyanation of phenanthrene in the presence of DCNB is much more efficient than when it is absent, and that the photocyanation in the absence of DCNB gradually accelerated with an increase in conversion,




Plots of yield of (3a) vs. irradiation time for the photocyanation of (1a) under oxygen in the presence (○) and absence of DCNB (●); solutions in acetonitrile-water of (1a) (0.2 mmol) and NaCN (4 mmol) [and DCNB (0.1 mmol)].

probably because of the photosensitisation effect of the product (CNP) formed.

The ensuing reactions of the cyanated radicals, $ArH-CN$, formed by nucleophilic attack of cyanide anions on the cation radicals of arenes appear to be im-

and/or polymerisation reactions, depending on their reactivities. Under O_2 , however, oxidative aromatisation of the radicals to (3) by O_2 and its related species appears to overcome the other reaction pathways. This might be the reason why the photocyanation is successful under O_2 even in cases where complex mixtures are formed under N_2 .


In the case of 1- and 2-methoxynaphthalene, effects of oxygen saturation are notable (Table 3). In either the



SCHEME 2

presence or absence of CNP (electron acceptor), irradiation under O_2 improved the yield of the cyanation product of 2-methoxynaphthalene compared with that obtained under N_2 . Moreover, photocyanation of 1-methoxynaphthalene in the absence of CNP under O_2 gave 4-methoxynaphthalene-1-carbonitrile and 1-cyanonaphthalene in the ratio 3 : 4, similar to that for photocyanation in the presence of CNP, while the latter product was much more predominantly formed than the

former by irradiation under N_2 . These observations suggest that molecular oxygen may act as an electron acceptor towards the excited singlet methoxynaphthalenes; electron transfer from excited singlet aromatic



compounds to O_2 has been reported.¹¹ The formation of 1-cyanonaphthalene under N_2 would arise, at least in part, from the direct nucleophilic attack of cyanide anion on the methoxylated carbon of 1-methoxynaphthalene.

In summary, it is noted that direct efficient cyanation of various arenes with sodium cyanide occurs upon irradiation in the presence of electron acceptors to give cyanation products in fairly good yields. Moreover, favourable effects of oxygen saturation were observed. Thus, the photocyanation described here provides a

TABLE 3
Photocyanation of 1- and 2-methoxynaphthalenes^a

Arene	Irradn. time/h	Products, yield (%) ^b					
		Under O_2				Under N_2	
		CNP					
1-Methoxynaphthalene	12	(3i) 20	(3c) 25	(3i) 12	(3c) 16	(3i) 6	(3c) 28
2-Methoxynaphthalene	14	(3j) 54		(3j) 50		(3j) 17	

^a For solutions in acetonitrile-water (3 ml; 9:1 v/v) of the arenes (0.3 mmol) and NaCN (3 mmol) [and CNP (0.07 mmol)].
^b G.l.c. yields based on the arenes used.

convenient, useful synthetic tool for the preparation of aromatic nitriles, a method which does not require the use of any such specific reagents as crown ethers,¹² phase-transfer catalysts,¹³ or ammonium cyanides,¹³ nor require high dilution, even in the case of anthracene.

EXPERIMENTAL

M.p.s were taken on a Yanagimoto hot stage. ¹H N.m.r. spectra were recorded for solutions in CCl_4 with Me_4Si as internal standard at 100 MHz on a JEOL JNM-PS-100

TABLE 4
Characterisation of the nitriles

Compound	M.p. (°C) (lit. m.p.)	¹ H N.m.r. (δ) in CCl_4	Found (%) [Calc. (%)]
(2a)	84—85	3.05 (d, 2 H), 3.88 (dd, 1 H), 7.0—7.4 (m, 4 H), 7.5—7.6 (m, 2 H)	C, 87.65; H, 5.2; N, 6.8 [C, 87.77; H, 5.40; N, 6.82]
(2c)	b.p. 93 at 0.1 mmHg	3.5 (m, 2 H), 3.9 (t, 1 H), 5.9 (dt, 1 H), 6.5 (dt, 1 H), 7.3—7.9 (m, 4 H)	C, 85.2; H, 5.6; N, 9.0 [C, 85.13; H, 5.85; N, 9.03]
(2g)	b.p. 70 at 0.01 mmHg	1.0 (d, 3 H), 2.2 (s, 3 H), 2.4—2.6 (m, 2 H) 2.8—3.1 (m, 1 H), 6.9—7.2 (m, 4 H)	
(3d)	193—194 ^a	2.7 (s, 3 H), 7.1—7.6 (m, 5 H), 7.9—8.2 (m, 1 H)	
(3e)	84—85	2.7 (s, 3 H), 7.2—7.9 (m, 5 H), 8.15 (d, 1 H)	
(3f)	(84.5—85.5) ^c 80—81	1.68 (s, 3 H), 3.76—4.16 (m, 4 H), 7.44 (t, 1 H), 7.64 (dd, 1 H), 7.76—8.06 (m, 2 H), 8.26 (br, s, 1 H)	C, 75.05; H, 5.35; N, 5.8 [C, 75.30; H, 5.48; N, 5.85]
(3f')	85	1.64 (s, 3 H), 3.7—4.16 (m, 4 H), 7.46—7.94 (m, 3 H), 7.96 (d, 1 H), 8.1 (br d, 1 H), 8.2 (br d, 1 H)	C, 75.0; H, 5.4; N, 5.85 [C, 75.30; H, 5.48; N, 5.85]
(3g)	82.5—83.5 (82—83) ^d	2.4 (s, 3 H), 3.6 (s, 3 H), 7.4—7.7 (m, 4 H), 8.0—8.1 (m, 1 H)	
(3h)	88 (86—87) ^e	2.5 (s, 3 H), 2.64 (s, 3 H), 7.2—7.5 (m, 3 H), 7.7 (d, 1 H), 7.95 (d, 1 H)	
(3i)	101—102 (102.5) ^f	4.0 (s, 3 H), 6.5 (d, 1 H), 7.3—7.7 (m, 3 H), 7.8—8.2 (m, 2 H)	
(3j)	96—97 (95—96) ^g	4.0 (s, 3 H), 7.0—8.0 (m, 6 H)	
(3k)	89—90 (89) ^h	3.86 (s, 3 H), 3.9 (s, 3 H), 6.44 (br s, 1 H), 6.50 (dd, 1 H), 7.44 (d, 1 H)	
(3l)	51—52 (50—52) ⁱ	2.2 (s, 3 H), 2.36 (s, 6 H), 6.7 (s, 2 H)	

^a M.p. of 4-Methylnaphthalene-1-carboxamide. ^b R. C. Fuson, B. C. McKusick, and F. W. Spangler, *J. Am. Chem. Soc.*, 1945, **67**, 597. ^c R. C. Fuson, C. H. McKeever, and L. C. Behr, *J. Am. Chem. Soc.*, 1941, **63**, 2774. ^d P. H. Gore, C. K. Thadani, and S. Thorburn, *J. Chem. Soc. (C)*, 1968, 2506. ^e P. H. Gore and M. Yusuf, *J. Chem. Soc. (C)*, 1971, 2586. ^f A. Fischer, M. A. Riddolls, and J. Vaughan, *J. Chem. Soc. (B)*, 1966, 106. ^g R. C. Fuson and D. H. Chadwick, *J. Org. Chem.*, 1948, **13**, 484. ^h G. Lohaus, *Chem. Ber.*, 1967, **100**, 2719. ⁱ R. C. Fuson and J. J. Denton, *J. Am. Chem. Soc.*, 1941, **63**, 654.

spectrometer or at 60 MHz on a JEOL JNM-60 spectrometer, i.r. spectra for solutions in CCl_4 on a Shimadzu IR-400, mass spectra on a Hitachi RMU-6E spectrometer, and g.l.c.-mass spectra on a Hitachi RMU-4 spectrometer. G.l.c. was carried out on a Shimadzu GC-3BF dual-column instrument with flame-ionisation detectors. Columns used were 75 cm \times 4 mm of 5% SE-30 on Celite 545, 5% Ucon Oil LB-550X on Celite 545, and 10% PEG-20M on Shimalite W.

Spectral grade acetonitrile was used without further purification. All the arenes and *p*-dicyanobenzene were used as received (Tokyo Kasei). 1-Cyanonaphthalene (Tokyo Kasei) was chromatographed on silica gel and then distilled *in vacuo*. 9-Cyanophenanthrene was prepared according to the literature method¹⁴ and recrystallised from methanol.

General Procedure for Photocyanation under N₂.—To a solution of sodium cyanide (36–120 mmol) in water (10 ml) was added acetonitrile (90 ml) containing the arene (3.6–10 mmol) and the electron acceptor (0.9–3 mmol). In the case of anthracene, 150 ml of the acetonitrile solution was used. The solution was introduced into a Pyrex vessel, N₂ was bubbled through for 20 min, and the solution was irradiated with an Eikosha PIH-300 high-pressure mercury lamp under cooling with water. The progress of the reaction was followed by g.l.c. To the irradiated solution was added brine (50 ml) and then diethyl ether (500 ml). After vigorous shaking, the ethereal layer was separated, washed three times with water, dried (MgSO_4), and then evaporated. The residue was chromatographed on silica gel (Merk, Art 7734, 70–230 mesh), using hexane and 10–30% benzene in hexane as the eluants. The results are summarised in Table 1.

General Procedures for Photocyanation under O₂.—A gentle stream of pure oxygen was bubbled through a solution in acetonitrile–water (100 ml) of the arene (6–10 mmol), the electron acceptor (0.92–5.0 mmol), and sodium cyanide (100 mmol) in a Pyrex vessel. The reaction vessel was then connected to a balloon filled with pure oxygen and

the solution was irradiated under cooling with water. After the arenes had been completely consumed, the photolysates were extracted with diethyl ether and treated in a manner similar to the work-up described above. In some cases, the products and electron acceptors were isolated without the use of column chromatography. The results are summarised in Table 2.

This work was supported by the Ministry of Education of Japan.

[0/914 Received, 16th June, 1980]

REFERENCES

- ¹ Part 33, M. Yasuda, C. Pac, and H. Sakurai, *Bull. Chem. Soc. Jpn.*, 1980, **53**, 502.
- ² K. Friedrich and K. Wallenfels in 'The Chemistry of the Cyano Group,' ed. Z. Rappoport, Interscience, New York, 1970, p. 67.
- ³ J. Cornelisse and E. Havinga, *Chem. Rev.*, 1975, **75**, 353; E. Havinga and J. Cornelisse, *Pure Appl. Chem.*, 1976, **47**, 1.
- ⁴ J. A. J. Vink, C. M. Lok, J. Cornelisse, and E. Havinga, *J. Chem. Soc., Chem. Commun.*, 1972, 710; S. Nilsson, *Acta Chem. Scand.*, 1973, **27**, 329.
- ⁵ K. Koyama, T. Suzuki, and S. Tsutsumi, *Tetrahedron Lett.*, 1965, 627; K. Yoshida and S. Nagase, *J. Am. Chem. Soc.*, 1979, **101**, 4268.
- ⁶ K. Mizuno, C. Pac, and H. Sakurai, *J. Chem. Soc., Chem. Commun.*, 1975, 553.
- ⁷ C. Pac, A. Nakasone, and H. Sakurai, *J. Am. Chem. Soc.*, 1977, **99**, 5806.
- ⁸ D. Rehm and A. Weller, *Isr. J. Chem.*, 1970, **8**, 259.
- ⁹ T. Hino, H. Masuhara, and N. Mataga, *Bull. Chem. Soc. Jpn.*, 1970, **49**, 394.
- ¹⁰ T. Asanuma, T. Gotoh, A. Tsuchida, M. Yamamoto, and Y. Nishijima, *J. Chem. Soc., Chem. Commun.*, 1977, 485.
- ¹¹ R. Potashnik, C. R. Goldshmidt, and M. Ottolenghi, *Chem. Phys. Lett.*, 1971, **9**, 424; Y. Taniguchi, A. Mioshi, N. Mataga, H. Shizuka, and T. Morita, *ibid.*, 1972, **15**, 223.
- ¹² R. Beugelmas, M.-T. Le Goff, J. Pusset, and G. Roussi, *J. Chem. Soc., Chem. Commun.*, 1976, 377.
- ¹³ R. Beugelmas, H. Ginsburg, A. Lecas, M.-T. Le Goff, and G. Roussi, *Tetrahedron Lett.*, 1978, 3271.
- ¹⁴ J. E. Callen, C. E. Dornfeld, and G. H. Coleman, *Org. Synth.*, Coll. Vol. III, 1958, p. 212.