

Benzylic Substitution of Benzyl Methyl Ether


By Ming Kuo Yeh, Department of Chemistry, Tamkang University, Tamsui, Taiwan

Treatment of benzyl methyl ether with *n*-butyl-lithium–tetramethylethylenediamine in hexane gave benzylic-lithiation, without *ortho*-lithiation or a Wittig type rearrangement. The benzylic-lithiated benzyl methyl ether was then allowed to react with benzophenone, 4-methoxybenzophenone, butyl bromide, ethyl phenylacetate, and 4-chlorobenzophenone.

THE Wittig rearrangement is well documented^{1,2} and occurs in ethers when in the presence of a strong base (*e.g.* sodamide or organolithium) [see equation (1) where R¹ and R² may be alkyl, aryl, or vinyl, and one of the




hydrogens may be replaced by an alkyl or aryl group]. Initially, the mechanism was thought to involve proton abstraction by base from the benzylic carbon, followed by either a concerted or an ion-pair elimination–addition mechanism (Scheme 1).¹ More recent evidence suggests




SCHEME 1

that the radical-pair mechanism² shown in Scheme 2 occurs after removal of the proton by the base. One of the radicals in the radical pair is a ketyl radical. There is evidence that, at least in some cases, the radical-pair mechanism accounts for only a portion of the product, and a concerted mechanism may also take place.³ The radical-pair mechanism has been suggested for cases where R² is alkyl or aryl and a [2,3] sigmatropic rearrangement when R² is allylic.




SCHEME 2

Following reports of *ortho*-lithiation of aromatic compounds such as that observed by Jones and Hauser⁴ [see equation (2)], Gschwend and Hamdan,⁵ and Meyer




and Seebach⁶ [see equation (3)], a parallel reaction with benzyl methyl ether was attempted in hexane. Tetramethylethylenediamine (TMEDA) was added to enhance the ability of *n*-butyl-lithium to extract the *ortho*-proton.



Studies showed that, with 1 or 2 equiv. of *n*-butyl-lithium, neither *ortho*-lithiation nor the usual Wittig rearrangement took place; removal of the benzylic proton resulted in the formation of a hexane-stable anion. Addition of *n*-butyl bromide to this and work-up gave 1-methoxy-1-phenylpentane in 85% yield. An equimolar proportion of *n*-butyl-lithium gave a yield of product > 2 mol equiv.

Other electrophiles gave similar results (see Table 1). The low yield of product from ethyl phenylacetate arises probably as a result of the acidity of the initial



product, 1-methoxy-1,3-diphenylacetone, which could undergo further condensation.

All the reported¹⁻⁶ solvents used for the Wittig rearrangements and related reactions have been moderately polar [*e.g.* hexane–tetrahydrofuran (THF), ether, ether–THF, and THF]. Use of a non-polar solvent, such as hexane with TMEDA either avoids or limits the Wittig rearrangement. It has been recently reported that carbanions derived from benzyl allyl ethers and thio-ethers undergo different rearrangements depending upon the position of metallation.⁷ Since the solvent used in these cases was THF–hexane, the above finding indicates that changing the reaction medium should also vary the ratio of different types of rearrangement products.

EXPERIMENTAL

M.p.s were determined on a Mel-Tem apparatus, and were uncorrected. I.r. spectra were measured on a Perkin-Elmer Model 237 or a Beckman Model 4250. ^1H N.m.r. spectra were recorded on a JEOL NM-60 HL spectrometer, using tetramethylsilane as the internal standard. Mass spectra were taken on a Hitachi GC-Mass M-52. Elementary analyses were carried out at the Microanalytical Laboratory, Chung-Shan Institute of Science and Technology. Bu^nLi in hexane was purchased from Tokyo Kasei Co., Japan, or E. Merck Co., Germany. Hexane was pre-dried over sodium, calcium hydride, or molecular sieve.

Lithiation of Benzyl Methyl Ether.—To a solution of

TABLE 1

Benzylic substitution of benzyl methyl ether		
Electrophile	Product	Yield (%)
Ph_2CO	$\text{PhCH}(\text{OMe})\text{C}(\text{OH})\text{Ph}_2$ (2)	76
BuBr	$\text{PhCH}(\text{OMe})\text{Bu}^a$ (3)	85
$\text{PhCO}\cdot\text{C}_6\text{H}_4\cdot\text{OMe}-p$	$\text{PhCH}(\text{OMe})\text{C}(\text{OH})(\text{C}_6\text{H}_4\text{OMe}-p)\text{Ph}^a$ (4)	65
$\text{PhCO}\cdot\text{C}_6\text{H}_4\text{Cl}-p$	$\text{PhCH}(\text{OMe})\text{C}(\text{OH})(\text{C}_6\text{H}_4\text{Cl}-p)\text{Ph}^a$ (5)	40
$\text{EtO}\cdot\text{CO}\cdot\text{CH}_2\text{Ph}$	$\text{PhCH}(\text{OMe})\text{C}(\text{O})\text{CH}_2\text{Ph}^a$ (6)	22

^a New compounds, the analytical data for which are given in Table 2.

TABLE 2
Analytical data for reaction products

Product	M.p. (°C) (B.p.)	m/e (M^+)	^1H N.m.r. ^a (δ)	ν_{max} (cm^{-1})	Elemental analysis Found (Calc.) (%)		
					C	H	Cl
(2)	137—137.5 ^b	304	3.26 (s, 4) 4.99 (s, 1) 6.90—7.70 (m, 15)	3 500 1 070 2 960	82.85 (82.76)	6.45 (6.57)	
(3)	(145—150)	178	0.91 (t, 3) 1.10—2.61 (m, 6) 3.95 (t, 1) 6.54—7.48 (m, 5)	1 460 1 100 3 495			
(4)	141.5—142		3.30 (s, 3) 3.70 (s, 3) 5.05 (s, 1) 6.50—7.71 (m, 14)	1 090 1 020 3 500	78.55 (78.45)	6.5 (6.53)	
(5)	155—155.5		3.20 (s, 3) 3.26 (s, 1) 4.90 (s, 1) 6.50—7.72 (m, 14)	1 100 1 070 2 980	73.5 (73.60)	5.55 (5.55)	10.65 (10.37)
(6)			3.26 (s, 3) 3.70 (s, 2) 4.71 (s, 1) 6.86—7.50 (m, 10)	1 732 1 095	80.2 (80.00)	6.55 (6.67)	

^a Solvents: CCl_4 for compounds (2) and (4); DCCl_3 for compounds (3), (5), and (6). ^b Lit., 139 °C (H. Gross and J. Freiberg, *Chem. Ber.*, 1966, **99**, 3260).

benzyl methyl ether (2.44 g, 0.02 mol) and TMEDA (2.32 g, 0.02 mol) in dry hexane (100 ml) at -10 °C was added, in drops, Bu^nLi (0.02 mol) in hexane *via* a rubber septum. After 3 h the solution gradually turned brownish red and was assumed to contain 0.02 mol of lithiated benzyl methyl ether (1).

General Procedure for the Reaction of (1) with Electrophiles.—To the mixture containing (1) (0.02 mol) at -10 °C was added, in drops, a solution of the electrophile (0.02 mol) in hexane (40—80 ml). The mixture was then stirred for 3 h, after which 6N HCl (30 ml) was added. The layers were separated and the aqueous layer was extracted with ether. The combined organic layers were dried and evaporated. The residue was then purified by distillation or recrystallization to give the benzylic-substituted benzyl

methyl ether. The yields and analytical data for the products are given in Tables 1 and 2, respectively.

[0/1537 Received, 8th October, 1980]

REFERENCES

- D. J. Cram, 'Fundamentals of Carbanion Chemistry,' Academic Press, New York, 1965, pp. 230—233.
- J. March, 'Advanced Organic Chemistry,' McGraw-Hill Book Co., New York, 1977, pp. 1015—1016.
- J. F. Garst and C. D. Smith, *J. Am. Chem. Soc.*, 1976, **98**, 1526.
- F. N. Jones and C. R. Hauser, *J. Org. Chem.*, 1962, **27**, 701.
- H. W. Gschwend and A. Hamdan, *J. Org. Chem.*, 1975, **40**, 2008, and references therein.
- N. Meyer and D. Seebach, *Angew. Chem., Int. Ed. Engl.*, 1978, **17**, 521.
- J. F. Biellmann, J. B. Ducep, and D. Schirlin, *Tetrahedron*, 1980, **36**, 1249, and references therein.