


Homochiral Syntheses of (-)-Dihydrocorynantheol

Masataka Ihara, Nobuaki Taniguchi, Ken Yasui, and Keiichiro Fukumoto*
 Pharmaceutical Institute, Tohoku University, Aobayama, Sendai 980, Japan

A Corynanthe-type indole alkaloid, (-)-dihydrocorynantheol (7), was enantioselectively synthesized from (+)-(4*S*,5*R*)-4-(2-benzyloxyethyl)-5-ethyl-3,4,5,6-tetrahydro-2-pyrone (3), derived from ethylmalonic acid. The synthetic compound (7) was identical in all respects with a sample prepared from yohimbine (8).

Recently, we developed asymmetric syntheses of chiral propane-1,3-diols by way of chiral half-esters of monoalkylmalonic acids, and discovered the diastereoselective formation of a single isomer of the menthyl half-ester (1) by crystallisation-induced



Scheme 1. Reagents: i, tryptamine; ii, MsCl, Et₃N; iii, KH, 18-crown-6; iv, POCl₃; v, NaBH₄; vi, H₂, PdCl₂.


asymmetric transformation.¹ The half-ester (1) was enantioselectively converted, through radical cyclisation of the (*Z*)- α,β -unsaturated ester (2), into the optically pure lactone (3), which was transformed into several ipecacuanha alkaloids.² As an extension of this work, the lactone (3) was further transformed into a Corynanthe-type indole alkaloid, (-)-dihydrocorynantheol (7).³⁻⁵ Preparation of the same enantiomer (7) was also carried out starting from natural yohimbine (8). We report here homochiral syntheses of (-)-dihydrocorynantheol (7).

Condensation of the lactone (3) with tryptamine gave the hydroxy amide (4), $[\alpha]_D^{25} + 5.64^\circ$ (CHCl₃), in 91% yield. Since reaction of compound (4) with phosphorus trichloride oxide did not produce the expected cyclised product (6), the amide (4) was transformed into the lactam (5), $[\alpha]_D^{24} + 16.97^\circ$ (CHCl₃) in 70% overall yield by successive methanesulfonylation followed by ring formation using potassium hydride in the presence of 18-crown-6. Treatment of the lactam (5) with phosphorus trichloride oxide in acetonitrile provided the iminium salt (6), which was reduced with sodium borohydride to the corresponding hexahydroquinolizidine, in 97% overall yield, as a mixture of two diastereoisomers. Catalytic hydrogenation of the mixture with palladium(II) dichloride under hydrogen atmosphere in a mixture of chloroform and methanol gave rise to debenzylation accompanied by isomerisation^{4a} to furnish (-)-dihydrocorynantheol (7), m.p. 178–181 °C; $[\alpha]_D^{23} - 20.41^\circ$ (CHCl₃) {lit.,^{3a} m.p. 181–183 °C; $[\alpha]_D^{27} - 19^\circ$ (CHCl₃)}, as a single isomer in 77% yield (Scheme 1).

The same enantiomer (7) of the alkaloid was readily synthesized from yohimbine (8) as follows. The yohimbone (9),⁶ derived from yohimbine (8), was first converted into the enamine, which was then treated with propane-1,3-dithiol di(toluene-*p*-sulphonate)⁷ in the presence of triethylamine. Treatment of the dithioacetal (10), $[\alpha]_D^{25} - 25.3^\circ$ (CHCl₃), obtained in 97% overall yield from ketone (9), with potassium hydroxide in refluxing 2-methylpropan-2-ol⁸ afforded the carboxylic acid, which was converted into the corresponding methyl ester (11), $[\alpha]_D^{26} + 25.3^\circ$ (CHCl₃), in 52% overall yield for two steps. Reduction of ester (11) with lithium aluminium hydride produced quantitatively the alcohol (12), $[\alpha]_D^{25} + 4.65^\circ$ (CHCl₃), which was further reduced with W-2 Raney nickel to afford (-)-dihydrocorynantheol (7) (Scheme 2). All physical properties of the product (7), obtained in 98% yield, were identical with those of the first specimen. Thus, homochiral syntheses of (-)-dihydrocorynantheol (7) were accomplished starting from both the key lactone (3) and yohimbine (8).

Experimental

General Methods.—M.p.s were measured on a Yanako micro melting-point apparatus and are uncorrected. IR spectra were recorded for CHCl₃ solutions on a Hitachi 260-10 spectrophotometer. NMR spectra were measured for CDCl₃ solutions



Scheme 2. Reagents: i, pyrrolidine; ii, TsS[CH₂]₃STs, Et₃N; iii, KOH; iv, MeOH, conc. H₂SO₄; v, LiAlH₄; vi, Raney Ni.

with JEOL JNM-PMX-60, JEOL-FX-90A, and JNM-GX-500 spectrometers. Chemical shifts are reported relative to internal SiMe₄. Mass spectra were taken on JEOL-JMS-O1SG-2 and JEOL-DX-300 spectrometers. Optical rotations were determined on a JASCO-DIP-340 polarimeter. All new compounds described in the Experimental section were homogeneous on TLC and HPLC.

(3*S*,4*R*)-(+)-3-(2-Benzoyloxyethyl)-4-hydroxymethyl-N-[2-(indol-3-yl)ethyl]hexanamide (4).—A mixture of the lactone (3)² (270.9 mg, 1.03 mmol) and tryptamine (412 mg, 2.57 mmol) in dry toluene (5 ml) was heated under reflux for 6 h. After removal of the solvent, the residue was heated at 130 °C for an additional 3 h and then diluted with toluene. After evaporation, silica gel column chromatography of the residue with CHCl₃-AcOEt (1:4, v/v) as eluant gave the amide (4) (396.7 mg, 91%) as an oil (Found: *M*⁺, 422.2549. C₂₆H₃₄N₂O₃ requires *M*, 422.2570); [α]_D²⁵ + 5.64° (c 4.56, CHCl₃); ν_{max} 3 550–3 150 (OH and NH) and 1 660 cm⁻¹ (C=O); δ_H(90 MHz) 0.71–1.02 (3 H, m, CH₂Me), 1.97–2.28 (2 H, m, CH₂CONH), 2.63–3.00 (3 H, m, CH₂Ar and OH), 3.29–3.71 (6 H, m, 2 × CH₂O and CH₂N), 4.40 (2 H, s, OCH₂Ph), 5.81–6.10 (1 H, m, NHCO), 6.90–7.65 (10 H, m, 5 × ArH and Ph), and 8.14–8.33 (1 H, br s, NH).

(4*S*,5*R*)-(+)-4-(2-Benzoyloxyethyl)-5-ethyl-N-[2-(indol-3-yl)ethyl]piperidin-2-one (5).—To a stirred solution of the above amide (4) (68.6 mg, 0.162 mmol) in dry benzene (5 ml) at 5 °C were added a solution of Et₃N (0.025 ml, 0.18 mmol) in dry benzene (0.25 ml) and a solution of methanesulphonyl chloride (0.013 ml, 0.17 mmol) in dry benzene (0.1 ml). After being stirred at the same temperature for 0.5 h, the reaction mixture was diluted with benzene, washed successively with saturated aq. NaHCO₃ (× 2), 1% HCl, and water, and then dried (Na₂SO₄),

and the solvent was evaporated off under protection from moisture. Water was completely removed by azeotropic distillation (× 2) using benzene. The resulting mesyl ester was used for the next reaction without further purification.

To a stirred suspension of 18-crown-6 (8.0 mg, 0.03 mmol) and KH (65 mg, 1.6 mmol) in dry 1,2-dimethoxyethane (DME) (5 ml) at 0 °C was added slowly a solution of the above mesyl ester in dry DME (2 ml). The mixture was stirred at room temperature for 0.5 h, poured into saturated aq. NH₄Cl, and extracted with CH₂Cl₂. The combined organic layers were washed with brine, dried (Na₂SO₄), and evaporated to give a residue. Silica gel column chromatography of the residue with benzene-acetone (85:15, v/v) as eluant afforded the lactam (5) (46.1 mg, 70% yield for two steps) as a viscous oil (Found: *M*⁺, 404.2473. C₂₆H₃₂N₂O₂ requires *M*, 404.2464); [α]_D²⁴ + 16.97° (c 2.52, CHCl₃); ν_{max} 1 620 cm⁻¹ (C=O); δ_H(500 MHz) 0.74 (3 H, t, *J* 7.6 Hz, CH₂Me), 2.07 (1 H, dd, *J* 8.7 and 16.3 Hz, CHHCON), 2.53 (1 H, dd, *J* 8.7 and 16.3 Hz, >CHHCON), 2.87 (1 H, dd, *J* 8.7 and 13.6 Hz, >CHCHHNCO), 3.03 (2 H, t, *J* 7.6 Hz, CH₂Ar), 3.16 (1 H, dd, *J* 5.4 and 13.6 Hz, CHCHHNCO), 3.40–3.75 (4 H, m, CH₂CH₂N and CH₂O), 4.49 (2 H, s, OCH₂Ph), 7.00–7.71 (10 H, m, 5 × ArH and Ph), and 8.10–8.22 (1 H, br s, NH).

(2*R*,3*R*,12*bS*)-(-)-3-Ethyl-2-(2-hydroxyethyl)-1,2,3,4,6,7,12,12*b*-octahydroindolo[2,3-*a*]quinolizine (7) (Dihydrocorynantheol).—(a) A mixture of the lactam (5) (15.1 mg, 0.0373 mmol) and freshly distilled POCl₃ (0.2 ml, 2.15 mmol) in dry MeCN (2 ml) was heated under reflux for 2.5 h under protection from moisture; during the reaction hydrogen chloride gas was evolved. After removal of the solvent under reduced pressure, the reagents were removed by azeotropic distillation with toluene (× 3) to give the iminium salt (6), which was used for the next reaction without further purification.

To a stirred solution of the above salt in distilled MeOH (2 ml) at 0 °C was added NaBH₄ (12 mg, 0.32 mmol). After being stirred at the same temperature for 1 h, the reaction mixture was poured into water, and the mixture was extracted with CHCl₃. The combined extracts were washed with brine, dried (K₂CO₃), and evaporated. Silica gel column chromatography of the residue with CHCl₃-MeOH (95:5, v/v) as eluant afforded quinolizidines (14 mg, 97%) as a yellowish oil.

A solution of the above product (14 mg, 0.036 mmol) in a mixture of distilled MeOH (2 ml) and distilled CHCl₃ (0.4 ml) was stirred with PdCl₂ (10 mg, 0.057 mmol) at room temperature under hydrogen (1 atm) for 40 min. The reaction mixture was poured into 10% aq. ammonium hydroxide. After extraction with CHCl₃, the combined extracts were washed with brine, dried (K₂CO₃), and evaporated to give a residue, which was subjected to silica gel column chromatography. Elution with CHCl₃-MeOH (96:4, v/v) afforded (-)-dihydrocorynantheol (7) (8.3 mg, 77%) as crystals, m.p. 178–181 °C (lit.,^{3a} 181–183 °C) (Found: *M*⁺, 298.2044. Calc. for C₁₉H₂₆N₂O: *M*, 298.2045); [α]_D²³ - 20.41° (c 0.25, CHCl₃) {lit.,^{3a} [α]_D²⁷ - 19° (c 1.02, CHCl₃)}; ν_{max} 3 270 (OH) and 2 800–2 700 cm⁻¹ (trans-quinolizidine bands); δ_H(500 MHz) 0.87–0.99 (3 H, m, CH₂Me), 2.56–2.78 (2 H, m, CH₂N), 2.98–3.25 (3 H, m, CH₂N and CHN), 3.70–3.84 (2 H, m, CH₂O), 7.04–7.49 (4 H, m, 4 × ArH), and 7.80–7.91 (1 H, br s, NH).

(b) To a solution of the dithioacetal (12) (see below) (12.3 mg, 0.0305 mmol) in acetone (5 ml) was added W-2 Raney nickel (1 g) under N₂. The mixture was heated under reflux for 18 h. The catalyst was removed by filtration, and the filtrate was evaporated under reduced pressure. The residue was chromatographed on silica gel with CHCl₃-MeOH (19:1, v/v) as eluant to give (-)-dihydrocorynantheol (7) (8.9 mg, 98%) as an oil, whose physical properties were identical with those of the compound prepared by method (a).

(-)-18,18-Trimethylenedithiopyhimban-17-one (**10**).—A solution of yohimbone (**9**)⁶ (662 mg, 2.25 mmol) and pyrrolidine (0.26 ml, 3.15 mmol) in dry benzene (30 ml) was heated under N₂ for 3 h in a flask equipped with a Dean–Stark trap for azeotropic removal of water. Removal of the solvent gave the enamine. A solution of the above enamine, propane-1,3-dithiol di(toluenep-sulphonate)⁷ (1.03 g, 2.48 mmol), and Et₃N (1 ml) in dry MeCN (30 ml) was heated under reflux for 14 h under N₂. After removal of the solvent, a mixture of 1.2M-HCl (5 ml) and CHCl₃ (5 ml) was added to the residue and then the mixture was stirred for 10 min at room temperature. The reaction mixture was basified with 10% aq. ammonia and extracted thoroughly with CHCl₃. The combined extracts were washed with water, dried (Na₂SO₄), and evaporated under reduced pressure. The residue was purified by column chromatography on silica gel with CHCl₃ as eluant to afford the dithioacetal (**10**) (868 mg, 97%) as an oil (Found: *M*⁺, 398.1479. C₂₂H₂₆N₂O₂S₂ requires *M*, 398.1486; [α]_D²⁵ -25.3° (c 0.64, CHCl₃); ν_{max} 3 470 (NH) and 1 705 cm⁻¹ (C=O); δ_H(90 MHz) 1.22–3.91 (21 H, m, 9 × CH₂ and 3 × CH), 6.91–7.56 (4 H, m, 4 × ArH), and 7.66–7.90 (1 H, br s, NH); *m/z* 398 (*M*⁺).

(2R,3S,12bS)-(+)-Methyl {3-[(1,3-Dithian-2-yl)methyl]-1,2,3,4,6,7,12,12b-octahydroindolo[2,3-a]quinolizin-2-yl}acetate (**11**).—To a solution of the dithioacetal (**10**) (554.5 mg, 1.39 mmol) in dry 2-methylpropan-2-ol (20 ml) was added KOH (782 mg, 13.9 mmol) and the mixture was heated under reflux for 2.5 h. After evaporation of the solvent, the residue was dissolved in MeOH (30 ml) and acidified with conc. H₂SO₄ (0.6 ml). The resulting mixture was stirred for 7 h at room temperature and for 3.5 h at 50 °C. After concentration, the residue was basified with 10% aq. ammonia and extracted thoroughly with CHCl₃. The extracts were washed with water, dried (Na₂SO₄), and evaporated. The residue was subjected to silica gel column chromatography with CHCl₃–AcOEt (19:1, v/v) as eluant to afford the ester (**11**) (312.6 mg, 52%) as a yellowish oil (Found: *M*⁺, 430.1728. C₂₃H₃₀N₂O₂S₂ requires *M*, 430.1749; [α]_D²⁶ +25.3° (c 0.64, CHCl₃); ν_{max} 3 470 (NH) and 1 730 cm⁻¹ (C=O); δ_H(90 MHz) 1.05–3.34 (21 H, m, 9 × CH₂ and 3 × CH), 3.73 (3 H, s, OMe), 4.03 (1 H, dd, *J* 4.5 and 9.8 Hz, CHS₂), 6.93–7.54 (4 H, m, 4 × ArH), and 7.70–7.93 (1 H, br s, NH); *m/z* 430 (*M*⁺).

(2R,3S,12bS)-(+)-3-[(1,3-Dithian-2-yl)methyl]-2-(2-hydroxyethyl)-1,2,3,4,6,7,12,12b-octahydroindolo[2,3-a]quinolizine (**12**).—To a suspension of lithium aluminium hydride (1.8 mg, 0.047 mmol) in dry tetrahydrofuran (THF) at 0 °C (2 ml) was added under N₂ a solution of the methyl ester (**11**) (27 mg, 0.062 mmol) in dry THF (3 ml). After being stirred for 30 min, the reaction mixture was treated by successive addition of 10% aq. ammonium chloride (0.1 ml) and CHCl₃. After being stirred for 1 h, the mixture was filtered through Celite, dried (Na₂SO₄), and evaporated. Purification of the residue by silica gel column chromatography with CHCl₃–MeOH (99:1, v/v) as eluant afforded the alcohol (**12**) (25 mg, 100%) as a yellowish oil

(Found: *M*⁺, 402.1794. C₂₂H₃₀N₂O₂S₂ requires *M*, 402.1799; [α]_D²⁵ +4.65° (c 0.43, CHCl₃); ν_{max} 3 480 (NH) and 3 300 cm⁻¹ (OH); δ_H(90 MHz) 1.10–3.27 (22 H, m, 9 × CH₂, 3 × CH, and OH), 3.51–3.88 (2 H, m, CH₂OH), 3.91–4.19 (1 H, m, CHS₂), 6.95–7.75 (4 H, m, 4 × ArH), and 8.22–8.42 (1 H, br s, NH); *m/z* 402 (*M*⁺).

Acknowledgements

This work was supported in part by a Grant in Aid for Scientific Research on Priority Areas, 01607002, Advanced Molecular Conversion from the Ministry of Education, Science and Culture of Japan. We thank Mr. K. Kawamura, Mrs. A. Satoh, Miss K. Mushiake, Miss M. Inada, and Miss N. Oikawa of this Institute for spectral measurements and preparation of the manuscript.

References

- 1 M. Ihara, M. Takahashi, N. Taniguchi, K. Fukumoto, and T. Kametani, *J. Chem. Soc., Chem. Commun.*, 1987, 619; M. Ihara, M. Takahashi, N. Taniguchi, K. Yasui, and K. Fukumoto, *J. Chem. Soc., Perkin Trans. 1*, 1989, 897.
- 2 M. Ihara, K. Yasui, N. Taniguchi, K. Fukumoto, and T. Kametani, *Tetrahedron Lett.*, 1988, **29**, 4963; M. Ihara, K. Yasui, N. Taniguchi, and K. Fukumoto, *J. Chem. Soc., Perkin Trans. 1*, 1990, 1469.
- 3 Isolation and structure determination: (a) B. Gilbert, L. D. Antonaccio, and C. Djerassi, *J. Org. Chem.*, 1962, **27**, 4702; (b) C. Vamvacas, W. V. Philipsborn, E. Schlittler, H. Schmid, and P. Karrer, *Helv. Chim. Acta*, 1957, **40**, 1793.
- 4 Synthesis of the racemate: (a) T. Kametani, N. Kanaya, H. Hino, S.-P. Huang, and M. Ihara, *J. Chem. Soc., Perkin Trans. 1*, 1981, 3168; (b) S. Takano, K. Masuda, and K. Ogasawara, *J. Chem. Soc., Chem. Commun.*, 1980, 887; (c) M. Ihara, N. Taniguchi, K. Fukumoto, and T. Kametani, *ibid.*, 1987, 1438; (d) B. Daniele, G. Lesma, G. Palmisano, and S. Tollari, *J. Chem. Soc., Perkin Trans. 1*, 1984, 1237; (e) S. Takano, K. Shibuya, M. Takahashi, S. Hatakeyama, and K. Ogasawara, *Heterocycles*, 1981, **16**, 1125; (f) T. Fujii, S. Yoshifuji, and H. Ito, *ibid.*, 1977, **7**, 149; (g) T. Fujii, M. Ohba, T. Tachinami, H. Miyajima, M. Koch, and E. Seguin, *ibid.*, 1986, **24**, 1215; (h) R. T. Brown, M. F. Jones, and M. Wingfield, *J. Chem. Soc., Chem. Commun.*, 1984, 847.
- 5 Synthesis of the (-)-enantiomer: T. Suzuki, E. Sato, K. Unno, and T. Kametani, *Chem. Pharm. Bull.*, 1986, **34**, 1584.
- 6 J. D. Albright and L. Goldman, *J. Org. Chem.*, 1965, **30**, 1107; *J. Am. Chem. Soc.*, 1967, **89**, 2416.
- 7 R. B. Woodward, I. J. Pachter, and M. L. Scheinbaum, *J. Org. Chem.*, 1971, **36**, 1137; S. Takano, K. Hiroya, and K. Ogasawara, *Chemistry Lett.*, 1983, 255.
- 8 J. A. Marshall and D. E. Seitz, *J. Org. Chem.*, 1974, **39**, 1814; S. Takano, M. Sasaki, H. Kanno, K. Shishido, and K. Ogasawara, *J. Org. Chem.*, 1978, **43**, 4169.

Paper 0/01221A
Received 20th March 1990
Accepted 23rd May 1990