

Oxocarbons and related compounds. Part 24.¹ Chlorosquarylation of indoles

Arthur H. Schmidt,* Stefan H. Thiel and Otfried Gaschler


Abteilung für Organische Chemie und Biochemie, Fachhochschule Fresenius, Kapellenstrasse 11-15, D-65193 Wiesbaden, Germany †

Chlorosquarylated indoles **3** and **4** have been synthesized from indoles **2** and squaric dichloride **1** and have been treated with nitrogen- and oxygen-nucleophiles.

A large number of indoles are distinguished by their biological activity. In particular, specifically substituted indoles serve as the starting material for the synthesis of natural products and also for pharmacologically active compounds.² As a consequence, there is a steady demand for novel synthetically versatile indole derivatives. Of fundamental importance in the search for such building blocks is the attachment of a polyfunctional subunit to the indole nucleus, with the capacity to undergo a great many, easy chemical transformations.³ We intended to achieve this by introducing the 2-chloro-3,4-dioxocyclobut-1-enyl group (chlorosquaryl group) to the indole nucleus. For this purpose we began an investigation into the reaction of 3,4-dichlorocyclobut-3-ene-1,2-dione (squaric dichloride) **1** with indoles.

Though squaric dichloride **1** has been used extensively for the acylation of arenes⁴ there is, surprisingly, only one mention in the literature of the extension of this reaction to indoles: a solution of squaric dichloride **1** and 1,2-dimethylindole **2a** (in excess) in dichloromethane was passed through a column of acidic alumina, thus affording 3,4-bis(1,2-dimethyl-3-indolyl)-cyclobut-3-ene-1,2-dione (0.67%).⁵

We have found that the reaction of the indoles **2a–f** with squaric dichloride **1** (1 equiv.) in an appropriate solvent affords the chlorosquarylated indoles **3** and **4** (Scheme 1, Table 1).




Scheme 1 Chlorosquarylation of indoles. For conditions see Table 1.

In agreement with reports on the acylation of indoles,⁶ chlorosquarylation takes place at C-3 of indoles **2a–c** (Table 1,

entries 1–4) with the formation of **3a–c**, and at C-2 of indoles **2d–f** (Table 1, entries 5–7), with the formation of **4a–c**. Depending on their specific substitution patterns, the chlorosquarylated indoles **3** and **4** exhibited different polarity and solubility properties. While **3a–c** and **4c** could be recrystallized easily, this was not the case with **4a** and **4b**. In an attempt to recrystallize **4b** from ethanol, the ethoxysquarylated indole **6** was obtained (*vide infra*).


On heating solutions of the chlorosquarylated indole **3b** in EtOH or THF with a primary amine (aniline) and a secondary amine (morpholine) for 30 min the aminosquarylated indoles **5a** and **5b** were obtained (Scheme 2). When **4b** was heated at



Scheme 2 Reagents and conditions: i, THF or EtOH, reflux, 30 min; ii, reflux, 3 min

reflux in EtOH for 3 min the ethoxysquarylated indole **6**, 3-ethoxy-4-(1-methyl-3-phenyl-1*H*-indol-2-yl)cyclobut-3-ene-1,2-dione, crystallized from the filtrate in analytically pure form (Scheme 2).

The chlorosquaryl unit's unique range of subsequent valuable synthetic transformations has been further demonstrated by the reaction of the chlorosquarylated indoles **3a–c** with sodium azide in methanol. Accompanied by the evolution of CO and N₂, the methyl indol-3-ylcyanoacetates **7a**⁷ (42%), **7b**⁷ (71%) and **7c** (66%) were obtained ‡ (Scheme 3). It is



Scheme 3 Reagents and conditions: i, acetone–MeOH, reflux 15–30 min

† New address: Abteilung für Organische Chemie und Biochemie, Fachhochschule Fresenius, Limburger Strasse 2, D-65510 Idstein, Germany.

‡ This is an extension of the reaction of 3-halogeno-4-phenylcyclobut-3-ene-1,2-dione with sodium azide. Phenylcyanoketene is generated which, in alcoholic solution, reacts further to give alkyl phenylcyanoacetates.⁸

Table 1 Chlorosquarylation of indoles

Entry	Indole	Solvent and conditions	Time (t/min)	Product ^a	Yield ^b (%)	Mp ^c (T/°C)
1	2a	Benzene, rt	45	3a	51	163–164 (decomp.)
2	2b	Benzene, rt	45	3b	26	169–170
3	2b	CHCl ₃ , rt	45	3b	21	168–169
4	2c	Benzene, reflux	120	3c	39	201–203 (decomp.)
5	2d	CH ₂ Cl ₂ , rt	720	4a	17	193–195
6	2e	Benzene, reflux	180	4b^d	75 ^e	195–197 (decomp.)
7	2f	Benzene, rt	45	4c	25	141–142

^a Compounds **3a–c**, **4a** and **4c** gave satisfactory elemental analyses and were characterized by IR, ¹H NMR, ¹³C NMR and mass spectroscopy.

^b Yields are of purified products, if not otherwise stated. ^c Uncorrected. ^d **4b** was purified by trituration with boiling light petroleum and did not give satisfactory elemental analyses. IR and mass spectra were taken and agreed with the structure. The structure of **4b** was confirmed by the preparation of the derivative **6** which gave satisfactory elemental analysis and was characterized by a complete set of spectra. ^e Yield of crude product.

noteworthy that methyl (5-methoxyindol-3-yl)cianoacetate has been successfully transformed into tryptamine analogues, which have proved to be potential antihypertensive drugs.⁹

Extension of this work to the chlorosquarylation of benzofurans and benzothiophenes is under investigation.

Experimental

3-Chloro-4-(1,2-dimethyl-1H-indol-3-yl)cyclobut-3-ene-1,2-dione **3a** (Table 1, Entry 1)

To a stirred solution of squaric dichloride **1** (1.74 g, 12.0 mmol) in benzene (30 cm³) was added a solution of 1,2-dimethylindole **2a** (2.00 g, 13.3 mmol) in benzene (30 cm³) over 10 min. When the addition was completed, the mixture was stirred for a further 45 min at room temperature. The reaction mixture was then filtered and the filtrate evaporated to dryness, leaving a dark green solid. This solid was recrystallized twice from CHCl₃–light petroleum (the first time with the aid of charcoal) to give **3a** as greenish crystals (1.83 g, 51%), mp 163–164 °C (decomp.) (Found: C, 64.4; H, 4.0; Cl, 13.6; N, 5.4. C₁₄H₁₀ClNO₂ requires C, 64.75; H, 3.88; Cl, 13.65; N, 5.39%); ν_{\max} (KBr)/cm⁻¹ 3020, 2920, 1780, 1750, 1600, 1535, 1490, 1440, 1400, 1100, 840 and 745; δ_{H} (400 MHz; CDCl₃) 2.78 (3 H, s), 3.74 (3 H, s), 7.24–7.32 (3 H, m) and 8.02 (1 H, d, *J* 7.8 Hz); δ_{C} (100.6 MHz; CDCl₃) 13.7, 30.4, 104.0, 110.0, 122.2, 122.9, 123.6, 125.5, 138.4, 146.3, 171.7, 187.5, 190.5 and 194.7; *m/z* (EI) 261 (M⁺, 11%), 259 (M⁺, 34), 205 (M⁺ – 2CO, 34) and (M⁺ – 2CO, 100).

Acknowledgements

The authors gratefully acknowledge financial support of this work from the Deutsche Forschungsgemeinschaft (DFG), Bonn-Bad Godesberg (Grant: Schm 309-6/1). They are also grateful to Dr G. Penzlin, Beilstein Institut, Frankfurt am Main, for his advice on nomenclature.

References

- Part 23, A. H. Schmidt, G. Kircher, St. Maus and H. Bach, *J. Org. Chem.*, in the press.
- (a) R. T. Brown, J. A. Joule and P. G. Sammes, *Comprehensive Organic Chemistry*, eds. D. H. R. Barton and W. D. Ollis, Pergamon Press, Oxford, 1979, vol. 4; (b) L. R. Smith, *Indoles*, Part II, Wiley-Interscience, New York, 1983; (c) J. E. Saxton, *Indoles*, Part IV, Wiley-Interscience, New York, 1983; (d) J. P. Kutney, *The Total Synthesis of Natural Products*, ed. J. Apsimon, Wiley-Interscience, New York, 1977, vol. 3.
- S. Blechert, R. Knier, H. Schroers and T. Wirth, *Synthesis*, 1995, 592.
- For selected references, see: (a) G. Maahs and P. Hegenberg, *Angew. Chem.*, 1966, **78**, 927; *Angew. Chem., Int. Ed. Engl.*, 1966, **5**, 888; (b) B. R. Green and W. Neuse, *Synthesis*, 1974, 46; (c) N. G. Argyropoulos, *Chem. Chron.*, 1986, **15**, 119; (d) P. M. Kazmaier, R. A. Burt and G. Baranyi, USP 4 624 904/1986, Xerox Corp., USA (*Chem. Abstr.*, 1987, **106**, 205165m); (e) Y. Akasaki, A. Tokida, K. Torigoe and A. Imai, Jap. P 62 249 952/1987, Fuji Xerox Co., Ltd., Japan (*Chem. Abstr.*, 1988, **109**, 170005s).
- M. Matsuoka, H. Soejima and T. Kitao, *Dyes Pigm.*, 1991, **16**, 309.
- (a) R. A. Jones, *Comprehensive Heterocyclic Chemistry*, eds. A. R. Katritzky and C. W. Rees, Pergamon Press, Oxford, 1984, vol. 4, ch. 3.05, p. 201; (b) R. J. Sundberg, *The Chemistry of Indoles*, Academic Press, New York, London, 1970.
- V. S. Rozhkov, Yu. I. Smushkevich and N. N. Suvorov, *Khim. Geterotsikl. Soedin.*, 1981, 372; *Chem. Heterocycl. Compd. (Engl. Trans.)*, 1981, **17**, 274; *Chem. Abstr.*, 1981, **95**, 80637n.
- (a) R. C. DeSelms, *Tetrahedron Lett.*, 1969, 1179; (b) A. H. Schmidt and W. Ried, *Tetrahedron Lett.*, 1969, 2431.
- L. G. Zepeda, M. Rojas-Gardida, M. S. Morales-Rios and P. Joseph-Nathan, *Tetrahedron*, 1989, **45**, 6439.

Paper 5/07570J

Received 20th November 1995

Accepted 11th January 1996