

Studies on tellurium-containing heterocycles. Part 10.¹ 2-Benzotelluropyrylium salts: first preparation and reactions with Grignard reagents


Haruki Sashida* and Kazuo Ohyanagi

Faculty of Pharmaceutical Sciences, Hokuriku University, Kanagawa-machi, Kanazawa
920-1181, Japan

The first simple preparation of 2-benzotelluropyrylium salts **4** from *o*-ethynylbenzyl bromides **1** in two steps, and the characteristic reaction of these salts **4** with benzylmagnesium bromide and other Grignard reagents are described.

The chemistry (structure and physical properties, syntheses and reactions) of not only the pyrylium salts² but also the thiopyrylium salts³ has been widely investigated and is now well established. Recently, extensive synthetic studies on their selenium⁴ and tellurium⁵ analogs have been undertaken, however the title compounds, the 2-benzotelluropyrylium salts, have not been prepared until now. We have previously focused on the synthesis of various tellurium- or selenium-containing heterocycles⁶ by an intramolecular cyclization of the tellurol or selenol moieties to an ethynyl group. In our continuing studies, we herein describe the successful synthesis of the 2-benzotelluropyrylium salts and their reactions with Grignard reagents.

The synthesis of the 2-benzotelluropyrylium salts **4** is shown in Scheme 1. In order to obtain isotellurochromenes **3**, the




Scheme 1 Reagents and conditions: i, NaHTe (1.2 equiv.), DMF, 0 °C, 1 h; ii, EtOH, 90 °C, 1–3 h; iii, Ph₃C⁺ BF₄⁻ (1 equiv.), MeNO₂, room temp., 30 min

precursors for the preparation of the telluropyrylium salts **4**, the conversion from *o*-ethynylbenzyl bromides **1** to the telluroles **2** was examined. Treatment of the benzyl bromides **1** with sodium hydrogen telluride⁷ in DMF at 0 °C, followed by addition of EtOH, and then heating at 90 °C, resulted in a direct ring closure to afford isotellurochromenes **3** (**3a**: 84% yield, **3b**: 64% yield) *via* the benzyltellurol intermediates **2**.[†] The

[†] The formation of compounds **2** was characterized by the isolation of bis(*o*-ethynylbenzyl) ditellurides, which were obtained by potassium ferricyanide oxidation of **2** before heating in ethanol.

isotellurochromenes **3** were produced by the 6-*endo-dig* ring closure of **2** at the sp carbon atom of the acetylenic moiety. The 5-*exo-dig* reaction products were not obtained. The isochromenes **3** were treated with triphenylcarbenium tetrafluoroboranuide (Ph₃C⁺ BF₄⁻) in MeNO₂ at room temperature to give the desired 2-benzotelluropyrylium tetrafluoroboranuides **4** (**4a**: 94% yield, **4b**: 89% yield) as stable pale green prisms. The salts **4** are the first synthetic examples of 2-benzotelluropyrylium salts. As expected, **4** have high reactivities and are easily decomposed on contact with water or alcohol. Thus, in order to examine the chemical behavior of the novel telluropyrylium salts **4** towards nucleophiles, the reaction of **4** with some Grignard reagents was carried out. Contrary to our expectations, treatment of the salts **4** with Grignard reagents, such as methyl-, ethyl- or phenyl-magnesium bromide (iodide) in THF at 0 °C resulted in a self-coupling reaction at the C-1 position to give 1,1'-bis(isotellurochromenyl) **5** as the sole characterized product in moderate yield (Scheme 2). However, when benzylmagnesium bromide was used as the Grignard reagent in Et₂O at 0 °C, 1-benzylisotellurochromenes **6** (**6a**: 48% yield, **6b**: 21% yield), normal coupling products, were obtained. In this case, no dimers **5** were obtained. It is already known that the reaction of telluraxanthylum salts⁸ with reducing agents, such as zinc powder or with nucleophiles, such as ammonium chloride, affords the telluraxanthyl dimers, produced *via* a free radical mechanism. This result clearly supports the present formation of **5** proceeding *via* a similar radical mechanism. In contrast, a similar reaction of their selenium analogs, the 2-benzoselenopyrylium salts, with any Grignard reagent gave the normal corresponding 1-substituted isoselenochromenes in good yields.

In addition, treatment of 1-benzyl-3-*tert*-butylisotellurochromene **6a** with Ph₃C⁺ BF₄⁻ gave 1-benzyltelluropyrylium tetrafluoroboranuide **7a** as the sole product. The ¹H NMR spectrum of the crude isolated telluropyrylium salt **7a** in CD₃CN showed two *tert*-butyl signals at δ 1.59 and 1.29 in a ratio of 2:3, indicating the presence of the salt **7a** and the benzyldiene compound **8a**. Treatment of the salt **7a** with base or by alumina column chromatography using methylene chloride as the eluent quantitatively gave **8a**. This finding suggests that the telluropyrylium salt **7a** exists in the solvent as an equilibrium mixture of **7a** and **8a**. The stereochemistry of the olefin moiety in **8a** was determined by a nuclear Overhauser enhancement (NOE) measurement. A NOE was observed between the benzyldiene 1'-H and the aromatic 8-H in the 400 MHz ¹H NMR spectrum of **8a**. Thus, the olefin moiety was determined to have (*Z*)-stereochemistry. In the case of a similar treatment of the 3-unsubstituted isotellurochromenes **6b** with Ph₃C⁺ BF₄⁻, the formation of salt **7b** and the benzyldiene compound **8b** was confirmed by the ¹H NMR spectrum in CD₃NO₂. However, both **7b** and **8b** were too unstable to isolate. The proposed mechanism for the formation of **8** is shown in Scheme 2.



Scheme 2 Reagents and conditions: i, EtMgBr (MeMgI, PhMgBr), THF, 0 °C, 30 min; ii, PhCH₂MgBr, Et₂O, 0 °C, 30 min; iii, Ph₃C⁺ BF₄⁻ (1 equiv.), MeNO₂, room temp., 2 h

Experimental

Melting points were measured on a Yanagimoto micro melting point hot stage apparatus and are uncorrected. IR spectra were determined with a Hitachi 270-30 spectrometer. Mass spectra (MS) and HRMS were recorded on a JEOL JMS-DX300 instrument. NMR spectra were determined with a JEOL EX-90A (90 MHz) or JEOL JNM-GSX 400 (400 MHz) spectrometer in CDCl₃ or CD₃CN using tetramethylsilane as internal standard and *J* values are given in Hz.

Isotellurochromene 3a

A solution of *o*-ethynylbenzyl bromide **1** (2.51 g, 10 mmol) in DMF (10 ml) was slowly added to a solution of sodium hydrogen telluride (12 mmol), which was freshly prepared from tellurium dust (1.53 g) and sodium borohydride (0.54 g) in DMF (40 ml) at 0 °C under an argon atmosphere. The reaction mixture was stirred under the conditions for 1 h. Ethanol (40 ml) was added to the reaction mixture, and then the whole mixture was heated at 90 °C with stirring for 1–3 h. After addition of water, the aqueous mixture was extracted with benzene. The organic extracts were washed with water and brine, dried (MgSO₄) and evaporated. The resulting residue was chromatographed on silica gel using *n*-hexane as eluent to give pure **3a**, yellow prisms, mp 63 °C (from *n*-hexane); δ_H(CDCl₃, 90 MHz) 1.29 (9H, s, Bu^t), 3.79 (2H, s, 1-H₂), 6.64 (1H, s, 4-H), 7.10–7.26 (4H, m, Ar-H) (HRMS *m/z* Calc. for C₁₃H₁₆Te: 302.0315. Found: 302.0313).

2-Benzotelluropyrylium salt 4a

Ph₃C⁺ BF₄⁻ (1.88 g, 5.5 mmol) was added to a stirred solution of the isochromene **3a** (1.51 g, 5 mmol) in dry MeNO₂ (10 ml) and the mixture was stirred at room temperature for 30 min. To the reaction mixture was added dry Et₂O to precipitate the telluropyrylium salt **4a**, pale green prisms, mp 101 °C (decomp.);

*v*_{max}(KBr)/cm⁻¹ 1054 (BF₄⁻); δ_H(CD₃CN, 400 MHz) 1.68 (9H, s, Bu^t), Ph-H [8.00 (1H, ddd, *J* 8.8, 7.2, 1.5), 8.34 (1H, br d, *J* 8.4), 8.47 (1H, ddd, *J* 8.4, 7.2, 1.5), 8.49 (1H, br d, *J* 8.8)], 9.14 (1H, s, 4-H), 13.22 (1H, s, 1-H); δ_C(CD₃CN, 100 MHz) 33.34 (q), 43.80 (s), 131.42 (d), 132.07 (d), 135.40 (d), 137.84 (d), 139.05 (d), 139.85 (s), 143.69 (s), 182.85 (s), 188.83 (s) (Anal. Calc. for C₁₃H₁₅BF₄Te: C, 40.49; H, 3.93. Found: C, 40.35; H, 3.75%).

1-Benzylisotellurochromene 6a

A solution of PhCH₂MgBr (4 mmol) in Et₂O (4 ml) was added to a suspended mixture of telluropyrylium salt **4a** (3 mmol) in Et₂O (10 ml) and the mixture was stirred at room temperature for 30 min. The reaction mixture was quenched by the addition of aqueous NH₄Cl, and extracted with Et₂O. The ethereal fractions were washed with brine, dried (MgSO₄) and evaporated. The residue was chromatographed on silica gel using hexane–CH₂Cl₂ (20:1) as eluent to give **6a**, yellow oil; δ_H(CDCl₃, 90 MHz) 1.31 (9H, s, Bu^t), 3.21 (2H, d, *J* 7.1, CH₂Ph), 3.96 (1H, br t, *J* 7.1, 1-H), 6.78 (1H, s, 4-H), 6.86–7.26 (9H, m, Ar-H); HRMS *m/z* Calc. for C₂₀H₂₂Te: 392.0785. Found: C, 392.0781.

Selected data for the telluropyrylium salt 7a. Yield 71%, pale green prisms, mp 105–108 °C (decomp.); *v*_{max}(KBr)/cm⁻¹ 1052 (BF₄⁻); δ_H(CD₃CN, 90 MHz) 1.55 (9H, s, Bu^t), 4.82 (2H, s, CH₂Ph), 7.22–7.59 (9H, m, Ph-H), 8.92 (1H, s, 4-H) (Anal. Calc. for C₂₀H₂₁BF₄Te: C, 50.49; H, 4.45. Found: C, 50.64; H, 4.47%).

Selected data for the benzylidene compound 8a. Yellow oil; δ_H(CDCl₃, 400 MHz) 1.24 (9H, s, Bu^t), 6.86 (1H, s, 4-H), 7.18–7.45 (8H, m, Ph-H), 7.48 (1H, s, 1'-H), 7.53 (1H, br d, *J* 7.8, 8-H); HRMS *m/z* Calc. for C₂₀H₂₀Te: 390.0628. Found: 390.0627.

References

- For part 9, see H. Sashida, *Heterocycles*, 1998, **48**, 631.
- (a) A. T. Balaban, A. Dinculescu, G. N. Dorofeenko, G. W. Fischer, A. V. Koblik, V. V. Mezhrtskii and W. Schroth, *Advances in Heterocyclic Chemistry: Pyrulium Salts: Synthesis, Reactions, and Physical Properties*, Supplement 2, ed. A. R. Katritzky, Academic Press, Inc., London, 1982.
- G. Doddi and G. Ercolani, *Advances in Heterocyclic Chemistry: Thiopyrylium, Selenopyrylium, and Telluropyrylium Salts*, vol. 60, ed. A. R. Katritzky, Academic Press, Inc., London, 1994, pp. 65–195.
- (a) M. Renson and P. Pirson, *Bull. Soc. Chim. Belg.*, 1966, **75**, 456; (b) A. Tadino, L. Christiaens and M. Renson, *Bull. Soc. Roy. Sci. Liege*, 1973, **42**, 129; (c) A. Tadino, L. Christiaens and M. Renson, *Bull. Soc. Roy. Sci. Liege*, 1973, **42**, 146.
- (a) I. D. Sadekov and V. Minkin, *Advances in Heterocyclic Chemistry: Six-Membered Heterocycles with a Tellurium Atom*, vol. 63, ed. A. R. Katritzky, Academic Press, Inc., London, 1995, pp. 1–60; (b) I. D. Sadekov, A. A. Ladatko, E. I. Sadekova, G. N. Dorofeenko and V. I. Minkin, *Khim. Geterotsikl. Soedin.*, 1981, 343 (*Chem. Abstr.*, 1981, **95**, 80 692b); (c) M. R. Detty and B. J. Murray, *J. Org. Chem.*, 1982, **47**, 5235; (d) M. R. Detty and B. J. Murray, *J. Am. Chem. Soc.*, 1983, **105**, 883; (e) M. Detty and H. R. Luss, *Organometallics*, 1986, **5**, 2250; (f) M. Detty, J. M. McKelvey and H. R. Luss, *Organometallics*, 1988, **7**, 1131.
- (a) H. Sashida, H. Kurahashi and Tsuchiya, *J. Chem. Soc., Chem. Commun.*, 1991, 802; (b) H. Sashida, K. Ito and T. Tsuchiya, *J. Chem. Soc., Chem. Commun.*, 1993, 1493; (c) H. Sashida, K. Ito and T. Tsuchiya, *Chem. Pharm. Bull.*, 1995, **43**, 19; (d) H. Sashida, K. Sadamori and T. Tsuchiya, *Synth. Commun.*, 1998, **28**, 713; (e) H. Sashida, *Synthesis*, 1998, 745; (f) H. Sashida and A. Kawamukai, *J. Heterocycl. Chem.*, 1998, **35**, 165.
- J. Liu, M. Qui and X. Zhou, *Synth. Commun.*, 1990, **20**, 2759.
- A. A. Bumber, E. P. Ivakhnenko, A. A. Ladatko, A. A. Arutyunyan, I. D. Sadekov and O. Yu. Okhlobistin, *Khim. Geterotsikl. Soedin.*, 1988, 1196 (*Chem. Abstr.*, 1989, **111**, 38 699h).

Paper 8/04036B

Received 29th May 1998

Accepted 29th May 1998