

Conversion of *N*-alkyldiisopropylamines into *N,N*-bis(5-chloro-3-oxo[1,2]dithiol-4-yl)amines

PERKIN

Susana Barriga,^a Lidia S. Konstantinova,^b Carlos F. Marcos,^a Oleg A. Rakitin,^b Charles W. Rees,^c Tomás Torroba,^d Andrew J. P. White^c and David J. Williams^c

^a Departamento de Química Orgánica, Facultad de Veterinaria, Universidad de Extremadura, 10071 Cáceres, Spain

^b N. D. Zelinsky Institute of Organic Chemistry, Academy of Sciences, Leninsky Prospect, 47, 117913 Moscow, Russia

^c Department of Chemistry, Imperial College of Science, Technology and Medicine, London, UK SW7 2AY


^d Departamento de Química, Facultad de Ciencias, Universidad de Burgos, 09001 Burgos, Spain

Received (in Cambridge) 11th June 1999, Accepted 12th July 1999

Triisopropylamine **1a** and S₂Cl₂, in the presence of DABCO and formic acid, give the bicyclic amine **2a**, and its *N*-dealkylated product **3**, rather than the tricyclic bisdithiolothiazine system **7** produced by Hünig's base **1b**. However, when S₂Cl₂ is in molar excess over DABCO, Hünig's base and related *N*-alkyldiisopropylamines **1c–g** also give the analogous bicyclic compounds **2b–g** together with the tricyclic compounds **7b–g**. The *N*-benzyl compound **2c** is debenzylated quantitatively to the secondary amine **3** by sulfuric acid. X-Ray crystallography of *N,N*-bis(dithioly)ethylamine **2b** shows the two dithiole rings to be steeply inclined to each other (85°) and the geometry at nitrogen to be slightly pyramidal. The crystal packing is dominated by S⋯O electrostatic and weak C=O⋯Cl charge-transfer interactions.

N-Ethyldiisopropylamine (Hünig's base) is a hindered non-nucleophilic base, commonly used as a deprotonating reagent in many organic reactions. In contrast to its generally assumed inertness, we have shown that it reacts with disulfur dichloride, S₂Cl₂, under very mild conditions to give bis[1,2]dithiolo[1,4]thiazines,¹ bis[1,2]dithiopyrroles,² and a dihydro[1,2]dithiolo[1,4]thiazine,³ in one-pot reactions. Under carefully controlled reaction conditions each of these heterocyclic systems can be obtained selectively.⁴ Only the isopropyl groups react with S₂Cl₂, the ethyl group being inert. *N*-(2-Chloroethyl)diisopropylamine reacts similarly,³ and even *N*-benzyldiisopropylamine reacts in the same way to give the *N*-benzyl bisdithiolothiazines and pyrroles and hence, by debenylation, the parent *N*-H compounds.⁵ To explore the limits of the reaction we have now included a third isopropyl group in the starting amine, subjecting the highly crowded triisopropylamine to the action of S₂Cl₂ and have obtained the first member of a new class of stable bis(1,2-dithiol-4-yl)amines. Furthermore we report how these bicyclic bis(dithioly)amines can be produced from other *N*-alkyldiisopropylamines in one-pot reactions with S₂Cl₂ under appropriate conditions.


Triisopropylamine⁶ **1a** was treated with S₂Cl₂ (15 equiv.) and 1,4-diazabicyclo[2.2.2]octane (DABCO) (12 equiv.) in 1,2-dichloroethane for 3 d at room temperature. Formic acid (30 equiv.) was then added and the mixture heated under reflux for 1.5 h since this treatment gives cleaner reactions by facilitating the hydrolysis of intermediate 3-chlorodithiolium salts to dithiol-3-ones.⁴ Products **2a**, orange solid, C₉H₇Cl₂NO₂S₄, mp 138–140 °C (12%) and **3**, orange solid, C₆HCl₂NO₂S₄, mp 144–145 °C (6%) were obtained after chromatography (Scheme 1). Structures **2a** and **3** were elucidated from mass spectrometry, HRMS, ¹H- and ¹³C-NMR and IR spectroscopy. When triisopropylamine **1a** was treated with an excess of S₂Cl₂ and DABCO in equimolecular proportions (16 equiv.), followed by treatment with formic acid, only **3** was isolated, in very small amounts; without the formic acid treatment no pure products



Scheme 1

could be isolated. It appears that for this amine an excess of S₂Cl₂ over DABCO is needed to furnish stable products. Under these conditions triisopropylamine starts to react like Hünig's base,⁴ following the sequence of dehydrogenation, displacement of chlorine from S₂Cl₂ and chlorination, up to formation of the intermediate **4** (cf. ref. 4) (Scheme 2). The next step in our proposed mechanism would be cyclization of **4** to **5**,⁴ but if we assume that these tricyclic dithiolium salts are highly delocalised and hence near-planar, the *N*-isopropyl intermediate **5** will be markedly more crowded than the *N*-ethyl analogue. Computational modelling⁷ shows that the planar structure **5** is sterically forbidden. Instead of ring closure to **5**, **4** could be chlorinated further by the excess of S₂Cl₂ to give **6** which would be converted by formic acid into the bicyclic *N*-isopropyl compound **2a** which, in turn, could be slowly dealkylated by the hot acid to give the secondary amine **3** (Scheme 2). Interestingly, there was no sign that the remaining isopropyl group in **2a** had reacted with S₂Cl₂ to give a third 1,2-dithiole ring. This could be a consequence of the electron-withdrawing effect of the two dichlorodithiolium rings (or the derived chlorodithiolone rings) which would suppress the oxidation of the third *N*-isopropyl group to the iminium ion (cf. Scheme 2). For the same reason one *N*-acetyl or *N*-cyano group protects diisopropylamine from reaction with S₂Cl₂ under the same conditions.⁴

We have proposed in Scheme 2 that bis-dithiolium salt **4** is the key intermediate which is diverted from further cyclisation by ring chlorination when DABCO is in deficiency with respect



Scheme 2

to disulfur dichloride. If this assumption is correct a similar diversion of the reaction of other *N*-alkyldiisopropylamines might also be observed under the same conditions to give hitherto undetected bicyclic products analogous to **2a**, and we now find this to be the case. Thus, *N*-ethyldiisopropylamine **1b** (1 equiv.) was treated with S_2Cl_2 (10 equiv.) and DABCO (6 equiv.) in 1,2-dichloroethane for 3 d at room temperature. Formic acid (20 equiv.) was then added and the mixture heated under reflux for 1.5 h. Chromatography afforded the bicyclic product **2b**, yellow solid, $C_8H_5Cl_2NO_2S_4$, mp 132–133 °C (19%) and the previously reported tricyclic product **7b**⁴ (10%) (Scheme 3). The structure of **2b** was determined from mass spectrometry, HRMS, ¹H- and ¹³C-NMR and IR spectroscopy and was confirmed by X-ray diffraction (see below). Analogous treatment of *N*-benzyldiisopropylamine **1c** with S_2Cl_2 (10 equiv.) and DABCO (6 equiv.) in 1,2-dichloroethane for 3 d at room temperature followed by addition of formic acid (20 equiv.) and heating under reflux for 1 h gave **2c**, yellow crystals, mp 164–165 °C (15%) in addition to the previously reported compound **7c**⁵ (27%) (Scheme 3). The structure of **2c** was fully supported by its spectroscopic properties. Treatment of a dilute solution of the *N*-benzyl compound **2c** in dichloromethane with concentrated sulfuric acid at 5–10 °C for 30 min gave a quantitative yield of the debenzylated compound **3**, identical with that from triisopropylamine **1a**.

Similarly, treatment of *N*-(2-chloroethyl)diisopropylamine **1d** with S_2Cl_2 , DABCO and formic acid (20 equiv.) as shown in Table 1 gave the bicyclic compound **2d**, yellow crystals, $C_8H_4Cl_3NO_2S_4$, mp 97–98 °C (7%) in addition to tricyclic compound **7d**³ (20%). In the same way the *N*-phenylthioethyl **1e**, phthalimidoethyl **1f**, and tetrachlorophthalimidoethyl **1g** derivatives of diisopropylamine, which were prepared from **1d**,⁸ were converted into the analogous bicyclic (*ca.* 20% yield) and






Fig. 1 The molecular structure of **2b**. Selected bond lengths (Å) are: S(1)–C(8A) 1.735(4), S(1)–S(2) 2.050(2), S(2)–C(3) 1.792(4), C(3)–O(3) 1.208(5), C(3)–C(3A) 1.456(6), C(3A)–C(8A) 1.347(6), C(3A)–N(4) 1.408(5), N(4)–C(4A) 1.405(5), N(4)–C(9) 1.472(5), C(4A)–C(7A) 1.347(6), C(4A)–C(5) 1.468(5), C(5)–O(5) 1.204(5), C(5)–S(6) 1.787(4), S(6)–S(7) 2.046(2), S(7)–C(7A) 1.727(4), C(7A)–Cl(7) 1.706(4), C(8A)–Cl(8) 1.701(4).



Scheme 3

tricyclic (*ca.* 15% yield) compounds (Scheme 3 and Table 1). These results are all explicable by the mechanism proposed in Scheme 2 where the bisdithiolium intermediate (*cf.* **4**) is either chlorinated further or (except for *N*-isopropyl) cyclised to a 1,4-thiazine.

The complex reactions of *N*-alkyldiisopropylamines with S_2Cl_2 provide new routes to novel heterocyclic structures by diversion of intermediates on the way to complete sulfuration of both isopropyl groups. We have now shown that chlorination by S_2Cl_2 and a deficiency of DABCO, and hydrolysis by formic acid, of these readily available tertiary amines give *N,N*-bis(1,2-dithiol-4-yl)amine **3** and its *N*-alkyl derivatives **2b–g** in one-pot, though low-yielding, reactions. Whilst 5-amino derivatives of 1,2-dithiol-3-ones and -thiones are readily available by displacement of chlorine from the activated 5-position,⁹ 4-amino isomers like **2** and **3** appear to be very rare.¹⁰

The X-ray analysis of **2b** confirmed the bisdithiole nature of the compound (Fig. 1), the two dithiole rings being inclined almost orthogonally (85° between their mean planes), this geometric arrangement being produced by torsional twists of 59 and 61° about the C(3A)–N(4) and N(4)–C(4A) bonds respectively. The geometry at the nitrogen centre is slightly pyramidalized with N(4) lying 0.16 Å out of the C(3A)–C(4A)–C(9) plane—the dithiole rings are inclined by 69° [C(3A)] and 50° [C(4A)] to this plane. The pattern of bonding within each ring reflects the formal double bond character of the C(3A)–C(8A)

Table 1 Reaction of amines with S₂Cl₂, DABCO and formic acid

Amine	Reagents			Formic acid/ mmol	Reflux time/min	Yields of products (%)						
	Amine/ mmol	S ₂ Cl ₂ / mmol	DABCO/ mmol			2a	3	7b ⁴	7c ⁵	7d ³	7e ⁷	7f ⁷
1a	5	75	60	150	90	2a	12	3	6			
1b	5	50	30	100	60	2b	19	7b⁴	10			
1c	5	50	30	100	60	2c	15	7c⁵	20			
1d	5	50	25	100	45	2d	7	7d³	20			
1e^a	5	50	30	100	45	2e	15	7e⁷	17			
1f	5	50	25	100	60	2f	20	7f⁷	15			
1g^a	5	50	25	100	45	2g	22	7g⁷	16			

^a Tetramethylammonium chloride (30 mg) was added to the reaction mixture.




Fig. 2 Part of one of the loosely-linked chains of molecules present in the crystals of **2b**. The intrachain S...O contacts are: a) 3.12, b) 3.06, c) 3.08 and d) 3.08 Å.

and C(4A)–C(7A) linkages [both 1.347(6) Å], though these distances are noticeably greater than that for an isolated C=C and indicate a degree of delocalisation within the ring system. Structural data for 1,2-dithiol-3-one ring systems are sparse and, with the exception of the fused ring systems that we have described previously,^{4,5} the current release of the Cambridge Crystallographic Database (October 1998) contains only two other examples^{11,12} of pendant 1,2-dithiol-3-one containing systems. Of these, the closer analogue is 4-methyl-5-(pyrazin-2-yl)-3*H*-1,2-dithiol-3-one where the pattern of bonding in the dithiole ring does not differ significantly from that observed here.

The packing of the molecules (Fig. 2) is dominated by electrostatic O...S interactions (ranging between 3.06 and 3.12 Å) and parallel stacking of the dithiole rings of symmetry related molecules; the mean interplanar separations are 3.73 [C(3A) ring to C(3A) ring] and 3.85 Å [C(4A) ring to C(4A) ring]. These interactions combine to form rather elegant twisted chains that extend in the crystallographic *c* direction. Adjacent C₁-related chains are oriented such that the carbonyl groups in one chain are directed towards the chlorine atoms in the next and *vice versa* (Fig. 3). The O...Cl distances, 3.03 and 3.04 Å, are significantly less than the sum of their van der Waals radii (*ca.* 3.2 Å) and clearly indicate some form of intermolecular interaction. If we ignore the possibility of significant intramolecular electron-transfer between the two dithiole rings, then the 4-amino-5-chloro-1,2-dithiole moiety can be represented by the resonance forms A–C in Scheme 4, with a minor contribution from D. The oxygen atom carries a partial negative charge,





Fig. 3 The C=O...Cl linking of adjacent centrosymmetrically related chains. The O...Cl distances are: e) 3.03 and f) 3.04 Å.



Scheme 4

explaining the 3-centre O...S interactions a to d in Fig. 2, which are clearly electrostatic. If a small positive charge were induced on the chlorine atom, as in contribution D, then the intermolecular O...Cl interactions e and f (Fig. 3) could also be electrostatic in nature, but the observed bond lengths do not indicate any marked shortening of either the C–Cl or the C(3A)–C(3) [C(4A)–C(5)] bond lengths. A survey of the literature reveals that short intermolecular C=O...Cl contacts are relatively commonplace (159 examples in the range 2.86–3.20 Å), though infrequently commented upon. However, two particular examples, in the structure of 2,3-dichlorobenzol-1,4-quinone,¹³ and perhaps more relevantly in that of 5-chloropyrimidin-2-one,¹⁴ this type of interaction has been attributed to “donor–acceptor” and “charge transfer” bonding

respectively. Examples of strong interactions of this type have been discussed in a review by Hassel and Rømming¹⁵ and are considered to involve a lone pair on the carbonyl and the d-orbitals of the halogen. We think it likely that the C=O...Cl contacts in the structure of **2b** are probably examples of weak interactions of this type.

Experimental

Melting points were determined on a Kofler hot-stage apparatus and are uncorrected. IR spectra were recorded on a Bruker Vector 200 A instrument in KBr pellets. ¹H NMR and ¹³C NMR spectra were recorded on a Bruker DRX 400 in CDCl₃ or *d*₅-pyridine solution. CH groups were distinguished by DEPT experiments. *J* values are given in Hz. Mass spectra were recorded on VG7070E and VG-AutoSpec instruments using electron impact ionisation. Light petroleum refers to the fraction bp 40–60 °C.

General procedure for the preparation of *N,N*-bis(5-chloro-3-oxo-1,2-dithiol-4-yl)alkylamines **2a–g**

Disulfur dichloride was added dropwise to a stirred solution of *N*-alkyldiisopropylamine **1a–g** and DABCO dissolved in 1,2-dichloroethane (150 ml) at –40 °C. The mixture was stirred for 15 min at –40 °C, then for 3 days at room temperature. Formic acid was then added dropwise at 5 °C and the mixture was refluxed for 45 min to 1.5 h. The reaction mixture was filtered through Celite and the solvent was removed in a rotary evaporator. Medium pressure liquid chromatography (MPLC) (silica gel Merck 60, light petroleum to CH₂Cl₂) of the residue afforded products **2a–g**, in addition to the previously reported products **7b–g**. The reaction conditions and yields of products are given in Table 1.

N,N-Bis(5-chloro-3-oxo-1,2-dithiol-4-yl)isopropylamine **2a**.

Orange solid (from light petroleum–CH₂Cl₂), mp 138–140 °C (Found M⁺, 358.8734. C₉H₇Cl₂NO₂S₄ requires *M*, 358.8737) (Found: C, 30.25; H, 2.0; N, 3.8. C₉H₇Cl₂NO₂S₄ requires C, 30.0; H, 2.0; N, 3.9%); δ_H (CDCl₃) 4.34 (heptet, 1H, *J* 7.0, CH), 1.21 (d, 6H, *J* 7.0, 2 × CH₃); δ_C (CDCl₃) 187.02 (C=O), 150.52 and 133.86 (2 × sp² tertiary C), 51.40 (CH), 21.45 (CH₃); ν_{max}/cm⁻¹ 2978, 2929, 1646 (C=O), 1524, 1382, 1260, 1123, 805, 714, 620; *m/z* 363 (M⁺ + 4, 16%), 361 (M⁺ + 2, 58), 359 (M⁺, 68), 344 (M – CH₃, 32), 323 (M – HCl, 74), 317 (M – C₃H₆, 29), 256 (M – C₃H₇SCO, 77), 225 (C₅HCl₂NOS₂, 37), 79 (CICS⁺, 94), 43 (C₃H₇⁺, 100).

N,N-Bis(5-chloro-3-oxo-1,2-dithiol-4-yl)amine **3**. Orange solid (from light petroleum–CH₂Cl₂), mp 144–145 °C (Found M⁺, 316.8201. C₆HCl₂NO₂S₄ requires *M*, 316.8267) (Found: C, 22.8; H, 0.35; N, 4.35. C₆HCl₂NO₂S₄ requires C, 22.65; H, 0.3; N, 4.4%); δ_H (CDCl₃) 5.54 (br s, 1H, NH); δ_C (CDCl₃) 185.52 (C=O), 139.65 and 129.36 (2 × sp² tertiary C); ν_{max}/cm⁻¹ 3302 (N–H), 1614 (C=O), 1575, 1550, 1453, 1283, 1094, 563; *m/z* 321 (M⁺ + 4, 24%), 319 (M⁺ + 2, 84), 317 (M⁺, 100), 256 (M – HSCO, 35) (Found M⁺, 255.8517. C₅Cl₂NO₂S₃ requires *M*, 255.8519), 225 (M – COS₂, 16) (Found M⁺, 224.8854. C₅HCl₂NO₂S₂ requires *M*, 224.8877), 218 (M – CHCl₂O, 17) (Found M⁺, 217.8866. C₅NOS₄ requires *M*, 217.8863), 210 (M – C₂ClOS, 20) (Found M⁺, 209.8909. C₄HClNOS₃ requires *M*, 209.8909), 79 (CICS⁺, 47) (Found M⁺, 78.9407. CCIS requires *M*, 209.8909).

N,N-Bis(5-chloro-3-oxo-1,2-dithiol-4-yl)ethylamine **2b**. Yellow solid (from light petroleum–CH₂Cl₂), mp 132–133 °C (Found M⁺, 344.8579. C₈H₅Cl₂NO₂S₄ requires *M*, 344.8502) (Found: C, 27.9; H, 1.5; N, 4.0. C₈H₅Cl₂NO₂S₄ requires C, 27.75; H, 1.5; N, 4.0%); δ_H (CDCl₃) 3.70 (q, 2H, *J* 7.1, CH₂), 1.18 (t, 3H, *J* 7.1, CH₃); δ_C (CDCl₃) 185.99 (C=O), 146.54 and

133.52 (2 × sp² tertiary C), 44.58 (CH₂, from DEPT), 14.77 (CH₃, from DEPT); ν_{max}/cm⁻¹ 2981, 2931, 2875, 1634 (C=O), 1535, 1252, 1122, 1047, 612; *m/z* 349 (M⁺ + 4, 6%), 347 (M⁺ + 2, 22), 345 (M⁺, 25), 309 (M – HCl, 9), 274 (M – HCl₂, 8), 218 (M – C₃H₅Cl₂O, 11), 210 (M – C₄H₄ClOS, 10), 79 (CICS⁺, 94). *Crystal data for 2b*: C₈H₅Cl₂NO₂S₄, *M* = 346.3, triclinic, *P* $\bar{1}$ (no. 2), *a* = 7.814(2), *b* = 8.211(1), *c* = 10.115(2) Å, *a* = 86.50(1), *β* = 88.77(2), *γ* = 85.46(1)°, *V* = 645.6(2) Å³, *Z* = 2, *D*_c = 1.781 g cm⁻³, μ(Cu-Kα) = 104.9 cm⁻¹, *F*(000) = 348, *T* = 293 K; yellow rhombs, 0.43 × 0.27 × 0.21 mm, Siemens P4/PC diffractometer, ω-scans, 2088 independent reflections. The structure was solved by direct methods and the non-hydrogen atoms refined anisotropically using full matrix least-squares based on *F*² to give *R*₁ = 0.047, *wR*₂ = 0.128 for 1784 independent observed absorption corrected reflections [*F*_o] > 4σ(*F*_o), 2θ ≤ 126°] and 155 parameters.†

N,N-Bis(5-chloro-3-oxo-1,2-dithiol-4-yl)benzylamine **2c**.

Orange solid (from light petroleum–CH₂Cl₂), mp 146–165 °C (Found M⁺, 406.8743. C₁₃H₇Cl₂NO₂S₄ requires *M*, 406.8737) (Found: C, 38.4; H, 1.8; N, 3.5. C₁₃H₇Cl₂NO₂S₄ requires C, 38.2; H, 1.7; N, 3.4%); δ_H (*d*₅-pyridine) 7.76 (d, *J* 7.3, 2H, ArH), 7.31 (t, *J* 7.7, 2H, ArH), 7.21 (t, *J* 7.4, 1H, ArH), 5.05 (s, 2H, CH₂); δ_C (*d*₅-pyridine) 187.26 (C=O), 148.07, 137.35 and 134.40 (3 × sp² tertiary C), 129.36, 128.74 and 128.28 (3 × ArH, from DEPT), 53.06 (CH₂, from DEPT); ν_{max}/cm⁻¹ 3053, 2934, 1658 (C=O), 1541, 1453, 1381, 1069, 955, 694, 560; *m/z* 411 (M⁺ + 4, 1%), 409 (M⁺ + 2, 6), 407 (M⁺, 6), 371 (M – HCl, 2), 91 (C₇H₇⁺, 100).

N,N-Bis(5-chloro-3-oxo-1,2-dithiol-4-yl)(2-chloroethyl)amine **2d**.

Yellow crystals (from light petroleum–CH₂Cl₂), mp 97–98 °C (Found M⁺, 378.8180. C₈H₄Cl₃NO₂S₄ requires *M*, 378.8190) (Found: C, 25.5; H, 1.15; N, 3.5. C₈H₄Cl₃NO₂S₄ requires C, 25.2; H, 1.1; N, 3.7%); δ_H (CDCl₃) 3.93 (t, *J* 6.8, 2H, CH₂), 3.75 (t, *J* 6.8, 3H, CH₂); δ_C (CDCl₃) 186.16 (C=O), 147.57 and 133.35 (2 × sp² tertiary C), 50.63 and 42.97 (2 × CH₂, from DEPT); ν_{max}/cm⁻¹ 2964, 1627 (C=O), 1547, 1533, 1078, 1007, 821; *m/z* 383 (M⁺ + 4, 7%), 381 (M⁺ + 2, 15), 379 (M⁺, 14), 347 (M⁺ + 4 – HCl, 16), 345 (M⁺ + 2 – HCl, 16), 343 (M⁺ – HCl, 62), 308 (M⁺ – HCl₂, 14), 160 (C₅H₃ClNOS⁺, 45), 79 (CICS⁺, 100).

N,N-Bis(5-chloro-3-oxo-1,2-dithiol-4-yl)(2-phenylthioethyl)amine **2e**.

Yellow solid (from light petroleum–CH₂Cl₂), mp 81–83 °C (Found M⁺, 452.8631. C₁₄H₉Cl₂NO₂S₅ requires *M*, 452.8614) (Found: C, 37.2; H, 2.1; N, 3.0. C₁₄H₉Cl₂NO₂S₅ requires C, 37.0; H, 2.0; N, 3.1%); δ_H (CDCl₃) 7.35 (dt, *J*_{ab} 8.0, *J*_{ac/ad} 1.5, 2H, Ph_{ortho}), 7.27 (td, *J*_{ba} 8.0, *J*_{be} 2.3, 2H, Ph_{meta}), 7.18 (tt, *J*_{cb} 7.3, *J*_{ca} 1.9, 1H Ph_{para}), 3.81 (t, *J* 7.9, 2H, CH₂), 3.17 (t, *J* 7.9, 2H, CH₂); δ_C (CDCl₃) 185.99 (C=O), 147.12, 135.20 and 133.31 (3 × sp² tertiary C), 129.47, 128.97 and 126.34 (3 × CH_{aromat}, from DEPT), 49.19 and 32.98 (2 × CH₂, from DEPT); ν_{max}/cm⁻¹ 3042, 3008, 2977, 2935, 1655 (C=O), 1582, 1542, 1478, 1441, 1069, 802, 740, 690; *m/z* 453 (M⁺, 1%), 346 (M⁺ + 2 – C₆H₅S, 26), 344 (M⁺ – C₆H₅S, 34), 79 (CICS⁺, 95), 77 (C₆H₅⁺, 100).

N,N-Bis(5-chloro-3-oxo-1,2-dithiol-4-yl)(2-phthalimidoethyl)amine **2f**.

Orange solid (from light petroleum–CH₂Cl₂), mp 162–163 °C (Found M⁺, 489.8747. C₁₆H₈Cl₂N₂O₄S₄ requires *M*, 489.8744) (Found: C, 39.3; H, 1.7; N, 5.6. C₁₆H₈Cl₂N₂O₄S₄ requires C, 39.1; H, 1.6; N, 5.7%); δ_H (CDCl₃) 7.83 (dd, *J*_{ab} 5.4, *J*_{ac} 3.1, 2H, PhthH_{ortho}), 7.70 (dd, *J*_{ba} 5.4, *J*_{ba} 3.1, 2H, PhthH_{meta}), 4.14 (t, *J* 5.9, 2H, CH₂), 3.91 (t, *J* 5.9, 2H, CH₂); δ_C (CDCl₃) 187.01 (C=O), 168.42 (N–C=O), 155.19, 146.56 and 145.31 (3 × sp² tertiary C), 134.36 and 123.72 (2 × CH_{aromat}),

† CCDC reference number 207/348. See <http://www.rsc.org/suppdata/p1/1999/2237> for crystallographic files in .cif format.

47.61 and 37.21 ($2 \times \text{CH}_2$); $\nu_{\text{max}}/\text{cm}^{-1}$ 1772 and 1720 (N–C=O), 1655 and 1638 (C=O), 1394, 1289, 1044; m/z 494 ($\text{M}^+ + 4$, 5%), 492 ($\text{M}^+ + 2$, 17), 490 (M^+ , 20), 347 ($\text{M}^+ + 4 - \text{phthH}$, 10), 345 ($\text{M}^+ + 2 - \text{phthH}$, 35), 343 ($\text{M}^+ - \text{phthH}$, 46), 334 ($\text{M}^+ + 4 - \text{phthCH}_2$, 7), 332 ($\text{M}^+ + 2 - \text{phthCH}_2$, 24), 330 ($\text{M}^+ - \text{phthCH}_2$, 29), 308 ($\text{M}^+ - \text{phthHCl}$, 11), 174 ($\text{phth-C}_2\text{H}_4^+$, 68), 160 (phthCH_2^+ , 76), 147 (phthH^+ , 37), 130 ($\text{phth}^+ - \text{O}$, 100).

***N,N*-Bis(5-chloro-3-oxo-1,2-dithiol-4-yl)[2-(tetrachlorophthalimido)ethyl]amine 2g.** Orange solid (from light petroleum– CH_2Cl_2), mp 244–245 °C (Found M^+ , 627.7106. $\text{C}_{16}\text{H}_4\text{Cl}_5\text{S}_4\text{N}_2\text{O}_4$ requires M , 627.7156) (Found: C, 27.9; H, 1.0; N, 3.4. $\text{C}_{16}\text{H}_4\text{Cl}_6\text{N}_2\text{O}_4\text{S}_4 + 2\text{CH}_2\text{Cl}_2$ requires C, 27.1, H, 1.0, N 3.5%); δ_{H} 4.0 (dd, J_{a} 7.0, J_{b} 5.0, 2H, CH_2), 3.82 (dd, J_{a} 7.0, J_{b} 5.0, 2H, CH_2); δ_{C} 186.35 (C=O), 163.39 (N–C=O), 147.91, 139.80, 132.97, 130.38 and 129.49 ($5 \times \text{sp}^2$ tertiary C), 46.82 and 36.82 ($2 \times \text{CH}_2$); $\nu_{\text{max}}/\text{cm}^{-1}$ 2933, 1779 and 1720 (N–C=O), 1666 and 1627 (C=O), 1548, 1534, 1397, 1355, 1070, 743; m/z 638 ($\text{M}^+ + 8$, 5%), 632 ($\text{M}^+ + 6$, 15), 630 ($\text{M}^+ + 4$, 14), 629 ($\text{M}^+ + 2 + \text{H}$, 21), 628 ($\text{M}^+ + 2$, 35), 626 (M^+ , 16), 347 ($\text{M}^+ + 4 - \text{Cl}_4\text{phthH}$, 25), 345 ($\text{M}^+ + 2 - \text{Cl}_4\text{phthH}$, 83), 343 ($\text{M}^+ - \text{Cl}_4\text{phthH}$, 100), 334 ($\text{M}^+ + 4 - \text{Cl}_4\text{phthCH}_2$, 17), 332 ($\text{M}^+ + 2 - \text{Cl}_4\text{phthCH}_2$, 58), 330 ($\text{M}^+ - \text{Cl}_4\text{phthCH}_2$, 71), 310 ($\text{M}^+ - \text{Cl}_4\text{phthSH}$, 27), 296 ($\text{Cl}_4\text{phthCH}_2^+$, 20), 266 ($\text{Cl}_4\text{phth}^+ - \text{O}$, 38), 79 (CICS^+ , 100).

***N,N*-Bis(5-chloro-3-oxo-1,2-dithiol-4-yl)amine 3 by debenzoylation.** *N,N*-Bis(5-chloro-3-oxo-1,2-dithiol-4-yl)benzylamine **2c** (110 mg, 0.27 mmol) was treated with concentrated sulfuric acid (0.6 ml) in dichloromethane (40 ml) at 5–10 °C for 30 min, then water (50 ml) was added and the organic layer separated, washed and dried (MgSO_4), affording *N,N*-bis(5-chloro-3-oxo-1,2-dithiol-4-yl)amine **3** (85 mg, 99%) identical with that described above.

Acknowledgements

We gratefully acknowledge financial support from the *Dirección*

General de Enseñanza Superior of Spain (DGES Project ref. PB96-0101), the *Consejería de Educación y Juventud y Fondo Social Europeo* (ref. IPR98CO17), the Royal Society, the Russian Foundation for Basic Research (grant no. 99-03-32984a) and MDL Information Systems (UK) Ltd, and we thank the Wolfson Foundation for establishing the Wolfson Centre for Organic Chemistry in Medical Science at Imperial College. We thank Mr R. Sheppard for the NMR experiments.

References

- 1 C. F. Marcos, C. Polo, O. A. Rakitin, C. W. Rees and T. Torroba, *Angew. Chem., Int. Ed. Engl.*, 1997, **36**, 281.
- 2 C. F. Marcos, C. Polo, O. A. Rakitin, C. W. Rees and T. Torroba, *Chem. Commun.*, 1997, 879.
- 3 C. F. Marcos, O. A. Rakitin, C. W. Rees, L. I. Souvorova, T. Torroba, A. J. P. White and D. J. Williams, *Chem. Commun.*, 1998, 453.
- 4 C. W. Rees, A. J. P. White, D. J. Williams, O. A. Rakitin, C. F. Marcos, C. Polo and T. Torroba, *J. Org. Chem.*, 1998, **63**, 2189.
- 5 C. F. Marcos, O. A. Rakitin, C. W. Rees, T. Torroba, A. J. P. White and D. J. Williams, *Chem. Commun.*, 1999, 29.
- 6 H. Bock, I. Goebel, Z. Havlas, S. Liedle and H. Oberhammer, *Angew. Chem., Int. Ed. Engl.*, 1991, **30**, 187.
- 7 SHELXTL PC, version 5.03, Siemens Analytical X-Ray Instruments, Inc., Madison, WI, 1994.
- 8 C. W. Rees, A. J. P. White, D. J. Williams, O. A. Rakitin, L. S. Konstantinova, C. F. Marcos and T. Torroba, *J. Org. Chem.*, 1999, **64**, 5010.
- 9 D. M. McKinnon, in *Comprehensive Heterocyclic Chemistry II*, ed. A. R. Katritzky, C. W. Rees and E. F. V. Scriven, Pergamon Press, 1996, vol. 3, p. 569.
- 10 R. F. C. Brown and I. D. Rae, *Aust. J. Chem.*, 1964, **17**, 447.
- 11 C. H. Wei, *Acta Crystallogr., Sect. C*, 1985, **41**, 1525.
- 12 P. K. Morgensen, O. Simonsen, C. E. Wainwright and A. E. Underhill, *Acta Crystallogr., Sect. C*, 1991, **47**, 1905.
- 13 B. Rees, *Acta Crystallogr.*, 1970, **B26**, 1304.
- 14 S. Furberg and J. Solbakk, *Acta Chem. Scand., Ser. A*, 1974, **28**, 435.
- 15 O. Hassel and C. Rømming, *Quart. Rev.*, 1962, **16**, 1.

Paper 9/04680A