

Fmoc/Acyl protecting groups in the synthesis of polyamide (peptide) nucleic acid monomers

Zoltán Timár,^a Lajos Kovács,^{*a} Györgyi Kovács^a and Zoltán Schmel^b

^a Department of Medicinal Chemistry, Dóm tér 8 and ^b Department of Pharmaceutical Analysis, Somogyi B. u. 4, Albert Szent-Györgyi Medical University, H-6720 Szeged, Hungary. Fax: +36-62-42 52 62; E-mail: kovacs@ovrisc.mdche.u-szeged.hu

Received (in Cambridge, UK) 29th September 1999, Accepted 10th November 1999

The chemical synthesis of polyamide (peptide) nucleic acid (PNA) monomers **22–25** has been accomplished using Fmoc [*N*-(2-aminoethyl)glycine backbone], anisoyl (adenine), 4-*tert*-butylbenzoyl (cytosine) and isobutyryl/diphenylcarbamoyl (guanine) protecting-group combinations, thus allowing oligomer synthesis on *both* peptide and oligonucleotide synthesizers. An alternative method for the preparation of (*N*⁶-anisoyladenine-9-yl)acetic acid **7** is described using partial hydrolysis of a dianisoylated derivative. Different methods were studied for guanine alkylation including (a) Mitsunobu reaction; (b) low-temperature, sodium hydride- and (c) *N,N*-diisopropylethylamine-mediated alkylation reactions to give preferentially *N*⁹-substituted derivatives. Empirical rules are proposed for differentiating *N*⁹/*N*⁷-substituted guanines based on their ¹³C NMR chemical-shift differences.


Introduction

Polyamide (or as originally referred: peptide) nucleic acids (PNA) are one of the most powerful analogues of oligonucleotides in terms of chemical and enzymic stability, double- and triple-helix formation, with potential applications in antisense diagnosis and therapeutics.^{1,2} In these compounds the entire sugar–phosphate backbone is replaced with an *N*-(2-aminoethyl)glycine moiety and the nucleobases are attached through an *N*-acetyl linkage.

The chemical synthesis of PNA mostly relies on the assembly of the protected *N*-(2-aminoethyl)glycine backbone and protected nucleobase-substituted acetic acid structural units followed by standard oligomerization protocols.^{3,4} The pioneering efforts of a Danish group resulted in the application of Boc (backbone) and Z (cytosine, adenine) or *O*-benzyl (guanine) protection.^{5,6} Later the Uhlmann group used a monomethoxytrityl (MMTr)/acyl (anisoyl, 4-*tert*-butylbenzoyl, isobutyryl/acyl) strategy.^{7,8} All these methods require the use of (strong) acidic conditions (*e.g.*, TFA, HF) in the oligomer construction and final cleavage from the support. The need for milder methods led to the employment of the Fmoc group for backbone protection and Z⁹ or MMTr groups¹⁰ for the nucleobases. The Fmoc group is a convenient alternative to acid-sensitive backbone-protecting groups (Boc, MMTr) and allows easy monitoring of the coupling process.¹¹ The combination Fmoc/acyl should also be feasible since the former group can be cleaved without affecting the more stable base-protecting acyl groups.^{12–14} Herein we report on our results concerning the use of Fmoc (backbone)/acyl (4-*tert*-butylbenzoyl for cytosine; anisoyl for adenine; isobutyryl/*N,N*-diphenylcarbamoyl for guanine) protecting groups in the synthesis of PNA monomers. The prior protecting-group combinations (with the exception of Fmoc/MMTr) were used *either* for peptide *or* oligonucleotide synthesis protocols. With biologically important PNA–DNA and PNA–peptide conjugates in mind our approach offers a substantial advantage over the existing ones since *both* oligomerization methodologies are possible with the same monomers. Beside this, use of the frequently applied urethane protecting groups (*e.g.*, Z) for nucleobases is not practical as in our experience the yields are often very low. This paper complements and details our preliminary account.¹⁵

Results and discussion


The choice of nucleobase-protecting groups was motivated by different considerations since uniform protection, though attractive, is not possible. Thymine does not require protection and our synthesis of the thymine monomer, starting from acid **1**, was based on the procedure of Thomson *et al.*⁹ The anisoylated cytosine derivative **2** was alkylated to give acid **3** (Scheme 1) but the solubilities of these substances were so low



Scheme 1 Reagents and conditions: a, (1) BrCH₂COOMe, K₂CO₃, DMF; (2) NaOH, then HCl; b, AnCl, py, 80 °C; c, 4-Bu^tC₆H₄COCl, Et₃N, DMF; d, NaH, DMF, BrCH₂COOEt; e, NaOH, aq. 1,4-dioxane, then HCl. An = 4-MeOC₆H₄CO.

in common solvents that we had to abandon this group. The 4-*tert*-butylbenzoyl group proved to be more rewarding; acid **5**⁸ was easily obtained *via* intermediate **4** and used later.


(*N*⁶-Anisoyladenine-9-yl)acetic acid **7**⁷ was prepared in 28% overall yield from adenine *via* the alkylation of compound **6** (Scheme 2). An improved overall yield (40%) was obtained when *tert*-butyl bromoacetate was used for the alkylation



Scheme 2 Reagents and conditions: a, 1.5 equiv. AnCl, py, 100 °C; b, NaH, DMF, BrCH₂COOMe; c, NaOH, then KHSO₄; d, NaH, DMF, BrCH₂COOBu^t; e, 50% (v/v) TFA–CH₂Cl₂, (±)-1,4-dithiothreitol; f, NaH, DMF, BrCH₂COOEt; g, 2.5 equiv. AnCl, py, 80 °C; h, NaOH, aq. MeOH, then HCl.

instead of methyl bromoacetate followed by acidolysis. In an alternative approach ethyl (adenin-9-yl)acetate **8**⁵ was anisoylated with excess of anisoyl chloride and the resulting *N*⁶,*N*⁶-dianisoylated derivative (not isolated) was subjected to partial hydrolysis to give acid **7** in an improved yield (70%; overall 45% from adenine).


The substitution of guanine is notorious for giving *N*⁹/*N*⁷-regioisomers.¹⁶ Although the 2-amino group is not really nucleophilic enough to interfere with many transformations, the poor solubility of unprotected guanine excludes its use in most reactions. The application of *N*²-acyl (acetyl, propionyl, isobutyryl, *etc.*)-protected derivatives increases the solubility but *N*²-acylation alone cannot solve the fundamental problem of the selectivity of alkylation.¹⁶ Constraining guanine from its dominant 6-lactam structure to lactim (enolate) form by different groups has a beneficial effect on the ratio of *N*⁹/*N*⁷-regioisomers. The most successful in this respect is the *N,N*-diphenylcarbamoyl protecting group¹⁷ which reportedly gives in some cases a 100:1 ratio in favour of the *N*⁹-regioisomer.^{18,19} To see how the introduction of this group alters the selectivity of alkylation, first 2-*N*-isobutyrylguanine **9**²⁰ was alkylated with *tert*-butyl bromoacetate in the presence of sodium hydride to afford a nearly 1:1 ratio of *N*⁹/*N*⁷-isomers (**11** and **12**, respectively) in 74% yield (Scheme 3). The selection of the isobutyryl



Scheme 3 Reagents and conditions: a, (Pr^tCO)₂O, DMF, 150 °C; b, Ac₂O, DMF, 100 °C; c, Ph₂NCOCl, EtNPr₂, py; d, NaH, DMF, 0 °C, BrCH₂COOBu^t. Ibu = Pr^tCO; Dpc = Ph₂NCO.

group was motivated by the fact that although its removal under basic conditions is more sluggish than that of other simple acyl groups (acetyl, propionyl)^{17,21} it confers steric hindrance on the 2-amino group and thus prevents unwanted alkylation/glycosylation on it.¹⁹

Next the *N,N*-diphenylcarbamoyl derivative **10**^{19,22} was chosen for alkylation studies under different conditions. Its transformation with *tert*-butyl glycolate²³ in the Mitsunobu




Scheme 4 Reagents and conditions: a, DIAD, Ph₃P, THF, HOCH₂COOBu^t; b, DIAD, 4-Me₂NC₆H₄PPh₂, THF, HOCH₂COOBu^t; c, DIAD, Bu₃P, THF, HOCH₂COOBu^t; d, NaH, –20 °C, DMF, BrCH₂COOBu^t; e, 8% (v/v) TFA–CH₂Cl₂, 1,3-(MeO)₂C₆H₄, 0 °C, 18 h; f, BrCH₂COOMe, EtNPr₂, DMF; g, NaOH, aq. 1,4-dioxane–MeOH, then HCl.

reaction^{24,25} provided the product (**13**, Scheme 4) with good regioselectivity; however, its purification was very difficult and it was contaminated with significant amounts of triphenylphosphine oxide. (4-Dimethylaminophenyl)diphenylphosphine,^{26,27} claimed to give a phosphine oxide which can be removed by acidic extraction,²⁸ proved to be unsatisfactory since the product was still contaminated with the corresponding phosphine oxide. Tributylphosphine, the oxide of which is water-soluble, gave a cleaner product but the yield was low (36%). In the next experiments sodium hydride-mediated alkylation with *tert*-butyl bromoacetate was used to obtain the desired compound. We noticed that at ambient temperature the relative proportion of *N*⁷-regioisomer was relatively high, while lowering the temperature favoured the formation of the desired *N*⁹-regioisomer. At –20 °C a clean reaction gave negligible amounts of the *N*⁷-isomer but the yield of *N*⁹-isomer was still low (40%). Acidolysis of ester **13** in dilute TFA–CH₂Cl₂ (0 °C; 18 h) removed the *N,N*-diphenylcarbamoyl (Dpc) group without affecting the *tert*-butyl ester functionality (→ **11**). Albeit there is some evidence for the lability of this group in 50% (v/v) TFA–CH₂Cl₂⁸ or in the presence of Lewis acids²⁹ it was surprising that the Dpc group was more sensitive towards acids than was the *tert*-butyl group. The latter was expected to cleave under similar conditions.^{14,30}

The application of *N,N*-diisopropylethylamine as a hindered base and methyl bromoacetate⁸ to circumvent premature cleavage of the Dpc group in the subsequent hydrolysis gave a 71% yield of the product **14**, of which 58% was available without chromatography, along with 15% of the *N*⁷-isomer **15**. Basic hydrolysis of ester **14** led to acid **16** in a clean transformation. The surprisingly high yield of the unwanted isomer **15** in the first reaction underlines the fact that even the sterically hindered Dpc protecting group is not sufficient to steer the reaction to complete regioselectivity. Thus the claim that the use of the Dpc group has solved the historic problem of regioselective *N*⁹-substitution of guanine^{18,19} seems to be restricted to the realm of glycosylation reactions, while alkylation transformations require further experimentation.


The coupling of nucleobase-substituted acetic acids **1**, **5**, **7**, **16** with the backbone unit **17**⁹ under standard peptide-coupling conditions afforded the PNA esters **18–21** which were subsequently acidolysed (TFA in dichloromethane) to give the PNA monomers **22–25** (Scheme 5). As expected from our previous experience (**13** → **11**, Scheme 4) in the latter reaction the Dpc protecting group was removed along with the *tert*-butyl group. It is clear that in this final deprotection step the protect-



Scheme 5 Reagents and conditions: a, HBTU, HOBT, DMF, EtNPr₂; b, 17–43% (v/v) TFA–CH₂Cl₂, 1,3-(MeO)₂C₆H₄.

ing groups of ester **21** (Fmoc, Ibu, Dpc, Bu^t) are not completely orthogonal and further research is required. Although the presence of the Dpc group is not essential in oligomer synthesis, it may confer better solubility on the guanine subunits.

UV spectroscopy is often used to locate substituents on nucleobases. However, this technique is not reliable in the case of guanines especially if only one regioisomer is available.³¹ Moreover, there are no data on the influence of lactam/lactim tautomerism of guanines on UV properties. Indeed, the UV spectra of *N*⁹/*N*⁷-regioisomers **11/12** (lactams) and **14/15** (lactims), respectively, in buffered ethanolic solutions (pH 0, 6 and 13) revealed that there are no significant differences which would justify the assignment based solely on UV data. Therefore the site of alkylation in guanine derivatives **13–15**, **26**³² was established in 2D NMR (HMQC^{33,34} and HMBC^{35,36}) experiments.



Scrutinizing the ¹³C NMR chemical-shift parameters of compounds **11–16**, **21**, **25**, **26** (Table 1) and a further 45 *N*⁹/*N*⁷-substituted guanines^{19,37–39} (altogether 54 compounds) show that δ_{C-5} is the most sensitive to the *N*⁹/*N*⁷-substitution pattern (Fig. 1, *N*⁹: 113.75–123.70 ppm; *N*⁷: 104.56–115.09 ppm; for regioisomers the difference [Δδ_{C-5} = δ_{C-5}(*N*⁹) – δ_{C-5}(*N*⁷)] is 7.86–9.82 ppm), insensitive to the lactam/lactim tautomerism (data not shown) and this signal alone can be of diagnostic value, especially if data for both regioisomers are available. However, due to the overlapping of chemical-shift ranges for regioisomers this parameter might not be sufficient for unambiguous assignment. Kjellberg and Johansson³⁷ suggested that δ_{C-1}, δ_{C-5} and δ_{C-8} could be used to differentiate *N*⁹/*N*⁷-substituted guanines. This was also corroborated, in part, by our findings; however, some further tendencies were also observed. The utility of differential parameters *a* = δ_{C-4} – δ_{C-5}, *b* = δ_{C-8} – δ_{C-5}, *c* = δ_{C-5} – δ_{C-1}, (Fig. 1) has been assessed and the following conclusions could be drawn:

1. The parameter *a* distinctly differs for the regioisomers {*N*⁹: 28.20–35.41 ppm; *N*⁷: 41.35–54.25 ppm; for regioisomers the difference [Δ*a* = *a*(*N*⁹) – *a*(*N*⁷)] is –9.93 to –20.53 ppm}, shows little variation for lactam/lactim tautomerism (data not shown) and presents no overlapping ranges. The diagnostic value of this observation is slightly diminished since δ_{C-4} usually cannot be simply identified without having recourse to more




Fig. 1 ¹³C NMR chemical-shift parameters of *N*⁹/*N*⁷-substituted guanines.

sophisticated assignment techniques (selective INEPT, HMQC, HMBC, etc.).

2. The parameter *b* shows similar characteristics {*N*⁹: 16.10–26.91 ppm; *N*⁷: 29.70–41.17 ppm; for regioisomers the difference [Δ*b* = *b*(*N*⁹) – *b*(*N*⁷)] is –11.68 to –16.15 ppm} and its utility is further enhanced by the fact that δ_{C-5} is unmistakable among the skeletal carbons and δ_{C-8} can simply be located in a *J*-modulated spin-echo experiment.

3. The parameter *c* can be clustered according to the nature of attached substituent rather than lactam/lactim tautomerism and it gives useful values for glycosylated derivatives {*N*⁹: 33.38–35.81 ppm; *N*⁷: 16.30–23.38 ppm; for regioisomers the difference [Δ*c* = *c*(*N*⁹) – *c*(*N*⁷)] is 10.61–12.79 ppm} while for (cyclo)alkylated compounds it is of less use {*N*⁹: 53.40–83.45 ppm; *N*⁷: 54.57–71.15 ppm; for regioisomers the difference [Δ*c* = *c*(*N*⁹) – *c*(*N*⁷)] is 10.68–13.35 ppm}. The identification of δ_{C-1}, involved in this parameter, often requires more sophisticated techniques.

As a conclusion it can be seen that the values *a*, *b* [both for (cyclo)alkyl and glycosylated derivatives] and *c* (for glycosylated derivatives) are useful for characterizing the *N*⁹/*N*⁷-substitution pattern of guanines. From a practical point of view the parameter *b* is the most convenient one for the reasons explained above. It is noteworthy that δ_{C-5} (118.81 ppm) and the parameters *a* (35.38 ppm) and *b* (26.06 ppm) for compound **10**¹⁹ are in good agreement with those for *N*⁹-substituted derivatives, suggesting that its dominant tautomer is 9*H* in DMSO-*d*₆ solution.

Further studies relating to the application of the above monomers in the preparation of PNA oligomers and our quest for novel combinations of protecting groups are in progress and will be reported in due time.

Experimental

General

The following abbreviations are employed: diisopropyl azodicarboxylate (DIAD); *N,N*-diisopropylethylamine (DIPEA); 2-(1*H*-benzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (HBTU); 1-hydroxybenzotriazole (HOBT); trifluoroacetic acid (TFA). Chemicals were purchased from Aldrich or Fluka. Sodium hydride refers to a 55% suspension in mineral oil. Anhydrous solvents were prepared as described.⁴⁰ Light petroleum refers to the fraction with distillation range 40–60 °C. Thymine derivatives **1**, **18** and **22** were prepared using the procedure of Thomson *et al.*⁹ *N*⁷-Benzylguanine hydrochloride **26**³² was prepared for comparison of its ¹³C NMR parameters with those of other guanine derivatives (see Table 1). Organic solutions were dried using magnesium sulfate and evaporated in Büchi rotary evaporators. TLC: Kieselgel 60 F₂₅₄ (Merck), visualization: UV light. Column chromatography: Kieselgel 60 (0.063–0.200 mm, Merck). Mp: Electrothermal IA 8103 apparatus. Elemental analysis: Perkin-Elmer CHN

analyzer model 2400. UV: PE Lambda 10 spectrometer, λ_{max} /nm (lg ϵ), sh: shoulder. IR: Bio-Rad FTS-60A (KBr pellets, ν_{max} /cm⁻¹; s, strong; m, medium; w, weak). NMR: Bruker Avance DRX 400 and 500 spectrometers (¹H: 400.13 MHz and 500.13 MHz; ¹³C: 100.62 MHz and 125.76 MHz, respectively), DMSO-d₆ solutions, δ (ppm), J (Hz). Spectral patterns: s, singlet; d, doublet; dd, double doublet; t, triplet; m, multiplet; br, broad; deut, deuterable. The superscripts *, # denote interchangeable assignments. For the 2D experiments (HMQC, HMBC) the standard Bruker software packages (INV4GSSW, INV4GSLRNDWS) were applied. For ¹³C NMR data of guanine derivatives see Table 1. Mass spectrometry: Finnigan MAT TSQ 7000, electrospray (ESI) and atmospheric pressure chemical ionization (APCI) techniques. IUPAC names: AutoNom™ 2.1 (as implemented in ChemDraw® 5.0). Statistical analysis of ¹³C NMR chemical-shift parameters of guanine derivatives (Fig. 1): Corel® Quattro® Pro 8.0, @ functions and SigmaPlot 4.01. Details are available upon request.

[4-(4-Methoxybenzoylamino)-2-oxo-1,2-dihydropyrimidin-1-yl]acetic acid 3

Cytosine (1.11 g, 10.0 mmol) suspended in pyridine (50 mL) was stirred at room temperature while 4-methoxybenzoyl chloride (2.56 g, 15.0 mmol) was added and the reaction mixture was stirred in an oil-bath at 80 °C. The cytosine rapidly dissolved, and then the product precipitated from the solution. After 2 h the mixture was evaporated *in vacuo* and coevaporated with methanol (2×). The residue was suspended in methanol and the mixture was filtered to afford compound 3 (2.28 g, 93%) as a white powder, mp > 260 °C. The product [4-methoxy-*N*-(2-oxo-1,2-dihydropyrimidin-4-yl)benzamide, 2] was insoluble in practically all solvents, therefore it was used without further purification; R_f 0.58 (CH₂Cl₂–MeOH 8:2); λ_{max} [0.20% (v/v) TFA in EtOH]/nm 216 (lg ϵ 4.00), 276 (4.35); ν_{max} /cm⁻¹ 3276w, 3150w, 3073w, 2994w, 2846w, 1714s, 1692s, 1621m, 1609m, 1578m, 1497s, 1406w, 1260s, 1189m, 801m.

The majority (2.20 g, 8.97 mmol) of this substance was suspended in anhydrous DMF (25 mL), anhydrous K₂CO₃ (1.24 g, 8.97 mmol) and methyl bromoacetate (0.86 mL, 8.97 mmol) were added and the mixture was stirred for 24 h. The reaction mixture was filtered, washed with DMF and the filtrate was evaporated. Water (9 mL) and 4 M HCl (0.4 mL) were added to the residue and stirred for 15 min. The ester was filtered off and then added to a mixture of aq. 2 M NaOH (6.5 mL, 13.0 mmol) and water (12 mL) and the reaction mixture was sonicated. The substance dissolved and after 30 min no starting material was present (TLC). The reaction mixture was acidified with 4 M HCl (3.6 mL, 14.4 mmol) and the precipitated substance was filtered off. Purification of the crude product (2.94 g) was attempted by recrystallization from methanol. 1 L of solvent was not enough to dissolve the above amount, and so 0.45 g was filtered off to give a white powder, TLC: single spot, mp 220 °C (darkens), 239 °C (decomp.). From the methanolic solution a second crop (0.86 g) was obtained as a white powder, TLC: single spot, mp 166 °C (darkens), 227 °C (decomp.). Overall yield: 1.31 g (48%); R_f 0.58 (MeCN–MeOH–AcOH 4:1:1) (Found: C, 55.5; H, 4.2; N, 13.7. Calc. for C₁₄H₁₃N₃O₅: C, 55.45; H, 4.3; N, 13.9%); λ_{max} (EtOH)/nm 208 (lg ϵ 4.58), 252 (4.55), 287 (4.27); ν_{max} /cm⁻¹ 3147w, 3076w, 1711s, 1631m, 1615w, 1503m, 1419w, 1251m, 1183m, 755w; δ_{H} (400 MHz) 3.84 (3 H, s, OCH₃), 4.56 (2 H, s, CH₂), 7.04 and 8.03 (2 × 2 H, 2 × d, J 8.9, ArH), 7.30 (1 H, d, J 7.2, H-5*), 8.08 (1 H, d, J 7.2, H-6*), 11.03 (1 H, br s, deut, OH[#]), 11.97 (1 H, br s, deut, NH[#]); m/z (ESI) 304 (44%, [M + H]⁺). This compound was mentioned by Breipohl *et al.*⁸ but not described in detail.

[6-(4-Methoxybenzoylamino)purin-9-yl]acetic acid 7

A. Alkylation of [6-(4-methoxybenzoylamino)purine] and subsequent acidolysis. To [6-(4-methoxybenzoylamino)purine] 6⁷

(6.47 g, 24.0 mmol) suspended in anhydrous DMF (120 mL) was added sodium hydride (1.05 g, 24.0 mmol) in portions and the mixture was stirred at room temperature for 30 min. *tert*-Butyl bromoacetate (3.9 mL, 26.4 mmol) was added dropwise and stirring was continued for 2 h. The mixture was evaporated *in vacuo* and the residue was suspended in a mixture of water (200 mL) and dichloromethane (200 mL). The resulting precipitate was filtered off and dried (5.66 g, 61%). The majority (5.50 g, 14.3 mmol) of this substance was stirred with 50% (v/v) TFA in dichloromethane (40 mL) and (±)1,4-dithiothreitol (0.20 g, 1.3 mmol) at room temperature for 4 h. The solution was evaporated *in vacuo* and the residue was coevaporated with EtOAc (5×). The residue was dissolved in 5% (w/v) aq. NaHCO₃ (150 mL), filtered and acidified with 10% (w/v) aq. NaHSO₄ (100 mL). The precipitated solid was filtered off (4.11 g, 88%), to give 7 mp 213 °C (darkens), 250 °C (decomp.). The characteristics (¹H NMR and mass spectra) of this substance were in good agreement both with the published values⁷ and with those of the substance prepared in procedure B.

B. Controlled hydrolysis of a dianisoylated derivative. Ethyl (adenin-9-yl)acetate 8⁵ (4.42 g, 20 mmol) was suspended in anhydrous pyridine (50 mL), heated to 80 °C for 30 min, then cooled to room temperature. 4-Methoxybenzoyl chloride (8.53 g, 50.0 mmol) was added in portions and the mixture was stirred for 18 h, then evaporated *in vacuo* and the residue was coevaporated with toluene (3×). The residue was dissolved in dichloromethane (70 mL), and the solution was washed with 10% (w/v) citric acid (2 × 30 mL), dried and evaporated *in vacuo*. The crude product (14.96 g) was dissolved in warm ethanol (100 mL), cooled to room temperature, 2 M aq. NaOH (30 mL) was added, and the solution was left at room temperature and checked from time to time by TLC. After 175 min more aq. NaOH (5 mL) was added. The reaction was stopped after 4 h by addition of 1 M HCl (35 mL), pH ≈ 5, and the solution was evaporated *in vacuo*. The crude product was recrystallized from methanol (1 L) to afford a white powder (4.58 g, 70%), mp 215 °C (darkens), 254 °C (decomp.) [lit.,⁷ 222–223 °C (decomp.)]. The ¹H NMR and mass spectra of this compound were in good agreement with the published values.⁷

(2-Isobutyrylamino-6-oxo-1,6-dihydropurin-9-yl)acetic acid *tert*-butyl ester 11 and (2-isobutyrylamino-6-oxo-1,6-dihydropurin-7-yl)acetic acid *tert*-butyl ester 12

N-(6-Oxo-6,9-dihydro-1*H*-purin-2-yl)isobutyramide 9^{19,22} (1.11 g, 5.0 mmol) was suspended in anhydrous DMF and the mixture was chilled to 0 °C. Sodium hydride (0.36 g, 8.25 mmol) was added and the mixture was stirred at 0 °C for 30 min. *tert*-Butyl bromoacetate (0.81 mL, 5.5 mmol) was added and the reaction was stopped after 2 h by addition of a small amount of solid CO₂ and methanol (2 mL). The reaction mixture was evaporated *in vacuo* and the residue was chromatographed using 0–5% (v/v) methanol in dichloromethane. Eluted first was the less polar *N*⁷-isomer 12, (0.43 g, 26%), second a mixture (in ≈ 1:1 ratio as judged by TLC and ¹H NMR) of *N*⁷- and *N*⁹-isomer (0.24 g, 14%), and third the pure *N*⁹-isomer 11 (0.56 g, 34%).

Ester 11: white powder, mp 204 °C (decomp., from EtOH); R_f 0.13 (CH₂Cl₂–MeOH 95:5) (Found: C, 53.9; H, 6.4; N, 21.1. Calc. for C₁₅H₂₁N₅O₄: C, 53.7; H, 6.3; N, 20.9%); λ_{max} [50% (v/v) 1 M HCl in EtOH, pH 0]/nm 206 (lg ϵ 4.26), 265 (4.23); λ_{max} [50% (v/v) phosphate buffer in EtOH, pH 6]/nm 260 (lg ϵ 4.17), 282sh (4.02); λ_{max} [50% (v/v) 0.1 M NaOH in EtOH, pH 13]/nm 216 (lg ϵ 4.39), 263 (4.06); ν_{max} /cm⁻¹ 3151w, 2980w, 2932w, 1753s, 1698m, 1673s, 1614m, 1562m, 1549m, 1483w, 1411m, 1233m, 1154m, 1143m, 795w; δ_{H} (500 MHz) 1.11 [6 H, d, J 6.8, (CH₃)₂CH], 1.40 (9 H, s, Bu[†]), 2.78 [1 H, pseudoquintet, (CH₃)₂CH], 4.88

Table 1 ^{13}C NMR chemical shifts of guanine derivatives (δ , ppm)^a

Compd.	Subst.	C-2	C-4	C-5	C-6	C-8	Other carbons
10 ^{b,c}	—	152.34	154.19	118.81	157.41	144.87	irrelevant
11 ^d	<i>N</i> ⁹	148.49	155.17	119.98	149.30	140.72	19.22 [(CH ₃) ₂ CH], 28.03 [(CH ₃) ₃ C], 35.01 [(CH ₃) ₂ CH], 45.17 (CH ₂ COO), 82.69 [(CH ₃) ₃ C], 167.03 (COOBu ^t), 180.56 (Pr ^t CO)
12	<i>N</i> ⁷	147.55	157.20	112.08	152.95	144.50	19.23 [(CH ₃) ₂ CH], 28.00 [(CH ₃) ₃ C], 35.06 [(CH ₃) ₂ CH], 48.26 (CH ₂ COO), 82.36 [(CH ₃) ₃ C], 167.20 (COOBu ^t), 180.32 (Pr ^t CO)
13 ^{e,f}	<i>N</i> ⁹	153.24	155.88	120.47	155.95	147.01	20.08 [(CH ₃) ₂ CH], 28.52 [(CH ₃) ₃ C], 35.18 [(CH ₃) ₂ CH], 45.82 (CH ₂ COO), 83.40 [(CH ₃) ₃ C], 127.76, 128.15, 130.26 (arom. Cs), 142.51 (arom. quaternary C), 151.03 (OCON), 167.33 (COOBu ^t), 175.97 (Pr ^t CO)
14 ^{e,f}	<i>N</i> ⁹	153.29	155.91	120.51	155.91	146.90	20.08 [(CH ₃) ₂ CH], 35.25 [(CH ₃) ₂ CH], 45.14 (CH ₂ COO), 53.48 (CH ₃ O), 127.79, 128.14, 130.26 (arom. Cs), 142.51 (arom. quaternary C), 151.03 (OCON), 168.85 (COOMe), 175.90 (Pr ^t CO)
15 ^f	<i>N</i> ⁷	149.53	164.63	112.19	141.28	150.44	19.20 [(CH ₃) ₂ CH], 34.26 [(CH ₃) ₂ CH], 47.50 (CH ₂ COO), 52.50 (CH ₃ O), 128.91, 129.04, 129.36 (arom. Cs), 141.28 (arom. quaternary C), 152.02 (OCON), 167.77 (COOMe), 174.94 (Pr ^t CO)
16 ^e	<i>N</i> ⁹	152.19	154.81	119.62	155.03	146.50	19.08 [(CH ₃) ₂ CH], 34.22 [(CH ₃) ₂ CH], 66.22 (CH ₂ COO), 126.90, 129.17, 129.37, 129.90 (arom. Cs), 141.49 (arom. quaternary C), 150.07 (OCON), 174.93 (Pr ^t CO, COOH)
21 ^{e,g}	<i>N</i> ⁹	152.01	154.72	119.40	155.05	146.31	18.98 [(CH ₃) ₂ CH], 27.47/27.50 [(CH ₃) ₃ C], 34.17 [(CH ₃) ₂ CH], 46.56 (OCH ₂ CH), 43.73/43.99, 47.09, 48.76, 49.96 (4 × CH ₂), 65.28 (OCH ₂ CH), 80.88/81.98 [(CH ₃) ₃ C], 119.88, 124.79/124.88, 126.81/126.85, 126.95, 127.40, 128.71, 129.20 (arom. Cs), 140.54, 141.45, 143.61/143.65 (arom. quaternary C), 150.00 (OCON), 155.13, 155.95/156.21, 166.24/166.74, 167.69/168.32 (4 × CO), 175.00 (Pr ^t CO)
25 ^{e,g}	<i>N</i> ⁹	147.79, 147.82	149.16, 149.24	119.49	154.79	140.45	18.76 [(CH ₃) ₂ CH], 34.58 [(CH ₃) ₂ CH], 46.66 (OCH ₂ CH), 43.86/44.00, 46.85/46.97, 47.76, 49.11 (4 × CH ₂), 65.43 (OCH ₂ CH), 120.01/120.04, 124.94/125.01, 126.95, 127.53 (arom. Cs), 140.64/140.68, 143.76/143.78 (arom. quaternary C), 156.07/156.30, 166.37/166.91, 170.28/170.72 (3 × CO), 180.05 (Pr ^t CO, COOH)
26 ^{d,f,h}	<i>N</i> ⁷	154.95	151.46	108.07	153.92	141.06	51.09 (CH ₂), 128.70, 129.15, 129.62 (arom. Cs), 136.38 (arom. quaternary C)

^a In DMSO-*d*₆; 125.76 MHz; *J*-modulated spin-echo experiments; for guanine numbering see ester **11**, Scheme 3. ^b Ref. 19. ^c The C-4, C-5, C-6, C-8 signals were observed only after adding trifluoroacetic acid. ^d The assignment of signals corresponding to C-2 and C-6 carbons is tentative. ^e The assignment of signals corresponding to C-4 and C-6 carbons is tentative. ^f Assignment based on HMQC and HMBC experiments. ^g Some signals were doubled due to the presence of rotamers. ^h *N*⁷-Benzylguanine hydrochloride, prepared according to Bridson *et al.*³²

(2 H, s, CH₂COO), 7.95 (1 H, s, H-8), 11.65 (1 H, br s, deut, NH), 12.10 (1 H, br s, deut, NH); *m/z* (ESI) 693 (20%, [2M + Na]⁺), 671 (55, [2M + H]⁺), 336 (100, [M + H]⁺).

Ester **12**: white powder, mp 202.5 °C (decomp., from EtOH); *R*_f 0.19 (CH₂Cl₂–MeOH 95:5) (Found: C, 53.65; H, 6.15; N, 21.1%); λ_{max} [50% (v/v) 1 M HCl in EtOH, pH 0]/nm 206 (lg ϵ 4.24), 263 (4.20); λ_{max} [50% (v/v) phosphate buffer in EtOH, pH 6]/nm 221 (lg ϵ 4.24), 265 (4.11), 282sh (3.98); λ_{max} [50% (v/v) 0.1 M NaOH in EtOH, pH 13]/nm 224 (lg ϵ 4.31), 269 (4.01); ν_{max} /cm^{−1} 3240w, 2981w, 2937w, 1741m, 1695s, 1677s, 1604s, 1535w, 1421w, 1390m, 1370m, 1238m, 1160m, 747w; δ_{H} (500 MHz) 1.11 [6 H, d, *J* 6.8, (CH₃)₂CH], 1.39 (9 H, s, Bu^t), 2.73 [1 H, pseudo-quintet, (CH₃)₂CH], 5.07 (2 H, s, CH₂COO), 8.11 (1 H, s, H-8), 11.55 (1 H, br s, deut, NH), 12.14 (1 H, br s, deut, NH); *m/z* (ESI) 693 (40%, [2M + Na]⁺), 671 (25, [2M + H]⁺), 358 (27, [M + Na]⁺), 336 (100, [M + H]⁺).

[6-Diphenylcarbamoyloxy-2-(isobutrylamino)purin-9-yl]acetic acid *tert*-butyl ester **13**

A. Mitsunobu reaction, general procedure. Compound **10**^{19,22} (1.00 g, 2.40 mmol) was suspended in anhydrous THF (50 mL) and the mixture was refluxed for 20 min to achieve partial dissolution of the starting material.²⁰ The suspension was cooled to room temperature, *tert*-butyl glycolate²³ (0.40 g, 3.0 mmol), the appropriate phosphine (3.19 mmol) and DIAD (0.62 mL, 3.19 mmol) were added dropwise, and the mixture was stirred at room temperature. The reaction mixture completely dissolved and became yellow coloured. After completion of the reaction (TLC) the solution was evaporated *in vacuo* and the residue was subjected to chromatographic purification.

A1. With triphenylphosphine.—Reaction time: 4 h at room temperature. Chromatography: 50–70% (v/v) ethyl acetate in light petroleum. Eluted first was the product **13** (0.31 g), slightly

contaminated with triphenylphosphine oxide. Further fractions were also obtained containing varying proportions of the product and triphenylphosphine oxide. The different, partly crystalline fractions were triturated with methanol upon which the product crystallized. This was filtered off and washed with light petroleum. The cleanest product (0.40 g, 31%), a white powder, melted at 183.2–185.5 °C. A further crystalline crop (0.39 g) containing the product and triphenylphosphine oxide (TLC) was also obtained; *R*_f 0.50 (CH₂Cl₂–MeOH 95:5) (Found: C, 63.5; H, 5.5; N, 15.7. Calc. for C₂₈H₃₀N₆O₅: C, 63.4; H, 5.7; N, 15.8%); λ_{max} (EtOH)/nm 205 (lg ϵ 4.60), 229 (4.53), 258sh (4.15), 279 (4.08); λ_{max} /cm^{−1} 3462w, 3346w, 2979w, 2934w, 1738s, 1715m, 1624m, 1587m, 1524m, 1449m, 1411m, 1305m, 1240m, 1187s, 1164s, 1056m, 758w, 700m; δ_{H} (500 MHz) 1.09 [6 H, d, *J* 6.8, (CH₃)₂CH], 1.43 (9 H, s, Bu^t), 2.87 [1 H, pseudo-quintet, *J* 6.8, (CH₃)₂CH], 5.03 (2 H, s, CH₂), 7.29–7.53 (10 H, m, ArH), 8.45 (1 H, s, H-8), 10.69 (1 H, br s, deut, NH); *m/z* (ESI) 557 (8%, [2Ph₃PO + H]⁺), 531 (100, [M + H]⁺).

A2. With 4-(dimethylamino)phenyl(diphenyl)phosphine.^{26,27}—Reaction time: 2.5 h at 0 °C. Work-up: the crude product was dissolved in dichloromethane (50 mL) and extracted successively with 4 M HCl (3 × 25 mL) and with 5% (w/v) aq. NaHCO₃ (50 mL). TLC revealed that most of the 4-(dimethylamino)phenyl(diphenyl)phosphine oxide remained in the organic phase. The organic phase was dried, and purified by column chromatography using 0–1% (v/v) methanol in dichloromethane. Methanolic trituration and filtration (light petroleum) afforded the product (0.42 g, 33%), mp 182.5–185.0 °C. The IR, ¹H NMR and mass spectra of this compound were in good agreement with those of the substance obtained in procedure **A1**.

A3. With tributylphosphine.—Reaction time: 1.5 h at 0 °C. Work-up: the crude product was dissolved in dichloromethane (50 mL) and extracted with water (3 × 25 mL) to remove

the tributylphosphine oxide. Chromatography: 0–1.5% (v/v) methanol in dichloromethane. Methanolic trituration and filtration (light petroleum) afforded the product (0.46 g, 36%), mp 182.2–184.8 °C. The IR, ¹H NMR and mass spectra of this compound were in good agreement with those of the substance obtained in procedure A1.

B. Low-temperature, sodium hydride-mediated alkylation. To compound **10**^{19,22} (1.40 g, 3.36 mmol) suspended in anhydrous DMF (20 mL) was added sodium hydride (0.16 g, 3.70 mmol) at room temperature. After 30 min the reaction mixture was chilled to –20 °C and maintained at this temperature. *tert*-Butyl bromoacetate (0.60 mL, 4.0 mmol) was added dropwise. After 2 h the reaction was stopped by addition of a small amount of solid CO₂ and methanol, and the mixture was evaporated *in vacuo* and coevaporated with toluene (2×). The residue was dissolved in a mixture of water and dichloromethane (20 mL each). The aqueous phase was extracted with dichloromethane (3 × 15 mL), and the combined organic phases were dried, and evaporated *in vacuo*. Chromatography: 0–1% (v/v) methanol in dichloromethane. Methanolic trituration and subsequent crystallization from methanol (15 mL) afforded the product (0.70 g, 40%) as a white powder, mp 183.1–184.5 °C. The IR, ¹H NMR and mass spectra of this compound were in good agreement with those of the substance obtained in procedure A1.

Hydrolysis of [6-diphenylcarbamoyloxy-2-(isobutrylamino)-purin-9-yl]acetic acid *tert*-butyl ester (**13** → **11**)

To ester **13** (0.210 g, 0.38 mmol) dissolved in anhydrous dichloromethane (6 mL) were added 1,3-dimethoxybenzene (0.070 mL, 0.53 mmol) and TFA (0.50 mL, 6.5 mmol) at 0 °C and the mixture was stirred for 18 h. The reaction mixture was diluted with dichloromethane (20 mL), and extracted with satd. aq. NaHCO₃ solution (3 × 10 mL) to remove the excess of acid. Chromatography: 0–10% (v/v) methanol in dichloromethane to give the lactam **11** (0.069 g, 55%) as an amorphous foam. The IR, ¹H NMR and mass spectra of this product were in good agreement with those of the substance obtained in a previous experiment (*vide supra*).

[6-Diphenylcarbamoyloxy-2-(isobutrylamino)purin-9-yl]acetic acid methyl ester **14** and [6-diphenylcarbamoyloxy-2-(isobutrylamino)purin-7-yl]acetic acid methyl ester **15** (*cf.* ref. 8)

To compound **10**^{19,22} (3.53 g, 8.47 mmol) suspended in anhydrous DMF (40 mL) was added DIPEA (2.87 mL, 16.74 mmol) and the mixture was briefly heated to 80 °C until a clear solution was obtained (10 min). The mixture was cooled to room temperature, methyl bromoacetate (0.87 mL, 9.32 mmol) was added, and the mixture was stirred for 20 h. The reaction mixture was evaporated *in vacuo* and the residue was coevaporated with methanol (3×). The partly crystalline material was suspended in methanol (40 mL) and added dropwise to water (120 mL) with vigorous stirring. The precipitate was filtered off (3.99 g, 96%) and recrystallized from EtOAc (190 mL) to afford the title products (2.01 g, 49%), mp 167.0–168.4 °C; from the mother liquor was obtained a further crop (0.37 g, 9%), mp 168.0–170.4 °C. The mother liquor was evaporated and chromatographed by using 1–4% (v/v) methanol in dichloromethane. Eluted first was ester **14** (0.54 g, 13%) then its regioisomer **15** (0.60 g, 15%). Overall yield of **14**: 2.92 g, 71%.

Ester **14**: white powder, mp 170.0–171.4 °C (from EtOAc); *R*_f 0.37 (CH₂Cl₂–MeOH 95:5) (Found: C, 61.35; H, 5.1; N, 17.4. Calc. for C₂₅H₂₄N₆O₅: C, 61.5; H, 4.95; N, 17.2%); λ_{max}[50% (v/v) 1 M HCl in EtOH, pH 0]/nm 207 (lg ε 4.50), 226 (4.53), 253sh (4.17), 279 (4.07); λ_{max}[50% (v/v) phosphate buffer in EtOH, pH 6]/nm 227 (lg ε 4.54), 259sh (4.14), 277 (4.08); λ_{max}[50% (v/v) 0.1 M NaOH in EtOH, pH 13]/nm 230 (lg ε 4.45),

276 (4.03); ν_{max}/cm^{–1} 3295w, 2996w, 2956w, 1756m, 1726s, 1681s, 1631m, 1599m, 1491m, 1384m, 1340s, 1233s, 1223s, 1192m, 1063m, 789w, 758w, 704w; δ_H (500 MHz) 1.08 [6 H, d, *J* 6.8, (CH₃)₂CH], 2.83 [1 H, pseudoquintet, *J* 6.8, (CH₃)₂CH], 3.73 (3 H, s, CH₃), 5.15 (2 H, s, CH₂), 7.31–7.49 (10 H, m, ArH), 8.43 (1 H, s, H-8), 10.64 (1 H, br s, deut, NH); *m/z* (ESI) 511 (25%, [M + Na]⁺), 489 (100, [M + H]⁺).

Ester **15**: amorphous foam; *R*_f 0.35 (CH₂Cl₂–MeOH 9:1) (Found: C, 61.3; H, 5.1; N, 17.0); λ_{max}[50% (v/v) 1 M HCl in EtOH, pH 0]/nm 207 (lg ε 4.59), 227 (4.56), 252sh (4.26), 283 (4.09); λ_{max}[50% (v/v) phosphate buffer in EtOH, pH 6]/nm 230 (lg ε 4.61), 256sh (4.17), 282 (4.02); λ_{max}[50% (v/v) 0.1 M NaOH in EtOH, pH 13]/nm 229 (lg ε 4.53), 287sh (3.90); ν_{max}/cm^{–1} 3438w, 2972w, 1753s, 1707m, 1639m, 1592w, 1493s, 1445m, 1301s, 1187s, 761m, 701m; δ_H (500 MHz) 1.11 [6 H, d, *J* 6.8, (CH₃)₂CH], 2.85 [1 H, pseudoquintet, *J* 6.8, (CH₃)₂CH], 3.56 (3 H, s, CH₃), 5.20 (2 H, s, CH₂), 7.28–7.48 (10 H, m, ArH), 8.53 (1 H, s, H-8), 10.57 (1 H, br s, deut, NH); *m/z* (ESI) 511 (40%, [M + Na]⁺), 489 (100, [M + H]⁺).

[6-Diphenylcarbamoyloxy-2-(isobutrylamino)purin-9-yl]acetic acid **16** (*cf.* ref. 8)

Ester **14** (2.24 g, 4.59 mmol) was suspended under sonication in a mixture of methanol (6 mL), 1,4-dioxane (24 mL) and water (12 mL). 1 M aq. NaOH (5 mL) was added and the mixture was stirred for 30 min. The pH of the mixture was brought to ≈6 by addition of 1 M HCl and the organics were evaporated off *in vacuo*. The solution was diluted with water (120 mL) and acidified to pH ≈3 by addition of 1 M HCl. The precipitate was filtered off, and washed with ice–water to give acid **16** (1.99 g, 91%), white powder, mp 156 °C (decomp.). Attempted recrystallization from EtOAc resulted in gel formation; *R*_f 0.40 (MeCN–MeOH–AcOH 8:1:1) (Found: C, 60.5; H, 4.4; N, 17.4. Calc. for C₂₄H₂₂N₆O₅: C, 60.75; H, 4.7; N, 17.7%); λ_{max}(EtOH)/nm 229 (lg ε 4.51), 260sh (4.11), 279 (4.06); ν_{max}/cm^{–1} 3379w, 1723s, 1627m, 1591m, 1522m, 1493m, 1441m, 1411m, 1307m, 1200s, 1187s, 760w, 701m; δ_H (500 MHz) 1.09 [6 H, s, (CH₃)₂CH], 2.83 [1 H, m, (CH₃)₂CH], 5.05 (2 H, s, CH₂), 7.29–7.50 (10 H, m, ArH), 8.46 (1 H, s, H-8), 10.65 (1 H, br s, deut, NH); *m/z* [ESI (CHCl₃ + MeOH)], 971 (8%, [2M + Na]⁺), 514 (8, [M + K]⁺), 497 (20, [M + Na]⁺), 475 (100, [M + H]⁺).

Esters 19–21

General procedure. To acid **5**,⁷ or **16** (2.0 mmol) dissolved in anhydrous DMF (20 mL) were added HOBt hydrate (0.61 g, 4.0 mmol) and HBTU (1.52 g, 4.0 mmol). Meanwhile ester **17**⁹ (1.95 g, 3.0 mmol for acids **5**, **7** and 1.30 g, 2.0 mmol for acid **16**) was suspended in dichloromethane (20/30 mL), extracted with satd. aq. NaHCO₃ (10 mL) and dried. The above dichloromethane solution of free **17** base and DIPEA (0.70 mL, 4.0 mmol) were added after 5 min and the reaction mixture was stirred at room temperature. Work-up: after evaporation of the solution *in vacuo*, the residue was dissolved in dichloromethane (30 mL), extracted with 1 M HCl (**19**: 3 × 10 mL) or with satd. aq. NaHSO₄ (**20**: 5 × 10 mL), and washed successively with satd. aq. NaHCO₃ (25 mL) and brine (25 mL). In the case of **21**, after evaporation of the reaction mixture the residue was triturated with EtOAc (5 mL) and filtered, followed by crystallization. The resulting crude products were purified chromatographically (**19**, **20**) or by crystallization (**21**).

({2-[4-(4-*tert*-Butylbenzoylamino)-2-oxo-1,2-dihydropyrimidin-1-yl]acetyl}-[2-(9H-fluoren-9-ylmethoxycarbonylamino)-ethyl]amino)acetic acid *tert*-butyl ester **19**.—Reaction time: 20 h at room temperature. Chromatography: 1–3% (v/v) methanol in dichloromethane, yield 0.99 g (70%), colourless oil. In a repeated experiment the extractive work-up was omitted and the crude product was triturated with methanol (2 mL) to yield a cleaner product (0.82 g, 58%), amorphous foam; *R*_f 0.49 (CH₂Cl₂–MeOH 95:5) (Found: C, 67.6; H, 6.4; N, 9.7. Calc. for

C₄₀H₄₆N₅O₇: C, 67.8; H, 6.5; N, 9.9%; λ_{\max} (EtOH)/nm 205 (lg ϵ 4.19), 265 (3.94), 288 (3.52), 300 (3.52); $\nu_{\max}/\text{cm}^{-1}$ 3450m, 3150w, 3070w, 2966w, 2936w, 1710s, 1670s, 1629m, 1563m, 1492s, 1408w, 1367s, 1254s, 1156s, 1113s, 851w, 759w, 740w; δ_{H} (500 MHz, rotamers) 1.29 (9 H, s, Bu^t), 1.40 (9 H, s, Bu^t), 1.46 (2 H, s, CH₂), 3.23 (2 H, m, CH₂), 3.44 (2 H, m, CH₂), 3.97 (3 H, s, CH₃O), 4.26 (1 H, m, CH), 4.68/4.90 (2 H, 2 s, CH₂), 6.27 (1 H, s, H-5*), 7.34 (2 H, dd, *J* 7.4 and 7.2, fluorenyl CH), 7.41 (2 H, dd, *J* 7.4 and 7.2, fluorenyl CH), 7.52 (2 H, d, *J* 8.3, 4-Bu^tC₆H₄CO), 7.83 (2 H, d, *J* 7.5, fluorenyl CH), 7.87 (2 H, d, *J* 7.5, fluorenyl CH), 7.98 (2 H, d, *J* 8.3, 4-Bu^tC₆H₄CO), 8.00 (1 H, s, H-6*); *m/z* (ESI) 730 (78%, [M + Na]⁺), 708 (100, [M + H]⁺).

([2-(9*H*-Fluoren-9-ylmethoxycarbonylamino)ethyl]-[2-[6-(4-methoxybenzoylamino)purin-9-yl]acetyl]amino)acetic acid tert-butyl ester **20**.—Reaction time: 1.5 h at room temperature. Chromatography: 0–20% (v/v) EtOAc in methanol, yield 1.20 g (85%), amorphous foam; *R*_f 0.42 (CH₂Cl₂–MeOH 95:5) (Found: C, 64.8; H, 5.4; N, 13.7. Calc. for C₃₈H₃₉N₇O₇: C, 64.7; H, 5.6; N, 13.9%); λ_{\max} (EtOH)/nm 206 (lg ϵ 4.78), 266 (4.41), 278 (4.41), 289 (4.42), 300sh (4.32); $\nu_{\max}/\text{cm}^{-1}$ 3065w, 2980w, 2943w, 1705m, 1670m, 1609m, 1586m, 1513w, 1458m, 1411w, 1252s, 1157m, 845s, 762m, 743m; δ_{H} (500 MHz, rotamers) 1.32 (9 H, s, Bu^t), 2.81 (2 H, s, CH₂), 3.25 (2 H, m, CH₂), 3.65 (2 H, m, CH₂), 3.94 (3 H, s, CH₃O), 4.50 (3 H, m, CH, CH₂), 5.25/5.44 (2 H, 2 s, CH₂), 7.16 (2 H, d, *J* 8.7, anisoyl CH), 7.39 (2 H, dd, *J* 7.4 and 7.2, fluorenyl CH), 7.49 (2 H, dd, *J* 7.4 and 7.2, fluorenyl CH), 7.76 (2 H, d, *J* 7.4, fluorenyl CH), 7.96 (2 H, d, *J* 7.4, fluorenyl CH), 8.13 (2 H, d, *J* 8.6, anisoyl CH), 8.40 (1 H, s, H-8*), 8.70 (1 H, s, H-2*), 11.10 (1 H, br s, NH); *m/z* (ESI) 706 (100%, [M + H]⁺).

([2-[6-Diphenylcarbamoyloxy-2-(isobutrylamino)purin-9-yl]acetyl]-[2-(9*H*-fluoren-9-ylmethoxycarbonylamino)ethyl]amino)acetic acid tert-butyl ester **21**.—Reaction time: 1.5 h at room temperature. The crude product was obtained as described above and was recrystallized from ethanol (750 mL) to give a white powder (1.22 g, 40%), mp 209.0–209.5 °C (decomp.); *R*_f 0.36 (CH₂Cl₂–MeOH 95:5) (Found: C, 66.3; H, 5.5; N, 12.9. Calc. for C₄₇H₄₈N₈O₈: C, 66.2; H, 5.7; N, 13.1%); λ_{\max} (EtOH)/nm 221 (lg ϵ 4.57), 228 (4.53), 256 (4.39), 266sh (4.38), 279sh (4.27), 289sh (4.07), 300 (3.81); $\nu_{\max}/\text{cm}^{-1}$ 3302m, 2979w, 1750m, 1732s, 1704s, 1656s, 1624w, 1591w, 1545m, 1493w, 1450m, 1192s, 1156m, 1055m, 762m, 697w; δ_{H} (500 MHz, rotamers) 1.06 [6 H, d, *J* 6.2, (CH₃)₂CH], 1.35/1.47 (9 H, 2 s, Bu^t), 2.83 [1.3 H, m, (CH₃)₂CH, NH], 3.52 (4 H, m, partly shielded by the water signal, 2 × CH₂), 3.95 (1 H, s, CH), 4.19–4.29 (2 H, m, CH₂), 4.32–4.35 (2 H, m, CH₂), 5.07/5.25 (2 H, 2 × s, guanyl CH₂), 7.22–7.48 (14 H, m, 2 × Ph, fluorenyl CH), 7.64 (2 H, d, *J* 7.3, fluorenyl CH), 7.86 (2 H, dd, *J* 7.9 and 7.9, fluorenyl CH), 8.30 (1 H, s, H-8), 10.57 (1 H, s, NH); *m/z* (ESI) 891 (1%, [M + K]⁺); 875 (5, [M + Na]⁺); 853 (100, [M + H]⁺).

([2-[4-(4-*tert*-Butylbenzoylamino)-2-oxo-1,2-dihydropyrimidin-1-yl]acetyl]-[2-(9*H*-fluoren-9-ylmethoxycarbonylamino)ethyl]amino)acetic acid **23**

To ester **19** (0.40 g, 0.56 mmol) dissolved in dichloromethane (20 mL) was added 1,3-dimethoxybenzene (0.19 mL, 1.45 mmol) followed by TFA (4.0 mL, 52.3 mmol) and the mixture was stirred for 6 h at room temperature. The solution was evaporated *in vacuo*, and the residue was coevaporated with acetonitrile (5×). The residue was triturated under diethyl ether, filtered and recrystallized from methanol (20 mL) to give a white powder (0.27 g, 73%), mp 197.4–199.0 °C (decomp.); *R*_f 0.86 (CH₂Cl₂–MeOH 6:4) (Found: C, 66.3; H, 5.7; N, 10.6. Calc. for C₃₆H₃₈N₅O₇: C, 66.2; H, 5.9; N, 10.7%); λ_{\max} (EtOH)/nm 205 (lg ϵ 4.88), 265 (4.68), 289 (4.25), 300 (4.27); $\nu_{\max}/\text{cm}^{-1}$ 3147w, 3067w, 2964w, 1708s, 1664s, 1610m, 1562m, 1490s, 1409w, 1367m, 1255m, 1113w, 853w, 760w, 741w; δ_{H} (500 MHz,

rotamers) 1.34 (9 H, s, Bu^t), 3.14/3.25 (1 H, 2 m, partly shielded by the water signal, CH₂), 3.38/3.47 (1.4 H, 2 m, partly shielded by the water signal, CH₂), 4.03/4.22 (2 H, 2 s, CH₂), 4.26 (1 H, m, CH), 4.31/4.36 (2 H, 2 d, *J* 6.7, CH₂), 4.70/4.89 (2 H, 2 s, cytosinyl CH₂), 7.27/7.41 (1 H, 2 d, *J* 7.5, H-5*), 7.33 (3 H, dd, *J* 7.5 and 6.8, fluorenyl CH, NH), 7.42 (2 H, d, *J* 6.8, fluorenyl CH), 7.53 (2 H, d, *J* 8.4, 4-Bu^tC₆H₄CO), 7.69 (2 H, dd, *J* 7.5 and 6.8, fluorenyl CH), 7.89 (2 H, d, *J* 7.5, fluorenyl CH), 7.94 (1 H, d, *J* 7.3, H-6*), 7.98 (2 H, dd, *J* 2.7, 8.4, 4-Bu^tC₆H₄CO), 11.20 (1 H, br s, NH[#]), 12.80 (1 H, br s, OH[#]); *m/z* (ESI) 652 (100%, [M + H]⁺).

([2-(9*H*-Fluoren-9-ylmethoxycarbonylamino)ethyl]-[2-[6-(4-methoxybenzoylamino)purin-9-yl]acetyl]amino)acetic acid **24**

To ester **20** (0.92 g, 1.30 mmol) dissolved in dichloromethane (20 mL) were added 1,3-dimethoxybenzene (0.23 mL, 1.82 mmol) and TFA (15.0 mL, 196.0 mmol) and the mixture was stirred for 6 h at room temperature. The solution was evaporated *in vacuo*, and the residue was coevaporated with acetonitrile (5×). The residue was dissolved in methanol (1 mL), diethyl ether (4.5 mL) was added, and the mixture was stored at 4 °C overnight. The resulting gum was triturated with diethyl ether, filtered and recrystallized from methanol (80 mL) to afford a white powder (0.59 g, 70%), mp 160.8–163.9 °C; *R*_f 0.76 (CH₂Cl₂–MeOH 6:4) (Found: C, 62.8; H, 4.65; N, 14.9. Calc. for C₃₄H₃₁N₇O₇: C, 62.9; H, 4.8; N, 15.1%); λ_{\max} (EtOH)/nm 206 (lg ϵ 4.74), 266 (4.43), 278 (4.43), 289 (4.44), 299sh (4.35); $\nu_{\max}/\text{cm}^{-1}$ 3440m, 3222w, 3102w, 3069w, 2978w, 2946w, 1713m, 1693w, 1647m, 1603m, 1582w, 1525m, 1500m, 1411w, 1252s, 1178m, 762m; δ_{H} (500 MHz, rotamers) 3.15/3.59 (2 H, 2 m, CH₂), 3.18 (2 H, s, CH₂), 3.86 (3 H, s, CH₃O), 4.03/4.10 (2 H, 2 m, CH₂), 4.22 (1 H, m, CH), 4.30/4.39 (2 H, 2 m, CH₂), 5.20/5.37 (2 H, 2 s, adenylyl CH₂), 7.08 (2 H, d, *J* 8.7, anisoyl CH), 7.29/7.42 (1 H, 2 br t, NH), 7.32 (2 H, dd, *J* 7.4 and 7.3, fluorenyl CH), 7.41 (2 H, dd, *J* 7.4 and 7.3, fluorenyl CH), 7.70 (2 H, d, *J* 7.4, fluorenyl CH), 7.88 (2 H, d, *J* 7.4, fluorenyl CH), 8.06 (2 H, d, *J* 8.7, anisoyl CH), 8.33 (1 H, s, H-8*), 8.62/8.67 (1 H, 2 s, H-2*); *m/z* (ESI) 650 (100%, [M + H]⁺).

([2-(9*H*-Fluoren-9-ylmethoxycarbonylamino)ethyl]-[2-(2-isobutrylamino-6-oxo-1,6-dihydropurin-9-yl)acetyl]amino)acetic acid **25**

To ester **21** (0.68 g, 0.79 mmol) suspended in dichloromethane (20 mL), was added 1,3-dimethoxybenzene (0.125 mL, 0.95 mmol) followed by TFA (7.34 mL, 95.3 mmol) and the mixture was stirred for 6 h at room temperature. The solution was evaporated *in vacuo*, and the residue was coevaporated with EtOAc (4×). The solid residue was triturated with EtOAc, and filtered (0.48 g, quant.), mp 202.0–206.0 °C and then recrystallized from ethanol (40 mL) to furnish a white powder (0.21 g, 45%), mp 208.8–210.6 °C (decomp.); from the mother liquor a further crop was obtained (0.026 g, 5%), mp 206.0–208.1 °C. Overall yield of the recrystallized product: 0.236 g, 50%. This reaction was repeated on a 3.0 mmol scale and afforded a quantitative yield of the crude acid **25** (1.80 g), mp 202.0–206.0 °C; *R*_f 0.16 (MeCN–MeOH–AcOH 8:1:1) (Found: C, 59.7; H, 5.3; N, 16.1. Calc. for C₃₀H₃₁N₇O₇: C, 59.9; H, 5.2; N, 16.3%); λ_{\max} (EtOH)/nm 205 (lg ϵ 4.79), 221sh (4.24), 256sh (4.45), 262 (4.48), 278sh (4.29), 289sh (4.13), 300 (4.03); $\nu_{\max}/\text{cm}^{-1}$ 3350w, 3131w, 3067w, 2965w, 2940w, 1693s, 1674m, 1610m, 1570m, 1542m, 1485w, 1411m, 1250m, 1154w, 757w, 743w; δ_{H} (500 MHz, rotamers) 1.11 [6 H, d, *J* 6.6, (CH₃)₂CH], 2.75 [1 H, pseudoquintet, *J* 6.6, (CH₃)₂CH], 3.13 (1 H, m), 3.35 (1.8 H, m, partly shielded by the water signal) and 3.49 (1.2 H, m, 2 × CH₂), 4.02/4.29 (2 H, 2 s, CH₂), 4.25 (1 H, m, CH), 4.32/4.38 (2 H, 2 d, *J* 5.8, 6.5, CH₂), 4.97/5.13 (2 H, 2 s, guanyl CH₂), 7.26/7.46 (1 H, 2 br t, NH*), 7.33 (2 H, m, fluorenyl CH), 7.41 (2 H, dd, *J* 7.2 and 7.2, fluorenyl CH), 7.68 (2 H, dd, *J* 7.5, 7.2, fluorenyl CH), 7.82 (1 H, s, H-8), 7.88 (2 H, d, *J* 7.5, fluorenyl

CH), 11.59/11.65 (1 H, 2 s, NH*), 12.07 (1 H, s, NH)*, 12.50 (1 H, br s, OH*); *m/z* (APCI) 602 (100%, [M + H]⁺).

Acknowledgements

This research has been supported by grants MKM 626, OTKA T 22551 and OTKA W 15521 (purchase of ChemProtect 1.0).

References

- 1 P. E. Nielsen, *Annu. Rev. Biophys. Biomol. Struct.*, 1995, **24**, 167.
- 2 P. E. Nielsen and G. Haaime, *Chem. Soc. Rev.*, 1997, **26**, 73.
- 3 B. Hyrup and P. E. Nielsen, *Bioorg. Med. Chem.*, 1996, **4**, 5.
- 4 E. Uhlmann, A. Peyman, G. Breipohl and D. W. Will, *Angew. Chem., Int. Ed.*, 1998, **37**, 2796.
- 5 K. L. Dueholm, M. Egholm, C. Behrens, L. Christensen, H. F. Hansen, T. Vulpus, K. H. Petersen, R. H. Berg, P. E. Nielsen and O. Buchardt, *J. Org. Chem.*, 1994, **59**, 5767.
- 6 L. Christensen, R. Fitzpatrick, B. Gildea, K. H. Petersen, H. F. Hansen, T. Koch, M. Egholm, O. Buchardt, P. E. Nielsen, J. Coull and R. H. Berg, *J. Pept. Sci.*, 1995, **1**, 175.
- 7 D. W. Will, D. Langner, J. Knolle and E. Uhlmann, *Tetrahedron*, 1995, **51**, 12069.
- 8 G. Breipohl, D. W. Will, A. Peyman and E. Uhlmann, *Tetrahedron*, 1997, **53**, 14671.
- 9 S. A. Thomson, J. A. Josey, R. Cadilla, M. D. Gaul, C. F. Hassman, M. J. Luzzio, A. J. Pipe, K. L. Reed, D. J. Ricca, R. W. Wiethe and S. A. Noble, *Tetrahedron*, 1995, **51**, 6179.
- 10 G. Breipohl, J. Knolle, D. Langner, G. Omalley and E. Uhlmann, *Bioorg. Med. Chem. Lett.*, 1996, **6**, 665.
- 11 L. A. Carpino, *Acc. Chem. Res.*, 1987, **20**, 401.
- 12 P. Kocienski, *Protecting Groups*, Georg Thieme Verlag, Stuttgart, 1994.
- 13 E. Sonveaux, in *Protocols for Oligonucleotide Conjugates, Methods in Molecular Biology*, ed. S. Agrawal, Humana Press, Totowa, 1994, vol. 26, pp. 1–71.
- 14 Hanessian Laboratory, *ChemProtect 1.0*, Department of Chemistry, Université de Montréal, Montréal, 1995.
- 15 L. Kovács, Z. Timár and B. Penke, presented at the *13th International Round Table. Nucleosides, Nucleotides and their Biological Applications*, Montpellier, France, 1998, Poster 78.
- 16 F. P. Clausen and J. Juhl-Christensen, *Org. Prep. Proced. Int.*, 1993, **25**, 375.
- 17 T. Kamimura, M. Tsuchiya, K. Koura, M. Sekine and T. Hata, *Tetrahedron Lett.*, 1983, **24**, 2775.
- 18 M. J. Robins, R. Zou, F. Hansske, D. Madej and D. L. J. Tyrrell, *Nucleosides, Nucleotides*, 1989, **8**, 725.
- 19 M. J. Robins, R. Zou, Z. Guo and S. F. Wnuk, *J. Org. Chem.*, 1996, **61**, 9207.
- 20 T. F. Jenny, K. C. Schneider and S. A. Benner, *Nucleosides, Nucleotides*, 1992, **11**, 1257.
- 21 H. Köster, K. Kulikowski, T. Liese, W. Heinkens and V. Kohli, *Tetrahedron*, 1981, **37**, 363.
- 22 R. Zou and M. J. Robins, *Can. J. Chem.*, 1987, **65**, 1436.
- 23 H. R. Kricheldorf and J. Kaschig, *Justus Liebigs Ann. Chem.*, 1976, **882**.
- 24 D. L. Hughes, *Org. React.*, 1992, **42**, 335.
- 25 D. L. Hughes, *Org. Prep. Proced. Int.*, 1996, **28**, 127.
- 26 S. Trippett and D. M. Walker, *J. Chem. Soc.*, 1961, 2130.
- 27 H. A. Brune, M. Falck, R. Hemmer, G. Schmidtberg and H. G. Alt, *Chem. Ber.*, 1984, **117**, 2791.
- 28 M. von Itzstein and M. Mocerino, *Synth. Commun.*, 1990, **20**, 2049.
- 29 Y. El-Kattan, G. Gosselin and J. L. Imbach, *J. Chem. Soc., Perkin Trans. 1*, 1994, 1289.
- 30 T. W. Greene and P. G. M. Wuts, *Protective Groups in Organic Synthesis*, John Wiley and Sons, New York, 2nd edn., 1991, p. 246.
- 31 A. Albert, in *Physical and Physicochemical Aids in Characterization and in Determination of Structure, Synthetic Procedures in Nucleic Acid Chemistry*, ed. W. W. Zorbach and R. S. Tipson, John Wiley and Sons, New York, 1973, vol. 2, pp. 47–123.
- 32 P. K. Bridson, G. Richmond and F. Yeh, *Synth. Commun.*, 1990, **20**, 2459.
- 33 L. Müller, *J. Am. Chem. Soc.*, 1979, **101**, 4481.
- 34 A. Bax, R. H. Griffey and B. L. Hawkins, *J. Magn. Reson.*, 1983, **55**, 301.
- 35 A. Bax and M. F. Summers, *J. Am. Chem. Soc.*, 1986, **108**, 2093.
- 36 A. Bax and D. Marion, *J. Magn. Reson.*, 1988, **78**, 186.
- 37 J. Kjellberg and N.-G. Johansson, *Tetrahedron*, 1986, **42**, 6541.
- 38 J. Boryski, *J. Chem. Soc., Perkin Trans. 2*, 1997, 649.
- 39 G. Lowe and T. Vilaivan, *J. Chem. Soc., Perkin Trans. 1*, 1997, 539.
- 40 D. D. Perrin and W. L. F. Armarego, *Purification of Laboratory Chemicals*, Pergamon Press, Oxford, 3rd edn., 1988.

Paper a907832k