

# Thermal reaction of diastereomeric benzocyclobutenols. Evidence for reversible opening of 1,2-dihydrobenzocyclobutenols to hydroxy-*o*-xylylenes

Etsuya Kawata, Masaichi Saito and Michikazu Yoshioka \*

Department of Chemistry, Faculty of Science, Saitama University, Shimo-okubo, Urawa, Saitama 338-8570, Japan


Received (in Cambridge, UK) 18th October 1999, Accepted 21st January 2000

Thermolysis of 1,2-dihydrobenzocyclobutenols **2a,b** or **3a–c** without solvent at 120 °C gave a 1 : 1 mixture of **2a–c** and **3a–c** together with 3-hydroxy-2,2-dimethyl-1-(*o*-methylphenyl)alkan-1-ones **1a–c** and *o*-methylisobutyrophenone **5**. Thermolysis of **2a** or **3a** under the same conditions but at 150 °C gave only **5**. Thermolysis of **2c** or **3c** in benzene-*d*<sub>6</sub> at 120 °C resulted in clean interconversion between **2c** and **3c**. Thermolysis of 1,2-dihydrobenzocyclobutenols **2d–f** or **3d–f** without solvent at 150 °C gave a mixture of **2d–f** and **3d–f** together with 3-acetoxy-2,2-dimethyl-1-(*o*-methylphenyl)alkan-1-ones **1d–f** and benzyl ketones **9a–c**.

It is well known that benzocyclobutenes undergo an electrocyclic ring opening of the four-membered ring to generate *o*-xylylenes. The *o*-xylylenes are so reactive that they react with various dienophiles to give [4 + 2] cycloadducts<sup>1</sup> or undergo dimerization.<sup>2</sup> The inter- and intramolecular cycloaddition reactions of *o*-xylylenes have been used in the synthesis of polycyclic ring systems.<sup>3</sup> Thermolysis of benzocyclobutenols affords hydroxy-*o*-xylylenes,<sup>4</sup> the geometry of which has been investigated by the analysis of the adduct with maleic anhydride or *N*-phenylmaleimide.<sup>4a</sup> The benzocyclobutenol preferentially opens to produce the (*E*)-dienol. Although the reaction of (*E*)-dienols in the presence of dienophiles has been widely studied,<sup>5</sup> there are few reports on recyclization of dienols generated from benzocyclobutenols. Sammes and co-workers reported that heating optically active 1-methyl-1,2-dihydrobenzocyclobutenol at 110 °C readily gave 2-methylacetophenone because the intermediary (*E*)-dienol underwent proton transfer faster than recyclization to produce acetophenone.<sup>6</sup> We report here the thermal interconversion of diastereomeric 1,2-dihydrobenzocyclobutenols **2** and **3**, which are prepared by photocyclization of 3-hydroxy- and 3-acetoxy-2,2-dimethyl-1-(*o*-methylphenyl)alkan-1-ones **1**.

## Results and discussion


Irradiation of the hydroxy ketone **1a** in methanol with Pyrex-filtered light gave two diastereomeric 1,2-dihydrobenzocyclobutenols **2a** and **3a** together with *trans*- and *cis*-cyclopropane-1,2-diols **4** (Scheme 1).<sup>6</sup> The 1,2-dihydrobenzocyclobutenols **2a** and **3a** could be isolated by column chromatography on silica. The configuration of **2a** was determined by X-ray crystallographic analysis to be (3*S*\*,1'*S*\*), so that **3a** had the (3*S*\*,1'*R*\*) configuration. The <sup>1</sup>H NMR spectrum of **2a** showed peaks due to two methylene protons of the four-membered ring as an AB quartet at δ 2.96 and 3.60 and two methyl singlets on C-2 at δ 0.78 and 1.13, whereas that of **3a** showed two methylene protons at δ 3.03 and 3.55 and two methyl singlets at δ 0.71 and 1.25. The two methylene signals of **2a** were further apart than those of **3a** and the two methyl singlets of **2a** were closer than those of **3a**. Irradiation of hydroxy ketones **1b,c** under the same conditions also gave two diastereomeric 1,2-dihydrobenzocyclobutenols **2b,c** and **3b,c**, the configurations of which could be



Scheme 1 Condition: i, hv.

assigned by comparison of their <sup>1</sup>H NMR spectra to those of **2a** and **3a**. Irradiation of acetoxy ketones **1d–f** under the same conditions also gave two diastereomeric benzocyclobutenols **2d–f** and **3d–f** in 95% yield. The configurations of these compounds were determined by the X-ray crystallographic analyses of **3d**, **2e**<sup>7</sup> and **2f**. Results of the photochemical reaction of hydroxy and acetoxy ketones **1a–f** are given in Table 1.

When the (3*S*\*,1'*S*\*)-1,2-dihydrobenzocyclobutenol **2a** was heated at 120 °C in a sealed glass tube and the progress of the reaction was monitored by <sup>1</sup>H NMR, the gradual disappearance of **2a** was observed together with the gradual formation of the (3*S*\*,1'*R*\*)-1,2-dihydrobenzocyclobutenol **3a** as well as with the formation of small amounts of the hydroxy ketone **1a** and *o*-methylisobutyrophenone **5** (Scheme 2). After 20 h, a thermal equilibrium between **2a** and **3a** was established, where the **2a** : **3a** ratio was 1 : 1. The yields of the mixture of **2a** and **3a**, the hydroxy ketone **1a** and *o*-methylisobutyrophenone **5** were 93, 2 and 5%, respectively. Heating the (3*S*\*,1'*R*\*)-1,2-dihydrobenzocyclobutenol **3a** under the same conditions for 20 h also gave the 1 : 1 mixture of **2a** and **3a** together with small amounts of **1a** and **5**. However, when **2a** or **3a** was heated under the same conditions but at 150 °C for 10 h, only 2-methylisobutyrophenone **5** was obtained. At 150 °C, the 1,2-dihydrobenzocyclobutenols **2a** and **3a** were converted completely into



Scheme 2 Conditions: i, 120 °C; ii, 150 °C.

the hydroxy ketone **1a** which further underwent retro-aldol cleavage to give **5**. The 1,2-dihydrobenzocyclobutenols **2b** and **3b** also underwent thermal interconversion at 120 °C. However, since the 1,2-dihydrobenzocyclobutenol **2c** has a melting point of 136 °C, it remained unchanged on heating at 120 °C without solvent. On the other hand, when a dilute benzene-*d*<sub>6</sub> solution of **2c** in an NMR tube was heated at 120 °C, the conversion into the diastereomeric isomer **3c** occurred cleanly without formation of either the hydroxy ketone **1c** or 2-methylisobutyrophenone **5**, though a long time was required to reach a thermal equilibrium between **2c** and **3c**. The compound **3c**, having a melting point below 120 °C, was converted into **2c** on heating at 120 °C without solvent. In this case, a large amount of the hydroxy ketone **1c** was formed after heating for 20 h. However, **3c** was converted cleanly into a 1:1 mixture of **2c** and **3c** on

Table 1 Photochemical reaction of *o*-tolyl ketones **1a–f**

Ketone <b>1</b>	Irradiation time/h <sup>a</sup>	Conversion (%) <sup>b</sup>	Yield (%) <sup>c</sup>		
			( <b>2</b> + <b>3</b> ) <sup>d</sup>	<b>4</b>	<b>2</b> : <b>3</b> Ratio <sup>e</sup>
<b>a</b>	2	66	17	43	4:1
<b>b</b>	4	74	47	—	3:2
<b>c</b>	5	50	37	—	4:3
<b>d</b>	15	97	95	—	3:2
<b>e</b>	15	100	95	—	6:5
<b>f</b>	15	94	95	—	1:1

<sup>a</sup> A solution of the ketone (600 mg) in methanol (160 cm<sup>3</sup>) was irradiated with a 100 W high-pressure mercury lamp through a Pyrex filter. Ketones **1a–c** were irradiated under ice-cooling. Ketones **1d–f** were irradiated at room temperature. <sup>b</sup> Based on the amount of consumed starting material. <sup>c</sup> Isolated yield based on converted starting material. <sup>d</sup> Sum yield of **2** and **3**. These compounds could be isolated by repeated chromatography. <sup>e</sup> Determined by <sup>1</sup>H NMR on the fractions of the mixture after chromatography.

Table 2 Thermal reaction of benzocyclobutenols **2a–f** and **3a–f**

Benzocyclobutenol	Solvent	Temp/°C	Time/h	Yield (%) <sup>a</sup>				<b>2</b> : <b>3</b> Ratio <sup>b</sup>
				<b>1</b>	( <b>2</b> + <b>3</b> )	<b>5</b>	<b>9</b>	
<b>2a</b>	None	120	20	2	93	5	—	1:1
<b>3a</b>	None	120	20	5	84	3	—	1:1
<b>2a</b>	None	150	10	—	—	88	—	—
<b>3a</b>	None	150	10	—	—	88	—	—
<b>2b</b>	None	120	20	7	71	—	—	1:1
<b>3b</b>	None	120	20	6	71	—	—	1:1
<b>2c</b>	C <sub>6</sub> D <sub>6</sub>	120	80	—	100	—	—	1:1
<b>3c</b>	None	120	20	63	16	6	—	1:1
<b>3c</b>	C <sub>6</sub> D <sub>6</sub>	120	80	—	100	—	—	1:1
<b>2d</b>	None	150	20	20	72	—	8	7:4
<b>3d</b>	None	150	20	9	68	—	11	7:4
<b>2e</b>	None	150	20	7	62	—	11	3:2
<b>3e</b>	None	150	20	18	60	—	11	3:2
<b>2f</b>	None	150	20	11	66	—	2	2:3
<b>3f</b>	None	150	20	20	66	—	7	2:3


<sup>a</sup> Isolated yield. <sup>b</sup> Estimated by <sup>1</sup>H NMR on the fractions after chromatography.

heating in benzene-*d*<sub>6</sub> at 120 °C. The acetoxy-substituted 1,2-dihydrobenzocyclobutenols **2d–f** and **3d–f** were also inter-converted into each other on heating at 150 °C without solvent, together with the formation of **1d–f** and benzyl ketones **9a–c**. After 20 h, a thermal equilibrium between **2d–f** and **3d–f** was established. Results of the thermal reaction of **2** and **3** are given in Table 2.

As already mentioned, the 1,2-dihydrobenzocyclobutenol undergoes selective thermal opening of the cyclobutene ring to form the (*E*)-dienol. The thermal interconversion of the diastereomeric 1,2-dihydrobenzocyclobutenols **2** and **3** probably proceeds via the (*E*)-dienol intermediate **6** because the (*E*)-dienol species is sufficiently long-lived to be able to cyclize to give epimers.<sup>8</sup> As shown in Table 2, the major reaction of the diastereomeric 1,2-dihydrobenzocyclobutenols **2** and **3** was the interconversion of diastereomers. Since the (*E*)-dienols formed from **2** and **3** are very congested because of the orientation of the bulky quarternary alkyl group, the possibility that β-hydroxy- and β-acetoxy ketones **1a–f** are formed from the (*Z*)-dienol cannot be ruled out. However, since the thermolysis of a dilute benzene-*d*<sub>6</sub> solution of **2c** or **3c** cleanly gave a 1:1 mixture of **2c** and **3c**, the 1,2-dihydrobenzocyclobutenols **2** and **3** would open selectively to the (*E*)-dienol because the (*Z*)-dienol is very short-lived and undergoes a rapid 1,5-sigmatropic hydrogen shift to give the ketone **1**.<sup>9</sup> The resulting (*E*)-dienol undergoes recyclization to give 1,2-dihydrobenzocyclobutenols **2** and **3** along with intermolecular hydrogen transfers to give hydroxy and acetoxy ketones **1**.<sup>10</sup> The lower yields of hydroxy ketones **1a,b** compared with **1c** in the thermolysis of **3a–c** without solvent at 120 °C may be due to a bulkier substituent in **3a,b** than in **3c**. The bulky substituent may prevent the approach of the initially formed (*E*)-dienols. The (*E*)-dienols formed from 1,2-dihydrobenzocyclobutenols having a relatively small substituent react with various dienophiles to give cycloadducts.<sup>5</sup> However, when the 1,2-dihydrobenzocyclobutenol **2a** or **3a** was heated at 120 °C in the presence of dimethyl acetylenedicarbonylate or maleic anhydride, no adduct of the (*E*)-dienol (**6** (*R*<sup>2</sup> = Pr<sup>i</sup>)) with the dienophile could be detected but the interconversion between **2a** and **3a** was observed, perhaps due to steric congestion of **6** (*R*<sup>2</sup> = Pr<sup>i</sup>) preventing the access of the dienophile.

The **2a–c**:**3a–c** ratios in the photochemical reaction of **1a–c** increased with increasing size of *R*<sup>2</sup>, whereas these ratios were 1:1 in the thermal reactions of **2a–c** and **3a–c** regardless of the size of *R*<sup>2</sup> (Table 1 and 2). It is well known that benzocyclobutenols are prepared from the (*E*)-dienols generated by the irradiation of *o*-tolyl ketones.<sup>11</sup> The photochemically generated (*E*)-dienols from **1a–c** in methanol must be solvated by meth-

anol. The solvated (*E*)-dienols may cyclize to **2** and **3** in a ratio that depends on the size of R<sup>2</sup>.<sup>12</sup> The **2d-f**:**3d-f** ratios in the thermal reaction of **2d-f** or **3d-f** were also different from those in the photochemical reaction of **1d-f**. In both thermal and photochemical reactions, the **2d-f**:**3d-f** ratios increased with increasing size of R<sup>2</sup>. Finally, the benzyl ketone **9** may be formed by a process involving homolytic cleavage between the aryl carbon and C-1 followed by hydrogen transfer (Scheme 3).<sup>13</sup>



Scheme 3 Condition: i, 150 °C.

## Experimental

Mps are uncorrected and bps are oven temperatures in Kugelrohr distillation. IR spectra were recorded on a Hitachi 270-50 spectrometer for solutions in CCl<sub>4</sub>. <sup>1</sup>H NMR spectra were obtained with a Bruker AC 200, a Bruker AC 300-P or a Bruker AM 400 spectrometer with CDCl<sub>3</sub> as a solvent. Tetramethylsilane was used as an internal standard and *J* values are given in Hz. <sup>13</sup>C NMR spectra were measured on a Bruker AC 200 or a Bruker AC 300-P spectrometer with CDCl<sub>3</sub> as a solvent. An Ushio 100 W high-pressure mercury lamp was used as an irradiation source. Starting compounds **1a-c** were prepared by the condensation of *o*-methylisobutyrophenone with the aldehyde according to previously described methods.<sup>14</sup> Compounds **1d-f** were prepared by refluxing **1a-c** in acetic anhydride in the presence of a catalytic amount of hydrochloric acid.

### General procedure for the photolysis of **1a-c**

A solution of **1** (600 mg) in methanol (160 cm<sup>3</sup>) was irradiated with a 100 W high-pressure mercury lamp through a Pyrex filter under argon for 2–15 h (see Table 1). The solvent was removed under reduced pressure, and the residue was chromatographed on silica gel [hexane–ethyl acetate (6:1 to 22:1)] to give 1,2-dihydrobenzocyclobutenols **2** and **3** and the cyclopropane-1,2-diol **4**. The physical properties of **4** have already been described in a previous paper.<sup>6</sup>

**(3*S*\*,1'*S*\*)-2-(1'-Hydroxy-1',2'-dihydrobenzocyclobuten-1'-yl)-2,4-dimethylpentan-3-ol 2a.** Mp 67–68 °C (from hexane) (Found: C, 77.1; H, 9.3. C<sub>15</sub>H<sub>22</sub>O<sub>2</sub> requires C, 76.9; H, 9.5%); ν<sub>max</sub>/cm<sup>-1</sup> 3400br (OH); δ<sub>H</sub> (300 MHz) 0.78 (3 H, s) and 1.13 (3 H, s) (1-H<sub>3</sub> and 2-Me), 0.99 (3 H, d, *J* 7) and 1.08 (3 H, d, *J* 7) (CHMe<sub>2</sub>), 2.00 (1 H, m, CHMe<sub>2</sub>), 2.96 (1 H) and 3.60 (1 H) (AB-pair, *J* 15, 2'-H<sub>2</sub>), 2.97 (1 H, br s) and 3.90 (2 H, m) (2 × OH and 3-H) and 7.1–7.3 (4 H, m, ArH); δ<sub>C</sub> (50 MHz) 16.3 (q), 16.9 (q), 21.7 (q) and 23.4 (q) (4 × Me), 29.3 (d, C-4), 42.4 (s, C-2), 43.7 (t, C-2'), 80.8 (d, C-3), 89.1 (s, C-1'), 121.9 (d), 123.5 (d), 127.1 (d), 129.2 (d), 142.5 (s) and 148.7 (s) (ArC).

**(3*S*\*,1'*R*\*)-2-(1'-Hydroxy-1',2'-dihydrobenzocyclobuten-1'-yl)-2,4-dimethylpentan-3-ol 3a.** Mp 74–75 °C (from hexane) (Found: C, 77.1; H, 9.3. C<sub>15</sub>H<sub>22</sub>O<sub>2</sub> requires C, 76.9; H, 9.5%); ν<sub>max</sub>/cm<sup>-1</sup> 3450br (OH); δ<sub>H</sub> (300 MHz) 0.71 (3 H, s) and 1.25

(3 H, s) (1-H<sub>3</sub> and 2-Me), 0.99 (3 H, d, *J* 7) and 1.06 (3 H, d, *J* 7) (CHMe<sub>2</sub>), 2.00 (1 H, m, CHMe<sub>2</sub>), 3.03 (1 H) and 3.55 (1 H) (AB-pair, *J* 15, 2'-H<sub>2</sub>), 3.01 (1 H, br s, OH), 3.78 (1 H, m, 3-H), 4.10 (1 H, br s, OH) and 7.1–7.3 (4 H, m, ArH); δ<sub>C</sub> (50 MHz) 16.4 (q), 17.3 (q), 21.4 (q) and 23.2 (q) (4 × Me), 29.5 (d, C-4), 42.5 (s, C-2), 43.1 (t, C-2'), 80.3 (d, C-3), 89.6 (s, C-1'), 121.9 (d), 123.5 (d), 127.1 (d), 129.2 (d), 142.5 (s) and 148.3 (s) (ArC).

**(3*S*\*,1'*S*\*)-2-(1'-Hydroxy-1',2'-dihydrobenzocyclobuten-1'-yl)-2-methylpentan-3-ol 2b.** Mp 96–97 °C (from hexane) (Found: C, 76.4; H, 9.2. C<sub>14</sub>H<sub>20</sub>O<sub>2</sub> requires C, 76.3; H, 9.2%); ν<sub>max</sub>/cm<sup>-1</sup> 3300br (OH); δ<sub>H</sub> (200 MHz) 0.75 (3 H, s) and 0.92 (3 H, s) (1-H<sub>3</sub> and 2-Me), 0.99 (3 H, t, *J* 7, 5-H<sub>3</sub>), 1.3–1.6 (2 H, m, 4-H<sub>2</sub>), 2.88 (1 H) and 3.53 (1 H) (AB-pair, *J* 15, 2'-H<sub>2</sub>), 3.74 (1 H, m, 3-H), 3.14 (1 H, br s) and 4.02 (1 H, br s) (2 × OH) and 7.1–7.3 (4 H, m, ArH); δ<sub>C</sub> (50 MHz) 11.3 (q), 16.8 (q) and 21.8 (q) (3 × Me), 24.6 (t, C-4), 41.3 (s, C-2), 43.5 (t, C-2'), 80.1 (d, C-3), 88.0 (s, C-1'), 121.9 (d), 123.2 (d), 126.8 (d), 128.9 (d), 142.4 (s) and 148.6 (s) (ArC).

**(3*S*\*,1'*R*\*)-2-(1'-Hydroxy-1',2'-dihydrobenzocyclobuten-1'-yl)-2-methylpentan-3-ol 3b.** Mp 104–105 °C (from hexane) (Found: C, 76.4; H, 9.2. C<sub>14</sub>H<sub>20</sub>O<sub>2</sub> requires C, 76.3; H, 9.2%); ν<sub>max</sub>/cm<sup>-1</sup> 3300br (OH); δ<sub>H</sub> (200 MHz) 0.69 (3 H, s) and 1.14 (3 H, s) (1-H<sub>3</sub> and 2-Me), 1.05 (3 H, t, *J* 7, 5-H<sub>3</sub>), 1.4–1.8 (2 H, m, 4-H<sub>2</sub>), 3.03 (1 H) and 3.57 (1 H) (AB-pair, *J* 15, 2'-H<sub>2</sub>), 3.65 (1 H, m, 3-H), 4.27 (1 H, br s) and 5.11 (1 H, br s) (2 × OH) and 7.1–7.3 (4 H, m, ArH); δ<sub>C</sub> (50 MHz) 11.1 (q), 16.6 (q) and 21.6 (q) (3 × Me), 24.8 (t, C-4), 41.7 (s, C-2), 42.9 (t, C-2'), 79.3 (d, C-3), 88.8 (s, C-1'), 121.9 (d), 123.5 (d), 127.1 (d), 129.1 (d), 142.4 (s) and 148.4 (s) (ArC).

**(2*S*\*,1'*S*\*)-3-(1'-Hydroxy-1',2'-dihydrobenzocyclobuten-1'-yl)-3-methylbutan-2-ol 2c.** Mp 136–137 °C (from hexane) (Found: C, 75.5; H, 8.7. C<sub>13</sub>H<sub>18</sub>O<sub>2</sub> requires C, 75.7; H, 8.8%); ν<sub>max</sub>/cm<sup>-1</sup> 3400br (OH); δ<sub>H</sub> (300 MHz) 0.62 (3 H, s) and 1.05 (3 H, s) (4-H<sub>3</sub> and 3-Me), 1.22 (3 H, d, *J* 7, 1-H<sub>3</sub>), 2.98 (1 H) and 3.63 (1 H) (AB-pair, *J* 15, 2'-H<sub>2</sub>), 3.69 (1 H, br s) and 3.55 (1 H, br s) (2 × OH), 4.19 (1 H, q, *J* 7, 2-H) and 7.1–7.3 (4 H, m, ArH); δ<sub>C</sub> (75 MHz) 15.5 (q), 18.4 (q) and 22.0 (q) (3 × Me), 41.4 (s, C-3), 43.9 (t, C-2'), 73.9 (d, C-2), 88.4 (s, C-1'), 121.9 (d), 123.5 (d), 127.1 (d), 129.3 (d), 142.4 (s) and 148.5 (s) (ArC).

**(2*S*\*,1'*R*\*)-3-(1'-Hydroxy-1',2'-dihydrobenzocyclobuten-1'-yl)-3-methylbutan-2-ol 3c.** Mp 84–85 °C (from hexane) (Found: C, 75.5; H, 8.7. C<sub>13</sub>H<sub>18</sub>O<sub>2</sub> requires C, 75.7; H, 8.8%); ν<sub>max</sub>/cm<sup>-1</sup> 3450br (OH); δ<sub>H</sub> (300 MHz) 0.62 (3 H, s) and 1.15 (3 H, s) (4-H<sub>3</sub> and 3-Me), 1.16 (3 H, d, *J* 7, 1-H<sub>3</sub>), 3.02 (1 H) and 3.55 (1 H) (AB-pair, *J* 15, 2'-H<sub>2</sub>), 3.85 (1 H, br s) and 4.20 (1 H, br s) (2 × OH), 4.07 (1 H, q, *J* 7) and 7.1–7.3 (4 H, m, ArH); δ<sub>C</sub> (50 MHz) 15.5 (q), 18.4 (q) and 21.7 (q) (3 × Me), 41.5 (s, C-3), 42.8 (t, C-2'), 73.2 (d, C-2), 88.8 (s, C-1'), 121.8 (d), 123.5 (d), 127.1 (d), 129.2 (d), 142.4 (s) and 148.3 (s) (ArC).

**(3*S*\*,1'*S*\*)-3-Acetoxy-2-(1'-hydroxy-1',2'-dihydrobenzocyclobuten-1'-yl)-2,4-dimethylpentane 2d.** Mp 72–73 °C (from hexane) (Found: C, 74.0; H, 8.9. C<sub>17</sub>H<sub>24</sub>O<sub>3</sub> requires C, 73.9; H, 8.8%); ν<sub>max</sub>/cm<sup>-1</sup> 3500br (OH) and 1720 (C=O); δ<sub>H</sub> (200 MHz) 0.90 (3 H, s) and 1.22 (3 H, s) (1-H<sub>3</sub> and 2-Me), 0.96 (3 H, d, *J* 7) and 1.01 (3 H, d, *J* 7) (CHMe<sub>2</sub>), 2.12 (3 H, s, COMe), 2.0–2.2 (1 H, m, CHMe<sub>2</sub>), 3.31 (1 H, br s, OH), 2.83 (1 H) and 3.59 (1 H) (AB-pair, *J* 15, 2'-H<sub>2</sub>), 5.18 (1 H, d, *J* 3, 3-H) and 7.1–7.3 (4 H, m, ArH); δ<sub>C</sub> (50 MHz) 17.7 (q), 18.1 (q), 21.1 (q), 21.3 (q) and 23.3 (q) (5 × Me), 28.7 (d, C-4), 43.2 (s, C-2), 43.4 (t, C-2'), 79.9 (d, C-3), 86.1 (s, C-1'), 122.1 (d), 123.3 (d), 127.0 (d), 129.1 (d), 143.0 (s) and 148.2 (s) (ArC) and 172.0 (s, C=O).

**(3*S*\*,1'*R*\*)-3-Acetoxy-2-(1'-hydroxy-1',2'-dihydrobenzocyclobuten-1'-yl)-2,4-dimethylpentane 3d.** Mp 67–68 °C (from hexane) (Found: C, 74.1; H, 8.8. C<sub>17</sub>H<sub>24</sub>O<sub>3</sub> requires C, 73.9; H,

8.8%);  $\nu_{\max}/\text{cm}^{-1}$  3500br (OH) and 1750 (C=O);  $\delta_{\text{H}}$  (200 MHz) 0.90 (3 H, s) and 1.14 (3 H, s) (1-H<sub>3</sub> and 2-Me), 0.97 (3 H, d, *J* 7) and 0.98 (3 H, d, *J* 7) (CHMe<sub>2</sub>), 2.14 (3 H, s, COMe), 2.1–2.3 (1 H, m, CHMe<sub>2</sub>), 2.67 (1 H, br s, OH), 2.99 (1 H) and 3.66 (1 H) (AB-pair, *J* 15, 2'-H<sub>2</sub>), 5.14 (1 H, d, *J* 2, 3-H) and 7.1–7.3 (4 H, m, ArH);  $\delta_{\text{C}}$  (50 MHz) 17.4 (q), 20.3 (q), 20.6 (q), 21.2 (q) and 23.2 (q) (5 × Me), 29.2 (d, C-4), 43.1 (s, C-2), 43.8 (t, C-2'), 81.0 (d, C-3), 86.9 (s, C-1'), 121.8 (d), 123.4 (d), 127.0 (d), 129.1 (d), 142.4 (s) and 148.7 (s) (ArC) and 172.0 (s, C=O).

**(3*S*\*,1'*R*\*)-3-Acetoxy-2-(1'-hydroxy-1',2'-dihydrobenzocyclobuten-1'-yl)-2-methylpentane 2e.** Mp 83–84 °C (from hexane) (Found: C, 73.3; H, 8.6. C<sub>16</sub>H<sub>22</sub>O<sub>3</sub> requires C, 73.3; H, 8.5%);  $\nu_{\max}/\text{cm}^{-1}$  3500br (OH) and 1720 (C=O);  $\delta_{\text{H}}$  (200 MHz) 0.82 (3 H, s) and 1.19 (3 H, s) (1-H<sub>3</sub> and 2-Me), 0.91 (3 H, t, *J* 7, 5-H<sub>3</sub>), 1.66 (2 H, m, 4-H<sub>2</sub>), 2.11 (3 H, s, COMe), 2.83 (1 H) and 3.58 (1 H) (AB-pair, *J* 15, 2'-H<sub>2</sub>), 3.55 (1H, br s, OH), 5.17 (1 H, dd, *J* 3 and 10, 3-H) and 7.1–7.3 (4 H, m, ArH);  $\delta_{\text{C}}$  (50 MHz) 10.8 (q), 17.7 (q), 21.1 (q) and 21.3 (q) (4 × Me), 22.8 (t, C-4), 42.5 (s, C-2), 43.1 (t, C-2'), 79.2 (d, C-3), 85.6 (s, C-1'), 122.1 (d), 123.2 (d), 126.9 (d), 129.0 (d), 142.8 (s) and 148.1 (s) (ArC) and 172.1 (s, C=O).

**(3*S*\*,1'*S*\*)-3-Acetoxy-2-(1'-hydroxy-1',2'-dihydrobenzocyclobuten-1'-yl)-2-methylpentane 3e.** Mp 72–73 °C (from hexane) (Found: C, 73.4; H, 8.5. C<sub>16</sub>H<sub>22</sub>O<sub>3</sub> requires C, 73.3; H, 8.5%);  $\nu_{\max}/\text{cm}^{-1}$  3500br (OH) and 1750 (C=O);  $\delta_{\text{H}}$  (200 MHz) 0.85 (3 H, s) and 1.08 (3 H, s) (1-H<sub>3</sub> and 2-Me), 0.88 (3 H, t, *J* 7, 5-H<sub>3</sub>), 1.67 (2 H, m, 4-H<sub>2</sub>), 2.11 (3 H, s, COMe), 2.79 (1 H, br s, OH), 2.96 (1 H) and 3.63 (1 H) (AB-pair, *J* 15, 2'-H<sub>2</sub>), 5.21 (1 H, dd, *J* 1 and 3, 3-H) and 7.1–7.3 (4 H, m, ArH);  $\delta_{\text{C}}$  (50 MHz) 10.9 (q), 19.7 (q), 20.3 (q) and 21.2 (q) (4 × Me), 23.7 (t, C-4), 42.5 (s, C-2), 44.2 (t, C-2'), 79.9 (d, C-3), 86.5 (s, C-1'), 122.0 (d), 123.4 (d), 127.0 (d), 129.2 (d), 142.7 (s) and 148.8 (s) (ArC) and 171.2 (s, C=O).

**(2*S*\*,1'*S*\*)-2-Acetoxy-3-(1'-hydroxy-1',2'-dihydrobenzocyclobuten-1'-yl)-3-methylbutane 2f.** Mp 79–80 °C (from hexane) (Found: C, 72.7; H, 8.2. C<sub>15</sub>H<sub>22</sub>O<sub>3</sub> requires C, 72.6; H, 8.1%);  $\nu_{\max}/\text{cm}^{-1}$  3630 (OH) and 1750 (C=O);  $\delta_{\text{H}}$  (200 MHz) 0.89 (3 H, s) and 1.13 (3 H, s) (4-H<sub>3</sub> and 3-Me), 1.25 (3 H, d, *J* 7, 1-H<sub>3</sub>), 2.04 (3 H, s, COMe), 2.88 (1 H) and 3.66 (1 H) (AB-pair, *J* 15, 2'-H<sub>2</sub>), 3.07 (1 H, br s, OH), 5.34 (1 H, q, *J* 7, 2-H) and 7.1–7.3 (4 H, m, ArH);  $\delta_{\text{C}}$  (50 MHz) 15.6 (q), 17.2 (q), 21.0 (q) and 21.6 (q) (4 × Me), 42.2 (s, C-3), 43.7 (t, C-2'), 74.0 (d, C-2), 85.8 (s, C-1'), 122.0 (d), 123.4 (d), 127.1 (d), 129.2 (d), 142.9 (s) and 148.6 (s) (ArC) and 171.0 (s, C=O).

**(2*S*\*,1'*R*\*)-3-Acetoxy-3-(1'-hydroxy-1',2'-dihydrobenzocyclobuten-1'-yl)-3-methylbutane 3f.** Bp 96–98 °C at 0.3 mmHg (Found: C, 72.6; H, 8.2. C<sub>15</sub>H<sub>22</sub>O<sub>3</sub> requires C, 72.6; H, 8.1%);  $\nu_{\max}/\text{cm}^{-1}$  3470br (OH) and 1740 (C=O);  $\delta_{\text{H}}$  (200 MHz) 0.87 (3 H, s) and 1.10 (3 H, s) (4-H<sub>3</sub> and 3-Me), 1.25 (3 H, d, *J* 7, 1-H<sub>3</sub>), 2.06 (3 H, s, COMe), 2.74 (1 H, br s, OH), 2.95 (1 H) and 3.64 (1 H) (AB-pair, *J* 15, 2'-H<sub>2</sub>), 5.22 (1 H, q, *J* 7, 2-H) and 7.1–7.3 (4 H, m, ArH);  $\delta_{\text{C}}$  (50 MHz) 16.0 (q), 19.4 (2q) and 21.4 (q) (4 × Me), 41.8 (s, C-3), 44.1 (t, C-2'), 75.2 (d, C-2), 86.2 (s, C-1'), 122.0 (d), 123.3 (d), 126.9 (d), 129.1 (d), 142.5 (s) and 148.6 (s) (ArC) and 170.3 (s, C=O).

#### Pyrolysis of 1',2'-dihydrobenzocyclobutenols 2a–f and 3a–f

1',2'-Dihydrobenzocyclobutenol 2a–c or 3a–c (100 mg) was sealed in an 8 mm diameter Pyrex tube under argon. The tube was heated at 120 °C for 20 h. The <sup>1</sup>H NMR analysis of the mixture revealed that interconversion occurred between 2a–c and 3a–c. The mixture was chromatographed on silica gel and eluted with a mixture of hexane and ethyl acetate (4:1 to 6:1) to give two isomeric 1,2-dihydrobenzocyclobutenols 2a–c and 3a–c, hydroxy ketone 1a–c and *o*-methylisobutyrophenone 5.

The same treatment of 1,2-dihydrobenzocyclobutenol 2d–f or 3d–f (100 mg) but at 150 °C for 20 h gave acetoxy ketone 1d–f, 1,2-dihydrobenzocyclobutenol 2d–f and 3d–f and 4-acetoxy-3,3-dimethyl-1-phenylbutan-2-one 9a–c. 1,2-Dihydrobenzocyclobutenol 2c or 3c (ca. 10 mg) in benzene-*d*<sub>6</sub> (ca. 0.5 cm<sup>3</sup>) was placed in a 5 mm diameter NMR tube and degassed by freeze–pump–thaw cycles. The tube was heated at 120 °C for 80 h. The <sup>1</sup>H NMR spectrum of the mixture showed only peaks due to a 1:1 mixture of 2c and 3c.

**4-Acetoxy-3,3,5-trimethyl-1-phenylhexan-2-one 9a.** Bp 97–98 °C at 0.4 mmHg (Found: C, 73.6; H, 8.8. C<sub>17</sub>H<sub>24</sub>O<sub>3</sub> requires C, 73.9; H, 8.8%);  $\nu_{\max}/\text{cm}^{-1}$  1740 (ester C=O) and 1720 (C=O);  $\delta_{\text{H}}$  (300 MHz) 0.86 (3 H, d, *J* 7) and 0.89 (3 H, d, *J* 7) (6-H<sub>3</sub> and 5-Me), 1.13 (3 H, s) and 1.22 (3 H, s) (3-Me<sub>2</sub>), 1.90 (1 H, d × sept, *J* 6 and 7, 5-H), 2.09 (3 H, s, COMe), 3.81 (1 H) and 3.88 (1 H) (AB-pair, *J* 16, 1-H<sub>2</sub>), 5.20 (1 H, d, *J* 6, 4-H) and 7.2–7.4 (5 H, m, ArH);  $\delta_{\text{C}}$  (75 MHz) 18.5 (q), 20.1 (q), 20.8 (q), 21.5 (q) and 22.7 (q) (5 × Me), 29.6 (d, C-5), 44.5 (t, C-1), 52.3 (s, C-3), 80.7 (d, C-4), 126.7 (d), 128.4 (d), 129.6 (d) and 134.3 (s) (ArC), 170.7 (s, OC=O) and 210.4 (s, C=O).

**4-Acetoxy-3,3-dimethyl-1-phenylhexan-2-one 9b.** Bp 86–88 °C at 0.3 mmHg (Found: C, 73.2; H, 8.5. C<sub>16</sub>H<sub>22</sub>O<sub>3</sub> requires C, 73.3; H, 8.5%);  $\nu_{\max}/\text{cm}^{-1}$  1740 (ester C=O) and 1720 (C=O);  $\delta_{\text{H}}$  (200 MHz) 0.86 (3 H, t, *J* 7, 6-H<sub>3</sub>), 1.16 (3 H, s) and 1.19 (3 H, s) (3-Me<sub>2</sub>), 1.4–1.5 (2 H, m, 5-H<sub>2</sub>), 2.07 (3 H, s, COMe), 3.82 (2 H, s, 1-H<sub>2</sub>), 5.25 (1 H, dd, *J* 5 and 8, 4-H) and 7.2–7.3 (4 H, m, ArH);  $\delta_{\text{C}}$  (50 MHz) 10.8 (q), 20.3 (q), 21.2 (q) and 23.5 (q) (4 × Me), 20.8 (t, C-5), 44.5 (t, C-1), 52.1 (s, C-3), 78.4 (d, C-4), 126.7 (d), 128.4 (d), 129.6 (d) and 134.4 (s) (ArC), 170.7 (s, ester C=O) and 210.1 (s).

**4-Acetoxy-3,3-dimethyl-1-phenylpentan-2-one 9c.** Bp 96–98 °C at 0.4 mmHg (Found: C, 72.7; H, 8.2. C<sub>15</sub>H<sub>22</sub>O<sub>3</sub> requires C, 72.6; H, 8.1%);  $\nu_{\max}/\text{cm}^{-1}$  1740 (ester C=O) and 1720 (C=O);  $\delta_{\text{H}}$  (200 MHz) 1.16 (3 H, d, *J* 7, 5-H<sub>3</sub>), 1.17 (3 H, s) and 1.19 (3 H, s) (3-Me<sub>2</sub>), 1.97 (3 H, s, COMe), 3.79 (2 H, s, 1-H<sub>2</sub>), 5.26 (1 H, q, *J* 7, 4-H) and 7.1–7.3 (5 H, m, ArH);  $\delta_{\text{C}}$  (50 MHz) 14.9 (q), 19.6 (q), 20.8 (q) and 21.0 (q) (4 × Me), 44.2 (t, C-1), 51.4 (s, C-3), 74.0 (d, C-4), 126.6 (d), 128.3 (d), 129.5 (d) and 134.2 (s) (ArC), 170.1 (s, ester C=O) and 210.1 (s, C=O).

#### Pyrolysis of 2a and 3a in the presence of dienophile

A solution of 100 mg of 2a or 3a and 2 equiv. of dimethyl acetylenedicarboxylate or maleic anhydride in 1 cm<sup>3</sup> of benzene was sealed in a glass tube. The tube was degassed by freeze–pump–thaw cycles and heated at 120 °C for 20 h. The solvent was removed and the residue was fractionated by chromatography on silica gel using a 4:1 mixture of hexane and ethyl acetate. The <sup>1</sup>H NMR spectrum of each of the fractions revealed that 2a and 3a were interconverted, while no adduct of the dienol arising from 2a or 3a with dienophile was detected.

#### Crystallographic analysis of 2a, 2f and 3d

Data were collected on a MAC Science DIP3000 diffractometer with Mo-K $\alpha$  radiation ( $\lambda = 0.71073$  Å) at 298 K and the structure was solved by direct methods. Crystal data for 2a: C<sub>15</sub>H<sub>22</sub>O<sub>2</sub>, *M* = 234.37, orthorhombic, *a* = 7.8840(5), *b* = 12.514(1), *c* = 14.297(1) Å, *V* = 1410.6(2) Å<sup>3</sup>, *Z* = 4, space group *P*2<sub>1</sub>2<sub>1</sub>,  $\mu = 0.067$  mm<sup>-1</sup>. The crystal used had dimensions of 0.5 × 0.5 × 0.4 mm. The final cycle of full-matrix least-squares refinement was based on 1747 observed reflections [*I* > 3 $\sigma$ (*I*)] and 242 variable parameters with *R*(*R*<sub>w</sub>) = 0.049 (0.062). Crystal data for 2f: C<sub>15</sub>H<sub>22</sub>O<sub>3</sub>, *M* = 248.32, monoclinic, *a* = 11.308(2), *b* = 9.460(1), *c* = 13.187(1) Å, *V* = 1402.4 (3) Å<sup>3</sup>, *Z* = 4, space group *P*2<sub>1</sub>/*n*,  $\mu = 0.075$  mm<sup>-1</sup>. The crystal used had dimensions of 0.3 × 0.3 × 0.28 mm. The final cycle of full-matrix least-squares refinement was based on 1920 observed

reflections [ $I > 3\sigma(I)$ ] and 243 variable parameters with  $R(R_w) = 0.053$  (0.067). Crystal data for **3d**:  $C_{17}H_{24}O_3$ ,  $M = 276.38$ , monoclinic,  $a = 10.106(1)$ ,  $b = 25.218(2)$ ,  $c = 12.298(2)$  Å,  $V = 3134.2(2)$  Å<sup>3</sup>,  $Z = 8$ , space group  $P2_1/n$ ,  $\mu = 0.074$  mm<sup>-1</sup>. The crystal used had dimensions of  $0.15 \times 0.15 \times 0.15$  mm. The final cycle of full-matrix least-squares refinement was based on 2941 observed reflections [ $I > 3\sigma(I)$ ] and 415 variable parameters with  $R(R_w) = 0.054$  (0.071).

CCDC reference number 207/396. See <http://www.rsc.org/suppdata/p1/a9/a908326j> for crystallographic files in .cif format.

## References

- 1 J. L. Charlton and M. M. Alauddin, *Tetrahedron*, 1987, **43**, 2873.
- 2 M. P. Cava and A. A. Deana, *J. Am. Chem. Soc.*, 1959, **81**, 4266; L. A. Errede, *J. Am. Chem. Soc.*, 1961, **83**, 949.
- 3 W. Oppolzer, *Synthesis*, 1978, 793.
- 4 (a) B. J. Arnold, P. G. Sammes and T. W. Wallace, *J. Chem. Soc., Perkin Trans. 1*, 1974, 409; (b) B. J. Arnold, P. G. Sammes and T. W. Wallace, *J. Chem. Soc., Perkin Trans. 1*, 1974, 415.
- 5 J. L. Charlton, G. L. Plourde, K. Koh and A. S. Secco, *Can. J. Chem.*, 1990, **68**, 2022; J. L. Charlton, K. Koh and G. L. Plourde, *Can. J. Chem.*, 1990, **68**, 2028; J. L. Charlton and S. Maddaford, *Can. J. Chem.*, 1993, **71**, 827; J. L. Charlton, S. Maddaford, K. Koh, S. Boulet and M. H. Saunders, *Tetrahedron: Asymmetry*, 1993, **4**, 645; J. L. Charlton, D. Bogucki and P. Guo, *Can. J. Chem.*, 1995, **73**, 1463; D. M. Coltart and J. L. Charlton, *Can. J. Chem.*, 1996, **74**, 88.
- 6 M. Yoshioka, S. Miyazoe and T. Hasegawa, *J. Chem. Soc., Perkin Trans. 1*, 1993, 2781.
- 7 K. Iida, E. Kawata, K. Komada, M. Saito, S. Kumakura and M. Yoshioka, *Acta Crystallogr., Sect. C*, 1998, **54**, 1938.
- 8 P. J. Wagner, D. Subrahmanyam and B.-S. Park, *J. Am. Chem. Soc.*, 1991, **113**, 709.
- 9 R. Haag, J. Wirz and P. J. Wagner, *Helv. Chim. Acta*, 1977, **60**, 2595.
- 10 K. Iida, M. Saito and M. Yoshioka, *J. Org. Chem.*, 1999, **64**, 7407.
- 11 M. Sobczak and P. J. Wagner, *Tetrahedron Lett.*, 1998, **39**, 2523; P. J. Wagner, M. Sobczak and B.-S. Park, *J. Am. Chem. Soc.*, 1998, **120**, 2488.
- 12 P. J. Wagner, *Acc. Chem. Res.*, 1971, **4**, 168.
- 13 O. L. Chapman, U.-P. E. Tsou and J. W. Johnson, *J. Am. Chem. Soc.*, 1987, **109**, 553.
- 14 M. Yoshioka, T. Suzuki and M. Oka, *Bull. Chem. Soc. Jpn.*, 1984, **57**, 1604; M. Yoshioka, K. Nishizawa, M. Arai and T. Hasegawa, *J. Chem. Soc., Perkin Trans. 1*, 1991, 541.

Paper a908326j