

# Novel photoadditions of tertiary amines to the $\alpha$ -position of $\alpha,\beta$ -unsaturated $\gamma,\delta$ -epoxy nitriles

PERKIN

Keitaro Ishii,\* Satoshi Hakamada, Mayumi Nagano, Masahiro Noji, Shigeo Sugiyama, Masashi Kotera and Masanori Sakamoto

Meiji Pharmaceutical University, 2-522-1 Noshio, Kiyose-shi, Tokyo 204-8588, Japan.

E-mail: ishiikei@my-pharm.ac.jp; Fax: +81 (0)424 958783; Tel: +81 (0)424 958783

Received (in Cambridge, UK) 14th November 2001, Accepted 7th May 2002

First published as an Advance Article on the web 29th May 2002

The photoreactions of  $\alpha,\beta$ -unsaturated  $\gamma,\delta$ -epoxy nitriles **1**, **2**, **13** and **16** with triethylamine give rise to novel 1 : 1  $\alpha$ -adducts (e.g., **5**) efficiently. After treatment with silica gel, the adducts undergo retro-Michael reaction leading to the corresponding  $\alpha$ -alkylidenenitrile derivatives (e.g., **3**). The epoxy nitrile **1** also reacts with various tertiary amines to afford  $\alpha$ -adducts. The reaction of **1** and the silylamine **24** gives mainly methylene derivative **22** and silylated compound **25** after treatment with silica gel. The reaction may proceed *via* single-electron-transfer from the amine to the excited epoxy nitrile.

## Introduction

Photoinduced electron-transfer reactions have received considerable interest recently in the discovery of new and synthetically useful chemical reactions.<sup>1</sup> For example, the photoadditions of tertiary amines to  $\alpha,\beta$ -unsaturated ketones and esters have been reported to afford 1 : 1  $\beta$ -adducts between carbonyl compounds and amines and reduced compounds.<sup>2</sup> Mariano *et al.* conducted intensive studies on the photoaddition using  $\alpha$ -trimethylsilylamine and on synthetic applications of the photoaddition to *N*-heterocyclic compounds.<sup>3,4</sup>

In a previous communication,<sup>5</sup> we have shown that direct irradiation of  $\alpha,\beta$ -unsaturated  $\gamma,\delta$ -epoxy nitrile **1**<sup>6</sup> with triethylamine (TEA) undergoes novel 1 : 1  $\alpha$ -addition and subsequent retro-Michael reaction leading to  $\alpha$ -alkylidenenitrile **3**. Similarly, the reaction of **2**<sup>7</sup> gives lactone **6** and  $\alpha$ -alkylidenenitrile **7** (Scheme 1). In this paper we describe the details of the photoreactions of the epoxy nitriles **1** and **2** with TEA, reactions of **1** with various tertiary amines, and studies on the scope and limitation of the addition for epoxy compounds. Furthermore, the addition of [(trimethylsilyl)methyl]diethylamine (TMSA) **24** to the epoxy nitrile **1** is also investigated.


## Results and discussion

### Photoadditions of epoxy nitriles with TEA


**Preparative irradiation of nitrile **1** in TEA.** The nitrile (*E*)-**1** was irradiated in TEA with a low-pressure mercury lamp through a quartz filter ( $\lambda = 254$  nm) at RT (conversion 51%) and subsequent silica gel column chromatography yielded (*Z*)-**1** (6%),  $\alpha$ -alkylidenenitrile **3** (26%) and reduced compound **4** (13%) (Scheme 1).<sup>†</sup> The structure of product **3** was deduced from the H–H and C–H COSY, phase-sensitive NOESY, and HMBC spectra. In the HMBC spectrum of **3**, the crosspeaks between H-3 and C(2) (<sup>2</sup>*J*), between H-3 and C(1) (<sup>3</sup>*J*) and between H-4 and C(2) (<sup>3</sup>*J*) are observed. The appearance of the crosspeak between H-3 and H-6' in the NOESY spectrum indicates *Z* orientation at the C(2) double bond.

The characteristic signals of **3** could not be observed in the <sup>1</sup>H NMR spectrum of the crude photoproduct of **1**. The TEA-adduct **5** was separated after basic aluminium oxide chromatography as a mixture of four diastereomers, which were treated with silica gel leading to **3** quantitatively. The result

<sup>†</sup> Yields for compounds throughout the rest of the paper are based on converted starting material.



Scheme 1 Reagents and conditions: i,  $\lambda = 254$  nm, Et<sub>3</sub>N, RT; ii, silica gel.




**Scheme 2** Reagents and conditions: i,  $(\text{EtO})_2\text{P}(\text{O})\text{CH}(\text{Me})\text{CN}$ , NaH, DMF; ii,  $\lambda = 254 \text{ nm}$ ,  $\text{Et}_3\text{N}$ , MeCN, RT; iii, silica gel; iv,  $(\text{EtO})_2\text{P}(\text{O})\text{CH}_2\text{CN}$ , NaH, DMF.

shows that **5** is transformed by acid-catalyzed retro-Michael reaction to the  $\alpha$ -alkylidenenitrile **3**.

**Preparative irradiation of nitrile 13 with TEA.** As the TEA-adduct **5** could not be separated in pure form, we attempted to obtain the TEA-adduct **14**, stable to silica gel. The epoxy nitrile **13** was prepared from isophorone oxide in 85% yield ( $E : Z \approx 2 : 3$ ) by the Horner–Emmons reaction. Preparative irradiation of **13** and TEA (30 equiv.) in acetonitrile with a low-pressure mercury lamp through a quartz filter ( $\lambda = 254 \text{ nm}$ ) at RT (conversion 65%) gave four adducts **14A–D** (20%) and two reduced compounds **15A–B** (16%) (Scheme 2). † One of the adducts **14A** was separated in *ca.* 80% purity and the structure was assured by its spectral data (see Experimental section).

**Preparative irradiation of nitrile 2 in TEA.** The nitrile ( $E$ )-**2** was irradiated in TEA with a low-pressure mercury lamp through a quartz filter ( $\lambda = 254 \text{ nm}$ ) at RT (conversion 80%) and subsequent silica gel chromatography afforded lactone **6** (10%),  $\alpha$ -alkylidenenitrile **7** (6%), reduced compounds ( $E$ )-**8** (5%) and ( $Z$ )-**8** (15%) and ketone **9** (6%) (Scheme 1). †

The structures of **6–8** were determined on the basis of spectral data. In particular, the  $^{13}\text{C}$  NMR spectrum of **6** shows a characteristic signal at  $\delta_{\text{C}} 165.9$  due to the  $\delta$ -lactone moiety, which is also evidenced by the IR band at  $1720 \text{ cm}^{-1}$ . As characteristic signals for the ethylidene moiety of **6**, the  $^1\text{H}$  NMR spectrum shows a doublet at  $\delta 1.91$  and a quartet-doublet at  $\delta 6.91$ , the  $^{13}\text{C}$  NMR spectrum has a quartet at  $\delta_{\text{C}} 13.8$  and a doublet at  $\delta_{\text{C}} 135.5$ . Furthermore, the ( $E/Z$ )-configuration at the ethylidene moiety was established by the significant NOE enhancement (6.9%) between 5-H and Me in the ethylidene group. The lactone **6** was presumably formed by cyclization of  $\alpha$ -alkylidenenitrile **10** (Fig. 1) possessing the ( $Z$ )-configuration



**Fig. 1**

of the exocyclic double bond at C(1') and by subsequent hydrolysis with silica gel.

Compound **7** could not be isolated in pure form, but the structure could be presumed by the spectral data of a 1 : 4 mixture of **7** and ( $Z$ )-**8**. The  $^1\text{H}$  NMR spectrum of **7** shows a double doublet at  $\delta 2.03$  and a quartet-doublet at  $\delta 6.23$ ; the  $^{13}\text{C}$  NMR spectrum has a quartet at  $\delta_{\text{C}} 17.0$  and a doublet at  $\delta_{\text{C}} 143.2$  due to the ethylidene moiety and two singlets at

$\delta_{\text{C}} 115.5$  and  $118.2$  due to C(1) and C(2). After treatment of **7** with silica gel, the formation of lactone **6** was not observed. Therefore, the ( $E/Z$ )-configuration of the exocyclic double bond at C(1') should be  $E$ . However, the ( $E/Z$ )-configuration at C(2) for **7** could not be determined.

The ( $E/Z$ )-configuration of **8** was deduced from the chemical shift for 2- $\text{H}_2$  and 3-H in the  $^1\text{H}$  NMR spectrum; the signals for 2- $\text{H}_2$  ( $\delta 3.49$  and  $3.71$ ) for ( $Z$ )-**8** and for 3-H ( $\delta 5.86$ ) for ( $E$ )-**8** appear at lower field than those for ( $E$ )-**8** ( $\delta 3.32$ ) and for ( $Z$ )-**8** ( $\delta 5.43$ ), respectively, due to the deshielding effects of the hydroxy group at C(2').

The mechanism for the formation of the ketone **9** in TEA solution is not yet clear, because **9** has been afforded by triplet sensitization reaction of **2** in earlier studies.<sup>6,7</sup>

We could not isolate the nitriles **11** and **12** (Fig. 1), which are mainly formed in the photoreaction of **2** in the presence of TEA (1 equiv.) in acetonitrile.<sup>6</sup> In the case of the photoreaction of **2** in TEA solution, the formation of **11** and **12** is suppressed because the SET process from TEA to the excited state **2\*** occurs (presumably) faster than the C( $\gamma$ )-C( $\delta$ ) bond cleavage.

**Preparative irradiation of nitrile 16 with TEA.** The nitrile **16** was prepared from *trans*-4,5-epoxyhexan-3-one in 49% yield ( $E : Z \approx 9 : 2$ ) by the Horner–Emmons reaction. A solution of the nitrile **16** ( $E : Z \approx 3 : 2$ ) and TEA (30 equiv.) in acetonitrile was irradiated with a low-pressure mercury lamp through a quartz filter ( $\lambda = 254 \text{ nm}$ ) at RT (conversion 92%) and subsequent silica gel column chromatography yielded  $\alpha$ -alkylidenenitriles ( $E$ )-**17** (14%) and ( $Z$ )-**17** (12%). † The reduced compound could not be observed.

The ( $E/Z$ )-configuration at C-3 for **17** was deduced from the chemical shift for 3- $\text{CH}_2\text{CH}_3$  in the  $^{13}\text{C}$  NMR spectrum; the signal ( $\delta_{\text{C}} 20.7$ ) for ( $E$ )-**17** appears at lower field than that for ( $Z$ )-**17** ( $\delta_{\text{C}} 29.4$ ) due to the steric interactions between 3-Et and 4- $\text{CH}(\text{OH})\text{Me}$ . Furthermore, the ( $E/Z$ )-configuration at C-3 was established by the significant NOE enhancements between 3- $\text{CH}_2\text{CH}_3$  and 5-H (8.5%) for ( $E$ )-**17** and between 3- $\text{CH}_2\text{CH}_3$  and 4-H (3.5%) for ( $Z$ )-**17**. The ( $Z$ )-configuration at the ethylidene moiety was determined also by the significant NOE enhancements between 3- $\text{CH}_2\text{CH}_3$  and 2-CH in the ethylidene group (6.6%) for ( $E$ )-**17** and between 5-H and 2-CH (2.2%) for  $Z$ -**17**.


In the order to test the viability of the reaction, epoxy ketones **18**<sup>8</sup> and **19**<sup>‡</sup> and epoxy nitrile **20**<sup>10</sup> (Fig. 2) were irradiated in the presence of TEA (30 equiv.) under the same reaction conditions. However, the TEA-adduct or  $\alpha$ -alkylidenenitrile derivative could not be obtained.

‡ The irradiations of  $\alpha,\beta$ -epoxy ketones in acetonitrile in the presence of 5 equiv. of  $\text{Et}_3\text{N}$  afford the corresponding  $\beta$ -hydroxy ketones.<sup>9</sup>


**Table 1** Analytical irradiations of **1** with TEA<sup>a</sup>

Entry	Concentration of TEA (mol dm <sup>-3</sup> ) [equiv.]	Solvent	Conversion (%)	Yields (%) <sup>†</sup> <sup>b</sup>	
				<b>3</b>	<b>4</b>
1	0.056 [1]	MeCN	56	trace	5 (6) <sup>c</sup>
2	0.282 [5]	MeCN	72	8	23 (3)
3	0.564 [10]	MeCN	85	19	20
4	1.13 [20]	MeCN	84	31	22
5	1.69 [30]	MeCN	95	45	14
6	2.26 [40]	MeCN	87	46	10
7	neat	—	49	27	14
8 <sup>d,e</sup>	0.564 [10]	MeCN	99	43	7
9 <sup>d</sup>	1.69 [30]	PrCN	51	36	12
10 <sup>d</sup>	1.69 [30]	Hexane	36	22	6

<sup>a</sup> A 0.0564 mol dm<sup>-3</sup> solution of **1** (*E/Z* = 2 : 1) in solvent was irradiated for 1.5 h at RT. <sup>b</sup> The yields were determined by <sup>1</sup>H NMR analysis using bis(trimethylsilyl)acetylene as internal standard, after stirring of the reaction mixture with silica gel in CHCl<sub>3</sub> for 2 h at RT. <sup>c</sup> Values in angle brackets are yields of cyclopropene **21**. <sup>d</sup> Irradiation for 1 h. <sup>e</sup> With 5 equiv. of Pr<sup>i</sup>NH<sub>2</sub>.

**Fig. 2**


**Analytical irradiations of **1** with TEA.** Photoreactions of nitrile **1** (*E/Z* = 2 : 1) in acetonitrile containing varying amounts of TEA (1, 5, 10, 20, 30 and 40 equiv.) or in TEA were performed, and results are shown in Table 1. The yield of **3** increased with increasing TEA concentration up to 40 equiv. In the low concentration range of TEA (entries 1 and 2) the cyclopropene **21**<sup>6</sup> (Fig. 3) was also formed. For the formation

**Fig. 3**

of **3** a high concentration of TEA is required. The results may suggest that TEA functions partially as a base in the reaction. Thus, isopropylamine (5 equiv.), possessing low electron donating character, was added to the solution of **1** and TEA (10 equiv.) in acetonitrile and the solution was irradiated. The yield of **3** (43%<sup>†</sup>; entry 8) was similar to that for the reaction using 30 equiv. of TEA (Entry 5). As the polarity of solvent decreases, the conversion and the yield of **3** decrease (entries 5, 9 and 10). These results may indicate that a single-electron transfer (SET) from TEA to the excited epoxy nitrile **1**<sup>\*</sup> occurs (Scheme 3). The resulting ion-radical intermediates (*e.g.*, **A**) are transformed into the reduced compound **4** and the adduct **5** *via* biradical intermediates (*e.g.*, **B**) in a manner similar to the reaction of enones and tertiary amines.<sup>2</sup> Direct irradiation of  $\alpha,\beta$ -unsaturated  $\gamma,\delta$ -epoxy nitriles and ketones (*e.g.*, **2** and **18**) ( $\lambda$  = 254 nm) in acetonitrile gave mainly the C( $\gamma$ )-C( $\delta$ ) bond-cleaved products (*e.g.*, **12**) and the cyclopropene (*e.g.*, **21**) *via* the carbene intermediate.<sup>7,8</sup> Therefore, in the low concentration range of TEA these compounds are formed faster than are the SET products (*e.g.*, **4** and **5**).

#### Photoadditions of nitrile **1** with various amines

The results of the photoreactions of the nitriles **1**, **2** and **16** with

**Scheme 3**

TEA prompted us to investigate the reactions of nitrile **1** and other amines. The nitrile **1** was irradiated in the presence of 30 equiv. of tertiary amines (dimethylethylamine, methyldiethylamine, tripropylamine, ethyldiisopropylamine, diethylethanolamine, 1-methylpyrrolidine, 1-methylpiperidine, tetramethylmethanediamine and tetramethylethylenediamine), and the results are summarized in Table 2. The reactivity for the addition reaction decreases in the order *N*-Me  $\approx$  *N*-Et > *N*-Pr > *N*-Pr<sup>i</sup> substituents owing (presumably) to the steric hindrance of the  $\alpha$ -carbon atom (entries 1–4). The  $\alpha$ -carbon atom in cyclic amines and in diamines has no reactivity (entries 6–9). Furthermore, the reactions of **1** with a secondary amine (diethylamine) and a primary amine (butylamine) were also performed under the same reaction conditions. However, the corresponding adducts could not be observed.

The structures of the  $\alpha$ -alkyldenitriles **22** and **23** (Fig. 3) were determined on the basis of their spectral data. In particular, the (*Z*)-orientation at the C(2) double bond in **23** was determined by comparison of the <sup>1</sup>H NMR chemical shifts of H-3 ( $\delta$  6.38) and H<sub>2</sub>-6' ( $\delta$  1.86 and 2.00) with those of **3** ( $\delta$  6.46 for H-3 and  $\delta$  1.87 and 1.98 for H<sub>2</sub>-6').

#### Photoaddition of nitrile **1** with TMSA **24**

**Preparative irradiation.** Since the reactions of the nitrile **1** and tertiary amine had given the  $\alpha$ -adducts in moderate yields, the photoreaction of **1** and TMSA **24**,<sup>11</sup> which is known as a stronger electron donor than TEA,<sup>§</sup> was investigated. A solution of the nitrile **1** (*E* : *Z* = 2 : 1) and **24** (10 equiv.) in acetonitrile was irradiated with a low-pressure mercury lamp through a quartz filter ( $\lambda$  = 254 nm) at RT (conversion 100%) and subsequent silica gel column chromatography yielded **3** (6%), **4** (23%), **22** (12%) and silylated compound **25** (21%) (Scheme 4).<sup>†</sup>

<sup>§</sup> Oxidation potentials of TMSA **24** and TEA are 0.57 V and 0.89 V *vs.* SCE, respectively.<sup>12</sup>

**Table 2** Irradiations of **1** with various tertiary amines<sup>a</sup>


Entry	Amine	Irradiation Time (h)	Conversion (%)	Yields (%) <sup>†b</sup>			
				3	22	23	4
1	EtNMe <sub>2</sub>	4.5	91	11	19	—	19
2	Et <sub>2</sub> NMe	5	94	22	13	—	20
3	Pr <sub>3</sub> N	3	76	—	—	12	28
4	Pr <sub>2</sub> <sup>i</sup> NEt	5	69	28	—	—	17
5	Et <sub>2</sub> N(CH <sub>2</sub> ) <sub>2</sub> OH	6.5	89	41	—	—	10
6	1-Methylpyrrolidine	3.5	81	—	2	—	20
7	1-Methylpiperidine	7	74	—	5	—	4
8	Me <sub>2</sub> NCH <sub>2</sub> NMe <sub>2</sub>	2.5	90	—	1	—	16
9	Me <sub>2</sub> N(CH <sub>2</sub> ) <sub>2</sub> NMe <sub>2</sub>	2.5	100	—	8	—	22

<sup>a</sup> A 0.0564 mol dm<sup>-3</sup> solution of **1** (*E/Z* = 2 : 1) in acetonitrile was irradiated ( $\lambda = 254$  nm) with 30 equiv. of amine at RT. <sup>b</sup> Isolated yields after silica gel flash-chromatography.

**Table 3** Irradiations of **1** with various concentrations of TMSA **24** in various solvents<sup>a</sup>

Entry	Concentration of TMSA (mol dm <sup>-3</sup> ) [equiv.]	Solvent	Conversion (%)	Yields (%) <sup>†b</sup>			
				3	22	25	4
1	-0.175 [5]	MeCN	41	3	7	10	8
2	0.350 [10]	MeCN	58	5	9	19	11
3	0.525 [15]	MeCN	51	8	10	20	8
4	-1.05 [30]	MeCN	53	4	6	12	5
5	0.350 [10]	0.1%MeOH–MeCN	60	8	13	19	7
6	0.350 [10]	0.7%MeOH–MeCN	89	5	11	11	8
7	0.350 [10]	1.4%MeOH–MeCN	81	5	11	10	7
8	0.350 [10]	10%MeOH–MeCN	74	4	8	1	6

<sup>a</sup> A 0.035 mol dm<sup>-3</sup> solution of **1** (*E/Z* = 2 : 1) was irradiated ( $\lambda = 254$  nm) for 0.5 h at RT. <sup>b</sup> The yields were determined by GLC analysis using octadecane as an internal standard, after stirring of the reaction mixture with silica gel in CHCl<sub>3</sub> for 2 h at RT.



**Scheme 4** Reagents and conditions : i,  $\lambda = 254$  nm, Et<sub>2</sub>NCH<sub>2</sub>TMS **24**, MeCN, RT; ii, silica gel, CHCl<sub>3</sub>, RT.

The structure of product **25** was determined on the basis of its spectral data. In particular, the (*Z*)-orientation at the C(2) double bond was determined by comparison of the <sup>1</sup>H NMR chemical shifts of H-3 ( $\delta$  6.57) and H<sub>2</sub>-6' ( $\delta$  1.85 and 2.03) with those of **3**.

We also confirmed the reaction of epoxy ketones **18**<sup>8</sup> and **19** and epoxy nitrile **20**<sup>10</sup> (Fig. 2) in the presence of 10 equiv. of TMSA **24** under the same reaction conditions. However, the TMSA-adducts could not be detected.

### Analytical irradiations

Photoreactions of nitrile **1** (*E/Z* = 2 : 1) in acetonitrile containing varying amounts of TMSA **24** (5, 10, 15 and 30 equiv.) (entries 1–4) and containing MeOH and **24** (entries 5–8) were performed, and results are shown in Table 3. The yield of **3**, **22** and **25** increased with increasing TMSA **24** concentration up to 0.525 mol dm<sup>-3</sup>. The yield of **25** decreased with increasing MeOH concentration. The probable reaction mechanism for the formation of  $\alpha$ -alkylidenenitriles **3**, **22** and **25** outlined in Scheme 5 appears to rationalize these results. SET between the excited nitrile **1**\* and TMSA **24** results in generation of the radical-ion-pair (e.g., **C**). The resulting cation radical **24**<sup>+</sup> undergoes  $\alpha$ -deprotonation (H<sub>a</sub><sup>+</sup> and H<sub>b</sub><sup>+</sup>) and nucleophile-assisted desilylation<sup>4</sup> leading to the adducts **26** and **27** and **28**, respectively. Finally, retro-Michael reaction gives the respective

olefins **25**, **3** and **22**. Compound **25** is preferentially formed in less polar aprotic solvents rather than in polar protic solvents. The results may show that the deprotonation of H<sub>a</sub><sup>+</sup> in **C** occurs favorably in a solvent cage (contact-ion-pair).<sup>4</sup>

In conclusion, the photoreactions of  $\alpha,\beta$ -unsaturated  $\gamma,\delta$ -epoxy nitriles **1**, **2**, **13** and **16** with TEA afford novel 1 : 1  $\alpha$ -adducts between the nitriles and TEA, and subsequent retro-Michael reaction gives  $\alpha$ -alkylidenenitriles (e.g., **3**). The nitrile **1** also reacts with various tertiary amines and TMSA **24**. The reaction is peculiar to  $\alpha,\beta$ -unsaturated  $\gamma,\delta$ -epoxy nitriles and may proceed *via* SET from the amine to the excited nitrile.

### Experimental

Mps and bps are uncorrected. Mps were measured with a Yanaco MP-3 apparatus and bps were measured with a Büchi

Kugel Rohr GKR-50 apparatus. UV spectra were recorded on a Hitachi 124 spectrometer and IR spectra on a Hitachi 215 spectrometer. NMR spectra were obtained with a JEOL JNM-EX270 (270 MHz; EX) or a JEOL JNM-GX400 (400 MHz; GX) spectrometer for samples in CDCl<sub>3</sub> solution using tetramethylsilane as internal standard, and *J*-values are given in Hz. Mass spectra (MS) and high-resolution MS (HRMS) were taken on a JEOL JMS-DX302 spectrometer. GLC was carried out on a Shimadzu GC-14A instrument (flame ionization detection) and a 30 m × 0.25 mm column of Rascot OV-1 was used. Column chromatography was performed with Nacalai silica gel 60 [230–400 mesh (SiO<sub>2</sub>)] or Merck aluminium oxide 90(basic) [70–230 mesh, activity III (Al<sub>2</sub>O<sub>3</sub>)].

An Eikosha 60 W low-pressure mercury lamp was used for irradiation. The photolysis solutions were purged with argon both before and during irradiation.

### Photoadditions of nitriles with triethylamine (TEA)

**Preparative irradiation of nitrile 1 in TEA.** A solution of (*E*)-**1** (700 mg, 3.99 mmol) in TEA (70 cm<sup>3</sup>) was irradiated with a low-pressure mercury lamp in a quartz test tube (conversion 51%) for 7 h at RT. After removal of the solvent, flash chromatography [SiO<sub>2</sub>; hexane–ethyl acetate (4 : 1)] of the residue afforded (*Z*)-**1** (21.7 mg, 6%),  $\alpha$ -alkylidene nitrile **3** (110 mg, 26%) and the alcohol **4** (48.2 mg, 13%). †

2-(3-Hydroxy-3,5,5-trimethylcyclohex-1-enyl)isocrotononitrile **3**, bp 130 °C at 0.22 mmHg (Found: M<sup>+</sup>, 205.1468. C<sub>13</sub>H<sub>19</sub>NO requires *M*, 205.1466);  $\lambda_{\max}$  (EtOH)/nm 245 (g/dm<sup>3</sup> mol<sup>-1</sup> cm<sup>-1</sup> 5000);  $\nu_{\max}$  (film)/cm<sup>-1</sup> 3420 (O–H) and 2190 (C≡N);  $\delta_{\text{H}}$  (EX) 1.02 and 1.05 (6 H, 2 s, 2 × 5'–CH<sub>3</sub>), 1.34 (3 H, s, 3'–CH<sub>3</sub>), 1.56 (1 H, d, *J* 14.2, 4'–H), 1.72 (1 H, dd, *J* 14.2 and 1.0, 4'–H), 1.72–1.82 (1 H, br s, OH), 1.87 and 1.98 (2 H, each d, *J* 16.5, 6'–H<sub>2</sub>), 2.12 (3 H, d, *J* 7.3, 4–H<sub>3</sub>), 6.09 (1 H, br s, 2'–H) and 6.46 (1 H, q, *J* 7.3, 3–H);  $\delta_{\text{C}}$  (EX) 17.5 (q, C-4), 27.8 and 30.8 (2 q, 2 × 5'–CH<sub>3</sub>), 30.1 (s, C-5'), 31.2 (q, 3'–CH<sub>3</sub>), 38.8 (t, C-6'), 50.0 (t, C-4'), 69.1 (s, C-3'), 115.7 (s, C-1), 119.0 (s, C-2), 130.8 (s, C-1'), 132.5 (d, C-2') and 139.9 (d, C-3); *m/z* 205 (M<sup>+</sup>, 25%), 190 (100), 163 (17), 148 (19), 139 (17), 134 (11) and 120 (13).

(3-Hydroxy-3,5,5-trimethylcyclohex-1-enyl)acetonitrile **4**, bp 110 °C at 0.3 mmHg (Found: M<sup>+</sup>, 179.1333. C<sub>11</sub>H<sub>17</sub>NO requires *M*, 179.1311);  $\nu_{\max}$  (film)/cm<sup>-1</sup> 3420 (O–H) and 2240 (C≡N);  $\delta_{\text{H}}$  (EX) 1.00 and 1.06 (6 H, 2 s, 2 × 5'–CH<sub>3</sub>), 1.30 (3 H, s, 3'–CH<sub>3</sub>), 1.53 and 1.68 (2 H, each d, *J* 14.2, 4'–H<sub>2</sub>), 1.65–1.75 (1 H, br s, OH), 1.76–1.90 (2 H, m, 6'–H<sub>2</sub>), 3.04 (2 H, br s, 2–H<sub>2</sub>) and 5.72 (1 H, d, *J* 1.0, 2'–H);  $\delta_{\text{C}}$  (EX) 25.7 (t, C-2), 27.7, 30.7 and 31.1 (3 q, 2 × 5'–CH<sub>3</sub> and 3'–CH<sub>3</sub>), 30.4 (s, C-5'), 42.1 (t, C-6'), 49.6 (t, C-4'), 69.1 (s, C-3'), 117.7 (s, C-1), 127.7 (s, C-1'), and 131.1 (d, C-2'); *m/z* 179 (M<sup>+</sup>, 3%), 164 (100), 139 (36), 123 (27), 100 (20) and 43 (28).

**Isolation of TEA-adduct 5.** A solution of **1** (1.16 g, 6.56 mmol) in TEA (116 cm<sup>3</sup>) was irradiated with a low-pressure mercury lamp in a quartz test-tube (conversion 61%) for 8 h at RT. After removal of the solvent, chromatography [Al<sub>2</sub>O<sub>3</sub>; hexane–ethyl acetate (2 : 1)] of the residue gave a mixture of four stereoisomers of compound **5** (635 mg, 57%). †

3-(Diethylamino)-2-(3-hydroxy-3,5,5-trimethylcyclohex-1-enyl)butyronitrile **5**, an oil (Found: M<sup>+</sup>, 278.2377. C<sub>17</sub>H<sub>30</sub>N<sub>2</sub>O requires *M*, 278.2358);  $\nu_{\max}$  (film)/cm<sup>-1</sup> 3410 (O–H) and 2230 (C≡N);  $\delta_{\text{H}}$  (EX) *inter alia* 2.33–2.67 (4 H, m, 2 × NCH<sub>2</sub>), 3.01–3.27 (2 H, m, 2- and 3–H) and 5.59 and 5.65 (1 H, 2 br s, 2'–H);  $\delta_{\text{C}}$  (EX) *inter alia* 39.6, 39.7, 40.2 and 40.6 (4 t, C-6'), 43.4, 43.5, 46.3 and 46.6 (4 d, C-2), 55.18, 55.22 and 55.5 (3 d, C-3), 69.1 and 69.2 (2 s, C-3'), 119.96 and 120.03 (2 s, C-1), 131.3, 131.4, 131.89 and 131.95 (4 s, C-1') and 131.8, 133.0 and 133.1 (3 d, C-2').

**Transformation of 5 into 3.** A solution of **5** (51.9 mg, 0.187 mmol) in chloroform (10 cm<sup>3</sup>) was stirred in the presence

of SiO<sub>2</sub> (1.0 g) for 2 h at RT. After filtration, <sup>1</sup>H NMR analysis of the reaction mixture showed that **5** was converted to  $\alpha$ -alkylidene nitrile **3** quantitatively.

**Preparation of nitrile 13.** Diethyl (1-cyanoethyl)phosphonate (7.64 g, 40.0 mmol) was added dropwise to a suspension of 60% NaH (1.60 g, 40.0 mmol) in dry DMF (56 cm<sup>3</sup>) at RT. After stirring of the mixture for 20 min at RT, isophorone oxide (5.53 g, 35.9 mmol) was added dropwise and stirring was continued at –50 °C for 20 min and for 30 min at RT; ice–water (53 cm<sup>3</sup>) was then added to the mixture, and the organic phase was extracted with diethyl ether. The continued extracts were washed with brine, dried over MgSO<sub>4</sub>, and concentrated *in vacuo*, giving a residue, which was subjected to flash chromatography [SiO<sub>2</sub>; hexane–ethyl acetate (9 : 1)] to afford the nitriles (*E*)-**13** (2.18 g, 32%) and (*Z*)-**13** (3.65 g, 53%).

(*E*)-2-[4,4,6-Trimethyl-7-oxabicyclo[4.1.0]heptan-2-ylidene]propiononitrile (*E*)-**13**, bp 85 °C at 0.25 mmHg (Found : C, 75.22; H, 8.99; N, 7.31. C<sub>12</sub>H<sub>17</sub>NO requires C, 75.35; H, 8.96; N, 7.32%);  $\nu_{\max}$  (film)/cm<sup>-1</sup> 2210 (C≡N);  $\delta_{\text{H}}$  (EX) 0.83, 0.97 and 1.39 (9 H, 3 s, 2 × 4'–CH<sub>3</sub> and 6'–CH<sub>3</sub>), 1.58 (1 H, dd, *J* 15 and 1, 1- or 5'–H), 1.87 (1 H, d, *J* 15.2, 5'–H), 2.08–2.10 (3 H, m with t character, *J* 1, 3–H<sub>3</sub>), 2.24–2.26 (2 H, m, 3'–H<sub>2</sub>) and 3.50 (1 H, s, 5'– or 1'–H);  $\delta_{\text{C}}$  (EX) 15.4, 24.7, 27.2 and 30.2 (4 q, 4 × CH<sub>3</sub>), 32.3 (s, C-4'), 41.1 and 42.9 (2 t, C-3' and C-5'), 56.8 (d, C-1'), 61.8 (s, C-6'), 108.3 (s, C-2), 119.0 (s, C-1) and 151.8 (s, C-2'); *m/z* 191 (M<sup>+</sup>, 100%), 176 (57), 148 (36), 135 (68), 118 (24), 107 (31) and 91 (21).

(*Z*)-2-[4,4,6-Trimethyl-7-oxabicyclo[4.1.0]heptan-2-ylidene]propiononitrile (*Z*)-**13**, bp 140 °C at 0.9 mmHg (Found : M<sup>+</sup>, 191.1312. C<sub>12</sub>H<sub>17</sub>NO requires *M*, 191.1310);  $\nu_{\max}$  (film)/cm<sup>-1</sup> 2210 (C≡N);  $\delta_{\text{H}}$  (EX) 0.79, 0.98 and 1.40 (9 H, 3 s, 2 × 4'–CH<sub>3</sub> and 6'–CH<sub>3</sub>), 1.58 (1 H, d, *J* 15, 5'–H), 1.88 (1 H, d, *J* 15, 5'–H), 1.97 (3 H, s, 3–H<sub>3</sub>), 2.03 (2 H, br s, 3'–H<sub>2</sub>) and 3.81 (1 H, s, 1'–H);  $\delta_{\text{C}}$  (EX) 16.4, 24.6, 27.0 and 30.7 (4 q, 4 × CH<sub>3</sub>), 32.5 (s, C-4'), 36.6 and 42.9 (2 t, C-3' and C-5'), 60.6 (d, C-1'), 61.7 (s, C-6'), 107.9 (s, C-2), 118.9 (s, C-1) and 152.9 (s, C-2'); *m/z* 191 (M<sup>+</sup>, 100%), 176 (48), 148 (32), 135 (64), 118 (23), 107 (31) and 91 (17).

**Preparative irradiation of nitrile 13 with TEA.** A solution of **13** (*E* : *Z* = 1 : 2) (736 mg, 3.85 mmol) and TEA (11.7 g, 11.6 mmol) in acetonitrile (100 cm<sup>3</sup>) was irradiated with a low-pressure mercury lamp in a quartz-test-tube (conversion 65%) for 2.5 h at RT. After removal of the solvent, flash chromatography [SiO<sub>2</sub>; hexane–ethyl acetate (4 : 1 → 1 : 1)] of the residue afforded adducts **14** (a mixture of four diastereomers, 142 mg, 20%), alcohols **15** [a mixture of two diastereomers (1 : 1), 77.7 mg, 16%]† and intractable materials (mainly polymers). Further purification of **14** and **15** by flash chromatography [SiO<sub>2</sub>; cyclohexane–ethyl acetate (3 : 2)] gave **14A** (contaminated with *ca.* 20% of **14B**) from **14**, and **15A** (contaminated with *ca.* 20% of **15B**) and **15B** (contaminated with *ca.* 30% of **15A**) from **15**.

3-(Diethylamino)-2-(3-hydroxy-3,5,5-trimethylcyclohex-1-enyl)-2-methylbutyronitrile **14A** (contaminated with *ca.* 20% of **14B**), an oil (Found : M<sup>+</sup> – CH<sub>3</sub>, 277.2275. C<sub>18</sub>H<sub>32</sub>N<sub>2</sub>O – CH<sub>3</sub> requires *M/Z*, 277.2280);  $\nu_{\max}$  (film)/cm<sup>-1</sup> 3460 (O–H) and 2220 (C≡N);  $\delta_{\text{H}}$  (EX) 0.99–1.06 (15 H, m, 4 × CH<sub>3</sub> and 4–H<sub>3</sub>), 1.30 and 1.50 (6 H, 2 s, 2 × CH<sub>3</sub>), 1.51–1.90 (5 H, m, OH, 4'–H<sub>2</sub> and 6'–H<sub>2</sub>), 2.36–2.49 (2 H, m, 2 × NCH), 2.36–2.49 (3 H, m, 2 × NCH and 3–H) and 5.96–5.98 (1 H, m, 2'–H);  $\delta_{\text{C}}$  (EX) 10.5, 14.1, 22.9, 27.0, 30.9 and 31.5 (7 q, 2 q at  $\delta_{\text{C}}$  14.1, 7 × CH<sub>3</sub>), 30.3 (s, C-5'), 37.8 and 44.3 (3 t, 2 t at  $\delta_{\text{C}}$  44.3, 2 × NCH<sub>2</sub> and C-6'), 49.8 (t, C-4'), 49.2 (s, C-2), 58.7 (d, C-3), 69.1 (s, C-3'), 122.8 (s, C-1), 129.9 (d, C-2') and 135.6 (s, C-1'); *m/z* 277 (M<sup>+</sup> – 15, 0.7%), 191 (0.7), 178 (16), 175 (15), 160 (15), 132 (11), 100 (100), 84 (11) and 56 (12).

2-(3-Hydroxy-3,5,5-trimethylcyclohex-1-enyl)-2-methylpropiononitrile **15A** (contaminated with *ca.* 20% of **15B**), bp 110 °C

at 0.35 mmHg (Found :  $M^+$ , 193.1470.  $C_{12}H_{19}NO$  requires  $M$ , 193.1467);  $\nu_{\max}$  (film)/ $cm^{-1}$  3520 (O–H) and 2230 (C≡N);  $\delta_H$  (EX) 1.01, 1.06 and 1.30 (9 H, 3 s, 3'-CH<sub>3</sub> and 2 × 5'-CH<sub>3</sub>), 1.42 (3 H, d,  $J$  7.3, 3-H<sub>3</sub>), 1.53 (1 H, d,  $J$  14, 4'-H), 1.62 (1 H, br s, OH), 1.68 (1 H, dt,  $J$  14 and 1.1, 4'-H), 1.83 (1 H, dd,  $J$  16.5 and 1.8, 6'-H), 1.89 (1 H, d,  $J$  16.5, 6'-H), 3.22 (1 H, q,  $J$  7.3, 2-H) and 5.68 (1 H, dd,  $J$  1.8 and 1.1, 2'-H);  $\delta_C$  (EX) 17.3, 27.4, 30.6 and 31.1 (4 q, 4 × CH<sub>3</sub>), 30.4 (s, C-5'), 32.6 (d, C-2), 40.1 (t, C-6'), 49.9 (d, C-4'), 69.1 (s, C-3'), 120.9 (s, C-1), 129.8 (d, C-2') and 133.4 (s, C-1');  $m/z$  193 ( $M^+$ , 2%), 178 (100), 160 (24), 139 (40), 106 (15), 91 (18) and 77 (12).

2-(3-Hydroxy-3,5,5-trimethylcyclohex-1-enyl)-2-methylpropionitrile **15B** (contaminated with *ca.* 30% of **15A**), an oil;  $\delta_H$  (EX) 1.01, 1.05 and 1.30 (9 H, 3 s, 3'-CH<sub>3</sub> and 2 × 5'-CH<sub>3</sub>), 1.42 (3 H, d,  $J$  7.3, 3-H<sub>3</sub>), 1.55 (1 H, d,  $J$  13.9, 4'-H), 1.69 (1 H, br s, OH), 1.68 (1 H, dd,  $J$  13.5 and 1.1, 4'-H), 1.82 and 1.89 (2 H, each d,  $J$  16.9, 6'-H<sub>2</sub>), 3.24 (1 H, q,  $J$  7.3, 2-H) and 5.67–5.69 (1 H, m, 2'-H);  $\delta_C$  (EX) 17.4, 27.5, 30.6 and 31.1 (4 q, 4 × CH<sub>3</sub>), 30.4 (s, C-5'), 32.6 (d, C-2), 40.0 (t, C-6'), 49.9 (d, C-4'), 69.1 (s, C-3'), 120.8 (s, C-1), 129.8 (d, C-2') and 133.3 (s, C-1').

**Preparative irradiation of nitrile 2 in TEA.** A solution of (*E*)-**2**' (1.22 g, 6.39 mmol) in TEA (122 cm<sup>3</sup>) was irradiated with a low-pressure mercury lamp in a quartz test-tube (conversion 80%) for 50 h at RT. After removal of the solvent, flash chromatography [SiO<sub>2</sub>; hexane–ethyl acetate (4 : 1)] of the residue afforded lactone **6** (110 mg, 10%),  $\alpha$ -alkylidenenitrile **7** (63.0 mg, 6%), alcohols (*E*)-**8** (54.0 mg, 5%) and (*Z*)-**8** (147 mg, 15%) and ketone **9**' (58.2 mg, 6%). †

4-(*E*)-Ethylidene-1,7,7-trimethyl-2-oxabicyclo[4.4.0]dec-5-en-3-one **6**, bp 125 °C at 0.2 mmHg (Found :  $M^+$ , 220.1460.  $C_{14}H_{20}O_2$  requires  $M$ , 220.1463);  $\nu_{\max}$  (film)/ $cm^{-1}$  1720 (C=O);  $\delta_H$  (EX) 1.18, 1.26 and 1.56 (9 H, 3 s, 1-CH<sub>3</sub> and 2 × 7-CH<sub>3</sub>), 1.39–2.08 (6 H, m, 8-, 9- and 10-H<sub>2</sub>), 1.91 (3 H, d,  $J$  7.6, 4 = CHCH<sub>3</sub>), 6.35 (1 H, s, 5-H) and 6.91 (1 H, qd,  $J$  7.6 and 1.0, 4 = CH);  $\delta_C$  (EX) 13.8 (q, 4 = CHCH<sub>3</sub>), 19.1 (t, C-9), 29.5, 29.8 and 30.9 (3 q, 3 × CH<sub>3</sub>), 36.3 (s, C-7), 39.5 and 39.9 (2 t, C-8 and C-10), 83.7 (s, C-1), 114.0 (d, C-5), 124.1 (s, C-4), 135.5 (d, 4 = CH), 149.3 (s, C-6) and 165.9 (s, C-3);  $m/z$  220 ( $M^+$ , 89%), 205 (43), 180 (22), 150 (100), 100 (47) and 96 (18).

2-[(*E*)-2-Hydroxy-2,6,6-trimethylcyclohexylidene]but-2-enenitrile **7** [contaminated with *ca.* 80% of (*Z*)-**8**], an oil;  $\delta_H$  (EX) *inter alia* 2.03 (3 H, dd,  $J$  7.3 and 1.3, 4-H<sub>3</sub>), 5.87 (1 H, br s, 2-CH=C) and 6.23 (1 H, qd,  $J$  7.3 and 1.7, 3-H);  $\delta_C$  (EX) *inter alia* 17.0 (q, C-4), 74.3 (s, C-2'), 115.5 and 118.2 (2 s, C-1 and C-2), 116.7 (d, 2-CH=C), 143.2 (d, C-3) and 157.2 (s, C-1').

3-[(*E*)-2-Hydroxy-2,6,6-trimethylcyclohexylidene]propionitrile (*E*)-**8**, bp 120 °C at 0.1 mmHg (Found:  $M^+$ , 193.1458.  $C_{12}H_{19}NO$  requires  $M$ , 193.1467);  $\nu_{\max}$  (film)/ $cm^{-1}$  3430 (O–H) and 2240 (C≡N);  $\delta_H$  (EX) 1.22, 1.29 and 1.39 (9 H, 3 s, 2'-CH<sub>3</sub> and 2 × 6'-CH<sub>3</sub>), 1.39–1.82 (7 H, m, 3', 4'- and 5'-H<sub>2</sub> and OH), 3.32 (2 H, d,  $J$  7.3, 2-H<sub>2</sub>) and 5.86 (1 H, t,  $J$  7.3, 3-H);  $\delta_C$  (EX) 18.1 and 18.2 (2 t, C-2 and C-4'), 29.6, 30.7 and 31.4 (3 q, 2 × 6'-CH<sub>3</sub> and 2'-CH<sub>3</sub>), 36.2 (s, C-6'), 39.1 and 40.9 (2 t, C-3' and C-5'), 73.8 (s, C-2'), 112.6 (d, C-3), 118.9 (s, C-1) and 156.3 (d, C-1');  $m/z$  193 ( $M^+$ , 10%), 178 (70), 160 (20), 153 (100), 135 (37), 108 (36), 100 (44), 95 (30) and 43 (33).

3-[(*Z*)-2-Hydroxy-2,6,6-trimethylcyclohexylidene]propionitrile (*Z*)-**8**, bp 120 °C at 0.15 mmHg (Found:  $M^+$ , 193.1468.  $C_{12}H_{19}NO$  requires  $M$ , 193.1467);  $\nu_{\max}$  (film)/ $cm^{-1}$  3420 (O–H) and 2240 (C≡N);  $\delta_H$  (EX) 1.09 and 1.16 (6 H, 2 s, 2 × 6'-CH<sub>3</sub>), 1.44 (3 H, s, 2'-CH<sub>3</sub>), 1.40–1.89 (7 H, m, 3', 4'- and 5'-H<sub>2</sub> and OH), 3.49 (1 H, dd,  $J$  18.1 and 7.6, 2-H), 3.71 (1 H, dd,  $J$  18.1 and 6.9, 2-H) and 5.43 (1 H, m with t-character,  $J$  7, 3-H);  $\delta_C$  (EX) 18.1 and 19.2 (2 t, C-2 and C-4'), 29.2, 31.4 and 31.9 (3 q, 2 × 6'-CH<sub>3</sub> and 2'-CH<sub>3</sub>), 37.5 (s, C-6'), 39.6 and 43.9 (2 t, C-3' and C-5'), 74.2 (s, C-2'), 113.6 (d, C-3), 120.2 (s, C-1) and 156.3 (d, C-1');  $m/z$  193 ( $M^+$ , 20%), 178 (100), 160 (61), 153 (90), 119 (39), 95 (49), 81 (40) and 43 (90).

**Irradiations of (*E*)-1 in acetonitrile containing varying amounts of TEA.** Aliquots of a 0.0564 mol dm<sup>-3</sup> solution of nitrile (*E*)-**1** in acetonitrile in the presence of TEA (1, 5, 10, 20, 30 and 40 equiv.) or in TEA were irradiated in quartz test-tubes under argon for 1.5 h at RT. Yields were determined by <sup>1</sup>H NMR (GX) analysis using bis(trimethylsilyl)acetylene as internal standard, after stirring of the reaction mixture with SiO<sub>2</sub> in CHCl<sub>3</sub> for 2 h at RT, and results are shown in Table 1.

**Preparation of nitrile 16.** Diethyl cyanomethylphosphonate (3.24 g, 18.3 mmol) was added dropwise to a suspension of 60% NaH (0.73 g, 18.3 mmol) in dry DMF (20 cm<sup>3</sup>) at 0 °C. After stirring of the mixture for 20 min at 0 °C, a solution of *trans*-4,5-epoxyhexan-3-one (1.90 g, 16.7 mmol) in DMF (6 cm<sup>3</sup>) was added dropwise. After stirring of the mixture for 2 h at RT, ice-water (24 cm<sup>3</sup>) was then added to the mixture, the organic phase was extracted with diethyl ether, and the combined extracts were subjected to the same work-up as used for the synthesis of **13**. Flash chromatography [SiO<sub>2</sub>; hexane–benzene–ethyl acetate (15 : 5 : 1)] afforded the nitriles (*E*)-**16** (875 mg, 40%) and (*Z*)-**16** (195 mg, 9%). (*E*,4*RS*,5*RS*)-3-Ethyl-4,5-epoxyhex-2-enenitrile (*E*)-**16**, bp 80 °C at 1 mmHg (Found:  $M^+$ , 137.0844.  $C_8H_{11}NO$  requires  $M$ , 137.0840);  $\nu_{\max}$  (CHCl<sub>3</sub>)/ $cm^{-1}$  2220 (C≡N) and 1620 (C=C);  $\delta_H$  (GX) 1.17 (3 H, t,  $J$  7.7, 3-CH<sub>2</sub>CH<sub>3</sub>), 1.40 (3 H, d,  $J$  5.1, 6-H<sub>3</sub>), 2.40 (1 H, dd,  $J$  17 and 7.7, 3-CH), 2.53 (1 H, dd,  $J$  17 and 7.7, 3-CH), 2.80 (1 H, qd,  $J$  5.1 and 1.8, 5-H), 3.14–3.15 (1 H, m with d-character,  $J$  1, 4-H) and 5.33 (1 H, s, 2-H);  $\delta_C$  (EX) 12.9 and 17.7 (2 q, 2 × CH<sub>3</sub>), 25.5 (t, 3-CH<sub>2</sub>), 58.5 and 58.6 (2 d, C-4 and C-5), 93.8 (s, C-2), 116.4 (s, C-1) and 165.4 (s, C-3);  $m/z$  137 ( $M^+$ , 21%), 122 (100), 108 (21), 94 (76), 79 (28) and 66 (69). (*Z*,4*RS*,5*RS*)-3-Ethyl-4,5-epoxyhex-2-enenitrile (*Z*)-**16** [contaminated with *ca.* 20% of (*E*)-**16**], an oil;  $\delta_H$  (GX) 1.07 (3 H, t,  $J$  7.3, 3-CH<sub>2</sub>CH<sub>3</sub>), 1.43 (3 H, d,  $J$  5.1, 6-H<sub>3</sub>), 1.98 (1 H, dqd,  $J$  17, 7.3 and 2, 3-CH), 2.13 (1 H, dqd,  $J$  17, 7.3 and 1.8, 3-CH), 3.09 (1 H, qd,  $J$  5.1 and 2.2, 5-H), 3.64–3.65 (1 H, m with d-character,  $J$  1, 4-H) and 5.32 (1 H, s, 2-H);  $\delta_C$  (EX) 11.6 and 17.6 (2 q, 2 × CH<sub>3</sub>), 23.6 (t, 3-CH<sub>2</sub>), 54.8 and 58.4 (2 d, C-4 and C-5), 96.5 (s, C-2), 116.1 (s, C-1) and 165.5 (s, C-3).

**Preparative irradiation of nitrile 16 with TEA.** A solution of **16** (*E* : *Z* = 3 : 2) (1.10 g, 8.02 mmol) and TEA (24.3 g, 241 mmol) in acetonitrile (76.5 cm<sup>3</sup>) was irradiated with a low-pressure mercury lamp in a quartz test-tube (conversion 92%) for 7 h at RT. After removal of the solvent, a solution of the residue in CHCl<sub>3</sub> (110 cm<sup>3</sup>) was stirred in the presence of SiO<sub>2</sub> (22 g) for 2 h at RT. After filtration, the filtrate was evaporated to give a residue, which was subjected to flash chromatography [SiO<sub>2</sub>; hexane–ethyl acetate (4 : 1)] to afford (*E*)-**17** (165 mg, 14%), (*Z*)-**17** (145 mg, 12%) and intractable materials (mainly polymers). † (*E*)-3-Ethyl-2-[(*Z*)-ethylidene]-5-hydroxyhex-3-enenitrile (*E*)-**17** [contaminated with *ca.* 45% of (*Z*)-**17**], an oil;  $\delta_H$  (GX) 1.03 (3 H, t,  $J$  7.3, 3-CH<sub>2</sub>CH<sub>3</sub>), 1.32 (3 H, d,  $J$  6.3, 6-H<sub>3</sub>), 1.91 (1 H, br s, OH), 2.10 (3 H, d,  $J$  7.1, 2-CHCH<sub>3</sub>), 2.30 and 2.37 (2 H, each dq,  $J$  14.3 and 7.3, 3-CH<sub>2</sub>), 4.66 (1 H, dq,  $J$  8.6 and 6.3, 5-H), 5.88 (1 H, d,  $J$  8.6, 4-H) and 6.50 (1 H, q,  $J$  7.1, 2-CH);  $\delta_C$  (GX) 13.7, 17.6 and 23.8 (3 q, 3 × CH<sub>3</sub>), 20.7 (t, 3-CH<sub>2</sub>), 64.5 (d, C-5), 116.0 and 118.4 (2 s, C-2 and C-1), 134.6 (d, C-4), 136.2 (s, C-3) and 140.4 (d, 2-CH). (*Z*)-3-Ethyl-2-[(*Z*)-ethylidene]-5-hydroxyhex-3-enenitrile (*Z*)-**17**, an oil (Found:  $M^+$ , 165.1154.  $C_{10}H_{15}NO$  requires  $M$ , 165.1083);  $\nu_{\max}$  (CHCl<sub>3</sub>)/ $cm^{-1}$  3370 (C–O) and 2180 (C≡N);  $\delta_H$  (GX) 1.03 (3 H, t,  $J$  7.6, 3-CH<sub>2</sub>CH<sub>3</sub>), 1.29 (3 H, d,  $J$  6.4, 6-H<sub>3</sub>), 1.58 (1 H, br s, OH), 2.11 (3 H, d,  $J$  7.0, 2-CHCH<sub>3</sub>), 2.23 (2 H, qd,  $J$  7.6 and 1.2, 3-CH<sub>2</sub>), 4.52 (1 H, dq,  $J$  9.2 and 6.4, 5-H), 5.45 (1 H, d,  $J$  9.2, 4-H) and 6.38 (1 H, q,  $J$  7.0, 2-CH);  $\delta_C$  (GX) 12.5, 17.4 and 23.7 (3 q, 3 × CH<sub>3</sub>), 29.4 (t, 3-CH<sub>2</sub>), 64.8 (d, C-5), 114.7 and 116.5 (2 s, C-2 and C-1), 133.1 (d, C-4), 137.4 (s, C-3) and 146.0 (d, 2-CH);  $m/z$  165 ( $M^+$ , 13%), 151 (41), 147 (9), 136 (33), 123 (100), 108 (30), 95 (25) and 77 (23).

### Photoadditions of nitrile **1** with various amines

By analogy with the photoreaction of **16** with TEA, the nitrile **1** was irradiated with 30 equiv. of an amine in acetonitrile, affording the corresponding adducts. The results are summarized in Table 2.

#### 2-(3-Hydroxy-3,5,5-trimethylcyclohex-1-enyl)acrylonitrile

**22**, bp 150 °C at 0.3 mmHg (decomposition) (Found:  $M^+$ , 191.1309.  $C_{12}H_{17}NO$  requires  $M$ , 191.1310);  $\nu_{\max}$  ( $CHCl_3$ )/ $cm^{-1}$  3590, 3470 (O–H) and 2220 (C≡N);  $\delta_H$  (EX) 1.03 and 1.06 (6 H, 2 s, 2 × 5'-CH<sub>3</sub>), 1.36 (3 H, s, 3'-CH<sub>3</sub>), 1.57 and 1.73 (2 H, each d,  $J$  14.2, 4'-H<sub>2</sub>), 1.66 (1 H, br s, OH), 1.87 (1 H, dd,  $J$  16.5 and 2, 6'-H), 2.01 (1 H, d,  $J$  16.5, 6'-H), 5.88 and 5.93 (2 H, each s, 3-H<sub>2</sub>) and 6.22 (1 H, br s, 2'-H);  $\delta_C$  (EX) 27.8, 30.9 and 31.1 (3 q, 3 × CH<sub>3</sub>), 30.1 (s, C-5'), 38.0 (t, C-6'), 49.5 (t, C-4'), 69.1 (s, C-3'), 117.1 (s, C-1), 124.6 (s, C-2), 126.9 (t, C-3), 130.8 (s, C-1') and 135.3 (d, C-2');  $m/z$  191 ( $M^+$ , 38%), 176 (100), 149 (27), 139 (41), 134 (29), 120 (26) and 106 (23).

(*Z*)-2-(3-Hydroxy-3,5,5-trimethylcyclohex-1-enyl)pent-2-enitrile **23**, bp 130 °C at 0.15 mmHg (Found:  $M^+$ , 219.1620.  $C_{14}H_{21}NO$  requires  $M$ , 219.1623);  $\nu_{\max}$  (film)/ $cm^{-1}$  3410 (O–H) and 2220 (C≡N);  $\delta_H$  (EX) 1.02 and 1.06 (6 H, 2 s, 2 × 5'-CH<sub>3</sub>), 1.12 (3 H, t,  $J$  7.6, 5-H<sub>3</sub>), 1.34 (3 H, s, 3'-CH<sub>3</sub>), 1.56 (1 H, d,  $J$  14.2, 4'-H), 1.72 (1 H, m with dt-character,  $J$  14.2 and 1, 4'-H), 1.75 (1 H, br s, OH), 1.86 (1 H, dd,  $J$  16.5 and 1.7, 6'-H), 2.00 (1 H, m with dt-character,  $J$  16.5 and 1, 6'-H), 2.50 (2 H, quintet,  $J$  7.6, 4-H<sub>2</sub>), 6.09 (1 H, br s, 2'-H) and 6.38 (1 H, t,  $J$  7.6, 3-H);  $\delta_C$  (EX) 13.2 (q, C-5), 25.3 (t, C-4), 27.9, 30.9 and 31.3 (3 q, 3 × CH<sub>3</sub>), 30.1 (s, C-5'), 38.8 (t, C-6'), 49.6 (t, C-4'), 69.1 (s, C-3'), 115.7 and 117.3 (2 s, C-1 and C-2), 130.8 (s, C-1'), 132.8 (d, C-2') and 146.5 (d, C-3);  $m/z$  219 ( $M^+$ , 31%), 204 (100), 177 (16), 148 (9), 139 (20) and 134 (10).

### Photoaddition of nitrile **1** with TMSA **24**

**Preparative irradiation.** A solution of **1** ( $E:Z = 2:1$ ) (470 mg, 2.65 mmol) and TMSA **24** (4.22 g, 26.5 mmol) in acetonitrile (71.6 cm<sup>3</sup>) was irradiated with a low-pressure mercury lamp in a quartz test-tube (conversion 100%) for 4 h at RT. After removal of the solvent, a solution of the residue in  $CHCl_3$  (97 cm<sup>3</sup>) was stirred in the presence of silica gel (1.9 g) for 2 h at RT. After filtration, the filtrate was evaporated to give the residue, which was subjected to flash chromatography [ $SiO_2$ ; benzene–ethyl acetate (9:1)] to afford **3** (6%), **4** (23%), **22** (12%) and **25** (21%). †

(*Z*)-2-(3-Hydroxy-3,5,5-trimethylcyclohex-1-enyl)-3-(trimethylsilyl)acrylonitrile **25**, mp 87–89 °C (Found:  $M^+$ , 263.1696.  $C_{15}H_{25}NOSi$  requires  $M$ , 263.1705);  $\nu_{\max}$  (film)/ $cm^{-1}$  3440 (O–H) and 2220 (C≡N);  $\delta_H$  (EX) 0.29 [9 H, s, Si(CH<sub>3</sub>)<sub>3</sub>], 1.04 and 1.06 (6 H, 2 s, 2 × 5'-CH<sub>3</sub>), 1.36 (3 H, s, 3'-CH<sub>3</sub>), 1.56 and 1.73 (2 H, each d,  $J$  14.2, 4'-H<sub>2</sub>), 1.70 (1 H, br s, OH), 1.85 (1 H, dd,  $J$  16.5 and 2.0, 6'-H), 2.03 (1 H, m with d-character,  $J$  16.5, 6'-H), 6.24 (1 H, br s, 2'-H) and 6.57 (1 H, s, 3-H);  $\delta_C$  (EX) –1.38 [3 q, Si(CH<sub>3</sub>)<sub>3</sub>], 27.9, 30.2 and 30.9 (3 q, 3 × CH<sub>3</sub>), 30.2 (s, C-5'), 38.4 (t, C-6'), 49.6 (t, C-4'), 69.3 (s, C-3'), 117.2 (s, C-1), 129.6 and 130.8 (2 s, C-2 and C-1'), 134.4 (d, C-2') and 145.5 (d,

C-3);  $m/z$  263 ( $M^+$ , 7%), 248 (18), 145 (86), 130 (100), 203 (17), 139 (24), 84 (16) and 73 (62).

**Irradiations with various concentrations of TMSA **24** in various solvents.** Aliquots of a 0.0564 mol dm<sup>-3</sup> solution of nitrile **1** ( $E/Z = 2:1$ ) in acetonitrile containing varying amounts of TMSA **24** (5, 10, 15 and 30 equiv.) and MeOH were irradiated in quartz test-tubes under argon for 0.5 h at RT. Yields were determined by GLC analysis using octadecane as internal standard, after stirring of reaction mixture with  $SiO_2$  in  $CHCl_3$  for 2 h at RT, and results are shown in Table 3.

### Acknowledgements

We thank the staff of the Analysis Center of Meiji Pharmaceutical University for performing the elemental analysis and measurements of GX-NMR (Miss S. Yoshioka) and mass spectra (Miss T. Koseki). We are also grateful to Miss A. Iizuka for her technical assistance. This work was partially supported by a Special Grant from Meiji Pharmaceutical University

### References

- 1 G. J. Kavarnos and N. J. Turro, *Chem. Rev.*, 1986, **86**, 401; J. Mattay, *Synthesis*, 1989, 233; J. Mattay and M. Vondenhof, *Top. Curr. Chem.*, 1991, **159**, 219; A. Albini, E. Fasani and M. Mella, *Top. Curr. Chem.*, 1993, **168**, 143; G. Pandey, *Top. Curr. Chem.*, 1993, **168**, 175; F. Müller and J. Mattay, *Chem. Rev.*, 1993, **93**, 99; J. Cossy, *Bull. Soc. Chim. Fr.*, 1994, **131**, 344; U. C. Yoon, P. S. Mariano, R. S. Givens and B. W. Atwater, in *Advances in Electron Transfer Chemistry*, ed. P. S. Mariano, JAI Press, Stamford, 1994, vol. 4, p. 117; F. D. Lewis, in *Advances in Electron Transfer Chemistry*, ed. P. S. Mariano, JAI Press, Stamford, 1996, vol. 5, p. 1; J. Cossy and J. P. Pete, in *Advances in Electron Transfer Chemistry*, ed. P. S. Mariano, JAI Press, Stamford, 1996, vol. 5, p. 141.
- 2 R. C. Cookson, J. Hudec and N. A. Mirza, *Chem. Commun.*, 1968, 180; N. J. Pienta and J. E. McKimmey, *J. Am. Chem. Soc.*, 1982, **104**, 5501; D. W. Smith and N. J. Pienta, *Tetrahedron Lett.*, 1984, **25**, 915; N. J. Pienta, *J. Am. Chem. Soc.*, 1984, **106**, 2704; S. Bertrand, N. Hoffmann, S. Humbel and J. P. Pete, *J. Org. Chem.*, 2000, **65**, 8690.
- 3 Y. T. Jeon, C.-P. Lee and P. S. Mariano, *J. Am. Chem. Soc.*, 1991, **113**, 8847.
- 4 U. C. Yoon and P. S. Mariano, *Acc. Chem. Res.*, 1992, **25**, 233.
- 5 K. Ishii, M. Kotera and M. Sakamoto, *J. Chem. Soc., Chem. Commun.*, 1994, 2465.
- 6 K. Ishii, D. Gong, R. Asai and M. Sakamoto, *J. Chem. Soc., Perkin Trans. 1*, 1990, 855.
- 7 K. Ishii, M. Abe and M. Sakamoto, *J. Chem. Soc., Perkin Trans. 1*, 1987, 1937.
- 8 B. Frei, H. Eichenberger, B. von Wartburg, H. R. Wolf and O. Jeger, *Helv. Chim. Acta*, 1977, **60**, 2968.
- 9 J. Cossy, A. Bouzide, S. Ibhi and P. Aclinou, *Tetrahedron*, 1991, **47**, 7775.
- 10 A. Jończyk, M. Fedoryński and M. Mąkosza, *Tetrahedron Lett.*, 1972, 2395.
- 11 E. Hasegawa, M. A. Brumfield and P. S. Mariano, *J. Org. Chem.*, 1988, **53**, 5435.
- 12 E. Hasegawa, K. Ishiyama, T. Fujita, T. Kato and T. Abe, *J. Org. Chem.*, 1997, **62**, 2396.