

## Electrophilic Aromatic Reactivities *via* Pyrolysis of Esters. Part 21.<sup>1</sup> $\sigma^+$ Values for Thiazole: the High Polarisability of Thiazole, and the Effect of Hydrogen Bonding on the Reactivity of *N*-Containing Heterocycles

Ryan August, Carole Davis, and Roger Taylor\*


School of Chemistry and Molecular Sciences, University of Sussex, Brighton BN1 9QJ, Sussex

The rates of gas-phase elimination of acetic acid from 1-arylethyl acetates (aryl = thiazol-2-yl, thiazol-4-yl, thiazol-5-yl, or phenyl) have been measured between 633.2 and 698.5 K. The relative rates of pyrolysis at 650 K are: thiazol-2-yl, 0.271; thiazol-4-yl, 0.491; and thiazol-5-yl, 1.104; coupled with the previously determined  $\rho$  factor for the reaction ( $-0.61$  at 650 K), this leads to the corresponding  $\sigma^+$  values 0.93, 0.505, and  $-0.07$ . The positional reactivity order, and the reactivity relative to benzene, are correctly predicted by  $\pi$ -electron density calculations. The reactivity of each position is substantially less than in solvolysis of the corresponding 1-aryl-1-chloroethanes, in contrast to the reactivity of thiophene which is closely similar in both pyrolysis and solvolysis reactions. Thiazole is thus much more polarisable than thiophene, and hence particularly susceptible to demands for resonance stabilisation of the respective transition states, the demand being less in the pyrolysis. The difference between the reactivities of the 2-position of thiazole and the corresponding  $\alpha$ -position of thiophene (1.72 sigma units) is substantially greater than the difference between the reactivities of the 4-position of thiazole and the  $\beta$ -position of thiophene (0.885 sigma units). This reflects the high 2,3- vs. 3,4-bond orders, and consequent variation in the deactivation by the (*ortho*) nitrogen atom. Nitrogen deactivates the 4-position more than it deactivates the 5-position, as expected. By contrast, in solvolysis the 5-position is deactivated more than the 4-position. This latter anomaly is attributed to hydrogen bonding, which is attenuated when the probe group is adjacent to the ring nitrogen atom. This explanation also accounts for the anomalously low reactivity of the 3- and 4-position of pyridine in determination of the electrophilic reactivity *via* solvolysis, as compared with gas-phase data, whereas the reactivity of the 2-position is the same in both reactions.


In previous parts of this series we have used the pyrolysis of 1-arylethyl acetates (1), a reaction which proceeds *via* the formation of a partial cation at the 1-carbon atom, and for which  $\rho$  is  $-0.66$  at 600 K ( $-0.61$  at 650 K), to determine electrophilic substituent constants for heterocycles. The particular advantage of the reaction (the first to be applied to determining heterocyclic reactivities by the Brown side-chain carbocation technique) is the absence of solvent, so that there are no problems associated either with protonation or with hydrogen bonding; this outweighs the disadvantage of the amount of synthetic work involved as compared with a conventional electrophilic substitution. So far the electrophilic reactivities of all positions of pyridine,<sup>2</sup> furan,<sup>3</sup> thiophene,<sup>3</sup> quinoline,<sup>4</sup> isoquinoline,<sup>5</sup> and benzo[*b*]thiophene,<sup>6</sup> and the 2- and 3-position of benzo[*b*]furan<sup>6</sup> have been determined by this method. The results for pyridine and thiophene, relevant to this study, are shown in terms of  $\sigma^+$  values in Scheme 1.

The results for thiophene are in excellent agreement with those determined from a number of other reactions,<sup>7</sup> the  $\sigma^+$  values being at the lower end of the spectrum of values obtained (the variation of data from all studies is *ca.* 0.05 sigma units), which reflects the smaller demand for resonance in the elimination as compared with some other reactions. The pyridine results are in excellent agreement with theoretical predictions,<sup>2</sup> but disagree with those from more recent studies [involving solvolysis of 1-aryl-1-chloropropanes (2)], which give the  $\sigma^+$  values shown in Scheme 2.<sup>8,9</sup>

It is noteworthy that whereas the results for the 2-position in gas-phase and solution studies are in good agreement (and in the latter show the smallest variation amongst the four media used), the values for the 3- and 4-position in the solution



Scheme 1.  $\sigma^+$  Values from pyrolysis of 1-arylethyl acetates



Scheme 2.  $\sigma^+$  Values from solvolysis of 1-aryl-1-chloropropanes (the ranges of values for pyridine arise from a range of solvents)

studies are both too positive, relative to the gas phase, by *ca.* 0.25 sigma units. We have previously suggested that this arises because of the very strong hydrogen bonding to which pyridine is subject, which provides additional electron withdrawal from the ring.<sup>10</sup> (The hydrogen chloride by-product in the solvolysis could also partially protonate the nitrogen.) When however the 1-chloropropyl substituent is at the 2-position, the nitrogen may be sufficiently shielded to inhibit hydrogen bonding (particularly in the transition state where the side-chain methyl



**Table.** Pyrolysis of compounds ArCH(Me)OAc

Ar	T/K	$10^3k/s^{-1}$	$\log(A/s^{-1})$	$E/kJ\ mol^{-1}$	Corr. coefft.	$k/k_0$ at 650 K
Phenyl	633.2	3.83	12.344	178.68	0.999 78	0
	655.1	12.3				
	671.2	27.8				
	688.1	58.4				
	696.5	83.5				
Thiazol-2-yl	633.2	1.03	12.444	187.09	0.999 84	0.271
	655.1	3.35				
	664.5	5.32				
	671.2	7.68				
	688.1	16.9				
Thiazol-4-yl	636.7	2.29	12.397	183.59	0.999 77	0.491
	651.4	5.02				
	664.5	9.69				
	678.8	18.7				
	698.5	50.1				
Thiazol-5-yl	633.2	4.30	12.240	176.97	0.999 78	1.104
	652.5	12.3				
	655.1	13.2				
	664.5	20.9				
	671.2	29.3				
	688.1	62.7				
	696.5	93.4				

situation therefore parallels exactly that found in pyridine. We do not rule out the possibility of intramolecular hydrogen bonding between nitrogen and the side-chain, which reduces opportunities for intermolecular bonding with the solvent in the solution studies.

### Experimental

**Kinetic Studies.**—The apparatus and general kinetic technique have been described in earlier parts of this series. The present work was carried out using a new reactor, the most significant feature of which is the gold-plated interior surface which minimises surface-catalysed reactions. Full details are described in a recent paper.<sup>18</sup>

Each compound showed excellent and reproducible kinetic behaviour, first-order to > 95% of reaction, and excellent Arrhenius plots (see Table for correlation coefficients). The reaction stoichiometry was  $2.0 \pm 0.03$  in  $10 \times t_{1/2}$  and only vinylthiazoles and acetic acid were detectable as reaction products. The vinylthiazoles are fairly stable; they underwent only very slow secondary decomposition which did not significantly interfere with the primary decomposition. To ensure maximum rigour in analysis of the data, we carried out new runs on 1-phenylethyl acetate, rather than use our existing data; this avoided any possible errors due to thermocouple ageing etc.

**1-(Thiazol-2-yl)ethyl Acetate.**—1-(Thiazol-2-yl)ethyl alcohol was prepared in 30% yield by the literature method,<sup>13</sup> except that 2-bromothiazole was the starting reagent. The alcohol (1 vol.) was heated under reflux with acetic anhydride (5 vol.) and pyridine (10 vol.) during 2 h. Normal work-up and fractional distillation yielded 1-(thiazol-2-yl)ethyl acetate (80%), b.p. 65 °C at 1.0 mmHg;  $n_D^{20}$  1.4970 (Found: C, 49.2; H, 5.3; N, 8.3.  $C_7H_9NO_2S$  requires C, 49.1; H, 5.3; N, 8.2%);  $\delta(CDCl_3)$  7.66 (1 H, d, H-5), 7.22 (1 H, d, H-4), 6.09 (1 H, q, CH), 2.02 (3 H, s,  $COCH_3$ ), and 1.62 (3 H, d,  $CH_3$ ).

**1-(Thiazol-4-yl)ethyl Acetate.**—4-Formylthiazole was prepared according to the literature method,<sup>13</sup> but the following modifications were incorporated to improve the yield.

(i) A substantially higher yield of 1,1,3-tribromoacetone can be obtained from the bromination of acetone,<sup>19</sup> if the crude product is fractionally distilled prior to recrystallisation from light petroleum, thereby removing most of the by-products.

(ii) The decomposition of 4-dibromomethyl-4,5-dihydro-4-hydroxythiazole hydrobromide with sulphuric acid must be continued until the solution is clear, indicating that all the hydrogen bromide has been evolved.

(iii) Heating 4-dibromomethylthiazole with aqueous ethanol does not give 4-formylthiazole, as described in the literature,<sup>20</sup> but instead a ca. 50:50 mixture of this product and the diethyl acetal. This mixture was converted into the aldehyde  $\delta(CDCl_3)$  10.13 (1 H, s, CHO), 8.98 (1 H, d, H-2), and 8.31 (1 H, d, H-5) by heating with 5% hydrochloric acid during 5 min.

1-(Thiazol-4-yl)ethyl alcohol was prepared in 80% yield, according to the literature method;<sup>13</sup>  $\delta(CDCl_3)$  8.59 (1 H, s, H-2), 7.08 (1 H, s, H-5), 4.82 (1 H, q, CH), and 1.35 (3 H, d,  $CH_3$ ). Acetylation as above then gave 1-(thiazol-4-yl)ethyl acetate (93%), b.p. 68 °C at 1 mmHg (Found: C, 49.3; H, 5.1; N, 8.4%);  $\delta(CDCl_3)$  8.79 (1 H, s, H-2), 7.27 (1 H, s, H-5), 6.15 (1 H, q, CH), 2.09 (3 H, s,  $CH_3$ ), and 1.65 (3 H, d,  $CH_3$ ).

**1-(Thiazol-5-yl) Acetate.**—1-(Thiazol-5-yl)ethyl alcohol was prepared from 2-chlorothiazole according to the literature method.<sup>13</sup> Acetylation as above gave 1-(thiazol-5-yl)ethyl acetate (55%), b.p. 33 °C at 0.15 mmHg (Found: C, 48.8; H, 5.4; N, 8.05%);  $\delta(CDCl_3)$  8.86 (1 H, s, H-5), 7.90 (1 H, s, H-2), 6.32 (1 H, q, CH), 2.02 (3 H, s,  $CH_3$ ), and 1.64 (3 H, d,  $CH_3$ ).

### References

- Part 20, M. A. Hossaini and R. Taylor, *J. Chem. Soc., Perkin Trans. 2*, 1982, 187.
- R. Taylor, *J. Chem. Soc.*, 1962, 4881.
- R. Taylor, *J. Chem. Soc. B*, 1968, 1397.
- R. Taylor, *J. Chem. Soc. B*, 1971, 2382.
- E. Glyde and R. Taylor, *J. Chem. Soc., Perkin Trans. 2*, 1975, 1783.
- H. B. Amin and R. Taylor, *J. Chem. Soc., Perkin Trans. 2*, 1978, 1053.
- R. Taylor, *J. Chem. Soc. B*, 1970, 1364.
- T. J. Broxton, G. L. Butt, L. W. Deady, S. H. Toh, R. D. Topsom, A. Fischer, and N. W. Morgan, *Can. J. Chem.*, 1973, 51, 1620.

- 9 D. S. Noyce, J. A. Virgilio, and B. Bartman, *J. Org. Chem.*, 1973, **38**, 2657.
- 10 R. Taylor, 'Specialist Periodical Report on Aromatic and Heteroaromatic Chemistry,' The Chemical Society, 1975, vol. 3, p. 222.
- 11 D. S. Noyce and J. A. Virgilio, *J. Org. Chem.*, 1973, **38**, 2660.
- 12 H. B. Amin and R. Taylor, *J. Chem. Soc., Perkin Trans. 2*, 1979, 624.
- 13 D. S. Noyce and S. A. Fike, *J. Org. Chem.*, 1973, **38**, 3316.
- 14 R. Taylor, 'Thiophene and Its Derivatives,' ed. S. Gronowitz, in 'Chemistry of Heterocyclic compounds,' Wiley, Chichester, in the press.
- 15 R. Phan-Tan-Luu, L. Bouscasse, E. Vincent, and J. Metzger, *Bull. Soc. Chim. Fr.*, 1969, 1149.
- 16 R. Taylor and G. G. Smith, *Tetrahedron*, 1963, **19**, 937.
- 17 C. Eaborn and D. R. M. Walton, *J. Organomet. Chem.*, 1965, **3**, 169; C. Eaborn, Z. S. Salih, and D. R. M. Walton, *J. Organomet. Chem.*, 1972, **36**, 47.
- 18 N. Al-Awadi and R. Taylor, *J. Chem. Soc., Perkin Trans. 2*, in the press.
- 19 C. Rappe, *Arkiv. Kemi*, 1963, **21**, 503.
- 20 H. Beganz and J. Rüger, *Chem. Ber.*, 1968, **101**, 3872.

Received 18th November 1985; Paper 5/2020