

# Kinetics and mechanism of phosphate-catalyzed hydrolysis of benzoate esters: comparison with nucleophilic catalysis by imidazole and *o*-iodosobenzoate

2 PERKIN

Omar A. El Seoud,<sup>\*a</sup> Marie-Françoise Ruasse<sup>b</sup> and Wagner A. Rodrigues<sup>a</sup>

<sup>a</sup> Instituto de Química, Universidade de São Paulo, C.P. 26077, 05513-970, São Paulo, S.P. Brazil. E-mail: elseoud@iq.usp.br; Fax: +55-11-3091-3874

<sup>b</sup> Institut de Topologie et de Dynamiques des Systèmes, Université de Paris VII-CNRS, 1 Rue Guy de la Brosse, Paris, F-75005, France

Received (in Cambridge, UK) 18th March 2002, Accepted 4th April 2002

First published as an Advance Article on the web 23rd April 2002

Phosphate-catalyzed hydrolysis of 2,4-dinitrophenyl 4-X-benzoate, and 3- or 4-Y-phenyl 3,5-dinitrobenzoates, where X and Y are substituents, has been studied spectrophotometrically. The following conclusions are based on catalytic rate constants, solvent kinetic isotope effect, detection of a mixed anhydride by FTIR, and application of the Hammett equation: (i) the catalytically active species is  $\text{HPO}_4^{2-}$ , (ii) the mechanism of catalysis is nucleophilic, the reaction proceeds *via* the irreversible formation of acyl phosphate, (iii) the formation of a tetrahedral species by attack of the  $\text{HPO}_4^{2-}$  on the ester CO group is rate limiting. Comparison of these results with our previous data on hydrolysis of benzoate esters with imidazole, or *o*-iodosobenzoate anion showed that: (i) the catalytic efficiency observed, *o*-iodosobenzoate > imidazole > phosphate, is due to a combination of steric and solvation effects, and the enhancement of the nucleophilicity of the former catalyst by the  $\alpha$ -effect, (ii) the nonlinear Brønsted-type plot between log (catalytic constant) and  $\text{p}K_a$  of the leaving 4-Y-phenol indicates that the rate-limiting step for *o*-iodosobenzoate changes as a function of changing Y, (iii) the reaction rate-limiting steps are different for the above-mentioned nucleophiles.

## Introduction

Acyl-transfers are important reactions in chemistry and enzymology.<sup>1-5</sup> Previously, we have studied the hydrolysis of benzoate esters of the general structure  $\text{X}-\text{C}_6\text{H}_4-\text{CO}_2-\text{C}_6\text{H}_4-\text{Y}$ , where X and Y are substituents, with two powerful nucleophiles, namely, imidazole, Imz, and *o*-iodosobenzoate anion, IBA. We now report on phosphate-catalyzed hydrolysis of the same benzoate esters. Phosphate acts as a general base in hydrolysis of esters with poor leaving groups (LG), *e.g.*, ethyl dichloroacetate.<sup>2</sup> The nucleophilic-catalysis pathway is favored, however, for esters with better LG, *e.g.*, phenyl carboxylates. Here we show that this is indeed the case for phosphate-catalyzed hydrolysis of the following esters: (i) 2,4-dinitrophenyl 4-X-benzoate (hereafter designated as series I), where X = methyl (I-a), hydrogen (I-b), chloro (I-c), cyano (I-d) and nitro (I-e), respectively, and 3- or 4-Y-phenyl 3,5-dinitrobenzoates (hereafter designated as series II), where Y = 4-methyl (II-a), hydrogen (II-b), 3-cyano (II-c), 4-cyano (II-d) and 4-nitro (II-e), respectively.

We show that:  $\text{HPO}_4^{2-}$  is the active catalytic species; the mechanism of catalysis is nucleophilic; the rate limiting step for

the first part of the reaction is attack of the catalyst on the ester CO group. We also compare nucleophilic catalysis by  $\text{HPO}_4^{2-}$ , Imz, and IBA.


## Experimental

### Synthesis


Reagents and solvents were purchased from Aldrich and Merck, and were purified as described elsewhere.<sup>6</sup> Series I was that available from a previous study,<sup>4b</sup> whereas series II was synthesized by reacting 3,5-dinitrobenzoyl chloride with the appropriate phenoxide under phase-transfer conditions.<sup>4a</sup> Esters II-a, II-b and II-e gave melting points that agree with literature values,<sup>7</sup> whereas esters II-c (mp 170–172 °C) and II-d (mp 225–226 °C) gave satisfactory elemental analysis, and the expected <sup>1</sup>H NMR spectra (acetone-*d*<sub>6</sub>, Bruker AC-200).

Tetrabutylammonium di- and/or mono-hydrogenphosphate were prepared by reacting phosphoric acid (85%) with aqueous tetrabutylammonium hydroxide (40%). The final solution pH was adjusted to 4.3 or 7.4 for the former and the latter compound, respectively. Dry salts were obtained by lyophilization.

The mixed carboxylic-phosphoric anhydrides were synthesized *in situ*,<sup>8b</sup> by reacting 1.0 mol of 4-X-benzoyl chloride (X = CN and NO<sub>2</sub>) with 1.05 mol of tetrabutylammonium dihydrogenphosphate in anhydrous acetonitrile, at room temperature. Reaction progress was followed by monitoring the disappearance of the ester  $\nu_{\text{C=O}}$  band and the appearance of the carbonyl group of the mixed anhydride (Perkin-Elmer 1750 FTIR spectrophotometer, 0.2 mm ClearTran cell, Wilmad Glass). We have obtained an excellent linear correlation (correlation coefficient,  $r = 0.997$ ) between  $\nu_{\text{C=O}}$  of these mixed anhydrides, 1700 (X = H),<sup>8c</sup> 1733 (X = 4-CN), 1741  $\text{cm}^{-1}$  (X = NO<sub>2</sub>) and the Hammett  $\sigma_p$  value of X. The same IR equipment and cell were employed in the detection of the produced acyl phosphoric



General structure of series I



General structure of series II

anhydride, *vide infra*, under the following experimental conditions: reaction medium, 65% CD<sub>3</sub>CN plus 35% of 0.1 M ((C<sub>4</sub>H<sub>9</sub>)<sub>4</sub>N)<sub>2</sub>HPO<sub>4</sub> in D<sub>2</sub>O, pD = 7.8, [I-e] = 0.05 M. The IR cell was kept at *ca.* 40 °C, and the spectrum of the reaction mixture was recorded as a function of time. The spectra of authentic samples were recorded in 65% CD<sub>3</sub>CN plus 35% D<sub>2</sub>O.

### Kinetic measurements

Kinetic measurements were carried out with Zeiss PM6KS or Beckman DU-70 Uv-vis spectrophotometers, fitted with thermostated cell holders whose temperature was kept constant at  $\pm 0.05$  °C (4000A digital thermometer, Yellow Springs Instruments). All runs were carried out in triplicate, under pseudo-first-order conditions, and were initiated by injecting the ester solution in anhydrous acetonitrile into the phosphate buffer, under the following experimental conditions: solvent, 10% CH<sub>3</sub>CN in water (v/v, N<sub>2</sub>-saturated); ionic strength, 1.0 M, maintained with NaCl; [phosphate buffer]<sub>total</sub> = 0.05 to 0.25 M; reaction pH = 7.8 to 8.2. We have verified that observed rate constants,  $k_{\text{obs}}$ , were independent of [ester] in the range  $1-6 \times 10^{-5}$  M, and used final [ester] =  $1-3 \times 10^{-5}$  M. The reaction mixture was homogenized with a micro-stirrer (type 338.004, Hellma). Reaction progress was followed by monitoring liberation of the appropriate phenol as a function of time, at the following  $\lambda$ : 362 nm, series I; 224, 227, 226, 270, and 400 nm for esters II-a to II-e, respectively. Plots of log (absorbance) *versus* time were rigorously linear over more than five half-lives, their slopes offered  $k_{\text{obs}}$ . For the same run, the relative standard deviation in  $k_{\text{obs}}$ , *i.e.*, ((standard deviation/ $k_{\text{obs}}$ )  $\times$  100) was  $\leq 0.2\%$ ; for triplicate runs, the difference between  $k_{\text{obs}}$  was  $\leq 1\%$ .


## Results and discussion

### Nature of the catalytically-active species

Independent of the reaction mechanism, the catalytically active species can be H<sub>2</sub>PO<sub>4</sub><sup>-</sup>, or HPO<sub>4</sub><sup>2-</sup>, or both. Hydrolysis of ester I-e was studied at 40 °C, at three pHs, where the buffer ratios, *i.e.*, [H<sub>2</sub>PO<sub>4</sub><sup>-</sup>]/[HPO<sub>4</sub><sup>2-</sup>] were: 80 : 20; 50 : 50; and 20 : 80, respectively. Second order, *i.e.*, catalytic rate constants,  $k_c$ , were calculated from plots of  $k_{\text{obs}}$  *versus* [H<sub>2</sub>PO<sub>4</sub><sup>-</sup>] and/or [HPO<sub>4</sub><sup>2-</sup>], respectively, giving: 0.004, 0.019, and 0.088 M<sup>-1</sup>s<sup>-1</sup> (H<sub>2</sub>PO<sub>4</sub><sup>-</sup>); 0.019, 0.020, and 0.020 M<sup>-1</sup>s<sup>-1</sup> (HPO<sub>4</sub><sup>2-</sup>). The constancy of the latter  $k_c$  indicates that HPO<sub>4</sub><sup>2-</sup> is the catalytically active species.

### Nature of catalysis by the buffer: general-base or nucleophilic?

Catalysis by the buffer (B/BH, B = HPO<sub>4</sub><sup>2-</sup>) can be general base (Scheme 1) or nucleophilic (Scheme 2). In all subsequent dis-




cussion, we concentrate on the part of the reaction that leads to formation of the phenoxide anion. The reason is that the nucleophilic catalysis pathway, if it applies, involves the formation of acyl phosphoric anhydride (*Int2*, Scheme 2) whose (slow) hydrolysis,<sup>8b</sup> is outside the scope of the present work.

As shown, we take the view that the reaction mechanism is stepwise,<sup>9-11</sup> involving the formation of a tetrahedral intermediate, *e.g.* *Int1* in Scheme 2, as given elsewhere for the reaction of acetate and benzoate esters with nucleophiles.<sup>9</sup>

The main difference between Schemes 1 and 2 is the formation of *Int2*, a carboxylic-phosphoric anhydride in the nucleophilic catalysis pathway. Formation of *Int2* does not involve direct participation of water, consequently, a small (secondary) kinetic solvent isotope effect is expected. Indeed,  $k_{\text{CH}_3\text{O}}/k_{\text{CD}_3\text{O}}$  for hydrolysis of esters I-d and I-e, at 40 °C were found to be 1.4 and 1.1, respectively, in agreement with a nucleophilic catalysis mechanism. For comparison,  $k_{\text{CH}_3\text{O}}/k_{\text{CD}_3\text{O}}$  were found to be (catalyst, ester, respectively): 1.4 and 1.3 (IBA, 4-nitrophenyl 4-nitrobenzoate, I-e), 1.3, 1.0, and 3.0 (Imz, I-e, 4-nitrophenyl acetate, ethyl dichloroacetate).<sup>14</sup> For the last ester, the catalysis is *general base*.<sup>1</sup>


We sought direct experimental evidence for the formation of *Int2*. FTIR was employed to follow changes in the  $\nu_{\text{C=O}}$  region for the phosphate-catalyzed hydrolysis of ester I-e, Fig. 1. Part



**Fig. 1** IR spectra of the  $\nu_{\text{C=O}}$  region. Part A shows peaks of authentic 2,4-dinitrophenyl 4-nitrobenzoate, Ester; 4-nitrobenzoyl phosphate, Intermediate; an equimolar mixture of potassium salts of 4-nitrobenzoic acid and 2,4-dinitrophenol, Products. Part B shows the dependence on time of the spectrum of I-e in the presence of phosphate buffer in CD<sub>3</sub>CN–D<sub>2</sub>O solvent, see Experimental for details.

A of this figure shows  $\nu_{\text{C=O}}$  for the following *authentic samples* in 65% CD<sub>3</sub>CN in D<sub>2</sub>O: ester I-e,  $\nu_{\text{C=O}} = 1762$  cm<sup>-1</sup>, the *initial* product if the catalysis were nucleophilic, *i.e.*, 4-nitrobenzoyl phosphate,  $\nu_{\text{C=O}} = 1741$  cm<sup>-1</sup>, and the *final* reaction products, *i.e.*, an equimolar mixture of the potassium salts of 4-nitrobenzoic acid and 2,4-dinitrophenol,  $\nu_{\text{C=O}} = 1722$  and 1709 cm<sup>-1</sup>, respectively. Part B of Fig. 1 shows changes in the same spectral region for phosphate-catalyzed hydrolysis of ester I-e after 10 minutes, 6 h and 18 h, respectively. A trace of *Int2* can be detected after 10 minutes, and is clearly seen after 6 hours, whereas the final reaction products are the only species present after 18 hours.<sup>12</sup> These changes show clearly the formation, and subsequent decomposition of *Int2*, 4-nitrobenzoyl phosphoric anhydride, in agreement with a nucleophilic catalysis mechanism.

Nucleophilic catalysis by phosphate buffers in aqueous solutions has been demonstrated for, *e.g.*, hydrolysis of 4-nitrophenyl acetate,<sup>13a</sup> phosphoryl transfer from phosphorylated pyridine monoanions,<sup>13b</sup> hydrolysis of guanosine 5'-phosphate 2-methylimidazole,<sup>13c</sup> hydrolysis of bis(4-nitrophenyl) methyl phosphonate,<sup>13d</sup> and esterolysis of 2,4-dinitrophenyl acetate in the presence of micellar benzyltrimethylammonium *n*-decyl phosphate.<sup>13e</sup> The first study had a very limited scope, and the




**Scheme 2** Nucleophilic catalysis by phosphate buffer.

question of the catalytically-active species has not been addressed.<sup>13a</sup> The second and third reactions are catalyzed by both buffer species, albeit more efficiently by  $\text{HPO}_4^{2-}$ ,<sup>13b,c</sup> the fourth reaction is catalyzed by  $\text{HPO}_4^{2-}$ , whereas the last reaction shows the formation of the mixed anhydride in a micellar solution.

### Details of the reaction mechanism

We now address the relative stabilities of *Int1* and *Int2*, respectively. Uv-vis spectra recorded during the hydrolysis of several esters showed sharp isosbestic points that lie between peaks of the ester and the phenol produced, e.g., at 295 and 242 nm for esters I-e and II-d, respectively. Additionally, identical rate constants were obtained when hydrolysis of I-c was studied by following either the disappearance of the ester, or the appearance of 4-chlorophenol. Both observations indicate that *Int1* does not accumulate, so that Scheme 2 can be simplified, as depicted in Scheme 3.



**Scheme 3** Simplified scheme for nucleophilic catalysis by phosphate buffer.

An obvious experiment to judge the reversibility of the reaction, *i.e.* whether  $k_{-c}$  is relevant, is to determine the (common ion) effect of the leaving phenol, added at the *start* of the run, on  $k_{\text{obs}}$ . Because  $\text{p}K_{\text{a}}$  of  $\text{HPO}_4^{2-}$  and the leaving phenol (e.g., 4-nitrophenol) are practically the same, their concentrations should be comparable ( $\geq 0.1 \text{ mol L}^{-1}$ ), in order to clearly see the effect of reversibility on  $k_{\text{obs}}$ . Experimentally, however, this phenol concentration cannot be employed because of its strong Uv-vis absorption.

Another approach is to investigate the reverse reaction, *i.e.*, that of *authentic* mixed anhydride with an appropriate phenoxide ion, in presence of a buffer, HB/B, whose B is a poor general base, and does not act as a nucleophile. We studied the reaction of equal amounts ( $4 \times 10^{-5} \text{ M}$ ) of 4-nitrobenzoyl phosphate and 2,4-dinitrophenolate anion (to give ester I-e), in the presence of  $0.05 \text{ mol L}^{-1}$  *N*-methylmorpholine buffer, at  $\text{pH} = 6.8$ . The Uv-vis spectrum of this mixture did not change as a function of time, during one hour. The same reaction was also investigated by FTIR, under the following conditions: solvent 65%  $\text{CD}_3\text{CN}$  in  $\text{D}_2\text{O}$  buffered with  $0.05 \text{ M}$  *N*-methylmorpholine ( $\text{p}D = 7.2$ ); [4-nitrobenzoyl phosphate] =  $0.025 \text{ M}$ ; [2,4-dinitrophenolate anion] =  $0.05 \text{ M}$ ,  $0.05 \text{ mm}$  path length ClearTran cell. The  $1850\text{--}1650 \text{ cm}^{-1}$  spectral region was periodically scanned during two hours; no ester appeared (at *ca.*  $1762 \text{ cm}^{-1}$ ). Both experiments clearly indicate that: (i) the mixed anhydride, *i.e.*, *Int2*, is stable under these reaction condi-

tions,<sup>8b,12</sup> (ii) the step of concern, *i.e.*, that leading to the formation of *Int2* plus the phenoxide ion is irreversible. That is, we need only be concerned with  $k_c$  of Scheme 3.

Table 1 shows the dependence of  $k_c$  of series I on temperature, along with the corresponding activation parameters. The Hammett plot ( $\log k_c$  versus  $\sigma_{\text{p}}$ )<sup>14</sup> gave an excellent straight line ( $r = 0.9981$ , sum of the squares of the residues,  $\Sigma Q = 0.068$ ) with  $\rho_{\text{X}} = 2.23$ . The reaction of series II was studied only at  $50^\circ\text{C}$ ; the following  $10^3 k_c$ ,  $\text{M}^{-1}\text{s}^{-1}$ , were calculated: 5.01, 7.08, 22.2, 51.3, and 97.3 for II-a ( $\text{Y} = 4\text{-CH}_3$ ), II-b ( $\text{Y} = \text{H}$ ), II-c ( $\text{Y} = 3\text{-CN}$ ), II-d ( $\text{Y} = 4\text{-CN}$ ) and II-e ( $\text{Y} = 4\text{-NO}_2$ ), respectively. For series II, the Hammett plot was satisfactory *only* when  $\sigma_{\text{p}}^-$  was employed ( $r = 0.9992$ ,  $\Sigma Q = 0.026$ ), giving  $\rho_{\text{Y}} = 0.89$ . The implications of these  $\rho$  for the rate-limiting step will be discussed in the next paragraph.

### Comparison of nucleophilic catalysis by phosphate, imidazole and the *o*-iodosobenzoate anion

Comparison of hydrolysis of these benzoate esters by  $\text{HPO}_4^{2-}$ , Phos, Imz, and IBA is instructive because: (i) all three reactions proceed by nucleophilic catalysis, *via* an irreversibly formed acylated catalyst, e.g., *Int2* for Phos, (ii) although the three catalysts have comparable  $\text{p}K_{\text{a}}$  (7.53, 7.05, and 7.20, for IBA, Imz, and Phos, respectively) their nucleophilic centers are different, nitrogen for Imz, and anionic oxygen for phosphate and IBA, respectively. An *outline* of the reactions of concern is depicted in Scheme 4 for an ester of the general formula  $\text{X-C}_6\text{H}_4\text{-CO}_2\text{C}_6\text{H}_4\text{-Y}$ , whereas relative rates, and differences between the activation parameters are listed in Table 2. Note that series I (2,4-dinitrophenyl 4-X-benzoates) is the *same* for the three catalysts, whereas series II for IBA and Imz, 4-Y-phenyl 2,4-dinitrobenzoates, is *different* from that employed with Phos, 3- or 4-Y-phenyl 3,5-dinitrobenzoates:

Imz and, specially, IBA are much more efficient nucleophiles than Phos (Table 2). This is due, in part, to differences in steric interactions, and of properties, e.g., solvation, of the nucleophiles. Steric hindrance to the attack on the ester CO group (to give *Int1*) is relevant to reactivity because of the presence of the *o*-nitro group in the phenol moiety.<sup>15</sup> This effect is specially small for IBA due to the favorable disposition of its nucleophilic (oxygen) center, which protrudes unhindered from the heterocyclic five-membered ring (the O-I-O<sup>-</sup> bond angle is  $165^\circ$ ).<sup>4b</sup> This, in part, is responsible for the favorable entropy of activation of the IBA-catalyzed reaction, as manifested by the corresponding *negative*  $T\Delta\Delta S^\ddagger$ , Table 2. Heat of solvation,  $\Delta H_{\text{solv}}^\circ$ , of the nucleophile contributes to the activation enthalpy since it must be de-solvated, at least partially, before attacking the ester CO group.  $\Delta H_{\text{solv}}^\circ$  are 17.9,<sup>4b</sup> 5.6<sup>16a</sup> and 250  $\text{kcal mol}^{-1}$ <sup>16b</sup> for IBA, Imz, and Phos, respectively. These data agree with the *positive* (*i.e.* favorable)  $\Delta\Delta H^\ddagger$  of Imz, and the fact that  $\Delta\Delta H^\ddagger$  for Imz is larger than the corresponding value for IBA. The higher  $\Delta H_{\text{solv}}^\circ$  of the latter nucleophile is more than compensated for by its enhanced nucleophilicity, due to the  $\alpha$ -effect.<sup>3</sup>

We now compare the rate limiting steps, RLS, involved. Our data indicate that these hydrolyses follow a stepwise mechanism, with RL formation, or breakdown of the corresponding

**Table 1** Dependence of the catalytic rate constants,  $k_c$ , on temperature and activation parameters for phosphate-catalyzed hydrolysis of ester series I, 2,4-dinitrophenyl 4-X-benzoates<sup>a</sup>

$T/^\circ\text{C}$	$10^4 k_c/\text{M}^{-1}\text{s}^{-1}$				
	X = CH <sub>3</sub>	X = H	X = Cl	X = CN	X = NO <sub>2</sub>
25.0	0.55	1.62	4.28	44.9	86.8
35.0	1.41	4.12	10.90	115.0	225.1
45.0	3.41	8.31	22.80	228.0	423.0
55.0	7.80	22.0	56.60	486.0	886.0


	Activation parameters <sup>b</sup>				
	X = CH <sub>3</sub>	X = H	X = Cl	X = CN	X = NO <sub>2</sub>
$\Delta H^\ddagger/\text{kcal mol}^{-1}$	16.6	16.0	15.9	14.6	14.2
$\Delta S^\ddagger/\text{cal K}^{-1}\text{mol}^{-1}$	-22.4	-22.3	-20.7	-20.1	-20.0
$\Delta G^\ddagger/\text{kcal mol}^{-1}$	23.3	22.6	22.1	20.6	20.2

<sup>a</sup> Reaction conditions: solvent, 10% acetonitrile in water, v/v; ionic strength = 1.0 M; [buffer]<sub>total</sub> = 0.05 to 0.25 M; reaction pH = 7.8 to 8.2. <sup>b</sup> The uncertainties in the activation parameters are  $\pm 0.1 \text{ kcal mol}^{-1}$  ( $\Delta H^\ddagger$  and  $\Delta G^\ddagger$ ) and  $\pm 0.5 \text{ cal K}^{-1}\text{mol}^{-1}$  ( $\Delta S^\ddagger$ ).

**Table 2** Comparison of the hydrolysis of series I, 2,4-dinitrophenyl 4-X-benzoates, at 25 °C, catalyzed by *o*-iodosobenzoate anion, IBA, imidazole, Imz, and HPO<sub>4</sub><sup>2-</sup>, Phos

X	$k_{c\text{-IBA}}/k_{c\text{-Phos}}^a$	$k_{c\text{-Imz}}/k_{c\text{-Phos}}^b$	$\Delta\Delta H^\ddagger^c$		$\Delta\Delta S^\ddagger^c$		$T\Delta\Delta S^\ddagger^c$	
			IBA	Imz	IBA	Imz	IBA	Imz
CH <sub>3</sub>	97363	4781	5.5	8.3	-4.1	11.4	-1.2	3.4
H	60864	2988	4.9	7.1	-5.4	7.7	-1.6	2.3
Cl	55070	2782	5.4	6.9	-3.9	7.3	-1.2	2.2
CN	39572	1604	4.5	6.3	-5.6	6.8	-1.7	2.0
NO <sub>2</sub>	31155	1072	4.3	5.3	-5.8	4.2	-1.7	1.3

<sup>a</sup> Reaction conditions: Phos,  $\mu = 1.0$ , solvent = 10% acetonitrile in water (v/v); IBA,  $\mu = 0.2$ , solvent = 14% acetonitrile in water (v/v).<sup>4b</sup> <sup>b</sup> Reaction conditions: Imz,  $\mu = 0.06$ , solvent = 10% acetonitrile in water (v/v).<sup>4a</sup> <sup>c</sup>  $\Delta\Delta H^\ddagger = \Delta H^\ddagger_{\text{Phos}} - \Delta H^\ddagger_{\text{Imz}}$ , kcal mol<sup>-1</sup>;  $\Delta\Delta S^\ddagger = \Delta S^\ddagger_{\text{Phos}} - \Delta S^\ddagger_{\text{Imz}}$ , cal K<sup>-1</sup> mol<sup>-1</sup>;  $T\Delta\Delta S^\ddagger$  is in kcal mol<sup>-1</sup>.

**Scheme 4** Nucleophilic catalysis by Phos, Imz, and IBA buffers, respectively.

tetrahedral intermediate (*Int1*, see Scheme 2), depending on the nucleophile and/or the strength of the LG. Structures of the relevant (*i.e.*, rate-limiting) transition states, TS, are shown in Fig. 2, followed by corroborating evidence.

The extents of bond formation (between the nucleophile and the ester CO group) and bond breaking (between the CO group and LG) can be probed by considering: Hammett  $\rho_X$  (series I) and  $\rho_Y$  (series II), Brønsted-type plots ( $\log k_c$  versus  $\text{p}K_a$  of the leaving phenol,  $\beta_{\text{LG}}$ ), and plots of  $\log k_c$  of the different nucleophiles. Examination of these data reveals the following:

(i) For Phos, the large magnitude of  $\rho_X - \rho_Y$  ( $= 1.34$ ) clearly indicates that the RLS is bond formation. For IBA and Imz, however, the corresponding differences are too small ( $\leq 0.12$ ) to allow an unambiguous conclusion with regard to the nature of the RLS. Therefore, we draw our conclusions mainly from the

above-mentioned Brønsted-type plots,  $\log k_c$ - $\log k_c$  plots, and the activation parameters.

(ii) A Brønsted-type plot between  $\log k_c$  IBA and  $\text{p}K_a$  of the leaving phenol exhibits a clear break (Fig. 3A), with  $\beta_{\text{LG}}$  changing from  $-0.78$  for the poor LG to  $-0.26$  for their good counterparts. Additionally, plots (not shown) of  $\log k_c$  IBA versus  $\log k_c$  Imz;  $\log k_c$  IBA versus  $\log k_c$  Phos; and  $\log k_c$  IBA versus  $\sigma^-$  are clearly nonlinear. We have employed the Hammett  $\sigma^-$  scale with IBA in order to be consistent with a Brønsted-type plot, and with the corresponding Hammett plots of the other two nucleophiles. The two straight lines of the  $\log k_c$  IBA versus  $\sigma^-$  plot yield  $\rho_Y$  that varies from 1.69 for poor LG, to 0.74 for their good counterparts. Therefore, there is a change in the RLS of IBA-catalyzed hydrolysis as a function of changing Y, the RLS being bond formation for




Fig. 2 Suggested structures for the rate-limiting transition states of nucleophile-catalyzed hydrolysis of benzoate esters.




Fig. 3 Part A shows the relationship between the catalytic rate constant for hydrolysis of 4-Y-phenyl 2,4-dinitrobenzoates in the presence of IBA at 45 °C,  $k_c$  IBA,<sup>4b</sup> and the  $pK_a$  of the leaving phenol. Part B shows the relationship between the catalytic rate constants for hydrolysis of 4-Y-phenyl 2,4-dinitrobenzoates in the presence of Imz at 40 °C,  $k_c$  Imz,<sup>4a</sup> and those for the hydrolysis of 4-Y-phenyl 3,5-dinitrobenzoates catalyzed by phosphate at 50 °C,  $k_c$  Phos. For the latter series, the rate constant for Y = 4-Cl was calculated from the Hammett plot.

good leaving phenols, and bond breaking for poor leaving ones.

(iii) In contrast, there is an excellent linear correlation between  $\log k_c$  Phos and  $\log k_c$  Imz ( $r = 0.9996$ ,  $\Sigma Q = 0.017$ , Fig. 3B), with a slope of 0.57. The corresponding Brønsted-type plots between  $\log k_c$  and  $pK_a$  of the leaving phenol are also linear ( $r = 0.9972$  and  $0.9984$ ,  $\Sigma Q = 0.085$  and  $0.036$  for Imz and Phos, respectively), with  $\beta_{LG}$  of  $-0.75$  and  $-0.40$  for Imz and Phos, respectively. The effect of Y on rates of hydrolysis of series II by these two nucleophiles is also well described by the  $\log k_c = \rho_Y \sigma_Y^-$  relationship, with  $\rho_Y = 1.59$  and  $0.89$  for Imz and Phos, respectively. Relative to Imz,  $\beta_{LG}$  and  $\rho_Y$  for the Phos-catalyzed reaction are markedly smaller; this indicates different magnitudes of bond breaking in their transition states, a conclusion that is consistent with a RL-formation and RL-breakdown of the tetrahedral intermediate for Phos and Imz, respectively.

(iv) In principle, the magnitude of bond formation between an attacking nucleophile and a reactant can be obtained from the slope,  $\beta_{Nucleophile}$ , of the Brønsted-type plot of  $\log k_c$  versus  $pK_a$  of the attacking species. Since distinct classes of nucleophiles were employed (oxygen and nitrogen),  $\beta_{Nucleophile}$  for the present reactions is not attainable by this approach. However,

$\rho_X$  is relevant to bond formation since it expresses the electronic demand on the X-substituted acyl moiety, in order to stabilize the partial negative charge on the acyl oxygen atom. Calculated  $\rho_X$  are 1.80, 1.71, and 2.23 for IBA, Imz, and Phos, respectively. These are consistent with a RL bond formation in the Phos-catalyzed reaction, with a demand on X larger than for the other two nucleophiles. Examination of the electrostatic interactions within the transition states shown in Fig. 2 corroborates this conclusion. For Phos, electrostatic repulsion is large in the dianionic TS, this being the reason for the large electronic demand on X,<sup>13b</sup> so that  $\rho_X$  is largest among the three nucleophiles.

(v) The activation entropies for hydrolysis of series I are consistent with the preceding discussion. As expected for bimolecular reactions, all  $\Delta S^\ddagger$  are negative but  $|\Delta S^\ddagger|_{Phos} < |\Delta S^\ddagger|_{Imz}$  in agreement with our previous conclusion regarding the importance of the bond formation (Phos), and bond breaking (Imz), steps respectively. The fact that  $|\Delta S^\ddagger|_{Phos} > |\Delta S^\ddagger|_{IBA}$  is unexpected, at least for good LG, and may reflect: (i) the above-mentioned favorable disposition of the IBA nucleophilic center, (ii) more extensive solvent reorganization required to solvate the dianionic TS of the Phos-catalyzed reaction.

## Conclusions

Phosphate-catalyzed hydrolysis of benzoate esters occurs via a stepwise mechanism, involving  $HPO_4^{2-}$  as a nucleophile, and the formation of acylated nucleophile, similar to Imz- and IBA-catalyzed hydrolysis of the same esters. The mechanism of catalysis and the nature of the RLS clearly depend on the structures of catalyst and substrate. Thus, increasing the strength of the LG from ethyl (in dichloroacetate)<sup>1</sup> to substituted phenyl (4-nitrophenyl acetate,<sup>1,13a</sup> series I and II) results in a change in the mechanism of catalysis by Phos and Imz from general base to nucleophilic. The weakest nucleophile, Phos, reacts with a RL-attack on the ester CO group, Imz reacts with RL phenol departure. The RLS of IBA, the strongest nucleophile, depends on the strength of the leaving phenol.

## Acknowledgements

We thank FAPESP for financial support, CNPq for a research fellowship to W. A. Rodrigues and a research productivity fellowship to O. A. El Seoud, E. B. Tada and C. Guizzo for their help during the preparation of this manuscript. This work has been carried out within a CAPES/COFECUB bilateral cooperation project.

## References

- M. L. Bender, R. J. Bergeron and M. Komiyama, *The Bioorganic Chemistry of Enzyme Catalysis*, Plenum, New York, 1984, pp. 45, 130.
- W. P. Jencks, *Catalysis in Chemistry and Enzymology*, Dover, New York, 1987, pp. 42, 163, 463.
- N. S. Isaacs, *Physical Organic Chemistry*, Longman, Essex, 1987, p. 375.
- (a) P. Menegheli, J. P. S. Farah and O. A. El Seoud, *Ber. Bunsenges. Phys. Chem.*, 1991, **95**, 1610; (b) O. A. El Seoud and M. F. Martins, *J. Phys. Org. Chem.*, 1995, **8**, 637.
- T. H. Fife, *Acc. Chem. Res.*, 1993, **26**, 325.
- D. D. Perrin, W. L. F. Armarego and D. R. Perrin, *Purification of Laboratory Chemicals*, 2nd edn, Wiley-Interscience, New York, 1988.
- (a) V. A. Izmail'skii and A. V. Belotsvetov, *J. Gen. Chem. USSR.*, 1941, **11**, 650 (*Chem. Abstr.*, 1942, **36**, 436); (b) R. C. Brown and R. E. Kremers, *J. Am. Pharm. Assoc.*, 1922, **11**, 607; (c) E. de B. Barnett and I. G. Nixon, *Chem. News*, 1924, **129**, 190.
- (a) W. P. Jencks and J. Carriuolo, *J. Am. Chem. Soc.*, 1961, **83**, 1743; (b) W. P. Jencks and G. Di Sabato, *J. Am. Chem. Soc.*, 1961, **83**, 4393.

- 4400; (c) G. Camici, G. Manao, G. Cappugi and G. Ramponi, *Experientia*, 1976, **32**, 535.
- 9 (a) C. D. Hubbard and J. F. Kirsch, *Biochemistry*, 1972, **11**, 2483; (b) E. A. Castro and G. B. Steinfert, *J. Chem. Soc., Perkin Trans. 2*, 1983, 453; (c) R. A. Moss, S. Swarp and S. Ganguli, *J. Chem. Soc., Chem. Commun.*, 1987, 860.
- 10 A. Williams, *Acc. Chem. Res.*, 1989, **22**, 387.
- 11 (a) S. Ba-Saif, A. K. Luthra and A. J. Williams, *J. Am. Chem. Soc.*, 1987, **109**, 6362; (b) E. Buncel, H. I. Um and S. S. Hoz, *J. Am. Chem. Soc.*, 1989, **111**, 971.
- 12 Preliminary (unpublished) results have shown that the hydrolysis of 4-nitrobenzoyl phosphate *in water* in the presence of 0.1 M phosphate buffer, at pH = 7.8, at 39 °C has a half-life of 28 minutes, compared with 3.04 minutes for the hydrolysis of I-d under more or less similar experimental conditions. Both reactions should be slowed down in 65% acetonitrile in water.
- 13 (a) C. J. O'Connor and R. G. Wallace, *Aust. J. Chem.*, 1984, **37**, 2559; (b) D. Herschlag and W. P. Jencks, *J. Am. Chem. Soc.*, 1989, **111**, 7587; (c) A. Kanavarioti and M. T. Rosenbach, *J. Org. Chem.*, 1991, **56**, 1513; (d) I. M. Kovach, A. J. Bennet, J. A. Bibbs and Q. Zhao, *J. Am. Chem. Soc.*, 1993, **115**, 5138; (e) V. G. Machado, C. A. Bunton, C. Zucco and F. Nome, *J. Chem. Soc., Perkin Trans. 2*, 2000, 169.
- 14 J. Shorter, *Pure Appl. Chem.*, 1994, **66**, 2451.
- 15 O. A. El Seoud, P. Menegheli, P. A. R. Pires and N. Z. Kiyan, *J. Phys. Org. Chem.*, 1994, **7**, 431.
- 16 (a) J. E. Delbene and I. Cohen, *J. Am. Chem. Soc.*, 1978, **100**, 5285; (b) D. W. Smith, *J. Chem. Educ.*, 1977, **54**, 540.