

Improved Syntheses of Tetrachloro-oxorhenium(vi) and Chlorotrioxorhenium(vii). Synthesis of Alkoxo- and Dialkylamido-rhenium Compounds. The Crystal and Molecular Structures of Di- μ -methoxy-tetramethoxy- μ -oxo-dioxodirhenium(vi) (*Re-Re*), Bis[lithium pentaisopropoxy-oxorhenate(vi)-lithiumchloride-tetrahydrofuran(1/1/2)], and *trans*-Tetraphenobis(trimethylphosphine)rhenium(iv) †

By Peter G. Edwards and Geoffrey Wilkinson,* Chemistry Department, Imperial College of Science and Technology, London SW7 2AY

Michael B. Hursthouse * and K. M. Abdul Malik, Chemistry Department, Queen Mary College, Mile End Road, London E1 4NS

Much improved synthetic procedures for ReOCl_4 , ReO_3Cl , ReOMe_4 , $\text{Li}_2[\text{ReMe}_6]$, and ReMe_6 are described. Interaction of ReOCl_4 with methanol in the presence of amine gives the unusual alkoxo-compound $(\text{MeO})_2\text{ORe}(\mu\text{-O})(\mu\text{-OMe})_2\text{ReO}(\text{OMe})_2$, (A), while LiOBu^t gives $\text{ReO}(\text{OBu}^t)_4$, and LiOPr^i gives an unusual dimeric species $\{\text{Li}[\text{ReO}(\text{OPr}^i)]_5\cdot\text{LiCl}(\text{thf})_2\}_2$, (B) (thf = tetrahydrofuran). The interaction with phenol in the presence of trimethylphosphine leads to the compound *trans*- $\text{Re}(\text{OPh})_4(\text{PMe}_3)_2$, (C), while with lithium bis(trimethylsilyl)amide a unique four-co-ordinate rhenium(v) compound, $\text{ReO}\{\text{N}(\text{SiMe}_3)_2\}_3$, is obtained. The crystal and molecular structures of (A)–(C) have been determined by X-ray crystallography. Compound (A) is monoclinic, space group $P2_1/n$, $a = 12.142(1)$, $b = 15.369(1)$, $c = 7.311(1)$ Å, $\beta = 90.22(1)^\circ$, and $Z = 4$. Compound (B) is triclinic, space group P_1 , $a = 14.602(5)$, $b = 13.363(3)$, $c = 9.244(6)$ Å, $\alpha = 98.39(4)$, $\beta = 102.35(5)$, $\gamma = 68.71(2)^\circ$, and $Z = 1$. Compound (C) is monoclinic, space group $P2_1/c$, $a = 10.079(2)$, $b = 10.527(3)$, $c = 14.443(3)$ Å, $\beta = 97.28(2)^\circ$, and $Z = 2$. The structures have been refined to R values of 0.059 (A), 0.069 (B) and 0.055 (C) using respectively 1 965, 4 061, and 1 797 observed intensities, measured on a diffractometer.

WITH the exception of the triply alkoxo-bridged carbonyl anions,¹ $[(\text{OC})_3\text{Re}(\mu\text{-OR})_3\text{Re}(\text{CO})_3]^-$ ($\text{R} = \text{Et}$ or Pr^i), other alkoxo-species have only one alkoxo-group and contain tertiary phosphine ligands² as in $\text{ReO}(\text{OEt})\text{-Cl}_2(\text{PPh}_3)_3$ or methyl and acetate groups as in the insoluble $\{\text{ReMe}(\text{OMe})(\text{O}_2\text{CMe})\}_n$.³ We have been unable to confirm the compounds said⁴ to have stoichiometry $\text{Re}(\text{OR})_3$ ($\text{R} = \text{Et}$ or Pr^i). Although we have as yet failed to isolate alkoxide or dialkylamido-compounds starting with ReCl_5 and $\text{ReCl}_4(\text{thf})_2$ (thf = tetrahydrofuran), we have isolated products from ReOCl_4 namely $\text{Re}_2\text{O}_3(\text{OMe})_6$, whose X-ray structure was determined, and $\text{ReO}(\text{OBu}^t)_4$, $\text{Li}[\text{ReO}(\text{OPr}^i)]_5$, and $\text{ReO}\{\text{N}(\text{SiMe}_3)_2\}_3$, which were described in a preliminary note;⁵ full details of this work are now given. We have subsequently⁶ synthesised the trirhenium cluster alkoxide $\text{Re}_3\text{Cl}_3(\text{OBu}^t)_6$ while trimethylsilyloxo-compounds with *N*-t-butylimido-groups $\text{Re}(\text{NBu}^t)_3(\text{OSiMe}_3)$ and $\text{Re}_3\text{O}_5(\text{NBu}^t)_4(\text{OSiMe}_3)_3$, the latter having bridging OSiMe_3 , O, and OReO_3 groups, have been recently described.⁷

We also describe improved syntheses of the starting material, ReOCl_4 , and of ReO_3Cl .

RESULTS AND DISCUSSION

Syntheses.—(1) *Tetrachloro-oxorhenium(vi) and chlorotrioxorhenium(vii)*. Methods for the synthesis of ReOCl_4 are (i) the oxidation of rhenium metal or rhenium halides in thionyl chloride⁸ or sulphuryl chloride;⁹ or (ii) the interaction of ReCl_5 with oxygen in a sealed tube at elevated temperature.¹⁰ These are impractical on a large scale and purification of the oxohalide can be tedious.

† No reprints available.

Rhenium pentachloride is readily made in large quantities in high yield¹¹ and we find that direct oxidation with dioxygen just below the melting point (261°C) in an open system at 1 atm ‡ pressure gives yields of up to 90% ReOCl_4 , an overall yield from rhenium of over 75%. The loss is due to formation of Re_3Cl_9 on thermal decomposition of ReCl_5 . Oxygenation at higher temperatures where the ReCl_5 burns gives lower yields. The ReOCl_4 is readily purified by vacuum distillation or sublimation and quantities ca. 25 g have been made in one run. The major contaminant in the product is chlorotrioxorhenium(vii) (per-rhenyl chloride, ReO_3Cl), but this is readily removed as it is a liquid (m.p. 4.5°C).

By direct oxidation of Re_3Cl_9 which again can be obtained in high yield from ReCl_5 ,¹¹ ReO_3Cl can be obtained in high yield. Thus, on heating Re_3Cl_9 in a stream of oxygen so that it inflames, the ReO_3Cl can be collected in a cold trap to give, after vacuum distillation, a colourless product in yields exceeding 80% based on rhenium metal. Early routes¹² to ReO_3Cl are tedious and difficult for large quantities while a recent procedure¹³ requires chlorine monoxide to react with rhenium halides.

(2) *Interaction of tetrachloro-oxorhenium(vi) with alcohols.* (a) *Methanol.* The interaction of ReOCl_4 with methanol and tertiary amine in diethyl ether leads to an orange diamagnetic complex that crystallises from light petroleum as oxygen- and moisture-sensitive rhombs. The compound can be sublimed with some decomposition at 90°C (10^{-3} Torr). Analytical and molecular-weight data indicate a stoichiometry Re_2O_3 -

‡ Throughout this paper: 1 atm = 101 325 Pa; 1 Torr = (101 325/760) Pa; 1 B.M. $\approx 9.27 \times 10^{-24}$ A m².

(OMe)₆, (A). Although the molecular ion is not observed, the mass spectrum shows the highest peak at *m/e* 342, corresponding to ReO₂(OMe)₄, while peaks at lower mass correspond to loss of CH₃ and O from this fragment.

The spectroscopic data (Table 1) are consistent with the structure (MeO)₂ORe(μ-O)(μ-OMe)₂ReO(OMe)₂, determined by X-ray study (Figure 1 and below). Thus the ¹H n.m.r. shows two singlets at δ 5.03 and 0.20 p.p.m. (area 2 : 1) indicating two types of MeO groups; the high-field peak is assigned to μ-OMe. The i.r. spectrum has bands expected for OMe groups [ν(CO) at 1 025 cm⁻¹], together with sharp bands at 968 and 982 cm⁻¹ assigned to ν(Re=O), and a band at 756 cm⁻¹ assigned to ν(Re-O-Re) as found in other species containing the O=Re-O-Re=O unit.¹⁴ Although a linear Re₂O₃ unit occurs in several rhenium complexes,¹⁵ the lower symmetry of the

shows alkoxy-bands (C-O, 1 115) and ν(Re-O) at 987 cm⁻¹.

The e.s.r. spectrum of a frozen benzene solution of the compound at 9.2 GHz consists of several fairly well resolved features spread over the region 0.15–0.64 T. It is typical of a rhenium(vi) complex with a very large hyperfine structure. It is very difficult to pick out the three sets of six lines expected for *S* = ½, *I* = 5/2 because, as with other rhenium(vi) spectra,¹⁶ the hyperfine lines are not equally spaced and their intensities vary markedly; they are thus easily masked by the 'forbidden' lines which abound. As yet a good interpretation of the spectrum is not feasible but the principal values of the *A* tensor for rhenium seem to be between about 0.04 and 0.06 cm⁻¹, while those of the *g* tensor are probably below 2.0. The high solubility of this salt in benzene indicates

TABLE 1
Spectroscopic data for the rhenium alkoxides and rhenium dialkylamide

Compound	¹ H n.m.r.		Infrared (cm ⁻¹)	
	δ/p.p.m. ^a	Assignment	ν(Re=O)	Other vibrations
Re ₂ O ₃ (OMe) ₆	5.03 (s, 2 H)	Re-OMe	982m, 986s	2 950m, 2 910s, 2 860w, 150m, 1 025s, 575s, 548s, 520m ^c
	0.20 (s, 1 H)	μ-(Re-OMe)	755w ^b	1
ReO(OBu ^t) ₄	1.15	Re-OCMe ₃	992m	1 380m, 1 358s, 1 230m, 1 167s, 1023w, 917s, 898s, 770m, 730w, 588m, 421m ^d
Li[ReO(OPr ⁱ) ₅] ₂ ·LiCl	1.35	Re-OCHMe ₂	987m	1 360m, 1 340w, 1 315w, 1 260m, 1 167m, 1 115s, 937s, 832s, 638m, 612s, 570m, 467w, 420w ^d
ReO[N(SiMe ₃) ₂] ₃	0.20	N-SiMe ₃	978	1 400m, 1 248m, 1 100w, 885s, 840s, 772m, 693w, 660m, 432w ^d

^a Relative to SiMe₄ δ = 0.0 p.p.m. ^b ν(Re-O). ^c In carbon disulphide. ^d In Nujol.

bent unit observed here explains the splitting of the ν(Re=O) band.

(b) *Isopropyl and t-butyl alcohols*. Using PrⁱOH brown solutions are obtained but no crystalline product could be isolated. *t*-Butyl alcohol also gives brown solutions, but from light petroleum low yields of very air-sensitive blue crystals were obtained. Similar blue crystals can be obtained in high yield, however, by interaction of ReOCl₄ with LiOBu^t, but these are thermally unstable above 0 °C, decomposing to give a black tar and a colourless liquid (mainly C₈ monounsaturated hydrocarbon plus a little Bu^tOH).

The complex is monomeric, and paramagnetic, showing a poorly resolved e.s.r. signal in frozen benzene, consistent with a monomeric, d¹, rhenium(vi) species. The ¹H n.m.r. spectrum shows a broadened singlet at δ 1.15 p.p.m. while the i.r. shows bands at 917 (C-O) and 992 (Re=O) cm⁻¹. It seems reasonable to suggest that this unstable compound is ReO(OBu^t)₄.

The reaction of LiOPrⁱ with ReOCl₄ leads to the isolation of green, light petroleum-soluble, paramagnetic crystals of stoichiometry Li[ReO(OPrⁱ)₅]₂·LiCl. Crystallisation from thf gives {Li[ReO(OPrⁱ)₅]₂·LiCl(thf)₂}₂, (B), whose structure has been determined by X-ray crystallography (Figure 3 and below). The conductivity in thf is very low (*ca.* 5 cm² ohm⁻¹ mol⁻¹), evidently due to close ion-pair formation. The complex reacts with moisture but not dry air. The ¹H n.m.r. spectrum has a broad signal at δ = 1.35 p.p.m., while the i.r.

strong ion pairing, but there is no sign of hyperfine interaction from Li⁺ (*I* = 3/2) in the spectrum of the anion.

(c) *Phenol*. The interaction of ReOCl₄ with LiOPh gives a purple solution from which no clean product could be isolated. However, the addition of trimethylphosphine gives a black powder insoluble in diethyl ether, benzene, and light petroleum and extraction of this with CH₂Cl₂ leads to a low yield of red, air-stable crystals. X-Ray diffraction studies confirm the structure as *trans*-Re(OPh)₄(PMe₃)₂ (C) (Figure 4 and below). The reduction to an octahedral rhenium(IV) complex is presumably due to the phosphine.

(3) *Interaction of ReOCl₄ with lithium dialkylamides*. With the exception of LiN(SiMe₃)₂, the interaction of lithium dialkylamides has so far not led to the isolation of well defined compounds. However, with LiN(SiMe₃)₂ a red, light petroleum-soluble, diamagnetic and volatile (sublimes 60 °C, 10⁻³ Torr) compound that crystallises from petroleum as red tetragonal plates is obtained. Analytical and spectroscopic data clearly show that the compound is ReO[N(SiMe₃)₂]₃ which has rhenium(v) in the previously unknown¹⁷ co-ordination of four. The ¹H n.m.r. spectrum has only a sharp singlet at δ = 0.2 p.p.m. due to CH₃ groups, while the i.r. spectrum has a band at 1 248 cm⁻¹ due to Si-C and one at 978 cm⁻¹ due to Re=O.

(4) *Synthesis of methylrhenium compounds*. The oxotetrachloride is a useful starting material for the synthesis of known rhenium(vi) alkyls, ReOMe₄,¹⁴ ReMe₆,

and $\text{Li}_2[\text{ReMe}_6]$,^{16,18} giving higher yields than the previous methods starting with $\text{ReOCl}_3(\text{PPh}_3)_3$, which were somewhat irreproducible and gave very variable yields. Thus, ReOMe_4 is obtained by action of a stoichiometric amount of methyl-lithium on ReOCl_4 at low temperatures followed by addition of water after warming. The use of excess of LiMe gives $\text{Li}_2[\text{ReMe}_6]$ directly and this in turn can be used to generate ReMe_6 by addition of water.

Structures.—(1) $\text{Re}_2\text{O}_3(\text{OMe})_6$. The structure of $\text{Re}_2\text{O}_3(\text{OMe})_6$, (A), is illustrated in Figure 1, which also indicates the atom labelling used. Interatomic distances and angles are given in Table 2. The molecule

TABLE 2

Intramolecular interatomic distances and angles for $\text{Re}_2\text{O}_3(\text{OMe})_6$ (A). The estimated standard deviations are given in parentheses

Bond lengths (Å)			
Re(2)—Re(1)	2.559(1)	O(1)—Re(1)	2.094(11)
O(2)—Re(1)	2.097(11)	O(3)—Re(1)	1.917(13)
O(4)—Re(1)	1.874(12)	O(5)—Re(1)	1.903(11)
O(6)—Re(1)	1.690(11)	O(1)—Re(2)	2.139(10)
O(2)—Re(2)	2.121(10)	O(3)—Re(2)	1.916(13)
O(7)—Re(2)	1.867(12)	O(8)—Re(2)	1.891(12)
O(9)—Re(2)	1.703(12)	C(1)—O(1)	1.436(17)
C(2)—O(2)	1.427(18)	C(4)—O(4)	1.499(20)
C(5)—O(5)	1.419(23)	C(7)—O(7)	1.476(20)
C(8)—O(8)	1.443(23)		
Bond angles (°)			
O(1)—Re(1)—Re(2)	53.6(3)	O(2)—Re(1)—Re(2)	53.1(3)
O(2)—Re(1)—O(1)	69.9(4)	O(3)—Re(1)—Re(2)	48.1(4)
O(3)—Re(1)—O(1)	73.4(5)	O(3)—Re(1)—O(2)	100.5(5)
O(4)—Re(1)—Re(2)	131.3(4)	O(4)—Re(1)—O(1)	100.9(5)
O(4)—Re(1)—O(2)	165.1(5)	O(4)—Re(1)—O(3)	87.5(6)
O(5)—Re(1)—Re(2)	127.8(4)	O(5)—Re(1)—O(1)	86.9(5)
O(5)—Re(1)—O(2)	84.2(4)	O(5)—Re(1)—O(3)	156.4(5)
O(5)—Re(1)—O(4)	83.6(5)	O(6)—Re(1)—Re(2)	102.2(5)
O(6)—Re(1)—O(1)	153.4(6)	O(6)—Re(1)—O(2)	86.9(6)
O(6)—Re(1)—O(3)	99.4(6)	O(6)—Re(1)—O(4)	104.4(7)
O(6)—Re(1)—O(5)	104.0(6)	O(1)—Re(2)—Re(1)	52.0(3)
O(2)—Re(2)—Re(1)	52.2(3)	O(2)—Re(2)—O(1)	68.6(4)
O(3)—Re(2)—Re(1)	48.1(4)	O(3)—Re(2)—O(1)	72.3(5)
O(3)—Re(2)—O(2)	99.6(5)	O(7)—Re(2)—Re(1)	130.7(4)
O(7)—Re(2)—O(1)	101.7(5)	O(7)—Re(2)—O(2)	165.5(5)
O(7)—Re(2)—O(3)	86.9(6)	O(8)—Re(2)—Re(1)	126.5(4)
O(8)—Re(2)—O(1)	87.1(5)	O(8)—Re(2)—O(2)	83.9(5)
O(8)—Re(2)—O(3)	155.8(5)	O(8)—Re(2)—O(7)	84.9(6)
O(9)—Re(2)—Re(1)	103.0(5)	O(9)—Re(2)—O(1)	152.7(6)
O(9)—Re(2)—O(2)	87.7(6)	O(9)—Re(2)—O(3)	99.8(6)
O(9)—Re(2)—O(7)	103.9(7)	O(9)—Re(2)—O(8)	104.2(6)
Re(2)—O(1)—Re(1)	74.4(3)	C(1)—O(1)—Re(1)	126.4(10)
C(1)—O(1)—Re(2)	124.8(10)	Re(2)—O(2)—Re(1)	74.7(3)
C(2)—O(2)—Re(1)	125.1(11)	C(2)—O(2)—Re(2)	124.7(11)
Re(2)—O(3)—Re(1)	83.8(5)	C(4)—O(4)—Re(1)	128.4(13)
C(5)—O(5)—Re(1)	131.3(10)	C(7)—O(7)—Re(2)	128.5(13)
C(8)—O(8)—Re(2)	130.7(12)		

contains two rhenium atoms bridged by one oxo- and two methoxy-groups and has the overall geometry of a facial bioctahedron with considerable distortion. It does not possess any crystallographic symmetry, but has an approximate mirror plane passing through the bridging oxygen atoms. The Re—O—Re bridge angles are all acute, with that involving the oxo-group *ca.* 10° wider than those with the methoxy-groups (83.8 *vs.* 74.4 and 74.7°). The acute angles at the bridging atoms and the short Re—Re distance (2.559 Å) are consistent with the presence of a Re—Re single bond, as required by the observed diamagnetism of the compound.




FIGURE 1 Molecular structure of $\text{Re}_2\text{O}_3(\text{OMe})_6$, (A), showing the atom-numbering

The structural parameters in the two ReO_6 octahedra are nearly the same, but each octahedron shows considerable distortions from idealised geometry. The $\text{O}_3\text{Re}(\text{O}_3)\text{ReO}_3$ bioctahedron retains the basic feature of eclipsing of the atoms occupying the corners of the two terminal triangles, but the triangle formed by the three bridging atoms is shifted so that although the Re—Re is still perpendicular to the $\text{O}(2) \cdots \text{O}(3)$ edge, it almost intersects it. The extent of this distortion is expressed by the dihedral angles between the planes $\text{Re}(1)\text{—O}(1)\text{—Re}(2)/\text{Re}(1)\text{—O}(2)\text{—Re}(2)$ 91.1°, $\text{Re}(1)\text{—O}(1)\text{—Re}(2)/\text{Re}(1)\text{—O}(3)\text{—Re}(2)$ 79.4°, and $\text{Re}(1)\text{—O}(2)\text{—Re}(2)/\text{Re}(1)\text{—O}(3)\text{—Re}(2)$ 11.7°, and is well demonstrated by the view chosen for Figure 2.




FIGURE 2 View of the $\text{Re}_2\text{O}_3(\text{OMe})_6$ molecule showing the eclipsing of the terminal ligands and staggering of the bridging oxo-group [O(3)] with the two terminal alkoxy-oxygen atoms [O(4) and O(7)]

One possible explanation for the distortion is the existence of a bent, off-centre, $\text{Re} \cdots \text{Re}$ interaction, stereochemically active in the space between O(6), O(9), and O(3). This would to some extent explain the uneven

by the oxygens O(1) and O(2) becoming planar. That this does not happen is probably a reflection of the strain which would occur in the bridges if the oxygen bridging orbitals were sp^2 rather than sp^3 hybridised.




FIGURE 3 Structure of $\{\text{Li}[\text{ReO}(\text{OPr}^i)_5] \cdot \text{LiCl}(\text{thf})_2\}_2$ (B), showing the atom numbering in the asymmetric unit. The primed atoms are generated by the centre of symmetry at $(0, \frac{1}{2}, \frac{1}{2})$

distribution of intramolecular $\text{O} \cdots \text{O}$ contacts (Table 3), where $\text{O}(3) \cdots \text{O}(6)$, $\text{O}(9)$, whilst still very short, are longer than $\text{O}(3) \cdots \text{O}(4)$, $\text{O}(7)$ and $\text{O}(2) \cdots \text{O}(6)$, $\text{O}(9)$.

In fact, one characteristic of this molecule is the con-

TABLE 3

Selected non-bonded short contacts (Å) in $\text{Re}_2\text{O}_3(\text{OMe})_6$

(a) Intramolecular

$\text{O}(1) \cdots \text{O}(2)$	2.40	$\text{O}(3) \cdots \text{C}(7)$	2.81
$\text{O}(1) \cdots \text{O}(3)$	2.40	$\text{O}(4) \cdots \text{O}(5)$	2.52
$\text{O}(1) \cdots \text{O}(4)$	3.06	$\text{O}(4) \cdots \text{C}(5)$	2.75
$\text{O}(1) \cdots \text{O}(5)$	2.75	$\text{O}(4) \cdots \text{O}(6)$	2.82
$\text{O}(1) \cdots \text{O}(7)$	3.11	$\text{O}(5) \cdots \text{O}(6)$	2.83
$\text{O}(1) \cdots \text{O}(8)$	2.78	$\text{O}(5) \cdots \text{C}(1)$	2.99
$\text{O}(2) \cdots \text{O}(3)$	3.09	$\text{O}(5) \cdots \text{C}(2)$	3.52
$\text{O}(2) \cdots \text{O}(5)$	2.69	$\text{O}(5) \cdots \text{C}(4)$	3.31
$\text{O}(2) \cdots \text{O}(6)$	2.62	$\text{O}(6) \cdots \text{C}(2)$	2.89
$\text{O}(2) \cdots \text{O}(8)$	2.69	$\text{O}(6) \cdots \text{O}(9)$	3.30
$\text{O}(2) \cdots \text{O}(9)$	2.67	$\text{O}(7) \cdots \text{O}(8)$	2.54
$\text{O}(2) \cdots \text{C}(1)$	2.85	$\text{O}(7) \cdots \text{O}(9)$	2.81
$\text{O}(3) \cdots \text{O}(4)$	2.62	$\text{O}(7) \cdots \text{C}(8)$	2.77
$\text{O}(3) \cdots \text{O}(6)$	2.75	$\text{O}(8) \cdots \text{O}(9)$	2.84
$\text{O}(3) \cdots \text{O}(7)$	2.60	$\text{O}(8) \cdots \text{C}(1)$	2.99
$\text{O}(3) \cdots \text{O}(9)$	2.77	$\text{O}(8) \cdots \text{C}(2)$	3.52
$\text{O}(3) \cdots \text{C}(4)$	2.86	$\text{O}(8) \cdots \text{C}(7)$	3.33

(b) Intermolecular

$\text{O}(1) \cdots \text{C}(2^{\text{I}})$	3.48	$\text{O}(6) \cdots \text{C}(8^{\text{III}})$	3.34
$\text{O}(5) \cdots \text{C}(4^{\text{II}})$	3.31	$\text{O}(8) \cdots \text{C}(7^{\text{II}})$	3.33
$\text{O}(5) \cdots \text{C}(8^{\text{III}})$	3.46	$\text{O}(9) \cdots \text{C}(4^{\text{V}})$	3.46
$\text{O}(6) \cdots \text{C}(1^{\text{IV}})$	3.06	$\text{O}(9) \cdots \text{C}(5^{\text{IV}})$	3.49

Symmetry code:

I $\frac{1}{2} + x, \frac{1}{2} - y, \frac{1}{2} + z;$	II $x, y, -1 + z;$
III $\frac{1}{2} - x, -\frac{1}{2} + y, -\frac{1}{2} - z;$	IV $-\frac{1}{2} + x, \frac{1}{2} - y, \frac{1}{2} + z;$
V $-\frac{1}{2} + x, \frac{1}{2} - y, -\frac{1}{2} + z$	

siderable intramolecular crowding, as evidenced by the values listed in Table 3. In particular, we note the contacts $\text{O}(2) \cdots \text{C}(1)$, $\text{O}(5) \cdots \text{C}(1)$, $\text{O}(8) \cdots \text{C}(1)$, $\text{O}(6) \cdots \text{C}(2)$, and $\text{O}(9) \cdots \text{C}(2)$ which could actually be relieved

The distribution of Re-O distances is as expected. The terminal $\text{Re}=\text{O}$ lengths (1.690, 1.703 Å) are similar to that found in the compound [acetone benzoylhydrazonido-(1-)-*N'*O]dichloro-oxo(triphenylphosphine)rhenium(v), 1.685 Å.¹⁹ The bridging Re-O-Re bonds are longer at 1.916, 1.917 Å and are similar to those involving the terminal OMe groups (1.867–1.903 Å). The bonds to the bridging methoxides are longest, at 2.094–2.139 Å.

The dimensions of the methoxide ligands are normal, with a mean C-O distance of 1.450 Å and, perhaps surprisingly, a fairly constant Re-O-C angle of 125–130°. The shorter intermolecular contacts (Table 3) involve oxygen atoms and indicate a fairly tight packing.

(2) $\{\text{Li}[\text{ReO}(\text{OPr}^i)_5] \cdot \text{LiCl}(\text{thf})_2\}_2$. The structure of this centrosymmetric dimer, (B), and the atom-numbering scheme are illustrated in Figure 3. Each half of the dimer contains an octahedral $[\text{ReO}(\text{OPr}^i)_5]^-$ ion in which three alkoxo-oxygen atoms [O(1), O(2), and O(3)] are involved in forming bridges with two Li^+ ions to produce Re-O-Li-O heterocycles in a manner similar to that found previously for two alkoxides of Cr^{III} and Fe^{III} .²⁰ One OPrⁱ group is also involved in forming a Li-O-Li-Cl ring.

Some of the more important bond lengths and angles are given in Table 4, and present interesting comparisons with those found for compound (A). Thus, $\text{Re}=\text{O}$ here is 1.707 Å; Re-O distances to the two terminal alkoxides are 1.898 and 1.899 Å, and to those alkoxides involved in Re-O-Li bridging, 1.987 and 2.023 Å. The Re-O bond to the unique four-co-ordinate oxygen atom is longest, at 2.083 Å. Even so, it is shorter than those to the three-

co-ordinate oxygens in (A). In neither compound is there any evidence of a *trans* influence weakening by the Re-O bond such as is found for other metal-oxo-functions.²¹ Nevertheless, in (B) there is the usual bending away of the 'equatorial' ligands from the Re=O group, with the metal atom lying some 0.23 Å above the equatorial O₄ plane towards the oxo-atom. Naturally, the disposition of the Re-O(2) and Re-O(3) bonds is also affected by the heterocycle formation.

The environment about the chloride ions is interesting in that it appears to be pyramidally three-co-ordinate. The 'empty' side of the co-ordination sphere is blocked by a second thf molecule, but there are no contacts to the chloride <4.0 Å. In fact this thf molecule is not bonded to any other portion of the complex and is rather poorly defined.

Dimensions of the OPrⁱ ligands are normal with average C-O and C-C distances of 1.425 and 1.517 Å respectively. Intermolecular contacts are all >3.71 Å and correspond to normal van der Waals packing.

TABLE 4

Selected intramolecular distances and angles for {Li[ReO(OPrⁱ)₂·LiCl(thf)₂]₂, (B)*

Bond lengths (Å)			
Re-O(1)	2.083(6)	Re-O(2)	1.987(8)
Re-O(3)	2.023(7)	Re-O(4)	1.898(8)
Re-O(5)	1.899(8)	Re-O(6)	1.707(9)
Li(1)-Cl	2.389(24)	Li(2)-Cl	2.422(18)
Li(2)-Cl'	2.379(25)	Li(1)-O(1)	1.976(23)
Li(2)-O(1)	1.995(22)	Li(2)-O(2)	1.919(21)
Li(1)-O(3)	1.855(22)	Li(1)-O(7)	1.959(21)
Bond angles (°)			
O(1)-Re-O(2)	77.8(3)	O(1)-Re-O(3)	75.7(3)
O(1)-Re-O(4)	89.2(3)	O(1)-Re-O(5)	91.0(3)
O(1)-Re-O(6)	163.1(4)	O(2)-Re-O(3)	91.5(3)
O(2)-Re-O(4)	165.8(3)	O(2)-Re-O(5)	88.7(3)
O(2)-Re-O(6)	92.0(4)	O(3)-Re-O(4)	90.7(3)
O(3)-Re-O(5)	166.3(3)	O(3)-Re-O(6)	91.4(4)
O(4)-Re-O(5)	85.9(3)	O(4)-Re-O(6)	101.9(4)
O(5)-Re-O(6)	102.3(4)	Li(1)-Cl-Li(2')	142.5(8)
Li(1)-Cl-Li(2)	69.5(7)	Li(2)-Cl-Li(2')	83.2(8)
O(1)-Li(1)-Cl	100.6(9)	O(1)-Li(2)-Cl	99.0(8)
O(3)-Li(1)-Cl	113.7(12)	O(2)-Li(2)-Cl	120.2(10)
O(1)-Li(1)-O(3)	82.2(7)	Cl'-Li(2)-Cl	96.8(9)
O(7)-Li(1)-Cl	104.6(10)	Cl'-Li(2)-O(1)	137.5(8)
O(7)-Li(1)-O(1)	132.3(15)	Cl'-Li(2)-O(2)	122.6(8)
O(7)-Li(1)-O(3)	121.5(12)	O(1)-Li(2)-O(2)	81.5(7)
Re-O(2)-C(2)	126.2(8)	Re-O(5)-C(5)	128.7(8)
Li(2)-O(2)-C(2)	130.5(10)	Li(2)-O(2)-Re	102.9(7)
Li(1)-O(3)-Re	104.2(8)	C(3)-O(3)-Re	123.8(7)
C(4)-O(4)-Re	131.6(8)	Li(1)-O(3)-C(3)	129.0(11)
Li(1)-O(1)-Re	97.9(7)	C(1)-O(1)-Re	121.4(7)
Li(2)-O(1)-Re	97.0(7)	Li(1)-O(1)-C(1)	126.9(9)
Li(2)-O(1)-Li(1)	87.3(9)	Li(2)-O(1)-C(1)	118.2(9)

* The primed atoms belong to the same dimer, and are generated by the centre of symmetry at (0, $\frac{1}{2}$, $\frac{1}{2}$).

(3) *trans*-Re(OPh)₄(PMe₃)₂. The molecular structure of (C) is shown in Figure 4, together with the atom numbering. Important interatomic distances and angles are given in Table 5. The monomeric molecules lie on crystallographic inversion centres, and the Re atom shows a distorted octahedral geometry with the phosphine ligands in *trans* positions. The Re-P distance (2.488 Å) is normal and comparable with those (2.472, 2.481, and 2.485 Å) in ReOCl₂(PPh₃)[Ph(O)CNCMe₂],¹⁹ ReCl₃-

(NMe)(PEtPh₂)₂,²² and ReCl₃(NC₆H₄COMe)(PEt₂Ph)₂,²³ while the mean Re-O distance (1.965 Å) is comparable with the mean Re-O(OPrⁱ) (terminal) distance (1.959 Å) in (B).

The ligand-geometry parameters are normal but some angular distortions are observed which can be attributed to interligand congestion. Thus, the Re-P-C(1) angle becomes considerably wider than the other two Re-P-C angles (122.5 vs. 108.9 and 114.6°) as a result of several




FIGURE 4 Molecular structure of Re(OPh)₄(PMe₃)₂, (C), showing the atom numbering in the asymmetric unit. The primed atoms are generated by the centre of symmetry at (0, 0, 0)

C(1)···Ph short contacts (Table 5). These contacts in turn widen the Re-O(1)-C(11) angle compared with the Re-O(2)-C(22) angle (141.3 vs. 131.9°). Also the octahedral P-Re-O(1') and P-Re-O(2') angles became larger than the P-Re-O(2) angles (97.4 and 96.3 vs. 82.6 and 83.7°). The intermolecular contacts (Table 5) are normal for van der Waals packing, with one contact of 3.57 and others greater than 3.70 Å.

EXPERIMENTAL

Microanalyses were by Butterworth Microanalytical Consultancy Ltd., Pascher, Bonn, and Imperial College Laboratories. Spectrometers: Perkin-Elmer R32 (¹H, 90 MHz), 597 and 325 (i.r.), and VG Micromass 7070 (mass spectra). Molecular weights were obtained cryoscopically and using an Hitachi-Perkin-Elmer 115 osmometer. Melting points were determined in capillaries sealed under nitrogen and magnetic moments in solution by Evans' method.

All operations except the syntheses of ReOCl₄ and ReO₃-Cl were carried out under dry oxygen-free nitrogen or *in vacuo*. All hydrocarbons and ethers were dried by refluxing

over sodium-benzophenone under nitrogen; dichloromethane and acetonitrile were dried by refluxing over calcium hydride under nitrogen, alcohols by refluxing over the corresponding magnesium alkoxide under nitrogen, triethylamine by refluxing over sodium under nitrogen, and phenol was recrystallised from dry high-boiling light petroleum. All solvents were distilled immediately prior to use. The light petroleum used had b.p. 40–60 °C.

(1) *Tetrachloro-oxorhenium*(vi).—In a distillation flask (100 cm³) with a U tube to act as a cold trap sealed to the side arm, was placed rhenium pentachloride (25 g), under N₂. A gas-inlet tube of a length such that oxygen could be passed directly onto the surface of the ReCl₅ was fitted in the neck of the distillation flask. The exhaust from the cold trap was connected to a bubbler and trap to avoid suck back and the U tube was cooled to –35 °C. Teflon sleeves were used in ground joints as ReOCl₄ reacts with greases.

TABLE 5

Important interatomic distances and angles for
Re(OPh)₄(PMe₃)₂, (C)

Bond lengths (Å)		Bond angles (°)	
P–Re	2.488(3)	O(1)–Re	1.939(9)
O(2)–Re	1.992(9)	C(1)–P	1.822(14)
C(2)–P	1.845(13)	C(3)–P	1.821(12)
C(11)–O(1)	1.336(13)	C(21)–O(2)	1.338(13)
C(12)–C(11)	1.394(14)	C(16)–C(11)	1.376(16)
C(13)–C(12)	1.344(15)	C(14)–C(13)	1.398(19)
C(15)–C(14)	1.387(18)	C(16)–C(15)	1.364(17)
C(22)–C(21)	1.364(16)	C(26)–C(21)	1.376(15)
C(23)–C(22)	1.412(17)	C(24)–C(23)	1.398(21)
C(25)–C(24)	1.357(22)	C(26)–C(25)	1.361(18)
Bond angles (°)		Bond angles (°)	
O(1)–Re–P	82.6(3)	O(2)–Re–P	83.7(2)
O(2)–Re–O(1)	88.7(3)	C(1)–P–Re	122.5(5)
C(2)–P–Re	108.9(5)	C(2)–P–C(1)	102.4(8)
C(3)–P–Re	114.6(5)	C(3)–P–C(1)	103.0(7)
C(3)–P–C(2)	103.4(7)	C(11)–O(1)–Re	141.3(8)
C(21)–O(2)–Re	131.9(7)	C(12)–C(11)–O(1)	122.6(10)
C(16)–C(11)–O(1)	119.4(10)	C(16)–C(11)–C(12)	118.0(11)
C(13)–C(12)–C(11)	120.7(12)	C(14)–C(13)–C(12)	120.9(13)
C(15)–C(14)–C(13)	119.0(13)	C(16)–C(15)–C(14)	118.9(14)
C(15)–C(16)–C(11)	122.4(13)	C(22)–C(21)–O(2)	120.2(11)
C(26)–C(21)–O(2)	121.2(11)	C(26)–C(21)–C(22)	118.5(11)
C(23)–C(22)–C(21)	120.7(12)	C(24)–C(23)–C(22)	118.4(13)
C(25)–C(24)–C(23)	120.0(15)	C(26)–C(25)–C(24)	120.3(16)
C(25)–C(26)–C(21)	122.0(14)		
Non-bonded short contacts ≤ 3.80 Å			
Intramolecular		Intermolecular	
C(1) ... C(11 ^I)	3.59	C(2) ... C(3 ^{III})	3.80
C(1) ... C(12 ^I)	3.46	C(2) ... C(12 ^{III})	3.75
C(1) ... C(22 ^I)	3.55	C(13) ... C(23 ^{IV})	3.80
C(3) ... C(21 ^I)	3.49	C(16) ... C(24 ^V)	3.80
C(3) ... C(22 ^I)	3.79	C(16) ... C(25 ^V)	3.71
C(3) ... C(26 ^I)	3.75	C(26) ... C(26 ^{VI})	3.57

Symmetry code:

I	–x, –y, –z;	II	–x, –½ + y, ½ – z;
III	–x, ½ + y, ½ – z;	IV	1 – x, –y, –z;
V	x, ½ – y, ½ + z;	VI	–x, 1 – y, –z

Oxygen was passed slowly over the rhenium chloride and the distillation flask was heated to 250 °C. Tetrachloro-oxorhenium began to reflux in the distillation flask and the oxygen flow was adjusted so that the oxohalide collected in the U tube, but was not flushed completely through the system. When no more oxohalide distilled the flask was allowed to cool and the apparatus was isolated under oxygen and transferred to a vacuum line. The more volatile ReO₃Cl was first condensed into a separate flask (cooled in liquid nitrogen) and the ReOCl₄ then collected. The product is quite pure after one distillation but can be further

purified by sublimation or trap-to-trap distillation. The initial product is satisfactory for syntheses. Yields after purification are typically ca. 19 g (80% based on ReCl₅).

(2) *Chlorotrioxorhenium*(vii).—In the apparatus described above was placed trirhenium nonachloride (25 g). The U tube was cooled to –35 °C and oxygen was passed slowly over the halide. The distillation flask was heated gently with a bunsen burner until the Re₃Cl₉ inflamed. The oxygen flow was adjusted so that the combustion was maintained without external heating and the chlorotrioxorhenium produced was not flushed through the system. The ReO₃Cl rapidly collects in the cold trap and freezes to a solid varying in colour from green-yellow to red. When all the Re₃Cl₉ had burned the flask was allowed to cool and the apparatus was isolated under oxygen and transferred to a vacuum line. The ReO₃Cl was distilled very slowly under vacuum into a separate flask cooled in liquid nitrogen. If the distillation is too rapid the product is coloured (yellow or red) and another distillation is necessary to obtain a colourless product, ca. 20 g (87% based on Re₃Cl₉).

(3) *Di-μ-methoxy-tetramethoxy-μ-oxo-dioxodirhenium*(vi) (Re–Re), (A).—A solution (red) of tetrachloro-oxorhenium(vi) (1 g) in diethyl ether (10 cm³) was added with vigorous stirring to a cooled (–78 °C) solution of methanol (5 cm³) and triethylamine (2 cm³) in diethyl ether (20 cm³). The violet solution was allowed to warm to ambient temperatures with stirring when a white precipitate formed. The solvent was removed *in vacuo* and the orange residue extracted with light petroleum (4 × 50 cm³). The orange solution was filtered, concentrated to ca. 75 cm³, and held at –20 °C overnight to give orange diamond-shaped crystals. A second crop of crystals was obtained by further concentration of the supernatant and cooling. The crystals were collected, washed with light petroleum (2 × 5 cm³), and dried under vacuum. Yield 0.64 g (74% based on ReOCl₄), m.p. 76–78 °C (Found: C, 12.7; H, 3.2; O, 23.5%; M 580. C₄H₁₂O₆Re requires C, 11.9; H, 3.0; O, 23.8%; M 606).

(4) *Oxotetra-t-butoxorhenium*(vi).—To a solution of ReOCl₄ (1 g, 2.9 mmol) in diethyl ether (20 cm³) cooled to –78 °C was added dropwise a solution of lithium t-butoxide [1.05 mol dm⁻³ in light petroleum (11.1 cm³), 11.6 mmol] with vigorous stirring. The resulting red solution was allowed to warm slowly to 10 °C, when it was green. The solvent was removed *in vacuo* and the residue extracted with light petroleum (3 × 20 cm³). The green solution was filtered, concentrated to ca. 20 cm³, and held at –20 °C overnight. The large blue crystals were collected, washed with cold light petroleum (2 × 5 cm³), and stored at –20 °C. Yield 1.2 g (84% based on ReOCl₄); decomp. ca. 0 °C, M 400 (calc. 494).

A small sample of the butoxide (ca. 0.1 g) was allowed to decompose at room temperature in a sealed flask with side tube for introducing samples into the mass spectrometer *via* a vacuum line. Both flasks were cooled in liquid nitrogen, evacuated, and connected in a closed system. The volatile products were then allowed to condense in the sample tube which was subsequently isolated from the vacuum line.

(5) *Lithium Pentaisopropoxo-oxorhenate*(vi)—*Lithium Chloride*, (B).—To a solution of ReOCl₄ (1 g, 2.9 mmol) in diethyl ether (20 cm³) cooled to –78 °C was added dropwise a solution of lithium isopropoxide [1 mol dm⁻³ in light petroleum (14.5 cm³), 14.5 mmol] with vigorous stirring. The red solution was allowed to warm slowly to room temperature when the colour had changed through brown to

pale green. The solvent was removed *in vacuo* and the green residue extracted with light petroleum ($5 \times 20 \text{ cm}^3$). The solution was filtered, concentrated to *ca.* 30 cm^3 , and held at -20°C overnight. The green needles were collected, washed with light petroleum, and dried *in vacuo*. Yield 1.1 g (69% based on ReOCl_4), decomp. $> 50^\circ \text{C}$ (Found: C, 33.2; H, 6.6; Cl, 5.8; O, 18.0. $\text{C}_{15}\text{H}_{35}\text{ClLi}_2\text{O}_6\text{Re}$ requires C, 33.0; H, 6.4; Cl, 6.4; O, 17.6%). Magnetic moment (298 K, Gouy method) μ_{eff} , *ca.* 0.9 B.M. Crystallisation from thf gave the solvate $\{\text{Li}[\text{ReO}(\text{OPr}^i)_5] \cdot \text{LiCl}(\text{thf})_2\}_2$, (B).

(6) *Tris[bis(trimethylsilyl)amido]oxorhenium(v)*.—To a solution of ReOCl_4 (1 g, 2.9 mmol) in diethyl ether (20 cm^3)

warm to room temperature when NaCl separated. Addition of trimethylphosphine (*ca.* 2 cm^3) caused a change in colour to red, and a brown insoluble precipitate was formed. After removal of solvent, the brown residue was extracted first with diethyl ether (discarded) and then with dichloromethane. Concentration of the dichloromethane solution gave diamond-shaped crystals in *ca.* 15% yield (Found: C, 47.8; H, 3.9; O, 9.7. $\text{C}_{30}\text{H}_{38}\text{O}_4\text{P}_2\text{Re}$ requires C, 48.0; H, 4.0; O, 9.7%).

(8) *Synthesis of Methylrhenium Compounds*.—(a) ReOMe_4 . To a solution of ReOCl_4 (1 g, 2.9 mmol) in diethyl ether (20 cm^3) at -100°C was added dropwise a cooled (-100°C)

TABLE 6

Crystal data and details of structure analyses

(i) Crystal Data	Compound (A) ^a	Compound (B) ^b	Compound (C) ^b
Formula	$\text{Re}_2\text{O}_3(\text{OME})_6$	$\{\text{Li}[\text{ReO}(\text{OPr}^i)_5] \cdot \text{LiCl}(\text{thf})_2\}_2$	$\text{Re}(\text{OPh})_4(\text{PMe}_3)_2$
<i>M</i>	606.6	1382.5	710.8
Crystal system	Monoclinic	Triclinic	Monoclinic
<i>a</i> /Å	12.142(1)	14.602(5)	10.079(2)
<i>b</i> /Å	15.369(1)	13.363(3)	10.527(3)
<i>c</i> /Å	7.311(1)	9.244(6)	14.443(3)
$\alpha/^\circ$	90	98.39(4)	90
$\beta/^\circ$	90.22(1)	102.35(5)	97.28(2)
$\gamma/^\circ$	90	68.71(2)	90
<i>U</i> /Å ³	1364.4	1637.1	1520.1
Space group	$P2_1/n$ (no. 14)	$P\bar{1}$ (no. 2)	$P2_1/c$ (no. 14)
<i>Z</i>	4	1	2
<i>D</i> _c /g cm ⁻³	2.953	1.402	1.553
<i>F</i> (000)	1104	704	710
Linear absorption coefficient, $\mu(\text{Mo-K}\alpha)/\text{cm}^{-1}$	179.7	38.5	41.5
Crystal size/mm	$0.33 \times 0.33 \times 0.08$	$0.30 \times 0.25 \times 0.22$	$0.24 \times 0.12 \times 0.08$
(ii) Data Collection			
Scan mode	$\omega/2\theta$	$\omega/2\theta$	$\omega/2\theta$
$\theta_{\text{min.}}, \theta_{\text{max.}}/^\circ$	1.5, 27	1.5, 24	1.5, 25
ω scan-width parameters <i>A, B</i> in $\omega = A + B \tan \theta$	0.85, 0.35	0.85, 0.35	0.85, 0.35
Horizontal aperture parameters <i>A, B</i> in aperture = $A + B \tan \theta$	4.0, 0.0	4.0, 0.0	4.0, 0.0
Total data	3212	4855	2894
Total unique data	2966	4761	2670
Observed data, $F_o > 4\sigma(F_o)$	1965	4061	1797
Crystal decay during data collection	5.8% (linear)	32.8% (linear)	None
(iii) Refinement			
No. of parameters	154	317	170
Weighting scheme coefficient <i>g</i> , in $w = 1/[\sigma^2(F_o) + g(F_o)^2]$	0.0022	0.0028	0.0012
Final $R = \Sigma \Delta F / \Sigma (F_o)$	0.059	0.069	0.055
$R' = (\Sigma w \Delta F^2 / \Sigma w F_o^2)^{1/2}$	0.072	0.086	0.071

^a Crystallographic measurements at 298 K. ^b Crystallographic measurements at 270 K.

cooled to -78°C was added with vigorous stirring a solution of $\text{LiN}(\text{SiMe}_3)_2$ [1 mol dm^{-3} in light petroleum (11.6 cm^3), 11.6 mmol]. The solution was allowed to warm slowly to ambient temperatures when it was bright red. The solvent was removed *in vacuo* and the residue extracted with light petroleum ($2 \times 20 \text{ cm}^3$), concentrated to *ca.* 5 cm^3 , and held at -20°C overnight. The red plates were collected and recrystallised from light petroleum to remove traces of $\text{LiN}(\text{SiMe}_3)_2$. The complex could also be purified by sublimation at 60°C (10^{-3} Torr). Yield of recrystallised material 1.25 g (63% based on ReOCl_4), m.p. $184\text{--}185^\circ \text{C}$ (Found: C, 31.5; H, 7.7; N, 6.0%; *M* 690. $\text{C}_{18}\text{H}_{34}\text{N}_3\text{OReSi}_6$ requires C, 31.7; H, 7.9; N, 6.2%; *M* 682).

(7) *trans-Tetraphenoxobis(trimethylphosphine)rhenium(IV)*, (C).—To ReOCl_4 (0.7 g) in diethyl ether (20 cm^3) at -78°C was added a solution of NaOPh (0.9 g, 4 equivalents) in thf (10 cm^3). The dark brown-purple solution was allowed to

solution of methyl-lithium [1.4 mol dm^{-3} in diethyl ether (8.3 cm^3), 11.6 mmol] with vigorous stirring. The solution was allowed to warm slowly (1 h) to ambient temperature when it was red. Degassed water (10 cm^3) was added dropwise with vigorous stirring, the solution cooled to -78°C and allowed to stand until excess of water had frozen, and filtered. The red-purple solution was evaporated to dryness at -40°C *in vacuo* and ReOMe_4 sublimed at 20°C (10^{-3} Torr); yield 0.51 g (67% based on ReOCl_4). Although the reaction is probably quantitative, loss of ReOMe_4 occurs on removal of solvent. This can be collected in a cold trap and more ReOMe_4 recovered by fractionation.

(b) $\text{Li}_2[\text{ReMe}_8]$. To a solution of ReOCl_4 (1 g, 2.9 mmol) in diethyl ether (20 cm^3) at -78°C was added dropwise with stirring a solution of methyl-lithium [1.4 mol dm^{-3} in diethyl ether (20.8 cm^3), 29 mmol]. The red solution was allowed to warm to, and held at, -10°C for 30 min, when

it was evaporated *in vacuo* and the residue extracted at -10°C with toluene ($3 \times 20\text{ cm}^3$). The solution was filtered, concentrated to *ca.* 15 cm^3 , and held at -78°C overnight. The thermally unstable red crystals were separated, washed with cooled light petroleum (10°C , $2 \times 5\text{ cm}^3$), and stored at -20°C ; yield 0.77 g (83% based on ReOCl_4).

(c) ReMe_6 . **CAUTION:** The red solution obtained from the reaction of ReOCl_4 with excess of methyl-lithium (above) was cooled to -78°C and degassed water (10 cm^3) was added dropwise with vigorous stirring. The solution was allowed to warm (until the ice melted) when it was green, re-cooled to -78°C , filtered, concentrated at -40°C *in vacuo* until green crystals appeared (*ca.* 5 cm^3), and then held at -78°C overnight. The crystals were collected and stored at -20°C . The compound may also be produced from $\text{Li}_2\text{-}[\text{ReMe}_6]$ by the same procedure and since it is potentially hazardous and should be treated with great caution it is better to prepare it from $\text{Li}_2[\text{ReMe}_6]$ in small quantities as required. In all operations with ReMe_6 , oxygen and moisture should be rigorously excluded.

(9) *X-Ray Crystallography.*—Crystals of compounds (A)–(C) were sealed under nitrogen in Lindemann capillaries. The determination of unit-cell constants and re-

TABLE 7

Fractional co-ordinates ($\text{Re} \times 10^5$, others $\times 10^4$) of the non-hydrogen atoms for $\text{Re}_2\text{O}_3(\text{OMe})_6$, (A)

	<i>x</i>	<i>y</i>	<i>z</i>
Re(1)	29 200(5)	20 564(5)	-511(8)
Re(2)	27 471(5)	37 146(5)	866(8)
O(1)	4 185(8)	2 959(7)	-483(14)
O(2)	2 502(8)	2 893(8)	-2 211(14)
O(3)	2 881(10)	2 852(8)	1 967(18)
O(4)	3 624(9)	1 265(9)	1 515(15)
O(5)	3 580(9)	1 373(9)	-1 920(13)
O(6)	1 589(9)	1 737(10)	-63(19)
O(7)	3 286(9)	4 513(9)	1 794(16)
O(8)	3 262(9)	4 517(9)	-1 675(15)
O(9)	1 359(10)	3 873(10)	119(22)
C(1)	4 844(13)	3 041(13)	-2 101(23)
C(2)	1 509(16)	2 848(14)	-3 257(30)
C(4)	3 669(16)	1 271(17)	3 563(25)
C(5)	4 256(17)	621(15)	-1 824(27)
C(7)	3 423(19)	4 402(16)	3 787(24)
C(8)	3 768(15)	5 360(14)	-1 456(30)

cording of intensity data were done using a Nonius CAD4 diffractometer and (Mo-K_α) radiation ($\lambda = 0.71069\text{ \AA}$) in a manner previously described in detail,²⁴ at room temperature (289 K) for (A) and at 270 K for both (B) and (C). The crystal data, important data-collection parameters, and refinement results are summarised in Table 6. The monoclinic space groups $P2_1/n$ for (A) and $P2_1/c$ for (C) were determined from systematic absences.

The centrosymmetric space group $P\bar{1}$ was confirmed for (B) by successful refinement. During data collection, compounds (A) and (B) suffered losses of 5.8 and 32.8% respectively in the intensities of their standard reflections, while (C) remained quite stable. All data sets were corrected for Lorentz, polarisation, and absorption effects, and those of (A) and (B) also for the crystal decay. The total number of reflections measured and those used in the refinement, $F_o > 4\sigma(F_o)$, are given in Table 6.

The rhenium atoms were determined from an *E* map in (A) and Patterson syntheses in both (B) and (C). The non-hydrogen atoms were then located and refined by successive electron-density syntheses and full-matrix least-

TABLE 8

Fractional co-ordinates ($\text{Re} \times 10^5$, others $\times 10^4$) of the non-hydrogen atoms for $\{\text{Li}[\text{ReO}(\text{OPr}^i)_5] \cdot \text{LiCl}(\text{thf})_2\}_2$, (B)

	<i>x</i>	<i>y</i>	<i>z</i>
Re	24 180(3)	17 657(3)	61 969(5)
Cl	1 086(2)	5 142(3)	4 568(4)
O(1)	1 839(5)	3 330(5)	7 139(8)
O(2)	1 116(6)	2 344(6)	4 860(9)
O(3)	3 019(6)	2 623(6)	5 300(9)
O(4)	3 502(5)	1 480(6)	7 816(10)
O(5)	1 773(6)	1 258(6)	7 357(9)
O(6)	2 824(7)	683(7)	4 973(10)
C(1)	1 705(9)	3 531(10)	8 626(14)
C(11)	648(11)	3 669(13)	8 767(17)
C(12)	2 001(17)	4 513(13)	9 306(21)
C(2)	642(11)	1 737(12)	3 788(18)
C(21)	12(15)	2 430(17)	2 541(20)
C(22)	15(23)	1 366(25)	4 554(32)
C(3)	3 463(9)	2 255(12)	3 961(15)
C(31)	2 724(15)	2 401(22)	2 595(20)
C(32)	4 224(20)	2 787(24)	4 030(25)
C(4)	4 431(9)	1 642(11)	8 052(15)
C(41)	4 706(12)	1 992(16)	9 648(18)
C(42)	5 163(12)	560(14)	7 507(24)
C(5)	2 211(11)	512(12)	8 485(16)
C(51)	2 650(14)	-627(12)	7 759(22)
C(52)	1 375(15)	697(18)	9 396(23)
O(7)	3 335(7)	4 904(8)	6 884(14)
C(71)	4 372(14)	4 530(17)	7 588(26)
C(72)	4 742(15)	5 472(18)	7 662(35)
C(73)	3 889(21)	6 361(20)	7 079(44)
C(74)	2 988(15)	6 044(16)	6 695(37)
Li(1)	2 497(16)	4 017(16)	6 131(30)
Li(2)	641(14)	3 835(15)	5 562(26)
O(8)	3 232(17)	7 291(33)	2 840(43)
C(81)	2 372(56)	8 176(33)	3 190(57)
C(82)	1 789(47)	8 094(60)	1 721(104)
C(83)	1 710(42)	7 341(50)	2 157(96)
C(84)	2 697(56)	6 776(41)	1 885(78)

squares refinement. In the final stages of refinement all the non-hydrogen atoms were assigned anisotropic temperature factors. None of the hydrogen atoms in any structure could be located from difference maps. These were ignored, except for the methine hydrogens in (B) and phenyl hydrogens in (C), which were inserted in idealised positions and included in the calculations of F_c . A common U_{iso} , refined to 0.08(2) and 0.09(2) \AA^2 respectively for the two groups of hydrogens in (B) and (C). The weighting scheme used was $w = 1/[\sigma^2(F_o) + g(F_o)^2]$, with the parameter g chosen to give nearly flat analyses of variances with

TABLE 9

Fractional co-ordinates ($\times 10^4$) of the non-hydrogen atoms for $\text{Re}(\text{OPh})_4(\text{PMe}_3)_2$, (C)

	<i>x</i>	<i>y</i>	<i>z</i>
Re	0	0	0
P	-1 594(3)	215(3)	1 172(2)
O(1)	1 271(9)	-281(7)	1 105(6)
O(2)	239(9)	1 863(8)	197(5)
C(1)	-3 395(14)	352(17)	842(11)
C(2)	-1 199(17)	1 681(14)	1 851(11)
C(3)	-1 457(14)	-1 023(13)	2 059(10)
C(11)	2 363(11)	-957(10)	1 406(7)
C(12)	2 629(11)	-2 136(11)	1 031(8)
C(13)	3 712(13)	-2 808(12)	1 379(9)
C(14)	4 621(15)	-2 326(14)	2 105(12)
C(15)	4 362(13)	-1 160(14)	2 493(9)
C(16)	3 237(13)	-512(13)	2 144(9)
C(21)	1 202(11)	2 662(10)	9(7)
C(22)	2 506(13)	2 420(12)	339(8)
C(23)	3 528(14)	3 265(14)	156(10)
C(24)	3 173(18)	4 355(15)	-372(11)
C(25)	1 869(19)	4 585(15)	-688(14)
C(26)	903(13)	3 761(11)	-489(9)

$(F_o/F_{\max})^{\frac{1}{2}}$ and $\sin \theta$. Final fractional atomic co-ordinates of the non-hydrogen atoms are given in Tables 7—9. Anisotropic temperature factors of these atoms, hydrogen-atom parameters, lists of observed and calculated structure factors, and a table containing all the bond lengths and angles in (B) have been deposited in Supplementary Publication No. SUP 22891 (39 pp.).*

Neutral-atom scattering factors were taken from refs. 25 (H), 26 (Re), and 27 (others), with those of the heavier elements being modified for anomalous dispersion, taking the $\Delta f'$ and $\Delta f''$ values from ref. 28. All calculations were performed on the ICL2980 computer at Queen Mary College using the programs SHELX²⁹ (structure determination and refinement) and PLUTO³⁰ (diagrams).

We thank the S.R.C. for a Studentship (to P. G. E.) and a Fellowship (to K. M. A. M.), and Dr. J. F. Gibson for assistance with e.s.r. measurements.

[0/355 Received, 4th March, 1980]

* For details see Notices to Authors No. 7, *J.C.S. Dalton*, 1979, Index issue.

REFERENCES

- ¹ A. P. Ginsberg and M. J. Hawkes, *J. Amer. Chem. Soc.*, 1968, **90**, 5930.
- ² N. P. Johnson, C. J. L. Lock, and G. Wilkinson, *J. Chem. Soc.*, 1964, 1054 and refs. therein.
- ³ R. A. Jones and G. Wilkinson, *J.C.S. Dalton*, 1978, 1063.
- ⁴ J. G. F. Druce, *J. Chem. Soc.*, 1937, 1407.
- ⁵ P. G. Edwards, G. Wilkinson, K. M. A. Malik, and M. B. Hursthouse, *J.C.S. Chem. Comm.*, 1979, 1158.
- ⁶ W. K. Wong and G. Wilkinson, unpublished work.
- ⁷ W. A. Nugent and R. L. Harlow, *J.C.S. Chem. Comm.*, 1979, 1105.
- ⁸ K. W. Bagnall, D. Brown, and R. Colton, *J. Chem. Soc.*, 1964, 3017; R. Colton, *Austral. J. Chem.*, 1965, **18**, 435.
- ⁹ D. A. Edwards and A. A. Woolf, *J. Chem. Soc. (A)*, 1966, 91.
- ¹⁰ A. Guest and C. J. L. Lock, *Canad. J. Chem.*, 1971, **49**, 603.
- ¹¹ R. Lincoln and G. Wilkinson, *Inorg. Synth.*, vol. 20, in the press.
- ¹² A. Brukl and K. Ziegler, *Chem. Ber.*, 1932, **65**, 916; C. J. Wolf, A. F. Clifford, and W. H. Joynton, *J. Amer. Chem. Soc.*, 1957, **79**, 4257.
- ¹³ K. Dehnicke and W. Liese, *Chem. Ber.*, 1977, **110**, 3959.
- ¹⁴ K. Mertis, D. H. Williamson, and G. Wilkinson, *J.C.S. Dalton*, 1975, 607.
- ¹⁵ See, for example, A. R. Middleton, A. F. Masters, and G. Wilkinson, *J.C.S. Dalton*, 1979, 542.
- ¹⁶ J. F. Gibson, G. M. Lack, K. Mertis, and G. Wilkinson, *J.C.S. Dalton*, 1976, 1492.
- ¹⁷ F. A. Cotton and G. Wilkinson, 'Advanced Inorganic Chemistry,' 4th edn., Wiley, New York, 1980, p. 884.
- ¹⁸ K. Mertis and G. Wilkinson, *J.C.S. Dalton*, 1976, 1488.
- ¹⁹ M. B. Hursthouse, S. A. A. Jayaweena, and A. Quick, *J.C.S. Dalton*, 1979, 279.
- ²⁰ M. Bochmann, G. Wilkinson, G. B. Young, M. B. Hursthouse, and K. M. A. Malik, *J.C.S. Dalton*, 1980, 901.
- ²¹ M. B. Hursthouse and M. Motevalli, *J.C.S. Dalton*, 1979, 1362.
- ²² D. Bright and J. A. Ibers, *Inorg. Chem.*, 1969, **8**, 703.
- ²³ D. Bright and J. A. Ibers, *Inorg. Chem.*, 1968, **7**, 1099.
- ²⁴ M. B. Hursthouse, R. A. Jones, K. M. A. Malik, and G. Wilkinson, *J. Amer. Chem. Soc.*, 1979, **101**, 4128.
- ²⁵ R. F. Stewart, E. R. Davidson, and W. T. Simpson, *J. Chem. Phys.*, 1965, **42**, 3175.
- ²⁶ D. T. Cromer and J. R. Waber in 'International Tables for X-Ray Crystallography,' Kynoch Press, Birmingham, 1974, vol. 4, p. 101.
- ²⁷ D. T. Cromer and J. B. Mann, *Acta Cryst.*, 1968, **A24**, 321.
- ²⁸ D. T. Cromer and D. Libermann, *J. Chem. Phys.*, 1970, **53**, 1891.
- ²⁹ SHELX Crystallographic Calculation Program, G. M. Sheldrick, University of Cambridge, 1976.
- ³⁰ PLUTO Program for Crystallographic Drawing, W. D. S. Motherwell, University of Cambridge.