

Additions and Corrections

Alkyl and Hydrido Derivatives of Tetrakis(trimethylphosphine)-osmium(II). X-Ray Crystal Structure of the Metallacycle $\text{Os}[(\text{CH}_2)_2\text{SiMe}_2](\text{PMe}_3)_4$ (1984, 877)

Torsten Behling, Gregory S. Girolami, Geoffrey Wilkinson, Richard E. Somerville, and Michael B. Hursthouse

Page 877, third line from the bottom in the left-hand column. The word 'required' should be replaced by 'equatorial'.

Page 878, Table 1. The following ^1H and $^{13}\text{C}\{-^1\text{H}\}$ n.m.r. data should be included.

Compound	^1H	Assignment	$^{13}\text{C}\{-^1\text{H}\}$
<i>fac</i> -OsCl(CH ₂ PMe ₂)(PMe ₃) ₃	0.07 (m)	Os-CH ₂	-20.4 (dd, 38.3, 9.5)
	0.62 (m)	Os-CH ₂	
	1.20 (dd, 9.8, 2.9)	PMe ₂	3.7 (d, 18.0)
	1.73 (dd, 10.4, 2.5)	PMe ₂	10.2 (d, 13.8)
	1.12 (d, 8.2)	PMe ₃	22.2 (d, 29.0)
	1.47 (dd, 7.8, 1.7)	PMe ₃	25.0 (m)
	1.57 (dd, 6.9, 1.0)	PMe ₃	

Page 878, Table 2. The following $^{31}\text{P}\{-^1\text{H}\}$ n.m.r. data should be included.

Compound	Spin system	Chemical shifts (p.p.m.)	Coupling constants (Hz)
<i>cis</i> -OsH ₂ (PMe ₃) ₄	A ₂ B ₂	$\delta_{\text{A}} = -47.9$ $\delta_{\text{B}} = -53.2$	$J_{\text{AB}} = 17.9$