

Synthesis and Characterization of Gold(III) Adducts and Cyclometallated Derivatives with 2-Substituted Pyridines. Crystal Structure of $[\text{Au}\{\text{NC}_5\text{H}_4(\text{CMe}_2\text{C}_6\text{H}_4)-2\}\text{Cl}_2]^{\dagger}$

Maria Agostina Cinellu,^a Antonio Zucca,^a Sergio Stoccoro,^a Giovanni Minghetti^{*,a}
Mario Manassero^{*,b} and Mirella Sansoni^b

^a Dipartimento di Chimica, Università di Sassari, via Vienna 2, I-07100 Sassari, Italy

^b Dipartimento di Chimica Strutturale e Stereochimica Inorganica, Università di Milano, Centro CNR, via Venezian 21, I-20133 Milano, Italy

The 2-benzylpyridines HL $[\text{NC}_5\text{H}_4(\text{CH}_2\text{Ph})-2$, $-(\text{CHMePh})-2$ and $(\text{CMe}_2\text{Ph})-2]$ reacted with $\text{AuCl}_3 \cdot 2\text{H}_2\text{O}$ or $\text{Na}[\text{AuCl}_4]$ to give adducts $[\text{Au}(\text{HL})\text{Cl}_3]$ or cyclometallated derivatives $[\text{AuLCl}_2]$. The metallation reaction involves direct activation of a C–H bond of the phenyl substituent. With the 2-alkylpyridines $\text{NC}_5\text{H}_4(\text{CMe}_3)-2$ and $\text{NC}_5\text{H}_4(\text{CH}_2\text{CMe}_3)-2$ (HL) only the tetrachloroaurate(III) salts $[\text{H}_2\text{L}][\text{AuCl}_4]$ were isolated. The structure of $[\text{Au}\{\text{NC}_5\text{H}_4(\text{CMe}_2\text{C}_6\text{H}_4)-2\}\text{Cl}_2]$ has been determined by X-ray diffraction. It reveals a six-membered ring having a boat-like conformation: there is a rather short interaction between one hydrogen of the Me group in axial position and the gold atom, $\text{Au} \cdots \text{H}$ 2.56(5) Å. The cyclometallated species react with PPh_3 to give cationic complexes $[\text{AuL}(\text{PPh}_3)\text{Cl}]^+$ and with 1,2-bis(diphenylphosphino)ethane (dppe) to yield $[\text{AuL}(\text{dppe})\text{Cl}]^+$. Under pressure (50 atm, 60 °C) of CO extrusion of the metal occurs. The organic products formed likely arise from insertion of CO into the Au–C bond. The reactivity of the cycloaurated species is compared with that of analogous palladium(II) derivatives described previously.

Most reviews which deal with cyclometallation reactions neglect gold chemistry, attention being focused on other d⁸ ions, e.g. Pd^{II}, Pt^{II} and Rh^I.¹

Nevertheless, a number of cycloaurated species involving nitrogen ligands has been described in previous years. Vicente *et al.*² have reported the synthesis, characterization and reactivity of many species arising from azobenzene and *N,N*-dimethylbenzylamines: all of them imply an aromatic C–H activation to give five-membered rings. The auration was not achieved directly: in most cases organomercury derivatives were used as arylating agents. Constable and co-workers³ have described gold(III) cyclometallated species with 6-(2-thienyl)- or 6-phenyl-2,2'-bipyridine, *i.e.* with potentially terdentate ligands, as well as with likewise substituted pyridines: some of these species were obtained by direct activation of an aromatic C–H bond. Very recently, the complex $[\text{Au}(\text{dpphen})\text{Cl}]^+$, where dpphen = 2,9-diphenyl-1,10-phenanthroline,⁴ has been reported.

Previously we have reported cyclometallated species of platinum(II) and palladium(II) with substituted pyridines and 2,2'-bipyridines both with benzyl and alkyl substituents.⁵ We have now extended our investigations to gold(III) and report the first results with 2-benzyl- and 2-alkyl-pyridines. The results for 2,2'-bipyridines will be described in a forthcoming paper.

With 2-benzylpyridines, activation of C(sp²)–H bonds was achieved directly. The structure in the solid state of one of these cycloaurated species was solved by X-ray diffraction: this is the first structure of a gold(III) six-membered C,N derivative.

Finally we deemed it worthwhile to study the reactivity of the cycloaurated derivatives with carbon monoxide. To the best of our knowledge, the reaction of organogold(III) derivatives with

CO has not been previously investigated. The behaviour of the gold species is compared with that of the palladium analogues.^{5d,6} A brief report on part of this work has been given.⁷

Results and Discussion

The 2-benzylpyridines (HL = $\text{NC}_5\text{H}_4\text{-CH}_2\text{Ph-2}$, $-\text{CHMePh-2}$ or $-\text{CMe}_2\text{Ph-2}$), and the 2-alkylpyridines (HL = $\text{NC}_5\text{H}_4\text{-CMe}_3\text{-2}$ and $-\text{CH}_2\text{CMe}_3\text{-2}$) were prepared from the appropriate cyanides, according to literature methods.⁸


The reaction of either $\text{AuCl}_3 \cdot 2\text{H}_2\text{O}$ or $\text{Na}[\text{AuCl}_4]$ with the 2-benzylpyridines was carried out in water at room temperature. With $\text{NC}_5\text{H}_4\text{-CH}_2\text{Ph-2}$ and $-\text{CHMePh-2}$ the reaction leads to the adducts $[\text{Au}(\text{HL})\text{Cl}_3]$ **1a** and **2a** in fairly good yields, *ca.* 75%. With $\text{NC}_5\text{H}_4(\text{CMe}_2\text{Ph})-2$, under the same conditions, only the salt $[\text{H}_2\text{L}][\text{AuCl}_4]$ **3s** was isolated (see Scheme 1). The formulation of **1a** and **2a** as $[\text{Au}(\text{HL})\text{Cl}_3]$ adducts is supported by microanalytical data, conductivity measurements, IR, NMR and mass spectra. In the IR spectra a strong absorption around 360 cm⁻¹ is assignable to a Au–Cl stretching mode, at a value comparable to that observed for other 1:1 adducts of AuCl_3 with heterocyclic ligands.⁹ In the ¹H NMR spectra all the resonances are shifted downfield upon co-ordination: in the aromatic region the spectra exhibit the expected nine resonances.

With $\text{NC}_5\text{H}_4\text{-CMe}_3\text{-2}$ and $-\text{CH}_2\text{CMe}_3\text{-2}$ only the salts $[\text{H}_2\text{L}][\text{AuCl}_4]$ **4s** and **5s** were isolated: so far, in spite of several attempts carried out under various conditions, adducts were not obtained.

The adducts **1a** and **2a** as well as the salt **3s** can be converted into the cycloaurated species $[\text{AuLCl}_2]$ **1m–3m** by warming their suspensions in aqueous acetonitrile (*ca.* 4:1). It is noteworthy that the metallated species can be obtained also from $\text{AuCl}_3 \cdot 2\text{H}_2\text{O}$ and the pyridines under reflux in water, *i.e.*

[†] Supplementary data available: see Instructions for Authors, *J. Chem. Soc., Dalton Trans.*, 1995, Issue 1, pp. xxv–xxx.

Non-SI unit employed: atm = 101 325 Pa.



Scheme 1 (i) $AuCl_3 \cdot 2H_2O$, water, room temperature (r.t.); (ii) MeCN-water, reflux

by direct activation of the C–H bond. Complexes **1m–3m** are isolated in good yields as white solids: they are not electrolytes and are slightly soluble in the most common solvents. The solubility in chlorinated solvents is sufficient to allow well resolved 1H NMR spectra to be obtained which unambiguously show the activation of an aromatic C–H bond. The strong deshielding of the H(6) proton, usually observed when a chlorine is in the proximity of the pyridine ring,¹⁰ is consistent with co-ordination of the nitrogen atom. The spectra indicate a boat conformation of the six-membered ring, as in the previously reported palladium derivatives.^{5d} No rapid inversion of the boat occurs on the NMR time-scale up to room temperature, as shown by the AB pattern displayed by the CH_2 resonance of complex **1m**, as well as by the two distinct resonances for the Me substituents in **3m**. The latter resonances are both at low field (δ 2.06 and 2.35, CD_2Cl_2) with respect to the free pyridine (δ 1.75, $CDCl_3$). The more deshielded one is likely to be associated with the methyl group pointing toward the metal, *i.e.* in a pseudo-axial position. In agreement with a boat conformation, two diastereomers are formed in the case of complex **2m** (*a* and *b*).

The structure of complex **3m** in the solid state has been solved by X-ray diffraction. It consists of the packing of $[AuCl_2]$ molecules separated by normal van der Waals interactions. An ORTEP¹¹ view of the molecule is shown in Fig. 1. Crystal data, selected bond distances and angles, and atomic coordinates are listed in Tables 1, 2 and 3, respectively. The gold atom displays an almost regular square-planar co-ordination, with a distortion in the N–Au–C(9) angle, 85.7(1)°. The bite of the six-membered cyclometallated ring is comparable with that observed in $[PtL(PPh_3)Cl]$ [$HL = NC_5H_4(CHMePh)-2$]^{5f} [$C-Pt-N$ 85.1(1)°], as well as in a number of six-membered N–Au–N rings [range 84.7(3)–86.8(2)°].¹² The bond lengths involving the gold atom are all normal. The Au–Cl(2) bond [2.390(1) Å] is much longer than Au–Cl(1) [2.282(1) Å] as expected owing to the great *trans* influence of the aryl carbon atom. The Au–C(9) bond length [2.021(3) Å] is identical with the Au–C(sp²) distance found in $[Au\{C_6H_3(N=NC_6H_4Me-4)-2-Me-5\}Cl_2]$, 2.021(5) Å,^{2m} and the Au–N bond length, 2.041(3) Å, is statistically coincident with Au–N(sp²), 2.031(6) Å, found in




Fig. 1 An ORTEP view of compound $[Au\{NC_5H_4(CMe_2C_6H_4)-2\}Cl_2]$ **3m**

Table 1 Crystallographic data for complex **3m**

Formula	$C_{14}H_{14}AuCl_2N$
<i>M</i>	464.2
Colour	Colourless
Crystal system	Triclinic
Space group	$P\bar{1}$
<i>a</i> /Å	8.379(1)
<i>b</i> /Å	8.712(2)
<i>c</i> /Å	11.533(2)
α /°	72.05(2)
β /°	79.18(1)
γ /°	62.09(2)
<i>U</i> /Å ³	706.8(3)
<i>Z</i>	2
<i>F</i> (000)	436
<i>D</i> _c /g cm ⁻³	2.181
Crystal dimensions/mm	0.17 × 0.23 × 0.48
μ (Mo–K α)/cm ⁻¹	107.5
Minimum, maximum transmission factors	0.68, 1.00
Scan mode	ω
ω -Scan width/°	1.2 + 0.35 tan θ
θ Range/°	3–27
Reciprocal space explored	+ <i>h</i> , ± <i>k</i> , ± <i>l</i>
Measured reflections	3087
Unique observed reflections [<i>I</i> > 3 σ (<i>I</i>)]	2829
Final <i>R</i> and <i>R'</i> ^a	0.025, 0.032
No. of variables	176
Goodness of fit ^b	1.41

^a $R = [\sum(|F_o - k|F_c|)/\sum F_o]$, $R' = [\sum w(F_o - k|F_c|)^2/\sum wF_o^2]^{\frac{1}{2}}$. ^b $[\sum w(F_o - k|F_c|)^2/(N_o - N_v)]^{\frac{1}{2}}$, where $w = 1/[\sigma(F_o)]^2$, $\sigma(F_o) = [\sigma^2(I) + (0.04I)^2]^{\frac{1}{2}}/2F_oL_p$, N_o is the number of observations and N_v the number of variables.

Table 2 Selected bond distances (Å) and angles (°) with estimated standard deviations (e.s.d.s) in parentheses for compound **3m**

Au–Cl(1)	2.282(1)	Au–Cl(2)	2.390(1)
Au–N	2.041(3)	Au–C(9)	2.021(3)
Cl(1)–Au–Cl(2)	91.6(4)	Cl(1)–Au–N	176.2(8)
Cl(1)–Au–C(9)	90.8(1)	Cl(2)–Au–N	91.9(9)
Cl(2)–Au–C(9)	177.4(1)	N–Au–C(9)	85.7(1)

Table 3 Fractional atomic coordinates with e.s.d.s in parentheses for the refined atoms of complex **3m**


Atom	x	y	z
Au	0.032 43(2)	-0.044 97(1)	0.308 93(1)
Cl(1)	-0.117 9(2)	0.253 5(1)	0.217 0(1)
Cl(2)	0.218 4(1)	0.009 6(1)	0.409 5(1)
N	0.161 8(4)	-0.315 3(4)	0.382 2(3)
C(2)	0.216 1(4)	-0.434 0(4)	0.313 4(3)
C(3)	0.300 9(6)	-0.616 1(5)	0.370 7(4)
C(4)	0.329 4(7)	-0.673 6(6)	0.493 8(5)
C(5)	0.272 7(7)	-0.551 5(7)	0.561 0(4)
C(6)	0.188 5(6)	-0.371 4(6)	0.502 7(4)
C(7)	0.179 1(4)	-0.363 9(4)	0.177 3(3)
C(8)	-0.025 9(4)	-0.243 5(4)	0.168 1(3)
C(9)	-0.115 6(5)	-0.102 6(5)	0.224 2(3)
C(10)	-0.302 4(6)	-0.004 2(6)	0.226 3(4)
C(11)	-0.403 8(5)	-0.042 5(7)	0.166 7(5)
C(12)	-0.317 2(5)	-0.174 1(6)	0.104 5(4)
C(14)	0.240 8(6)	-0.520 5(5)	0.119 2(4)
C(15)	0.287 3(5)	-0.257 0(5)	0.109 4(4)
H(151)	0.263(6)	-0.168(6)	0.142(5)
H(152)	0.256(6)	-0.189(7)	0.032(5)
H(153)	0.415(6)	-0.321(6)	0.122(5)

the cation $[\text{Au}(\text{dmp})(\text{NC}_9\text{H}_6\text{O})]^+$ [dmp = 2-(dimethylamino-methyl)phenyl].^{2b} The six-membered metallacycle is in a boat conformation, with atoms N, C(2), C(8) and C(9) essentially coplanar [maximum deviations from their best plane $-0.016(4)$ Å for C(2) and $+0.015(4)$ Å for C(8)]. This best plane forms dihedral angles with planes C(2)–C(7)–C(8) and N–Au–C(9) of $44.6(3)$ and $38.2(2)^\circ$, respectively. The three hydrogen atoms bonded to C(15) are in a staggered conformation with respect to the C(7)–C(2), C(7)–C(8), and C(7)–C(14) bonds. As a consequence atom H(151) is in a pseudo-axial position with respect to the metal co-ordination plane. The Au...H(151) distance, $2.56(5)$ Å, is rather short, well below the value reported as upper limit for long-range metal–hydrogen interactions in platinum(II) chemistry.¹³ Bond lengths and angles within the bidentate ligand are normal, with the two aromatic rings strictly planar and forming a dihedral angle of $63.2(2)^\circ$ with each other.

Reactivity of Cyclometallated Derivatives 1m–3m.—To evaluate the stability of the cycloaurated ring, we investigated first the reaction of compounds **1m–3m** with triphenylphosphine. The reaction (molar ratio $\text{PPh}_3:\text{Au} = 1:1$ or $2:1$) carried out in the presence of NaBF_4 gives the cationic species $[\text{AuL}(\text{PPh}_3\text{Cl})][\text{BF}_4^-]$ **1p–3p**. The spectroscopic data are consistent with **1p–3p** being cyclometallated species. In particular, the ^1H NMR spectra suggest that the rigid boat-like conformation of the ring is maintained: in agreement, diastereomers (*a* and *b*) are observed in the case of complex **2p**. Of the two possible geometrical isomers, *trans*-P–Au–N or *trans*-P–Au–C, the IR and NMR spectra, taken together, support a *trans*-P–Au–N arrangement. Indeed in the IR spectra the Au–Cl stretching vibrations are observed at values consistent with a chlorine *trans* to a carbon atom^{2c} (see Experimental section) and in the ^1H NMR spectra the H(6) protons are strongly deshielded (Table 4).

No further reaction occurs on addition of an excess of PPh_3 to complex **1p** (^{31}P NMR criterion): thus it seems that these gold(III) derivatives are rather stable towards both ring opening and reduction to gold(I) species.

At variance, the nitrogen atom can be displaced from the metal by reaction with the potentially chelating diphosphine $(\text{Ph}_2\text{PCH}_2)_2$ (dppe) (see Scheme 2). Reaction of complexes **1m** and **2m** with dppe leads to the cationic derivatives $[\text{AuL}(\text{dppe})\text{Cl}][\text{BF}_4^-]$ **1d** and **2d**. The two sharp resonances



Scheme 2 (i) PPh_3 , BF_4^- ; (ii) dppe , BF_4^- ; (iii) AgBF_4 ; (iv) $\text{HBF}_4 \cdot \text{Et}_2\text{O}$

observed in the $^{31}\text{P}\{-^1\text{H}\}$ NMR spectra at room temperature are consistent with square-planar species: the difference between their chemical shifts suggests that the phosphorus atoms are *trans* to ligands having markedly different *trans* influence, such as Cl and C. In the ^1H NMR spectra, the single resonances observed for the protons of the substituents at the benzylic carbon atom indicate a non-co-ordinated nitrogen atom (see Table 4). In agreement, complex **1d** was easily protonated by reaction with HBF_4 . The co-ordination of the pyridine ring can be restored and the chloride ligand removed by means of silver tetrafluoroborate to give complexes **1d'** and **2d'**.

Although the chemistry of gold with carbon monoxide is rather sparse,¹⁴ we deemed it worth investigating the reaction of the cyclometallated species **1m–3m** with CO in order to compare their behaviour with that of homologous palladium(II) and platinum(II) derivatives.^{5d,f,6} Under mild conditions, *e.g.* bubbling CO at room temperature through a solution of **1m** in CH_2Cl_2 or in $\text{CH}_2\text{Cl}_2\text{–EtOH}$, no reaction occurs: the starting complex is almost quantitatively recovered, only trace amounts of gold being formed. Under more vigorous conditions, 50 atm of CO, 60°C in $\text{CH}_2\text{Cl}_2\text{–EtOH}$ solution (*ca.* 4:1), a mirror of gold and bright-yellow organic molecules are formed (see Scheme 3). The organic derivatives were separated by chromatography on silica gel and fully characterized by mass and NMR (^1H , $^{13}\text{C}\{-^1\text{H}\}$) spectra (see Table 5). From **1m**, three carbonylated species were obtained, I–III. Compound II, identical to that obtained as the only product from $[\{\text{Pd}(\text{L})\text{Cl}\}_2]$ [$\text{HL} = \text{NC}_5\text{H}_4(\text{CH}_2\text{Ph})\text{–}2$] and CO under comparable conditions,⁶ is likely to be the first product of the reaction. Its oxidative chlorination, brought about by gold(III) chloride, can easily explain the formation of compound I. Indeed, in a separate experiment II was converted into I upon treatment with AuCl_3 . The C–C coupling, involved in the formation of compound III, may arise from the activation of a C–Cl bond. It is worth noting that both chlorination and C–C coupling of the ligand have previously been observed in the reaction of $\text{Na}[\text{AuCl}_4]$ with 2-(2-thienyl)pyridine.¹⁵ When the benzylic carbon atom bears a substituent, as in complex **2m**, one only product, 11-methylbenzo[*b*]quinolizin-6-one IV is obtained, as in the case of the palladium analogue.^{5d}

The formation of the species I–IV implies insertion of CO into the metal–carbon bond, followed by nucleophilic intramolecular attack of the nitrogen atom on an acyl intermediate. Consistent with the existence of this intermediate, in the case of **3m** the expected ester V is obtained. As far as we know, insertion of CO into gold(III)–carbon bonds is unprecedented. The analogies observed in the behaviour of

Table 4 Proton NMR data^a

Compound	Solvent	H(6)	Other aromatics	CH	CH ₂	CH ₃	Other
NC ₅ H ₄ -(CH ₂ Ph)-2	CDCl ₃	8.55 (d)	7.6–7.05		4.16 (s)		
1a	CDCl ₃	8.69 (d)	8.0–7.25		4.73 (s)		
1m	CD ₂ Cl ₂	9.27 (d)	8.1–7.1		H(A) 4.56 (d); H(B) 4.08 (d) [<i>J</i> (AB) = 15]		
1p	CDCl ₃ CD ₂ Cl ₂	8.95 (d) 9.10 (d)	8.0–6.4 8.1–6.5		4.37 (br s) H(A) 4.58 (d); H(B) 4.29 (d) [<i>J</i> (AB) = 15]		
1d	CDCl ₃	8.15 (d)	7.9–6.75		3.94 (s)		3.45 (m), 2.95 (m) CH ₂ of dppe
1d'	CD ₂ Cl ₂	8.33 (d)	8.25–7.3		4.36 (s)		3.38 (m), 2.87 (m) CH ₂ of dppe
1d''^b	CD ₂ Cl ₂	8.70 (d)	8.3–6.8		4.10 (br)		2.2 (br m, 1 H), 3.4–4.5 (br m, 3 H) CH ₂ of dppe
NC ₅ H ₄ -(CHMePh)-2	CDCl ₃	8.58 (d)	7.6–7.0	4.28 (q, 7.0)		1.70 (d, 7.0)	
2a	Me ₂ SO	8.48 (d)	7.7–7.1	4.24 (q, 7.0)		1.58 (d, 7.0)	
2m^c	CDCl ₃	8.66 (d)	8.0–7.3	5.30 (q, 7.2)		1.86 (d, 7.2)	
	CDCl ₃	<i>a</i> [2] <i>b</i> [1]	9.40 (d) 9.30 (d)	8.1–7.1	4.43 (q, 7.2) 4.68 (q, 7.2)	2.16 (d, 7.2) 1.94 (d, 7.2)	
	Me ₂ SO	<i>a</i> [3] <i>b</i> [1]	9.21 (d) 9.16 (d)	8.3–7.0	4.84 (q, 7.5) 4.73 (q, 7.0)	1.98 (d, 7.5) 1.86 (d, 7.0)	
2p^c	CDCl ₃	<i>a</i> [2] <i>b</i> [1]	9.15 (t) ^d 9.11 (t) ^d	8.0–6.4	4.80 (q, 7.4) 4.67 (q, 6.9)	2.14 (d, 7.4) 1.99 (d, 6.9)	
2d	CD ₂ Cl ₂	8.18 (d)	7.9–6.9	4.04 (q, 6.9)		1.21 (d, 6.9)	3.60 (m), 2.86 (m) CH ₂ of dppe
2d'	CD ₂ Cl ₂	8.68 (d)	8.3–6.8	1.11 (d, 7.2)		4.46 (q, 7.2)	2.57 (m, 1 H), 4.2–3.7 (m, 3H) CH ₂ of dppe
NC ₅ H ₄ -(CMe ₂ Ph)-2	CDCl ₃	8.57 (d)	7.6–7.1			1.75 (s)	
3s	(CD ₃) ₂ CO	8.94 (d)	8.9–7.3			1.99 (s)	
3m	CD ₂ Cl ₂	9.39 (d)	8.1–7.0			2.35 (s), 2.06 (s)	
3p	CDCl ₃	9.32 (t) ^d	8.15–6.45			2.36 (s), 2.11 (s)	
NC ₅ H ₄ -(CMe ₃)-2	CDCl ₃	8.55 (d)	7.6–7.05			1.36 (s)	
4s	CDCl ₃	8.86 (d)	8.7–8.1			1.62 (s)	
	(CD ₃) ₂ CO	8.96 (d)	8.8–8.1			1.63 (s)	
NC ₅ H ₄ -(CH ₂ CMe ₃)-2	CDCl ₃	8.53 (d)	7.6–7.1		2.68 (s)	0.94 (s)	
5s	CDCl ₃	8.93 (br m)	8.5–7.8		3.10 (s)	1.08 (s)	

^a Spectra recorded at room temperature, chemical shifts in ppm from internal SiMe₄, coupling constants in Hz. ^b At –40 °C, CH₂ (pyridine) AB system, δ 3.91 (d, 1 H) and 4.04 (d, 1 H), *J*(AB) = 17.0 Hz; CH₂ (dppe), δ 2.11 (m, 1 H), 3.39 (m, 1 H), 4.05 (m, 1 H) and 4.47 (m, 1 H). ^c Compounds **2m** and **2p** are present in solution as diastereomers, *a* and *b* (see text), integral ratios are in square brackets. ^d Due to coupling with phosphorus.


palladium(II) and gold(III) cyclometallated derivatives suggests the potential reactivity of the latter species, which is still largely unexplored.

Experimental

General.—The pyridines HL were prepared according to literature methods;⁷ AuCl₃·2H₂O and Na[AuCl₄]·2H₂O were obtained from Aldrich and Johnson Matthey respectively. Elemental analyses were performed with a Perkin-Elmer Elemental Analyser 240B by Mr. A. Canu (Dipartimento di Chimica, Università di Sassari). Conductivities were measured with a Philips PW 9505 conductimeter. Infrared spectra were recorded with a Perkin-Elmer 983 spectrophotometer using Nujol mulls, ¹H, ¹³C-¹H and ³¹P-¹H NMR spectra with a Varian VXR 300 spectrometer operating at 299.9, 75.4 and 121.4 MHz, respectively. The ¹H and ¹³C-¹H NMR data are collected in Tables 4 and 5; chemical shifts are given in ppm relative to internal tetramethylsilane (¹H) and external 85% H₃PO₄ (³¹P). Mass spectra were obtained with a VG 7070 instrument operating under FAB, with 3-nitrobenzyl alcohol as supporting matrix, or electron-impact (EI) conditions.

Preparations.—[Au{NC₅H₄(CH₂Ph)-2}Cl₃] **1a**. An aqueous solution (20 cm³) of AuCl₃·2H₂O (1.55 mmol) was added to pure NC₅H₄(CH₂Ph)-2 (0.262 g, 1.55 mmol): the resulting yellow suspension was stirred for 3 d at room temperature. The solid product was filtered off and air dried to give complex **1a** (0.540 g, 74%). Recrystallization from acetone–diethyl ether yielded an orange crystalline product: m.p. 149–150 °C [Found: C, 30.8; H, 2.3; N, 2.9%; *M*⁺ 471 (EI). C₁₂H₁₁AuCl₃N requires C, 30.5; H, 2.3; N, 2.9%; *M* 471]; $\tilde{\nu}_{\max}/\text{cm}^{-1}$ 1601s, 1563m, 1485s, 1477s, 1441vs, 1160m, 1105m, 766vs, 740vs, 731vs, 695s, 474s, 458s and (Au–Cl) 362vs; *m/z* 471 (*M*⁺) and 435 (*M* – HCl) (EI).

[Au{NC₅H₄(CH₂C₆H₄)-2}Cl₂] **1m**. **Method (a)**. A suspension of complex **1a** (0.471 g, 1 mmol) in MeCN (5 cm³) and water (20 cm³) was refluxed until the precipitate became white (ca. 8 h). The solid product was filtered off and air dried to give **1m** (0.400 g, 92%). Recrystallization from dichloromethane–diethyl ether yielded the analytical sample: m.p. 244–245 °C [Found: C, 33.2; H, 2.3; N, 3.1%; *M*⁺ 435 (EI). C₁₂H₁₀–AuCl₂N requires C, 33.0; H, 2.3, N, 3.2%; *M* 435]; $\tilde{\nu}_{\max}/\text{cm}^{-1}$ 1607m, 1563m, 1479s, 1025s, 779m, 751vs, (Au–Cl) 358vs and 294vs.

Scheme 3 (i) 50 atm CO, 60 °C, CH₂Cl₂-EtOH; (ii) K₂CO₃, CH₂Cl₂Table 5 Proton and ¹³C-¹H NMR data for compounds I-IV^a

Compound	I ^b (R = Cl)	II ^b (R = H)	III ^c	IV ^{c,d} (R = Me)
H ¹	7.88 (m)	7.19 (d)	6.89 (d)	7.57 (d)
H ²	7.16 (dd)	6.90 (dd)	6.89 (d)	6.99 (dd)
H ³	6.70 (dd)	6.56 (dd)	6.73 (m)	6.60 (dd)
H ⁴	8.89 (d)	8.77 (d)	9.02 (d)	8.89 (d)
H ⁷	8.66 (d)	8.57 (d)	8.73 (m)	8.69 (d)
H ⁸	7.58 (dd)	7.45 (dd)	7.53 (m)	7.53 (dd)
H ⁹	7.86 (m)	7.67 (dd)	7.56 (m)	7.79 (dd)
H ¹⁰	8.15 (d)	7.60 (d)	7.15 (m)	7.91 (d)
CH ₃				2.57 (s)
H ¹¹		6.76 (s)		
C ¹			123.8	122.3
C ²			127.8	125.5
C ³			112.4	111.2
C ⁴			127.1	126.6
C ⁷			128.8	128.5
C ⁸			125.7	124.8
C ⁹			133.1	132.1
C ¹⁰			124.5	122.7
C ¹¹		101.0		
CH ₃				12.1

^a Room temperature, CDCl₃, chemical shifts in ppm from internal SiMe₄. ^b ¹³C NMR, aromatics: I, δ 111.9, 122.3, 123.0, 125.8, 126.7, 128.0, 128.5 and 133.2; II, δ 111.9, 125.1, 125.6, 125.9, 126.1, 128.2 and 132.3. ^c Assignments based on correlation (COSY) and heteronuclear correlation (HETCOR) experiments. ^d Data taken from ref. 5(d).

Method (b). An aqueous solution (20 cm³) of AuCl₃·2H₂O (1 mmol) was added to pure NC₅H₄(CH₂Ph)-2 (0.169 g, 1 mmol); the resulting yellow suspension was refluxed until the precipitate became white (*ca.* 8 h). The solid product was filtered off and air dried to give complex **1m** (0.350 g, 80%).

[Au{NC₅H₄(CHMePh)-2}Cl₃]**2a**. An aqueous solution (20 cm³) of AuCl₃·2H₂O (1.55 mmol) was added to pure NC₅H₄(CHMePh)-2 (0.283 g, 1.55 mmol); the resulting yellow suspension was stirred for 3 d at room temperature. The solid product was filtered off and air dried to give complex **2a** (0.564

g, 75%). Recrystallization from acetone–diethyl ether yielded an orange crystalline product: m.p. 169–170 °C [Found: C, 32.3; H, 2.7; N, 2.8%; M^+ 485 (EI). $C_{13}H_{13}AuCl_3N$ requires C, 32.1; H, 2.7; N, 2.9%; M 485]; $\tilde{\nu}_{max}/cm^{-1}$ 1605vs, 1565m, 1482vs, 1432vs, 1161vs, 1124m, 1053m, 792m, 760vs, 700vs, 585s, 545vs and (Au–Cl) 359vs; m/z 485 (M^+) and 449 ($M - HCl$).

[$Au\{NC_5H_4(CHMeC_6H_4)_2\}Cl_2$] **2m**. A suspension of complex **2a** (0.200 g, 0.41 mmol) in MeCN (3 cm³) and water (15 cm³) was refluxed until the precipitate became white (*ca.* 7 h). The solid product was filtered off and air dried to give the analytical sample (0.165 g, 90%), m.p. 277–278 °C [Found: C, 34.9; H, 2.7; N, 3.0%; M^+ 449 (EI). $C_{13}H_{12}AuCl_2N$ requires C, 34.7; H, 2.7; N, 3.1%; M 449]; $\tilde{\nu}_{max}/cm^{-1}$ 1603vs, 1557m, 1477s, 1449vs, 1164w, 1025s, 797s, 764vs, 756vs and (Au–Cl) 352vs and 294vs.

[$HNC_5H_4(CMe_2Ph)_2$][$AuCl_4$] **3s**. *Method (a)*. An aqueous solution (40 cm³) of $AuCl_3 \cdot 2H_2O$ (2 mmol) was added to pure $NC_5H_4(CMe_2Ph)_2$ (0.394 g, 2 mmol): the resulting yellow suspension was stirred for 3 d at room temperature. The solid product was filtered off and air dried to give complex **3s** (0.628 g, 58%). Recrystallization from acetone–diethyl ether yielded an orange crystalline product: m.p. 134–135 °C [Found: C, 32.0; H, 3.0; N, 2.6%; M^+ 198 (FAB). $C_{14}H_{16}AuCl_4N$ requires C, 31.3; H, 3.0; N, 2.6%; $M(H_2L^+)$ 198]; $\Lambda_M(5 \times 10^{-4} mol dm^{-3}, acetone)$ 160 $\Omega^{-1} cm^2 mol^{-1}$; $\tilde{\nu}_{max}/cm^{-1}$ 3242vs, 3176vs, 3131vs, 1609vs, 1536s, 1494m, 1240m, 771vs, 759vs, 741s, 701vs, 566m, 541m and (Au–Cl) 357vs.

Method (b). To a chloroform solution (10 cm³) of anhydrous $AuCl_3$ (0.080 g, 0.26 mmol) was added a chloroform solution (10 cm³) of $NC_5H_4(CMe_2Ph)_2$: the resulting orange-yellow solution was stirred for 1 h at room temperature and then concentrated to small volume. Addition of diethyl ether gave complex **3s** (0.102 g, 73%).

Method (c). To a solution of $NC_5H_4(CMe_2Ph)_2$ (0.078 g, 0.39 mmol) in ethanol (5 cm³) was added a solution of $Na[AuCl_4] \cdot 2H_2O$ (0.156 g, 0.39 mmol) in the same solvent (15 cm³): the resulting solution was stirred for 12 h at room temperature and then evaporated to dryness. The residue was taken up with dichloromethane, filtered and concentrated to small volume: addition of diethyl ether gave complex **3s** (0.130 g, 62%).

[$Au\{NC_5H_4(CMe_2C_6H_4)_2\}Cl_2$] **3m**. A suspension of complex **3s** (0.462 g, 0.86 mmol) in MeCN (5 cm³) and water (25 cm³) was refluxed until the precipitate became white (*ca.* 8 h). The solid product was filtered off and air dried to give complex **3m** (0.334 g, 84%). Recrystallization from dichloromethane–diethyl ether yielded the analytical sample: m.p. 283–284 °C [Found: C, 35.7; H, 3.0; N, 3.1%; M^+ 463 (FAB). $C_{14}H_{14}AuCl_2N$ requires C, 36.2; H, 3.0; N, 3.0%; M 463]; $\tilde{\nu}_{max}/cm^{-1}$ 1601s, 1579m, 1557m, 1479s, 1390m, 1368s, 1025vs, 771vs, 757s, 745vs, 720s, 569s and (Au–Cl) 353vs and 294vs.

[$HNC_5H_4(CMe_3)_2$][$AuCl_4$] **4s**. An aqueous solution (20 cm³) of $AuCl_3 \cdot 2H_2O$ (1.87 mmol) was added to pure $NC_5H_4(CMe_3)_2$ (0.253 g, 1.87 mmol): the resulting yellow suspension was stirred for 3 d at room temperature. The solid product was filtered off and air dried to give complex **4s** (0.563 g, 63%). Recrystallization from dichloromethane–diethyl ether yielded a bright yellow crystalline product: m.p. 201–202 °C [Found: C, 22.6; H, 2.9; N, 2.9%; M^+ 136 (FAB). $C_9H_{14}AuCl_4N$ requires C, 22.7; H, 2.9; N, 2.9%; $M(H_2L^+)$ 136]; $\Lambda_M(5 \times 10^{-4} mol dm^{-3}, acetone)$ 168 $\Omega^{-1} cm^2 mol^{-1}$; $\tilde{\nu}_{max}/cm^{-1}$ 3268vs, 3195vs, 1624s, 1610vs, 1532s, 1276s, 1243m, 1097m, 766vs, 739s, 541vs and (Au–Cl) 358vs.

The same product was obtained from $Na[AuCl_4]$ and $NC_5H_4(CMe_3)_2$ in aqueous or in ethanolic solution, at room temperature or at reflux.

[$HNC_5H_4(CH_2CMe_3)_2$][$AuCl_4$] **5s**. An aqueous solution (20 cm³) of $AuCl_3 \cdot 2H_2O$ (1.68 mmol) was added to pure $NC_5H_4(CH_2CMe_3)_2$ (0.251 g, 1.68 mmol): the resulting yellow suspension was stirred for 3 d at room temperature. The solid was filtered off and air dried to give a first crop of an orange-

yellow product; a second crop was obtained by evaporation of the mother-liquor. The two crops were combined and crystallized from acetone–diethyl ether to give the analytical sample (0.370 g, 45%): m.p. 124–125 °C [Found: C, 24.0; H, 3.1; N, 2.9%; M^+ 150 (FAB). $C_{10}H_{16}AuCl_4N$ requires C, 24.5; H, 3.3; N, 2.9%; $M(H_2L^+)$ 150]; $\Lambda_M(5 \times 10^{-4} mol dm^{-3}, acetone)$ 160 $\Omega^{-1} cm^2 mol^{-1}$; $\tilde{\nu}_{max}/cm^{-1}$ 3245s, 3222s, 3173s, 3123s, 1625m, 1607s, 1529m, 1227w, 1169w, 782m, 755s and (Au–Cl) 359vs.

Reactions of the Cyclometallated Derivatives 1m–3m.—With PPh₃. General procedure. To a suspension of [$AuLCl_2$] (1 mmol) in acetone (20 cm³) was added solid PPh_3 (1 or 2 mmol) and $NaBF_4$ (excess). After a few minutes a solution was obtained; this was stirred for 1–4 h at room temperature and then evaporated to dryness. The residue was taken up with dichloromethane, filtered and concentrated to small volume: addition of diethyl ether gave a white solid of [$AuL(PPh_3)_2Cl$][BF_4] **1p–3p**, yield 80–90%. Recrystallization from dichloromethane–diethyl ether gave the analytical sample.

[$Au\{NC_5H_4(CH_2C_6H_4)_2\}(PPh_3)Cl$][BF_4] **1p**: m.p. 142 °C (decomp.) [Found: C, 48.4; H, 3.7; N, 1.7%; M^+ 662 (FAB). $C_{30}H_{25}AuBClF_4NP$ requires C, 48.1; H, 3.3; N, 1.9%; $M[AuL(PPh_3)Cl^+]$ 662]; $\Lambda_M(5 \times 10^{-4} mol dm^{-3}, acetone)$ 160 $\Omega^{-1} cm^2 mol^{-1}$; $\tilde{\nu}_{max}/cm^{-1}$ 1608m, 1558w, 1480s, 1059vs (br), 534vs, 515vs, 501vs and (Au–Cl) 305s; $\delta_p(CD_2Cl_2)$ 33.9.

[$Au\{NC_5H_4(CHMeC_6H_4)_2\}(PPh_3)Cl$][BF_4] **2p**: m.p. 205–207 °C [Found: C, 48.9; H, 3.3; N, 1.9%; M^+ 676 (FAB). $C_{31}H_{28}AuBClF_4NP$ requires C, 48.7; H, 3.7; N, 1.8%; $M[AuL(PPh_3)Cl^+]$ 676]; $\Lambda_M(5 \times 10^{-4} mol dm^{-3}, acetone)$ 164 $\Omega^{-1} cm^2 mol^{-1}$; $\tilde{\nu}_{max}/cm^{-1}$ 1610m, 1585w, 1565w, 1490s, 1055vs (br), 782w, 770m, 746vs, 721vs, 712s, 698s, 688s, 555s, 536vs, 513vs, 499vs and (Au–Cl) 313vs; $\delta_p(CDCl_3)$ 33.6 (s, P_a) and 32.8 (s, P_b) (a : b = 2 : 1).

[$Au\{NC_5H_4(CMe_2C_6H_4)_2\}(PPh_3)Cl$][BF_4] **3p**: m.p. 184–185 °C [Found: C, 49.3; H, 3.7; N, 1.7%; M^+ 690 (FAB). $C_{32}H_{29}AuBClF_4NP$ requires C, 49.4; H, 3.7; N, 1.8%; $M[AuL(PPh_3)Cl^+]$ 690]; $\Lambda_M(5 \times 10^{-4} mol dm^{-3}, acetone)$ 160 $\Omega^{-1} cm^2 mol^{-1}$; $\tilde{\nu}_{max}/cm^{-1}$ 1601s, 1570w, 1554w, 1480vs, 1051vs (br), 782s, 748vs, 731s, 719s, 691vs, 571m, 533vs, 510vs and (Au–Cl) 316s; $\delta_p(CDCl_3)$ 33.2.

With Ph₂PCH₂CH₂PPh₂ (dppe). General procedure. To a suspension of [$AuLCl_2$] (0.5 mmol) in acetone (20 cm³) was added solid dppe (0.5 mmol) and $NaBF_4$ (excess). After a few minutes a solution was obtained; this was stirred for 6 h at room temperature and then evaporated to dryness. The residue was taken up with dichloromethane, filtered and concentrated to small volume: addition of diethyl ether gave a white solid of [$AuL(dppe)Cl$][BF_4] **1d** and **2d**.

[$Au\{NC_5H_4(CH_2C_6H_4)_2\}(dppe)Cl$][BF_4] **1d**: yield 87%, m.p. 82–84 °C [Found: C, 51.4; H, 4.0; N, 1.7%; M^+ 798 (FAB). $C_{38}H_{34}AuBClF_4NP_2$ requires C, 51.5; H, 3.8; N, 1.6%; $M[AuL(dppe)Cl^+]$ 798]; $\Lambda_M(5 \times 10^{-4} mol dm^{-3}, acetone)$ 164 $\Omega^{-1} cm^2 mol^{-1}$; $\tilde{\nu}_{max}/cm^{-1}$ 1587s, 1564w, 1103vs, 1057vs (br), 996vs, 734vs, 691vs, 529vs and (Au–Cl) 327w; $\delta_p(CDCl_3)$ 37.1 [1 P, d, $J(PP)$ 54, P(A)] and 26.3 [1 P, d, $J(PP)$ 54 Hz, P(X)].

[$Au\{NC_5H_4(CHMeC_6H_4)_2\}(dppe)Cl$][BF_4] **2d**: yield 89%, m.p. 100–102 °C [Found: C, 52.2; H, 3.9; N, 1.6%; M^+ 812 (FAB). $C_{39}H_{36}AuBClF_4NP_2$ requires C, 52.6; H, 4.0; N, 1.6%; $M[AuL(dppe)Cl^+]$ 812]; $\Lambda_M(5 \times 10^{-4} mol dm^{-3}, acetone)$ 160 $\Omega^{-1} cm^2 mol^{-1}$; $\tilde{\nu}_{max}/cm^{-1}$ 1585m, 1566w, 1104vs, 1058vs (br), 996s, 730vs, 692s, 530vs (br) and (Au–Cl) 328m; $\delta_p(CD_2Cl_2)$ 35.4 [1 P, d, $J(PP)$ 57, P(A)] and 25.4 [1 P, d, $J(PP)$ 57 Hz, P(X)].

Reactions of Complex 1d.—With $HBF_4 \cdot Et_2O$. To a solution of complex **1d** (0.075 g, 0.085 mmol) in dichloromethane (15 cm³) was added $HBF_4 \cdot Et_2O$ ($5 \times 10^{-2} cm^3$, *ca.* 0.36 mmol) with stirring: the solution became slightly opalescent. It was stirred for 1 h at room temperature and then filtered and concentrated to small volume: addition of diethyl ether gave the white solid

[Au{HNC₅H₄(CH₂C₆H₄)-2}(dppe)Cl][BF₄]₂ **1d'** (0.070 g, 85%); m.p. 126–128 °C {Found: C, 46.0; H, 3.6; N, 1.4%; M^+ 797 (FAB). C₃₈H₃₅AuB₂ClF₈NP₂ requires C, 46.9; H, 3.6; N, 1.4%; M [Au(HL)(dppe)Cl²⁺] 799}; $\Lambda_M(5 \times 10^{-4} \text{ mol dm}^{-3}$, acetone) 210 $\Omega^{-1} \text{ cm}^2 \text{ mol}^{-1}$; $\tilde{\nu}_{\text{max}}/\text{cm}^{-1}$ 3500vs (br), 1620s, 1585m, 1545m, 1060vs (br), 740vs, 700vs and (Au–Cl) 325w; $\delta_p(\text{CD}_2\text{Cl}_2)$ 35.0 [1 P, d, $J(\text{PP})$ 55, P(A)] and 24.3 [1 P, d, $J(\text{PP})$ 55 Hz, P(X)].

With AgBF₄. To a solution of complex **1d** (0.133 g, 0.15 mmol) in acetone (15 cm³) was added a solution of AgBF₄ (0.030 g, 0.15 mmol) in acetone (10 cm³), with stirring. Silver chloride was filtered off and the solution evaporated to dryness; the residue was taken up with dichloromethane, filtered and concentrated to small volume. Addition of diethyl ether gave a pale yellow product [Au{NC₅H₄(CH₂C₆H₄)-2}(dppe)][BF₄]₂ **1d'** (0.120 g, 85%); m.p. 175 °C (decomp.) {Found: C, 48.4; H, 3.7; N, 1.5%; M^+ 763 (FAB). C₃₈H₃₄AuB₂F₈NP₂ requires C, 48.7; H, 3.6; N, 1.5%; M [AuL(dppe)²⁺] 763}; $\Lambda_M(5 \times 10^{-4} \text{ mol dm}^{-3}$, acetone) 210 $\Omega^{-1} \text{ cm}^2 \text{ mol}^{-1}$; $\tilde{\nu}_{\text{max}}/\text{cm}^{-1}$ 1609s, 1586m, 1568m, 1059vs (br), 996vs, 845m, 817m, 732vs and 692vs; $\delta_p(\text{CD}_2\text{Cl}_2)$: 36.6 [1 P, d, $J(\text{PP})$ 24, P(A)] and 26.7 [1 P, d, $J(\text{PP})$ 24 Hz, P(X)]; m/z 763 (M^+) and 850 ($[M + \text{BF}_4]^+$) (EI).

*Reaction of [Au{NC₅H₄(CHMeC₆H₄)-2}(dppe)Cl][BF₄]₂ **2d** with AgBF₄.*—To a solution of complex **2d** (0.092 g, 0.102 mmol) in acetone (15 cm³) was added a solution of AgBF₄ (0.020 g, 0.102 mmol) in acetone (10 cm³), with stirring. Silver chloride was filtered off and the solution evaporated to dryness; the residue was taken up with dichloromethane, filtered and concentrated to small volume. Addition of diethyl ether gave a white product [Au{NC₅H₄(CHMeC₆H₄)-2}(dppe)][BF₄]₂ **2d'** (0.076 g, 78%); m.p. 190 °C (decomp.) {Found: C, 48.8; H, 3.8; N, 1.5%; M^+ 777 (FAB). C₃₉H₃₆AuB₂F₈NP₂ requires C, 49.2; H, 3.8; N, 1.5%; M [AuL(dppe)²⁺] 777}; $\Lambda_M(5 \times 10^{-4} \text{ mol dm}^{-3}$, acetone) 210 $\Omega^{-1} \text{ cm}^2 \text{ mol}^{-1}$; $\tilde{\nu}_{\text{max}}/\text{cm}^{-1}$ 1606s, 1585m, 1568m, 1052vs (br), 823w, 729vs and 692vs; $\delta_p(\text{CD}_2\text{Cl}_2)$ 32.9 [1 P, d, $J(\text{PP})$ 12, P(A)] and 23.7 [1 P, d, $J(\text{PP})$ 12 Hz, P(X)]; m/z 777 (M^+) and 865 ($[M + \text{BF}_4]^+$) (FAB).

Reactions with Carbon Monoxide.—General procedure. A Pyrex bottle containing a suspension of [AuLCl₂] **1m–3m**, equipped with a magnetic stirrer bar, was placed in a stainless-steel autoclave. After removal of the air, the vessel was pressurized with CO (50 atm), heated at 60 °C and stirred for several hours. At the end of the reaction the CO was released and the mixture was worked up.

[Au{NC₅H₄(CH₂C₆H₄)-2}Cl₂] **1m** in CH₂Cl₂–EtOH. A suspension of complex **1m** (0.150 g, 0.345 mmol) in CH₂Cl₂–EtOH (4:1, 25 cm³) was subjected to the general procedure described above for 24 h, after which the mixture consisted of a bright yellow solution and metallic gold. The solution was filtered and evaporated to dryness (0.067 g). The organic product was a mixture (TLC) from which three pure compounds were isolated by chromatography on a column (58 × 2 cm) of silica gel (Merck, 70–230 mesh ASTM), using dichloromethane (**I** and **II**) and acetone (**III**) as eluents. First band eluted (R_f 0.28 on silica gel plates): **I** (0.010 g, 0.042 mmol, 12%) [Found: M^+ 229 (FAB). C₁₃H₈ClNO requires M 229]; $\tilde{\nu}_{\text{max}}/\text{cm}^{-1}$ (CO) 1675vs and 1630s, 1605s, 1563s, 1544w and 1522s; m/z 229 (100, M^+), 201 ($M - \text{CO}$), 166 ($M - \text{CO} - \text{Cl}$) and 139 ($M - \text{CO} - \text{Cl} - \text{HCN}$) (EI). Second band eluted (R_f 0.17): **II** (0.003 g, 0.015 mmol, 4.4%) [Found: M^+ 195 (FAB). C₁₃H₉NO requires M 195]; $\tilde{\nu}_{\text{max}}/\text{cm}^{-1}$ (CO) 1665vs and 1628vs, 1610s, 1589w, 1569m and 1524m; m/z 195 (100, M^+) and 167 ($M - \text{CO}$) (EI). Third band eluted (R_f 0.09): **III** (0.054 g, 0.14 mmol, 81%), m.p. 130 °C (decomp.) [Found: M^+ 388 (FAB). C₂₆H₁₄N₂O₂ requires M 388]; $\tilde{\nu}_{\text{max}}/\text{cm}^{-1}$ (CO) 1652vs and 1628vs, 1603s, 1565m and 1520s; m/z 388 (100, M^+), 359 ($M - \text{HCO}$) and 195 ($\frac{1}{2}M + \text{H}$) (EI).

[Au{NC₅H₄(CHMeC₆H₄)-2}Cl₂] **2m** in CH₂Cl₂–EtOH. A

suspension of complex **2m** (0.112 g, 0.25 mmol) in CH₂Cl₂–EtOH (4:1, 25 cm³) was subjected to the general procedure for 24 h, after which the mixture consisted of an intense yellow solution and metallic gold. The solution was filtered and evaporated to dryness, the residue was taken up with diethyl ether, filtered and evaporated to dryness to yield the analytical sample **IV** (0.0454 g, 87%), m.p. 198–200 °C [Found: M^+ 209 (EI). Calc. for C₁₄H₁₁NO: M 209]; $\tilde{\nu}_{\text{max}}/\text{cm}^{-1}$ (CO) 1652vs and 1632vs, 1606s, 1568s, 1548w and 1524s (CH₂Cl₂).

[Au{NC₅H₄(CMe₂C₆H₄)-2}Cl₂] **3m** in CH₂Cl₂–EtOH. A suspension of complex **3m** (0.180 g, 0.388 mmol) in CH₂Cl₂–EtOH (4:1, 25 cm³) was subjected to the general procedure for 20 h, after which the mixture consisted of a pale yellow solution and metallic gold. The solution was filtered and evaporated to dryness, the residue was taken up with ethanol and filtered to separate a small amount of the starting material; the solution was evaporated to dryness, taken up with dichloromethane and neutralized with K₂CO₃. After filtration, the solution was evaporated to dryness to yield the analytical sample **V** (0.078 g, 74%) [Found: $M + \text{H}^+$ 270 (FAB). C₁₇H₁₉NO₂ requires M 269]; $\tilde{\nu}_{\text{max}}/\text{cm}^{-1}$ (CO) 1719s and 1658w, 1585m and 1563m (CH₂Cl₂); $\delta_{\text{H}}(\text{CDCl}_3)$ 8.52 [1 H, ddd, $J(\text{H}^3, \text{H}^6)$ 1.0, $J(\text{H}^4, \text{H}^6)$ 2.0 and $J(\text{H}^5, \text{H}^6)$ 4.8, H⁶], 7.67 [1 H, dd, $J(\text{H}^4, \text{H}^6)$ 1.3 and $J(\text{H}^5, \text{H}^6)$ 8.0, H⁶], 7.55 [1 H, ddd, $J(\text{H}^3, \text{H}^4)$ 8.1, $J(\text{H}^4, \text{H}^5)$ 7.5 and $J(\text{H}^4, \text{H}^6)$ 2.0, H⁴], 7.47 [1 H, ddd, $J(\text{H}^3, \text{H}^5)$ 1.7, $J(\text{H}^4, \text{H}^5)$ 7.6 and $J(\text{H}^5, \text{H}^6)$ 8.0, H⁵], 7.42 [1 H, dd, $J(\text{H}^3, \text{H}^4)$ and $J(\text{H}^3, \text{H}^5)$ 1.7, H³], 7.27 [1 H, td, $J(\text{H}^3, \text{H}^4)$ 7.6, $J(\text{H}^4, \text{H}^5)$ 7.6 and $J(\text{H}^4, \text{H}^6)$ 1.3, H⁴], 7.13 [1 H, dt, $J(\text{H}^3, \text{H}^4)$ 8.1, $J(\text{H}^3, \text{H}^5)$ 1.0 and $J(\text{H}^3, \text{H}^6)$ 1.0, H³], 7.07 [1 H, ddd, $J(\text{H}^4, \text{H}^5)$ 7.5, $J(\text{H}^3, \text{H}^5)$ 1.0 and $J(\text{H}^5, \text{H}^6)$ 4.8, H⁵], 3.72 [2 H, q, $J(\text{HH})$ 7.2, OCH₂], 1.80 (6 H, s, 2 CH₃) and 1.07 [3 H, t, $J(\text{HH})$ 7.2 Hz, OCH₂CH₃] (COSY); $\delta_{\text{C}}(\text{CDCl}_3)$ 169.8 (1 C, C² or CO), 168.4 (1 C, C² or CO), 148.4 (1 C, C⁶), 147.1 (1 C, C¹), 135.8 (1 C, C⁴), 133.1 (1 C, C²), 130.4 (1 C, C⁵), 129.5 (1 C, C³), 127.2 (1 C, C⁶), 126.0 (1 C, C⁴), 121.2 (1 C, C³), 120.5 (1 C, C⁵), 60.7 (1 C, OCH₂), 46.6 [1 C, C(CH₃)₂], 30.7 (2 C, 2 CH₃) and 13.8 (1 C, CH₂CH₃) (HETCOR); m/z 268 ($M - \text{H}$), 224 ($M - \text{OCH}_2\text{CH}_3$) and 196 (100, $M - \text{CO}_2\text{CH}_2\text{CH}_3$) (EI).

X-Ray Data Collection and Structure Determination.—Crystal data and other experimental details are summarized in Table 1. The diffraction experiment was carried out on an Enraf-Nonius CAD-4 diffractometer at room temperature using Mo-K α radiation ($\lambda = 0.71073 \text{ \AA}$) with a graphite-crystal monochromator in the incident beam. The calculations were performed on a PDP 11/73 computer using the SDP package¹⁶ and the physical constants tabulated therein. Periodic monitoring of three standard reflections revealed a crystal decay of about 4.9% (on intensities) at the end of data collection. The diffracted intensities were corrected for Lorentz, polarization, decay and absorption effects (empirically).¹⁷ Scattering factors and anomalous dispersion corrections were taken from ref. 18. The structure was solved by Patterson and Fourier methods and refined by full-matrix least squares, minimizing the function $\sum w(F_o - k|F_c|)^2$. Anisotropic thermal parameters were refined for all the non-hydrogen atoms. All the hydrogen atoms were detected on the Fourier maps; those bonded to atom C(15) were included in the least-squares refinement, whereas the remainder were placed in their ideal positions (C–H 0.97 Å, B 1.20 times that of the carbon atom to which they are attached). In the final least-square cycles an extinction parameter, g , was included [according to the formula $|F_c|(\text{corrected}) = |F_c|/(1 + gI_c)$], which refined to the value $1.14(4) \times 10^{-6}$. The final Fourier map showed a maximum residual of $1.4(2) \text{ e \AA}^{-3}$ at 0.90 \AA from the gold atom. The final atomic coordinates are listed in Table 3.

Additional material available from the Cambridge Crystallographic Data Centre comprises H-atom coordinates, thermal parameters and remaining bond lengths and angles.

Acknowledgements

Financial support from Ministero dell'Università e della Ricerca Scientifica e Tecnologica (MURST) (40%) and Consiglio Nazionale delle Ricerche (CNR) is gratefully acknowledged.

References

- I. Omae, *Organometallic Intramolecular-coordination Compounds*, Elsevier, Amsterdam, 1986; E. C. Constable, *Polyhedron*, 1984, **3**, 1037; J. Dehan and M. Pfeffer, *Coord. Chem. Rev.*, 1976, **18**, 327; M. I. Bruce, *Angew. Chem., Int. Ed. Engl.*, 1977, **16**, 73; A. D. Ryabov, *Chem. Rev.*, 1990, **90**, 403; G. R. Newkome, W. E. Puckett, V. K. Gupta and G. E. Kiefer, *Chem. Rev.*, 1986, **86**, 451.
- (a) J. Vicente and M. T. Chicote, *Inorg. Chim. Acta*, 1981, **54**, L259; (b) J. Vicente, M. T. Chicote and M. D. Bermudez, *Inorg. Chim. Acta*, 1982, **63**, 35; (c) J. Vicente, M. T. Chicote and M. D. Bermudez, *J. Organomet. Chem.*, 1984, **268**, 191; (d) J. Vicente, M. T. Chicote, M. D. Bermudez, X. Solans and M. Font-Altaba, *J. Chem. Soc., Dalton Trans.*, 1984, 557; (e) J. Vicente, M. T. Chicote, M. D. Bermudez and M. Garcia-Garcia, *J. Organomet. Chem.*, 1985, **295**, 125; (f) J. Vicente, M. T. Chicote, M. D. Bermudez, P. G. Jones and G. M. Sheldrick, *J. Chem. Res.* 1985, (S) 72; (g) J. Vicente, M. T. Chicote, M. D. Bermudez and M. J. Sanchez-Santano, *J. Organomet. Chem.*, 1986, **310**, 401; (h) J. Vicente, M. T. Chicote, M. D. Bermudez, P. G. Jones, C. Fittschen and G. M. Sheldrick, *J. Chem. Soc., Dalton Trans.*, 1986, 2361; (i) J. Vicente, M. T. Chicote, M. D. Bermudez, M. J. Sanchez-Santano and P. G. Jones, *J. Organomet. Chem.*, 1988, **354**, 381; (j) J. Vicente, M. D. Bermudez, M. T. Chicote and M. J. Sanchez-Santano, *J. Chem. Soc., Dalton Trans.*, 1990, 1945; (k) J. Vicente, M. D. Bermudez, J. Escribano, M. P. Carrillo and P. G. Jones, *J. Chem. Soc., Dalton Trans.*, 1990, 3083; (l) J. Vicente, M. D. Bermudez, M. T. Chicote and M. J. Sanchez-Santano, *J. Organomet. Chem.*, 1990, **381**, 285; (m) J. Vicente, M. D. Bermudez, M. P. Carrillo and P. G. Jones, *J. Chem. Soc., Dalton Trans.*, 1992, 1975; (n) J. Vicente, M. D. Bermudez and J. Escribano, *Organometallics*, 1991, **10**, 3380; (o) J. Vicente, M. D. Bermudez, M. P. Carrillo and P. G. Jones, *J. Organomet. Chem.*, 1993, **456**, 305; (p) J. Vicente, M. D. Bermudez and F. J. Carrion, *Inorg. Chim. Acta*, 1994, **220**, 1.
- E. C. Constable, R. P. G. Henney and T. A. Leese, *J. Organomet. Chem.*, 1989, **361**, 277; E. C. Constable and T. A. Leese, *J. Organomet. Chem.*, 1989, **363**, 419; E. C. Constable, R. P. G. Henney, T. A. Leese and D. A. Tocher, *J. Chem. Soc., Dalton Trans.*, 1990, 443.
- C.-W. Chan, W.-T. Wong and C.-M. Che, *Inorg. Chem.*, 1994, **33**, 1266.
- (a) G. Minghetti, M. A. Cinellu, S. Gladiali, G. Chelucci, F. Demartin and M. Manassero, *J. Organomet. Chem.*, 1986, **307**, 107; (b) M. A. Cinellu, S. Gladiali and G. Minghetti, *J. Organomet. Chem.*, 1989, **363**, 401; (c) G. Minghetti, M. A. Cinellu, S. Stoccoro, G. Chelucci and A. Zucca, *Inorg. Chem.*, 1990, **29**, 5137; (d) G. Minghetti, M. A. Cinellu, S. Stoccoro and A. Zucca, *Gazz. Chim. Ital.*, 1992, **122**, 455; (e) S. Stoccoro, G. Chelucci, M. A. Cinellu, A. Zucca and G. Minghetti, *J. Organomet. Chem.*, 1993, **450**, C15; (f) G. Minghetti, A. Zucca, S. Stoccoro, M. A. Cinellu, M. Manassero and M. Sansoni, *J. Organomet. Chem.*, 1994, **481**, 195; (g) G. Minghetti, M. A. Cinellu, S. Stoccoro, A. Zucca and M. Manassero, *J. Chem. Soc., Dalton Trans.*, 1995, 777.
- M. A. Cinellu, G. Minghetti, S. Stoccoro, A. Zucca and G. Chelucci, I Congresso Gruppo Interdivisionale Chimica Organometallica, CO.GI.CO. '92, Camerino, 1992, Abstracts 22.
- M. A. Cinellu, G. Minghetti, S. Stoccoro, A. Zucca and M. Manassero, XVIth International Conference on Organometallic Chemistry, University of Sussex, 1994, Abstracts P. 105.
- U. Azzena, G. Chelucci, G. Delogu, S. Gladiali, M. Marchetti, F. Soccolini and C. Botteghi, *Gazz. Chim. Ital.*, 1986, **116**, 307; C. Botteghi, G. Chelucci, G. Chessa, G. Delogu, S. Gladiali and F. Soccolini, *J. Organomet. Chem.*, 1986, **304**, 217.
- G. Minghetti, M. L. Ganadu, C. Foddai, M. A. Cinellu, F. Cariati, F. Demartin and M. Manassero, *Inorg. Chim. Acta*, 1984, **86**, 93; J. Shamir, A. Givan, L. Canovese, L. Cattalini, P. Uguagliati and M. L. Tobe, *J. Raman Spectrosc.* 1993, **24**, 233.
- P. K. Byers and A. J. Canty, *Organometallics*, 1990, **9**, 210.
- C. K. Johnson, ORTEP, Report ORNL-5138, Oak Ridge National Laboratory, Oak Ridge, TN, 1976.
- P. K. Byers, A. J. Canty, L. M. Engelhart, J. M. Patrick and A. H. White, *J. Chem. Soc., Dalton Trans.*, 1985, 981; P. K. Byers, A. J. Canty, N. J. Minchin, J. M. Patrick, B. W. Skelton and A. H. White, *J. Chem. Soc., Dalton Trans.*, 1985, 1183; A. J. Canty, N. J. Minchin, P. C. Healy and A. H. White, *J. Chem. Soc., Dalton Trans.*, 1982, 1795; G. A. Annibale, L. Canovese, L. Cattalini, G. Natile, M. Biagini-Cingi, A. M. Manotti-Lanfredi and A. Tiripicchio, *J. Chem. Soc., Dalton Trans.*, 1981, 2280.
- P. S. Pregosin and F. Wombacher, *Magn. Reson. Chem.*, 1991, **29**, S106 and refs. therein.
- F. Calderazzo, *Angew. Chem., Int. Ed. Engl.*, 1977, **16**, 299; *J. Organomet. Chem.*, 1990, **400**, 303 and refs. therein.
- E. C. Constable and L. R. Sousa, *J. Organomet. Chem.*, 1992, **427**, 125.
- B. A. Frenz and Associates, SDP PLUS, Version 1.0, Enraf-Nonius, Delft, 1980.
- A. C. T. North, D. C. Phillips and F. S. Mathews, *Acta Crystallogr., Sect. A*, 1968, **24**, 351.
- International Tables for X-Ray Crystallography*, Kynoch Press, Birmingham, 1974, vol. 4.

Received 14th March 1995; Paper 5/01597I