

Preparation and crystal structure of $[(\text{AlBr}_3)_2(\text{Se}_2\text{N}_2)]$, the first example of a main-group element adduct of diselenium dinitride

Paul F. Kelly* and Alexandra M. Z. Slawin

Department of Chemistry, Loughborough University, Loughborough LE11 3TU, UK

The room temperature reaction of AlBr_3 with Se_4N_4 in CH_2Br_2 generated $[(\text{AlBr}_3)_2(\text{Se}_2\text{N}_2)]$ as an air-sensitive, yellow solid; X-ray crystallography confirmed the presence of the neutral Se_2N_2 ligand, bridging *via* two Al–N bonds.

The recent report of the preparation of the first examples of adducts of diselenium dinitride, Se_2N_2 ,¹ added weight to the fast developing conclusion that the chemistry of selenium–nitrogen systems may well, ultimately, prove to be as diverse as that of their sulfur counterparts.² The adducts in question, salts of $[\text{Pd}_2\text{Cl}_6(\text{Se}_2\text{N}_2)]^{2-}$, form in the high-temperature reaction of Se_4N_4 with simple halogen-bridged palladium salts, and appear to be stable both to air and friction (in marked contrast to Se_4N_4 which is a particularly pernicious explosive). Here we report on the preparation of the first example of a main-group metal adduct of diselenium dinitride.

It has long been known that Se_4N_4 reacts with AlBr_3 to generate the S_2N_2 adduct $[(\text{AlBr}_3)_2(\text{S}_2\text{N}_2)]$,³ we have now found that the analogous reaction of Se_4N_4 may be performed. Thus treatment of a suspension of Se_4N_4 (**CAUTION:** severe explosive!) in CH_2Br_2 with solid AlBr_3 (molar ratio 1:5), followed by thorough stirring, resulted in the immediate formation of a dark red, clear solution which was allowed to stand overnight. During this time a yellow crystalline solid was deposited; this was separated by decanting the mother liquor, washed with CH_2Br_2 (20 cm^3) and dried *in vacuo*.

The crystal structure of this product confirms it to be $[(\text{AlBr}_3)_2(\text{Se}_2\text{N}_2)]$ (see Fig. 1), revealing the presence of a neutral Se_2N_2 unit bridging the two AlBr_3 moieties *via* two Al–N bonds;† *i.e.* isostructural with $[(\text{AlBr}_3)_2(\text{S}_2\text{N}_2)]$.⁵ The resulting $\text{Al}_2\text{Se}_2\text{N}_2$ unit is effectively planar [the largest deviation from planarity being at N(1) 0.06 Å]; this plane also contains one of the bromine atoms on each aluminium. As has been noted many times for adducts of S_2N_2 ,⁶ there is a significant interaction between one of the bromines and the chalcogen–nitrogen ring; in this case it can be quantified by

† Crystal data. $\text{Al}_2\text{Br}_6\text{N}_2\text{Se}_2$, $M = 719.32$, monoclinic, space group $P2_1/n$ (no. 14), $a = 7.613(1)$, $b = 9.940(2)$, $c = 9.552(2)$ Å, $\beta = 110.41(2)^\circ$, $U = 677.5(2)$ Å³, $Z = 2$, $D_c = 3.53$ g cm⁻³, $\mu(\text{Cu-K}\alpha) = 290.63$ cm⁻¹, $F(000) = 636.0$, dimensions 0.1 × 0.1 × 0.03 mm, $T = 298$ K. All measurements were made on a Rigaku AFC7S diffractometer with graphite-monochromated Cu-K α ($\lambda = 1.54178$ Å) radiation and employing the ω - 2θ scan technique to a maximum 2θ value of 120.1°. An empirical absorption correction was applied which resulted in transmission factors ranging from 0.387 to 1.00. The structure was solved by direct methods⁴ and expanded using Fourier techniques with all atoms refined anisotropically on F (apart from Al and N which were refined isotropically) to give $R = 0.074$ for 669 observed reflections [$I > 3\sigma(I)$]. Maximum shift/error in final cycle 0.10. It should be noted that the air sensitivity of the crystals invariably led to a degree of decomposition upon mounting. Atomic coordinates, thermal parameters and bond lengths and angles have been deposited at the Cambridge Crystallographic Data Centre (CCDC). See Instructions for Authors, *J. Chem. Soc., Dalton Trans.*, 1996, Issue 1. Any request to the CCDC for this material should quote the full literature citation and the reference number 186/262.

the fact that the N(1)–Al(1)–Br(1) angle is 94.2°, which is some 15° smaller than the average of the corresponding angles to the other bromines. The average Se–N bond length in $[(\text{AlBr}_3)_2(\text{Se}_2\text{N}_2)]$ is effectively identical to that in $[\text{Pd}_2\text{Cl}_6(\text{Se}_2\text{N}_2)]^{2-}$.

The IR spectrum of the crystalline product shows an excellent correlation with that of $[(\text{AlBr}_3)_2(\text{S}_2\text{N}_2)]$, in particular the two strong bands which occur in the latter at 870 and 733 cm^{-1} are seen at 787 and 658 cm^{-1} in the selenium product (and retain the same intensity ratio). The yield of $[(\text{AlBr}_3)_2(\text{Se}_2\text{N}_2)]$ prepared in this manner, though not particularly high (typically 30 mg from 56 mg Se_4N_4 which equates to 14% based upon selenium) is sufficient to allow investigation into its chemical properties. Such investigations are currently ongoing and two points have emerged so far. The first of these is its severe air sensitivity; while aerial degradation of the S_2N_2 adducts of aluminium halides takes place over a number of seconds, samples of $[(\text{AlBr}_3)_2(\text{Se}_2\text{N}_2)]$ appear to decompose effectively instantaneously, turning a dark brown-black in colour and then lightening over a matter of minutes to orange (a colour consistent with the ultimate formation of Se_4N_4). It is worth noting that while this air sensitivity makes the product a lot harder to handle than the aforementioned palladium adducts (which are perfectly air stable), the fact that the reaction is performed at ambient temperature (rather than the 100 °C required in the latter case) greatly reduces the risk of Se_4N_4 detonation.

The other observation noted is that, unlike $[(\text{AlBr}_3)_2(\text{S}_2\text{N}_2)]$, $[(\text{AlBr}_3)_2(\text{Se}_2\text{N}_2)]$ does not undergo halogen exchange in CH_2Cl_2 . Thus we cannot, unfortunately, prepare a pure sample of $[(\text{AlCl}_3)_2(\text{Se}_2\text{N}_2)]$ in this manner. Attempts to prepare the latter by direct reaction of AlCl_3 with Se_4N_4 appear to be only partially successful. The IR spectrum of the off-white product of this reaction suggests that $[(\text{AlCl}_3)_2(\text{Se}_2\text{N}_2)]$ is indeed present but is significantly contaminated with an unidentified by-product.




Fig. 1 Crystal structure of $[(\text{AlBr}_3)_2(\text{Se}_2\text{N}_2)]$. Selected bond distances (Å) and angles (°): Al(1)–N(1) 1.92(2), N(1)–Se(1) 1.77(1), N(1)–Se(1*) 1.81(1); N(1)–Al(1)–Br(1) 94.2(5), N(1)–Al(1)–Br(2) 112.7(5), N(1)–Al(1)–Br(3) 108.1(5), Se(1)–N(1)–Se(1*) 97.6(7), N(1)–Se(1)–N(1*) 82.4(7), Se(1)–N(1)–Al(1) 122.4(8), Se(1*)–N(1)–Al(1) 139.9(8)

The isolation of $[(\text{AlBr}_3)_2(\text{Se}_2\text{N}_2)]$ in this manner represents a new class of reaction for Se_4N_4 . Previous reactions have involved either transition-metal species or selenium cations;² this is the first example of reaction with a main-group Lewis acid. As such it only serves to fuel the growing conclusion that a substantial body of chemistry of selenium–nitrogen systems awaits elucidation and suggests that, in view of the extensive range of S_2N_2 adducts that have been prepared, many more examples of adducts of Se_2N_2 should be possible.

References

- 1 P. F. Kelly and A. M. Z. Slawin, *Angew. Chem., Int. Ed. Engl.*, 1995, **34**, 1758.
- 2 T. M. Klapötke, in *The Chemistry of Inorganic Ring Systems; Studies in Inorganic Chemistry*, ed. R. Steudel, Elsevier, Amsterdam, 1992, **14**, 409; T. Chivers, *Main Group Chemistry News*, 1993, **1**, 6.
- 3 H. W. Roesky, J. Anhaus and W. S. Sheldrick, *Inorg. Chem.*, 1984, **23**, 75.
- 4 TEXSAN, Crystal Structure Analysis Package, Molecular Structure Corporation, Houston, TX, 1985 and 1992.
- 5 U. Thewalt and K. Holl, *Z. Naturforsch., Teil B*, 1984, **39**, 145.
- 6 See for example, U. Thewalt and M. Burger, *Angew. Chem., Int. Ed. Engl.*, 1982, **21**, 634.

Received 16th August 1996; Communication 6/05731D