

Dipalladium complexes with a bridging arylyene or diarylyene ligand. Synthesis and CNR' and CO insertion into the Pd–C bonds

Yong-Joo Kim,^{*,†,a} Sang-Won Song,^a Sang-Chul Lee,^a Soon-W. Lee,^b Kohtaro Osakada^c and Takakazu Yamamoto^c

^a Department of Chemistry, Kangnung National University, Kangnung 210-702, Korea

^b Department of Chemistry, Sung Kyun Kwan University, Natural Science Campus, Korea

^c Research Laboratory of Resources Utilization, Tokyo Institute of Technology, 4259 Nagatsuta, Midori-ku, Yokohama 226-8503, Japan


The complexes *trans*-[PdEt₂L₂] (L = PMe₃ or PEt₃) reacted with 4,4'-dibromobiphenyl, 4,4'-diiodobiphenyl, 1,4-diiodobenzene, di(4-bromophenyl) ether, di(4-chlorophenyl) sulfone, 9,10-dibromoanthracene and 4,4'-dibromooctafluorobiphenyl to give various bimetallic palladium(II) complexes with a bridging arylyene or diarylyene ligand, *trans,trans*-[L₂XPdRPaXL₂] (L = PMe₃ or PEt₃; R = C₆H₄C₆H₄, C₆H₄, C₆H₄OC₆H₄, C₆H₄SO₂C₆H₄, C₁₄H₈ or C₆F₄C₆F₄; X = Br, Cl or I). The structure of *trans,trans*-[(Et₃P)₂BrPdC₆H₄C₆H₄PdBr-(PEt₃)₂] has been determined by X-ray crystallography. The complex *trans,trans*-[(Me₃P)₂BrPdC₆H₄C₆H₄PdBr-(PMe₃)₂] reacts with *tert*-butyl, cyclohexyl and phenyl isocyanides to give the iminoacylpalladium complexes *trans,trans*-[(Me₃P)₂BrPdC(=NR')C₆H₄C₆H₄C(=NR')PdBr(PMe₃)₂] (R' = Bu^t, C₆H₄ or Ph) through insertion of the isocyanide into the two aryl–palladium bonds. Reactions of *trans,trans*-[L₂IPdRPaIL₂] (R = C₆H₄C₆H₄, L = PEt₃; R = C₆H₄, L = PMe₃ or PEt₃) with CO (1 atm) proceeded smoothly to give *trans,trans*-[(Et₃P)₂IPdCOC₆H₄C₆H₄PdI(PEt₃)₂] and *trans,trans*-[L₂IPdCOC₆H₄PdIL₂] (L = PMe₃ or PEt₃), respectively, through insertion of CO into one of the aryl–palladium bonds.

Dinuclear transition metal complexes with a bridging arylyene or diarylyene ligand, L_nMRML_n or L_nMRRML_n, possess a unique structure containing the π-conjugated ligand between two metal centres and have a rigid rod-like arrangement of the molecule. Dipalladium complexes with diarylyene ligands have been used as the precursor of transition metal-containing macrocycles which can include various guest molecules.¹ Analogous dinuclear complexes of Ni and Pd would show unique chemical properties such as insertion of small molecules into the metal–carbon bonds since they have two aryl–nickel² or palladium³ bonds with higher reactivity than aryl–platinum bonds.⁴ Oxidative addition of arylyene dihalides to zerovalent complexes of group 10 metals is expected to provide a convenient synthetic route of the above complexes. We have previously reported the synthesis of several arylyene bridged dinickel(II) complexes⁵ in connection with the reaction mechanism of dehalogenative polycondensation of dibromoarenes promoted by nickel(0) complexes.⁶ Stang and his co-workers⁷ have applied the bis-oxidative addition to synthesis of diarylyene bridged dipalladium and diplatinum complexes. These reactions are more easily performed than an alternative metathesis reaction of dichloro complexes of Group 10 metals with arylating reagents.⁸ In this paper we will describe the preparation of dinuclear palladium complexes with a bridging arylyene ligand *via* oxidative addition of dihalogenoarylyene compounds and their reaction with isocyanides and CO resulting in insertion into the palladium–aryl bonds.

Results and Discussion

Preparation and structures of arylyene bridged dipalladium complexes

Reactions of *trans*-[PdEt₂(PMe₃)₂] {or *trans*-[PdEt₂(PEt₃)₂]} with 4,4'-dibromobiphenyl, 4,4'-diiodobiphenyl, 1,4-diiodo-



Scheme 1

benzene, di(4-bromophenyl) ether, di(4-chlorophenyl) sulfone, 9,10-dibromoanthracene and 4,4'-dibromooctafluorobiphenyl in a 2:1 ratio at 55 °C in thf or toluene give the corresponding dinuclear complexes *trans,trans*-[L₂XPdRPaXL₂] (Scheme 1). The reactions are believed to involve oxidative addition of two C–Cl, C–Br or C–I bonds of the substrate to two molecules of [Pd(CH₂=CH₂)L₂] (L = PMe₃ or PEt₃) generated through

[†] E-Mail: yjkim@knusun.kangnung.ac.kr

Non-SI unit employed: atm = 101 325 Pa.

Table 1 Yields and analytical data of the complexes

Complex		Colour	Yield (%)	Analysis ^a (%)		
				C	H	N
1	$[(\text{Me}_3\text{P})_2\text{BrPdC}_6\text{H}_4]_2$	Off-white	52	35.22 (34.77)	5.35 (5.35)	
2	$[(\text{Me}_3\text{P})_2\text{IPdC}_6\text{H}_4]_2$	Colorless	53	31.07 (31.23)	5.01 (4.80)	
3	$[(\text{Et}_3\text{P})_2\text{BrPdC}_6\text{H}_4]_2$	Colorless	62	43.04 (43.35)	6.76 (6.87)	
4	$[(\text{Et}_3\text{P})_2\text{IPdC}_6\text{H}_4]_2$	Colorless	52	40.01 (39.62)	6.53 (6.28)	
5	$[(\text{Me}_3\text{P})_2\text{Pd}]_2\text{C}_6\text{H}_4$	Colorless	68	25.89 (25.52)	4.81 (4.76)	
6	$[(\text{Et}_3\text{P})_2\text{Pd}]_2\text{C}_6\text{H}_4$	Colorless	53	35.51 (35.49)	6.42 (6.35)	
7	$[(\text{Me}_3\text{P})_2\text{BrPdC}_6\text{H}_4]_2\text{O}$	Colorless	58	34.31 (34.11)	5.49 (5.25)	
8	$[(\text{Me}_3\text{P})_2\text{ClPdC}_6\text{H}_4]_2\text{SO}_2^b$	Colorless	34	35.64 (35.84)	5.61 (5.51)	
9	$[(\text{Me}_3\text{P})_2\text{BrPd}]_2\text{C}_{14}\text{H}_8$	Pale green	62	36.84 (36.60)	5.09 (5.20)	
10	$[(\text{Me}_3\text{P})_2\text{BrPdC}_6\text{F}_4]_2$	Colorless	69	29.95 (29.62)	3.87 (3.73)	
11	$[\text{PdBr}(\text{C}_6\text{H}_4\text{C}_6\text{H}_4\text{Br})(\text{PMe}_3)]_2$	Colorless	57	37.59 (37.89)	4.61 (4.59)	
12	$[\text{PdBr}(\text{C}_{14}\text{H}_8\text{Br})(\text{PMe}_3)]_2$	Pale green	69	40.64 (40.40)	4.50 (4.41)	
13	$[(\text{Me}_3\text{P})_2\text{BrPd}(\text{CNBu}^t)\text{C}_6\text{H}_4]_2$	Pale green	68	41.02 (41.03)	6.26 (6.28)	2.18 (2.81)
14	$[(\text{Me}_3\text{P})_2\text{BrPd}(\text{CNC}_6\text{H}_{11})\text{C}_6\text{H}_4]_2$	Pale yellow	85	43.47 (43.57)	6.43 (6.35)	2.63 (2.67)
15	$[(\text{Me}_3\text{P})_2\text{BrPd}(\text{CNPh})\text{C}_6\text{H}_4]_2$	Pale yellow	89	43.91 (44.08)	5.49 (5.26)	2.77 (2.71)
16	$[(\text{Me}_3\text{P})_2\text{IPd}(\text{CNBu}^t)\text{C}_6\text{H}_4]_2$	Yellow	60	33.07 (33.19)	5.77 (5.77)	2.46 (2.76)
17	$[(\text{Et}_3\text{P})_2\text{IPd}]_2\text{COC}_6\text{H}_4\text{C}_6\text{H}_4$	Yellow	58	39.70 (39.54)	6.33 (6.12)	
18	$[(\text{Me}_3\text{P})_2\text{IPd}]_2\text{COC}_6\text{H}_4$	Pale yellow	98	25.91 (26.08)	4.26 (4.61)	
19	$[(\text{Et}_3\text{P})_2\text{IPd}]_2\text{COC}_6\text{H}_4$	Yellow	84	35.83 (35.69)	6.31 (6.18)	

^a Required values are given in parentheses. ^b S 4.47 (3.99)%.




Fig. 1 An ORTEP¹¹ drawing of complex **3** showing the atom-labeling scheme and 50% probability thermal ellipsoids. Symmetry-equivalent atoms are generated by the crystallographic inversion center located midway between C(4) and C(4*). Selected bond distances and angles: Pd–C(1) 2.008(8), Pd–P(2) 2.298(3), Pd–P(1) 2.307(3), Pd–Br(1) 2.5254(14) and C(4)–C(4*) 1.50(2) Å; C(1)–Pd–P(2) 89.2(3), P(1)–Pd–Br(1) 90.53(9), C(1)–Pd–P(1) 89.9(2), C(2)–C(1)–C(6) 116.0(8), P(2)–Pd–P(1) 177.2(1), C(2)–C(1)–Pd 121.2(7), C(1)–Pd–Br(1) 179.2(3), C(6)–C(1)–Pd 112.8(7) and P(2)–Pd–Br(1) 90.39(9)°

β -hydrogen elimination of an ethyl ligand of $[\text{PdEt}_2\text{L}_2]$ ($\text{L} = \text{PMe}_3$ or PET_3) and ensuing reductive elimination of ethane.⁹ Yields and analytical and spectroscopic data of the complexes are summarized in Tables 1 and 2. Complexes **2**, **4** and **5** are obtained also from the reaction of diiodoaromatic compounds with $[\text{Pd}(\text{CH}_2=\text{CHPh})(\text{PMe}_3)]_2$ at room temperature. The aromatic dihalides, including sterically demanding 9,10-dibromoanthracene⁵ and a dichloroarylene compound, undergo smooth oxidative addition of both carbon–halogen bonds in the molecule to give the desired dinuclear complexes in moderate yields. A *trans* configuration around the Pd is supported by the ¹H and ¹³C NMR signals of PMe_3 or PET_3 ligands showing apparent triplets (PCH_3) or quintet (CH_3 or PET_3) arising from virtual coupling.¹⁰ Complexes **1–4**, **7** and **8** show the ¹³C NMR signals due to *ipso*- and *ortho*-carbons in the arylene ligand as triplets [$J(\text{CP}) = 4–7$ Hz], revealing that the two PR_3 ligands are magnetically equivalent.

Fig. 1 shows the molecular structure of complex **3** determined by X-ray crystallography. The molecule contains a crystallographic inversion centre at the midpoint of the C–C bond between two phenylene rings. The co-ordination sphere of each Pd can be described as a slightly distorted square plane with

average atomic displacement of 0.022 Å. The two phenyl rings are essentially planar and lie nearly perpendicular to the molecular plane, defined by Pd, P(1), P(2), Br(1) and C(1), with a dihedral angle of 82.2(1)°. This structure is similar to reported analogues of Ni and Pd, *trans,trans*- $[(\text{Et}_3\text{P})_2\text{BrNiC}_6\text{H}_4\text{C}_6\text{H}_4\text{NiBr}(\text{PET}_3)_2]$ ⁵ and *trans,trans*- $[(\text{Et}_3\text{P})_2\text{IPtC}_6\text{H}_4\text{C}_6\text{H}_4\text{PtI}(\text{PET}_3)_2]$ ⁷

Reactions of 4,4'-diiodobiphenyl and of 1,4-diiodobenzene giving complexes **2**, **4** and **5** proceed smoothly even at room temperature, whereas similar reactions of 4,4'-dibromobiphenyl and 9,10-dibromoanthracene with the palladium complex in a 1:2 molar ratio at room temperature give the mononuclear complexes *trans*- $[\text{PdBr}(\text{RBr-}p)(\text{PMe}_3)_2]$ ($\text{R} = \text{C}_6\text{H}_4\text{C}_6\text{H}_4$ **11** or C_{14}H_8 **12**). These results imply the easier oxidative addition of the C–I than the C–Br bond to the palladium(0) center, which is consistent with the previous observations that the relative reactivity of carbon(aryl)–halogen bonds toward oxidative addition to PdL_4 ($\text{L} =$ tertiary phosphine) decreases in the order $\text{C–I} > \text{C–Br} > \text{C–Cl}$.¹² The C–Br bond of **11** and **12** remaining intact could react with palladium(0) complexes at elevated temperature since the reactions in Scheme 1 were completed at 55 °C. In order to obtain further insight on the pathway of reactions we have examined the reaction of **11** with a palladium(0) complex $[\text{Pd}(\text{CH}_2=\text{CHPh})(\text{PMe}_3)_2]$, which has a similar structure to that of $[\text{Pd}(\text{CH}_2=\text{CH}_2)(\text{PMe}_3)_2]$ at 55 °C. Complex **1** was formed in 55% yield as shown in Scheme 2, suggesting stepwise oxidative addition of the two C–Br bonds in the substrate to the palladium(0) centers in the reactions in Scheme 1.


Reaction of CNR and CO with the diarylene bridged dipalladium complexes

The diarylene bridged dipalladium complexes prepared readily react with isocyanides and CO to give several new dinuclear complexes. Reactions of **1** with *tert*-butyl, cyclohexyl and phenyl isocyanides in a 1:2 molar ratio at room temperature proceed smoothly to afford *trans*- $[(\text{Me}_3\text{P})_2\text{BrPdC}(\text{=NR}')\text{C}_6\text{H}_4\text{C}_6\text{H}_4\text{C}(\text{=NR})\text{PdBr}(\text{PMe}_3)_2]$ **13–15** in 68–89% yields, as shown in Scheme 3. Complexes **13–15** do not undergo further insertion of the isocyanide into the Pd–CNR bonds at room temperature. Their IR spectra show a characteristic $\nu(\text{C}=\text{N})$ band at 1530–1590 cm^{-1} . The ¹³C–¹H NMR peaks due to the C=N carbons are observed at δ 176–184. The ¹H and ¹³C NMR spectra clearly

Table 2 The NMR data of the complexes (δ , J/Hz)

Complex	$^1\text{H}^a$	R	$^{13}\text{C}\{-^1\text{H}\}^b$	$^{31}\text{P}\{-^1\text{H}\}^c$
1	1.23 (36 H, CH_3) ^d	7.26 (ABq, 8 H)	14.4, ^d 125.6, 135.7, 135.8 (t, $J = 5$), 152.8 (t, $J = 6$)	-18.1
2	1.31 (36 H, CH_3) ^d	7.26 (ABq, 8 H)	15.8, ^d 125.6, 135.4 (t, $J = 5$), 135.8, 155.9 (t, $J = 6$)	-19.9
3	1.10 (36 H, CH_3) ^e	7.26 (ABq, 8 H)	8.2, 14.7, ^d 125.3, 135.1, 136.5 (t, $J = 4$), 152.7 (t, $J = 6$)	12.3
4	1.61 (m, 24 H, CH_2) ^e			
	1.08 (36 H, CH_3) ^e	7.27 (ABq, 8 H)	8.2, 16.0, ^d 125.3, 135.21, 136.5 (t, $J = 4$), 155.2	13.0
	1.67 (m, 24 H, CH_2)			
5	1.30 (36 H, CH_3) ^d	6.90 (s, 4 H)	16.2, ^d 136.0, 149.7	-20.5
6	1.08 (36 H, CH_3) ^e	6.86 (s, 4 H)	8.2, 15.9, ^d 136.4, 148.7	11.8
7	1.22 (36 H, CH_3) ^d	6.70 (d, 4 H)	14.4, ^d 118.7, 135.7 (t, $J = 5$), 146.0 (t, $J = 6$)	-17.5
		7.12 (d, 4 H)		
8	1.12 (36 H, CH_3) ^d	7.36 (d, 4 H)	13.3, ^d 125.3, 136.1 (t, $J = 5$), 137.4, 164.5 (t, $J = 7$)	-14.8
		7.55 (d, 4 H)		
9	0.98 (36 H, CH_3) ^e	7.34 (m, 4 H)	<i>f</i>	-16.2
	1.67 (m, 24 H, CH_2)	8.68 (m, 4 H)		
10	1.36 (36 H, CH_3) ^d		15.0 ^{d,g}	-13.8
11	1.12 (18 H, CH_3) ^d	7.28 (m, 4 H)	14.4, ^d 120.6, 125.9, 128.1, 131.8, 134.4, 136.1 (t, $J = 5$), 140.2, 155.0 (t, $J = 6$)	-18.0
		7.46 (m, 4 H)		
12	0.99 (18 H, CH_3) ^d	7.47 (m, 2 H)	14.7, ^d 118.1, 124.1, 126.9, 128.0, 130.6, 133.5, 137.5 (t, $J = 4$), 164.3	-15.7
		7.53 (m, 2 H)		
		8.46, 8.82 (d, 4 H)		
13	1.32 (36 H, CH_3) ^d	7.60 (d, 4 H)	15.1, ^d 31.1, 57.3, 126.2, 130.8, 140.5, 144.1, 176.2 (C=N)	-19.5
		8.28 (d, 4 H)		
14	1.38 (36 H, CH_3) ^d	7.59 (d, 4 H)	15.4, ^d 24.6, 25.7, 34.0, 68.5, 126.4, 129.8, 140.7, 142.8, 182.2 (C=N)	-17.8
		8.21 (d, 4 H)		
15	1.25 (36 H, CH_3) ^d	7.71 (d, 4 H)	15.0, ^d 121.1, 125.4, 126.6, 128.7, 130.6, 141.3, 143.9, 151.2, 184.0 (C=N)	-18.8
		8.39 (d, 4 H)		
16	1.39 (36 H, CH_3) ^d	8.14 (s, 4 H)	16.4, ^d 31.2, 58.1, 129.9, 145.2, 179.1 (C=N)	-21.7
17	1.08 (36 H, CH_3) ^e	7.34 (m, 4 H)	8.06, 8.18, 15.9, ^d 16.2, ^d 125.8, 126.0, 129.9, 133.6, 137.0, 140.7, 144.9,	10.7
	1.70 (18 H, CH_2)	7.69 (d, 2 H)	159.1, 234.4 (CO)	13.0
	1.86 (6 H, CH_2)	8.18 (d, 2 H)		
18	1.28 (18 H, CH_3) ^d	7.42 (d, 2 H)	16.0, ^d 16.1, ^d 128.3, 135.6 (t, $J = 5$), 138.6 (t, $J = 15$), 169.7 (t, $J = 5$), 235.4 (CO)	-19.4
	1.36 (18 H, CH_3) ^d	7.57 (d, 2 H)		-21.7
19	1.07 (36 H, CH_3) ^e	7.36 (d, 2 H)	8.01, 8.14, 16.1, ^d 16.2, ^d 136.3 (t, $J = 5$), 138.6, 168.6, 233.8 (CO)	10.0
	1.67 (18 H, CH_2)	7.84 (d, 2 H)		12.5
	1.86 (6 H, CH_2)			

^a Obtained in CDCl_3 at 25 °C. Peak positions were referenced to internal SiMe_4 . The ^1H NMR peaks not included are as follows: **13**, 1.59 (s, CH_3 , 18 H); **14**, 1.43 (m, 4 H, CH_2), 1.74 (m, 10 H, CH_2) and 1.86 (m, 4 H, CH_2); **15**, 7.20 (t, 2 H, C_6H_5), 7.43 (t, 4 H, C_6H_5) and 7.78 (d, 2 H, C_6H_5); **16**, 1.61 (s, CH_3 , 18 H). ^b Obtained in CDCl_3 at 25 °C. Peak positions were referenced to internal SiMe_4 . ^c Obtained in CDCl_3 at 25 °C. Peak positions were referenced to external 85% H_3PO_4 . ^d Observed as an apparent triplet due to virtual coupling. ^e Observed as an apparent quintet due to virtual coupling. ^f ^{13}C NMR spectrum not obtained due to low solubility of the complex. ^g Precise assignment of the peaks is not feasible due to severe splitting caused by C-F coupling.



indicate the symmetrical structure of the molecules formed through insertion of the isocyanide into the two aryl-palladium bonds.

Reaction of complex **5** with 2 mol equivalents of *tert*-butyl isocyanide gives a neutral complex with symmetrical structure,


trans,trans-[(Me_3P)₂IPdC(=NBu^t)C₆H₄C(=NBu^t)PdI(PMe₃)₂]**16** after recrystallization of the product. The crude product before purification shows an IR peak at 2160 cm^{-1} assigned to $\nu(\text{C}\equiv\text{N})$ of the isocyanide co-ordinated to a Pd. These results suggest that the reaction gives not only **16** but also cationic complexes such as [(Me_3P)₂(CNR')PdC₆H₄CPd(CNR')(PMe₃)₂]₂ or [(Me_3P)₂IPd(C=NR')C₆H₄Pd(CNR')I(PMe₃)₂] ($\text{R}' = \text{Bu}^t$) which may be regarded as the intermediate for formation of **16**.

Isocyanides which are isoelectronic to CO have been known to undergo facile insertion into Pd-C bonds.¹³ Bimetallic palladium complexes with a bridging dialkynyl ligand have been reported to undergo successive insertion of two or more molecules of aromatic isocyanide into a palladium-alkynyl bond giving the dinuclear product with unsymmetrical structure.¹⁴ A bimetallic platinum complex with a bridging thienylene group also reacts with aryl isocyanides to give unsymmetric or symmetric iminoacylplatinum complexes at much higher temperature.¹⁵ Reactions in Scheme 3 occur smoothly at room temperature, which implies facile insertion of the isocyanide into the palladium-aryl bond.


Reactions of CO (1 atm) with complexes **4**, **5** and **6** at room temperature result in insertion of a CO molecule into a Pd-C bond to give the arylpalladium complexes **17**, **18** and **19**, respectively, as shown in Scheme 4. The products are obtained as yellow solids and are characterized by spectroscopic meas-

urement and elemental analyses. The IR spectra show a strong $\nu(\text{CO})$ absorption in the range 1618–1630 cm^{-1} which is comparable with the $\nu(\text{CO})$ of a benzoyl group co-ordinated to palladium.³ The ^{13}C NMR signals at δ 234–235 support the single insertion of CO into the palladium–aryl bond. Two triplets for PMe_3 (**18**) and two triplets and two singlets for PEt_3 (**17** and **19**) in the ^{13}C NMR spectra as well as two singlets in the ^{31}P - $\{^1\text{H}\}$ NMR spectra support the presence of unequivalent phosphine ligands due to the unsymmetrical molecule structure. The presence of the aryl group in the complex at one Pd prevents further reaction of CO with the other Pd. This co-operation effect of the bimetallic system enabled synthesis of the unsymmetrical bimetallic palladium complexes with a bridging COR ligand.

As described above, one can easily prepare arylene-bridged dinuclear complexes from the reaction of diethylpalladium(II) complexes with various dihalogenoaromatic compounds. The sulfonyl group of $\text{ClC}_6\text{H}_4\text{SO}_2\text{C}_6\text{H}_4\text{Cl}$ facilitated oxidative addition of both C–Cl bonds in the molecule, whereas oxidative addition of aryl chloride with palladium(0) complexes is not common.¹⁶ The arylene bridged dinuclear palladium complexes undergo insertion of small molecules such as isocyanides and CO into their Pd–C bonds at room temperature; CO reacts with one of bonds, while insertion of isocyanide occurs at both sides of the arylene ligand.



Scheme 3



Scheme 4

Experimental

General, materials and measurements

All manipulations of the air-sensitive compounds were performed under a nitrogen or argon atmosphere with use of a standard Schlenk technique. Solvents were distilled from sodium–benzophenone and stored under a nitrogen atmosphere. The complexes $\text{trans}[\text{PdEt}_2\text{L}_2]$ (L = PMe_3 or PEt_3) were prepared according to the literature methods.⁹ 4,4'-Dibromobiphenyl, 4,4'-diiodobiphenyl, 1,4-diiodobenzene, di(4-bromophenyl) ether, di(4-chlorophenyl) sulfone, 9,10-dibromoanthracene and 4,4'-dibromooctafluorobiphenyl were obtained from Aldrich Co. and further purified by recrystallization.

Elemental analyses were carried out by analytical laboratories at the Tokyo Institute of Technology in Japan and the Basic Science Institute in Korea. The IR spectra were recorded on a Hitachi 270-30 spectrophotometer, NMR (^1H , ^{13}C - $\{^1\text{H}\}$ and ^{31}P - $\{^1\text{H}\}$) spectra on Bruker 300 and 500 MHz spectrometers. Chemical shifts were referred to internal SiMe_4 or to external 85% H_3PO_4 .

Preparations

Complexes 1–10. To a Schlenk flask containing $\text{trans}[\text{PdEt}_2(\text{PMe}_3)_2]$ (0.707 g, 2.23 mmol) at 0 °C were added 4,4'-dibromobiphenyl (0.348 g, 1.12 mmol) and thf (5 cm^3) in that order. The mixture was heated at 55 °C for 1 h to give an orange solution. Stirring at room temperature for 2 h resulted in a reddish yellow solid, which was filtered off, washed with Et_2O (2 cm^3) and recrystallized from CH_2Cl_2 –hexane to give $\text{trans},\text{trans}[(\text{Me}_3\text{P})_2\text{BrPdC}_6\text{H}_4\text{C}_6\text{H}_4\text{PdBr}(\text{PMe}_3)_2]$ **1** as slightly gray crystals (0.481 g, 52%). Complexes **2–10** were prepared analogously.

Complexes **2**, **4** and **5** were alternatively prepared by the following method. To a Schlenk flask containing $\text{trans}[\text{PdEt}_2(\text{PMe}_3)_2]$ (0.352 g, 1.11 mmol) at 0 °C were added styrene (0.462 g, 4.44 mmol) and thf (3 cm^3) in that order. The mixture was heated at 55 °C for 1 h to give a homogeneous yellow solution. This was cooled to room temperature and 4,4'-diiodobiphenyl (0.451 g, 1.11 mmol) added. Stirring for 2 h at room temperature resulted in white solids which were filtered off, washed with Et_2O (2 $\text{cm}^3 \times 2$), hexane (2 cm^3) and recrystallized from CH_2Cl_2 –hexane to give white solid **2** (0.543 g, 53%).

Complexes 11 and 12. To a Schlenk flask containing $\text{trans}[\text{PdEt}_2(\text{PMe}_3)_2]$ (0.354 g, 1.12 mmol) at 0 °C were added styrene (0.581 g, 5.60 mmol) and thf (4 cm^3) in that order. The mixture was heated at 55 °C for 1 h to give a homogeneous yellow solution. This was cooled to room temperature and 4,4'-dibromobiphenyl (0.350 g, 1.12 mmol) added. Stirring for 2 h at room temperature resulted in white solids which were filtered off, washed with Et_2O (2 $\text{cm}^3 \times 2$), hexane (2 cm^3) and recrystal-

lized from CH₂Cl₂-hexane to give white solid **11** (0.362 g, 57%). Complex **12** was prepared analogously.

Reaction of [Pd(CH₂=CHPh)(PMe₃)₂] with complex **11.** To a Schlenk flask containing *trans*-[PdEt₂(PMe₃)₂] (0.127 g, 0.40 mmol) at 0 °C were added styrene (0.168 g, 1.60 mmol) and thf (4 cm³) in that order. The mixture was heated at 55 °C for 1 h to give a yellow solution. This was cooled to room temperature and complex **11** (0.230 g, 0.40 mmol) added. The reaction mixture was stirred at 55 °C for 1 h, and the resulting reddish purple solids were filtered off, washed with hexane, Et₂O, and recrystallized to give analytically pure **1** (0.186 g, 55%).

Reactions of isocyanides with complexes **1 and **5**.** To a CH₂Cl₂ (3 cm³) solution of complex **1** (0.203 g, 0.245 mmol) was added *tert*-butyl isocyanide (46 mg, 0.563 mmol) at room temperature. Stirring the mixture for 2 h resulted in pale green solids which were filtered off, washed with hexane, and recrystallized from CH₂Cl₂-hexane to give pale green crystals of *trans*-[(Me₃P)₂BrPdC(=N^tBu)⁺C₆H₄C₆H₄C(=N^tBu)⁻]PdBr(PMe₃)₂] **13** (0.165 g, 68%). IR (KBr): ν(C=N)/cm⁻¹ 1586. Reactions of cyclohexyl isocyanide and of phenyl isocyanide with **1** were carried out analogously to give *trans*-[(Me₃P)₂BrPdC(=NC₆H₁₁)⁺-C₆H₄C₆H₄C(=NC₆H₁₁)⁻]PdBr(PMe₃)₂] **14** and *trans*-[(Me₃P)₂BrPdC(=NPh)⁺C₆H₄C₆H₄C(=NPh)⁻]PdBr(PMe₃)₂] **15**, respectively. IR (KBr): ν(C=N)/cm⁻¹ 1576 (**14**), 1534 (**15**).

Reactions of *tert*-butyl isocyanide with complex **5** were carried out analogously to give *trans,trans*-[(Me₃P)₂IPdC(=N^tBu)⁺C₆H₄C(=N^tBu)⁻]PdI(PMe₃)₂] **16** as yellow crystals (yield 60%). IR (KBr): ν(C=N)/cm⁻¹ 1590.

Reactions of CO with complexes **4, **5** and **6**.** Complex **5** (0.227 g, 0.268 mmol) was dissolved in CH₂Cl₂ (3 cm³) at room temperature. After evacuation of the system CO (1 atm) was introduced and the initial colorless solution immediately turned to yellow. After stirring for 3 h at room temperature the solvent was evaporated under reduced pressure to give a yellow residue, which was recrystallized from CH₂Cl₂-hexane to give yellow crystals of *trans,trans*-[(Me₃P)₂IPdC(O)C₆H₄PdI(PMe₃)₂] **18** (0.230 g, 98%). IR (KBr): ν(C=O)/cm⁻¹ 1620. Reactions of CO with **4** and **6** were carried out analogously to give *trans,trans*-[(Et₃P)₂IPdC(O)C₆H₄C₆H₄PdI(PET₃)₂] **17** (58%) and *trans,trans*-[(Et₃P)₂IPdC(O)C₆H₄PdI(PET₃)₂] **19**, respectively. IR (KBr): ν(C=O)/cm⁻¹ 1630 (**17**), 1618 (**19**).

Crystallography

Crystal data. C₃₆H₆₈Br₂P₄Pd₂ **3**, *M* = 997.40, monoclinic, space group *P*2₁/*n* (no. 14), *a* = 14.225(4), *b* = 18.720(5), *c* = 9.101(3) Å, β = 107.31(2)°, *U* = 2314(1) Å³, *T* = 298 K, *Z* = 2, μ(Mo-Kα) = 2.7 mm⁻¹, 2968 reflections measured, 2774 unique (*R*_{int} = 0.018), from which 2766 with *I* > 2σ(*I*) were used in refinement. Final *R*₁ and *wR*₂ values are 0.0587 and 0.1326.

Full-matrix least squares refinement was carried out with Pd, Br, and carbons of the aryl group anisotropic and with PET₃ carbons isotropic. Many carbon atoms of the PET₃ ligands showed severe disorder, and determination of reasonable occupancy of the possible positions of the ethyl groups was not feasible. Hydrogens were located in calculated positions.

CCDC reference number 186/974.

See <http://www.rsc.org/suppdata/dt/1998/1775/> for crystallographic files in .cif format.

Acknowledgements

This work was supported by the Basic Science Research Institute Program (NO. BSRI-96-3440), Korean Ministry of Education. Y.-J. K. is grateful also to a bilateral exchange program between JSPS (Japan Society for the Promotion of Science) and KOSEF (Korea Science and Engineering Foundation).

References

- 1 J. Manna, J. A. Whiteford, P. J. Stang, D. C. Muddiman and R. D. Smith, *J. Am. Chem. Soc.*, 1996, **118**, 8731; P. J. Stang, N. R. Persky and J. Manna, *J. Am. Chem. Soc.*, 1997, **119**, 4777; J. Manna, C. J. Kuehl, J. A. Whiteford, P. J. Stang, D. C. Muddiman, S. A. Hofstadler and R. D. Smith, *J. Am. Chem. Soc.*, 1997, **119**, 11 611; P. J. Stang and B. Olenyuk, *Acc. Chem. Res.*, 1997, **30**, 502.
- 2 E. Carmona, P. Palma, M. Paneque, M. L. Poveda, E. Gutiérrez-Puebla and A. Monge, *J. Am. Chem. Soc.*, 1996, **108**, 6424; E. Carmona, E. Gutiérrez-Puebla, J. M. Marín, A. Monge, M. Paneque, M. L. Poveda and C. Ruiz, *J. Am. Chem. Soc.*, 1989, **111**, 2883; C. Amatore and A. Jutand, *J. Am. Chem. Soc.*, 1911, **113**, 2819; K. Osakada, R. Sato and T. Yamamoto, *Organometallics*, 1994, **13**, 4645.
- 3 B. Crociani, M. Nicolini and T. Boschi, *J. Organomet. Chem.*, 1971, **33**, C81; B. Crociani, M. Nicolini and R. L. Richards, *J. Organomet. Chem.*, 1976, **104**, 259; F. Ozawa, T. Sugimoto, Y. Yuasa, M. Santra, T. Yamamoto and A. Yamamoto, *Organometallics*, 1984, **3**, 683; F. Ozawa, T. Sugimoto, T. Yamamoto and A. Yamamoto, *Organometallics*, 1984, **3**, 692.
- 4 G. K. Anderson and R. J. Cross, *J. Chem. Soc., Chem. Commun.*, 1978, 819; *J. Chem. Soc., Dalton Trans.*, 1979, 1246; 1980, 712; R. J. Cross and J. Gemmill, *J. Chem. Soc., Dalton Trans.*, 1981, 2317; M. Kubota, D. A. Phillips and J. E. Jacobsen, *J. Coord. Chem.*, 1980, **10**, 125; G. K. Anderson, H. C. Clark and J. A. Davies, *Organometallics*, 1982, **1**, 64.
- 5 Z.-H. Zhou and T. Yamamoto, *J. Organomet. Chem.*, 1991, **414**, 119; Y.-J. Kim, R. Sato, T. Maruyama, K. Osakada and T. Yamamoto, *J. Chem. Soc., Dalton Trans.*, 1994, 943.
- 6 T. Yamamoto, A. Morita, Y. Miyazaki, T. Maruyama, H. Wakayama, Z.-H. Zhou, Y. Nakamura, T. Kanbara, S. Sasaki and K. Kubota, *Macromolecules*, 1992, **25**, 1214; T. Yamamoto, T. Maruyama, Z.-H. Zhou, T. Ito, T. Fukuda, Y. Yoneda, F. Begum, T. Ikeda, S. Sasaki, H. Takezoe, A. Fukuda and K. Kubota, *J. Am. Chem. Soc.*, 1994, **116**, 4832.
- 7 J. Manna, C. J. Kuehl, J. A. Whiteford and P. J. Stang, *Organometallics*, 1997, **16**, 1897.
- 8 W. D. Müller and H. A. Brune, *Chem. Ber.*, 1986, **19**, 759.
- 9 F. Ozawa, T. Ito, Y. Nakamura and A. Yamamoto, *J. Organomet. Chem.*, 1979, **168**, 375; F. Ozawa, T. Ito and A. Yamamoto, *J. Am. Chem. Soc.*, 1980, **102**, 6457; F. Ozawa, T.-I. Son, S. Ebina, K. Osakada and A. Yamamoto, *Organometallics*, 1992, **11**, 171.
- 10 P. S. Pregosin and R. W. Kunz, *³¹P and ¹³C NMR of Transition Metal Phosphine Complexes*, Springer, Heidelberg, 1979, pp. 65–68.
- 11 C. K. Johnson, ORTEP, Report ORNL-5138, Oak Ridge National Laboratory, Oak Ridge, TN, 1976.
- 12 G. W. Parshall, *J. Am. Chem. Soc.*, 1974, **96**, 2360; P. Fitton and E. A. Rick, *J. Organomet. Chem.*, 1971, **28**, 287; D. H. Gerlach, A. R. Kane, G. W. Parshall, J. P. Jenson and E. L. Muetterties, *J. Am. Chem. Soc.*, 1971, **93**, 3543; M. Hidai, T. Kashiwagi, T. Ikeuchi and Y. Uchida, *J. Organomet. Chem.*, 1971, **30**, 385.
- 13 Y. Yamamoto and H. Yamazaki, *Coord. Chem. Rev.*, 1972, **8**, 225; Y. Yamamoto, T. Tanase, T. Yanai, T. Asano and K. Kobayashi, *J. Organomet. Chem.*, 1993, **456**, 287; B. Crociani, M. Sala, A. Polo and G. Bombieri, *Organometallics*, 1986, **5**, 1369; J. G. P. Delis, P. G. Aibel, P. W. N. M. van Leeuwen, K. Vrieze, N. Veldman and A. L. Spek, *J. Chem. Soc., Chem. Commun.*, 1995, 2233; Y.-J. Kim, J.-Y. Lee, D.-H. Kim and S.-W. Lee, *Bull. Korean Chem. Soc.*, 1996, **17**, 663.
- 14 H. Ogawa, T. Joh and S. Takahashi, *J. Chem. Soc., Chem. Commun.*, 1988, 561; K. Onitsuka, H. Ogawa, T. Joh, S. Takahashi, Y. Yamamoto and H. Yamazaki, *J. Chem. Soc., Dalton Trans.*, 1991, 1531; K. Onitsuka, T. Joh and S. Takahashi, *J. Organomet. Chem.*, 1994, **464**, 247; K. Onitsuka, K. Yanai, F. Takei, T. Joh and S. Takahashi, *Organometallics*, 1994, **13**, 3862; F. Takei, K. Yanai, K. Onitsuka and S. Takahashi, *Angew. Chem., Int. Ed. Engl.*, 1996, **35**, 1554.
- 15 S. Kotani, K. Shiina and K. Sonogashira, *J. Organomet. Chem.*, 1992, **429**, 403; K. Onitsuka, K. Murakami, K. Matsukawa, K. Sonogashira, T. Adachi and T. Yoshida, *J. Organomet. Chem.*, 1995, **490**, 117.
- 16 P. M. Maitlis, P. Espinet and M. J. H. Russell, in *Comprehensive Organometallic Chemistry*, eds. G. Wilkinson, F. G. A. Stone and E. W. Abel, Pergamon, Oxford, 1982, vol. 6, pp. 311–313.

Received 3rd February 1998; Paper 8/00925B