

Butenynyl complexes of iron(II) containing the tripodal tetraphosphine ligand P(CH₂CH₂PMe₂)₃

Leslie D. Field,* Barbara A. Messerle,* Ronald J. Smernik, Trevor W. Hambley and Peter Turner

School of Chemistry, University of Sydney, Sydney, NSW 2006, Australia.
E-mail: L.Field@chem.usyd.edu.au

Received 8th April 1999, Accepted 16th June 1999

The preparation and characterisation of iron(II) η^3 -but-1-en-3-yn-2-yl complexes [Fe(η^3 -RC \equiv C–C=CHR)L]⁺ [L = P(CH₂CH₂PMe₂)₃, R = Ph **1**; R = Bu^t **2**; R = *p*-HC \equiv CC₆H₄ **3**] is reported. The phenyl substituted butenynyl complex **1** was prepared by the reaction of FeCl₂L **5**, FeH(Cl)L **6** or [FeH(H₂)L]⁺ **7** with phenylacetylene in alcohol solvent. The coordinated but-1-en-3-yn-2-yl fragment is bound as a σ -vinyl/ π -acetylenic ligand. In solution, complex **1** exists as a pair of equilibrating isomers (**1a** and **1b**) which differ in the anchoring mode of the butenynyl ligand *i.e.* depending on whether the π -bound acetylenic group is *cis* or *trans* to the apical phosphorus of L in the octahedral coordination sphere. Assignment of the relative stereochemistry of **1a** and **1b** was achieved by analysis of the 2D NOESY spectrum. Exchange peaks in the NOESY spectrum also provided information on the mechanism of exchange between **1a** and **1b**. The crystal structure of **1** showed the solid state structure to be that of the major solution state isomer **1a** (π -bound acetylenic group is *cis* to the apical phosphorus of L). Complex **1** catalyses the stereospecific head-to-head dimerisation of phenylacetylene to *Z*-1,4-diphenylbut-1-en-3-yne.

Introduction

The reaction of metal complexes with terminal alkynes to afford butenynyl complexes is an important C–C bond forming reaction. In a number of cases, the free butenyne is subsequently liberated, effectively completing the catalytic cycle of head-to-head dimerisation of the alkyne.^{1,2} As part of a wider study examining the dimerisation (and oligomerisation) of acetylenes, we have examined a range of metal complexes where alkyne coupling occurs. Octahedral complexes containing tripodal tetradentate ligands appear particularly suited to alkyne dimerisation since the two sites available for alkyne coordination are constrained by the nature of the ligand to be mutually *cis*, facilitating the coupling of the alkyne fragments. Indeed, both the formation of butenynyl complexes and catalytic head-to-head coupling of terminal alkynes have been reported for complexes containing the tripodal tetraphosphine ligands P(CH₂CH₂PPh₂)₃ (L¹) **8**² and P(CH₂CH₂CH₂PMe₂)₃ (L²) **9**.³

The preparation and characterisation of a number of butenynyl complexes containing bi- and tetra-dentate phosphine ligands have been reported. Iron(II) dihydrogen hydrido complexes containing the bidentate phosphines 1,2-bis(dimethylphosphino)ethane (dmpe) and 1,2-bis(diethylphosphino)ethane (depe) react with terminal alkynes to form bis(acetylide) complexes in good yield.⁴ The bis(acetylide) complexes are protonated to initially form metal vinylidene complexes which rearrange and couple to form complexes containing coordinated butenyne. For some bis(acetylide) complexes, methanol employed as a reaction solvent is sufficiently acidic to carry out the protonation, in others, a stronger acid (*e.g.* trifluoroacetic acid) is required.⁵ The dichloro complex FeCl₂(dmpe)₂ also reacts with phenylacetylene in the presence of hexafluorophosphate to give the corresponding butenynyl complex.⁶

Reaction of the ruthenium dihydrogen hydrido complex [RuH(H₂)L]⁺ with terminal alkynes proceeds *via* two isolable intermediates, the σ -alkenyl complex [Ru(CH=CHR)L]⁺ and the σ -alkynyl complex [Ru(C \equiv CR)L]⁺, to eventually give the corresponding butenynyl complex [Ru(η^3 -RC \equiv C–C=CHR)-

L]⁺.^{2a-c,e} Reaction of the osmium dinitrogen hydrido complex [OsH(N₂)L]⁺ with terminal alkynes also yields butenynyl complexes.^{2d} The reaction proceeds *via* the vinylidene hydride complex [OsH(C=CHR)L]⁺ and one equivalent of RCH=CH₂ is produced as the reaction progresses. Butenynyl complexes of osmium can also be prepared directly from the dichloro complex OsCl₂L.^{2d}


The reaction of the iron complexes [FeH(X)L]⁺ (X = N₂, H₂) with terminal alkynes affords σ -alkenyl complexes [Fe(HC=CHR)L]⁺ and σ -alkynyl complexes [Fe(C \equiv CR)L]⁺, but these do not react further to give the corresponding butenynyl complexes.⁷

We have recently reported⁸ an efficient and high-yielding synthesis of the tetradentate ligand P(CH₂CH₂PMe₂)₃ (L) **4**, which has facilitated investigation of iron complexes containing **4**.⁹ Here, we report the synthesis of the iron(II) butenynyl complexes, [Fe(η^3 -RC \equiv CC=CHR)L]⁺ [R = Ph **1**, Bu^t **2**, *p*-HC \equiv CC₆H₄ **3**] and their characterisation by NMR spectroscopy and X-ray crystallography.

Results and discussion

Preparation of butenynyl complexes

The iron(II) diphenylbutenynyl complex [Fe(η^3 -PhC \equiv C–C=CHPh)L]⁺ **1** was prepared by the reaction of phenylacetylene with the dichloro complex **5**, the chloro hydrido complex **6** or the dihydrogen hydrido complex **7** in alcohol solvent (Scheme 1). In the reactions of phenylacetylene with **6** and **7**, styrene was observed as a by-product.



Scheme 1




Fig. 1 Schematic representation of cross peaks observed between L and butenylnyl ligands in the NOESY spectrum of the major diphenylbutenylnyl isomer **1a**. (a) Cross peaks between the vinylic proton and Me_A, backbone protons H¹ and H², and between the *ortho* protons of Ph² and Me_A indicate that Ph² lies near to the central phosphorus P_C; (b) cross peaks between the *ortho* proton of Ph¹ and Me_A, Me_B and Me_C, indicate that Ph¹ lies near the terminal phosphorus P_U.





Fig. 2 Schematic representation of cross peaks observed between L and butenylnyl ligands in the NOESY spectrum of the minor diphenylbutenylnyl isomer **1b**. Cross peaks between the vinylic proton and Me_C, indicate that Ph² lies near the terminal phosphorus P_U. Cross peaks between the *ortho* protons on Ph¹ and the ligand backbone protons H¹ and H², and also Me_A indicate that Ph¹ lies near to the central phosphorus P_C.

Reaction of phenylacetylene with **6** and **7** was rapid and no intermediates were detected (by ³¹P NMR) in either case. Reaction of **5** with phenylacetylene was significantly slower, and a number of (uncharacterised) intermediate complexes were visible if the course of the reaction was followed by ³¹P NMR. The addition of an ethanol solution of sodium tetraphenylborate to an ethanol solution of **1** resulted in the immediate precipitation of the red tetraphenylborate salt of **1**. The tetrafluoroborate salt of **1** was also prepared by the addition of sodium tetrafluoroborate, however, the BF₄⁻ salt was more soluble in methanol and ethanol, resulting in incomplete precipitation. Both the tetrafluoroborate and tetraphenylborate salts of **1** were soluble in acetone.

The ³¹P NMR spectrum of **1** (BPh₄⁻ salt in acetone) at room temperature contains sharp resonances at δ 171.9 (t, P_C), 64.4 (t, P_U) and 49.5 (dd, P_T). For the purposes of identifying the phosphorus nuclei of the coordinated ligand L, mutually *trans* terminal phosphorus nuclei were labelled P_T, the central phosphorus from which the three arms radiate was labelled P_C and the remaining terminal phosphorus was labelled P_U (see Figs. 1 and 2 for a labelled diagram). In complex **1**, no spin-spin coupling was detected between the terminal phosphines, P_T and P_U, in contrast with most complexes of L which have been characterised.⁹ A smaller set of broadened resonances arising from a second, minor product, was observed at δ 165.4 (P_C), 78.9 (P_U) and 54.1 (P_T). At 233 K both species gave rise to sharp resonances in the ³¹P NMR spectrum. The major species gave rise to signals at δ 171.7 (t, P_C), 65.1 (t, P_U) and 50.5 (dd, P_T); the minor species gave rise to resonances at δ 164.8 (dt, P_C), 79.3 (dt, P_U) and 55.4 (dd, P_T). The coupling constant between P_T and P_U for the minor species was also small (12.9 Hz). The ratio of major to minor species was *ca.* 11 : 1 at 233 K. The broadness


of the resonances of the minor species at 300 K is due to exchange between the two products at this temperature. The two species were identified (*vide infra*) as the isomeric complexes **1a** and **1b** which differ in the orientation of the butenylnyl ligand with respect to the rest of the molecule. For convenience, the exchanging mixture of isomeric butenylnyl complexes **1a** and **1b** is referred to as **1**.



Reaction of the chloro hydrido complex FeH(Cl)L **6** with *tert*-butylacetylene afforded the corresponding butenylnyl complex **2**. In contrast to **1**, only one set of resonances was observed in the 233 K ³¹P NMR spectrum of **2**. This may be due to the absence of appreciable quantities of the minor product corresponding to **1b**, or fast exchange between isomers even at this temperature. Reaction of the chloro hydrido complex FeH(Cl)L **6** with 1,4-diethynylbenzene also afforded the corresponding butenylnyl complexes **3a** and **3b**. The ³¹P NMR spectrum of **3a,b** at both 300 and 233 K were similar to those of the phenylbutenylnyl complexes, with two products observed at 233 K in the ratio of approximately 8 : 1. The ³¹P NMR spectra of **3** and **1** are almost identical and the major and minor isomers of **3** would therefore correspond to those of the phenylbutenylnyl complex.

Assignment of stereochemistry in butenylnyl complexes

Isomeric butenylnyl complexes have been observed for the ruthenium and osmium systems with L. In these cases no exchange is apparent in the 300 K ³¹P NMR spectrum. As discussed by Bianchini *et al.*,^{2b} octahedral complexes of L offer two different sites for the unsymmetrical bidentate butenylnyl ligand and hence two isomers are possible, with the triple bond *trans* to either P_C or *trans* to P_U. There is also the possibility of *E/Z* isomers about the butenylnyl double bond, resulting in a total of four possible isomers (I–IV). A *trans* stereochemistry



about the double bond of the ruthenium^{2b} and osmium^{2d} complexes of L was assigned (structures I and II) on the basis of the size of the long range coupling constant ³J_{C(2)-H(C4)}.¹⁸

The relative stereochemistry of the butenylnyl ligand with respect to L and the stereochemistry about the double bond for **1a** and **1b** was determined using 2D NMR COSY and

NOESY experiments. These spectra were recorded on the tetrafluoroborate salt (to avoid complication of ^1H spectra by resonances of the tetraphenylborate counter ion). Both the major and the minor isomers exhibited strong cross peaks between the vinylic proton and the relevant (see below) protons of L in the NOESY spectrum and were hence assigned a *Z* configuration about the butenyne double bond. The major isomer **1a** was found to have structure I whilst the minor isomer **1b** was assigned structure II.

In complex **1a**, strong NOESY cross peaks from the *ortho* protons on Ph¹ to Me_A, Me_B and Me_C indicated that the triple bond was *trans* to P_C. Cross peaks from the olefinic proton, to Me_A, H¹ and H² and from the *ortho* protons on Ph² to Me_A confirmed this assignment (Fig. 1). The expected cross peak from the vinylic proton H to the *ortho* proton of Ph² was obscured by the stronger cross peak between the *ortho* and *meta* protons on Ph² (δ 7.57 (Ph² H_{meta}), 7.56 (Ph² H_{ortho})).

The stereochemistry of the minor isomer **1b** was also determined from the NOESY spectrum and a full spectral assignment of the ^1H NMR spectrum of **1b** was achieved despite the large excess of **1a**. The ^1H resonances of L were assigned using the same methodology as was used for other octahedral complexes of L.⁹ A strong NOESY cross peak between the resonance of the vinylic proton and Me_C indicated that the double bond is located *trans* to P_C. This stereochemistry was confirmed by the presence of strong NOESY cross peaks between the resonances of the *ortho* protons on Ph¹ and H¹ and H² on the backbone of the PP₃ ligand (Fig. 2).

Exchange between isomers **1a** and **1b**

Exchange processes on a timescale similar to the mixing time of the NOESY experiment (usually 1–10 s) give rise to exchange cross peaks in the NOESY spectrum. These cross peaks are easily differentiated from the ‘real’ NOESY cross peaks in small molecules as they are characteristically of opposite sign. The NOESY spectrum of the diphenylbutenyne complexes **1a** and **1b** acquired at 303 K contains exchange peaks as well as the expected NOESY cross peaks.

The ^1H NMR spectrum of **1** (303 K, 600 MHz) contained resonances due to the two isomers **1a** and **1b**, although the resonances for the minor isomer **1b** were significantly broadened by exchange. The NOESY spectrum acquired at 303 K indicated that exchange processes were occurring within as well as between the two isomeric complexes. Each of the isomers **1a** and **1b** has three methyl resonances and exchange peaks are present between all of the methyl resonances of L (a six site exchange). Exchange peaks were also observed within and between the methylene resonances of L of both isomers.

In the aromatic region of the 2D NOESY spectrum acquired at 303 K (Fig. 3), more specific exchange was observed. An exchange peak was observed between the vinylic proton of the major isomer and the corresponding vinylic proton in the minor isomer. Exchange peaks were also observed between the alkyne-bound phenyl group in the major isomer (Ph¹) and the alkyne-bound phenyl group in the minor isomer (Ph^{1'}), and between the corresponding alkene-bound phenyl groups Ph² and Ph^{2'} (Fig. 4).

The most probable mechanism for the exchange involves the decoordination of the acetylene of the butenyne ligand to give complex **10** (where the butenyne is coordinated by the σ -bond to the vinylic carbon) followed by rotation around the metal–carbon bond and re-coordination of the alkyne to give the other stereoisomer (Scheme 2). There is ample precedent for the existence of η^1 -bound butenyne complexes, including X-ray crystal structures.^{1a,b}

X-Ray crystallography of $[\text{FeL}(\eta^3\text{-PhC}\equiv\text{C}-\text{C}=\text{CHPh})]\text{BPh}_4$ **1a**

Diffraction quality crystals were obtained by slow evaporation of a saturated acetone solution of the tetraphenylborate salt of




Fig. 3 Aromatic region of the NOESY spectrum (600 MHz, 303 K, acetone- d_6) of $[\text{Fe}(\eta^3\text{-PhC}\equiv\text{C}-\text{C}=\text{CHPh})]^+$ **1** (BPh_4 salt). Negative cross peaks are represented by dashed contours, positive peaks (due to exchange) are represented as solid contours.





Fig. 4 Schematic representation showing the exchange observed in the 303 K NOESY spectrum between H-*o* of **1a** and H'-*o* of **1b**, Ph¹ of **1a** and Ph^{1'} of **1b**, and Ph² of **1a** and Ph^{2'} of **1b**.



1. The structure determination shows the solid state structure to be the same as that determined for the major isomer **1a** by 2D NMR methods (Fig. 5). The crystal data parameters are summarised in Table 1. Selected bond lengths and angles are listed in Tables 2 and 3.

Comparison of the structure of **1a** with that of the triphenylphosphine chloro complex $[\text{FeCl}(\text{PPh}_3)_3]\text{BPh}_4$ **11**^{9a} shows the Fe–L fragment to be similar for both complexes, despite the different co-ligands (Table 2). Fe–P bond lengths




Fig. 5 ORTEP plot (25% thermal ellipsoids, non-hydrogen atoms) of $[\text{Fe}(\eta^3\text{-PhC}\equiv\text{C-C=CHPh})\text{L}]^+$ **1a** (BPh_4 salt).

Table 1 Crystallographic data for $[\text{Fe}(\eta^3\text{-PhC}\equiv\text{C-C=CHPh})\text{L}]^+$ **1a** (BPh_4 salt)

Empirical formula	$\text{C}_{52}\text{H}_{61}\text{BFeP}_4$
Formula weight	876.61
Crystal system	Triclinic
Space group	$P\bar{1}$
Z	2
Lattice Parameters:	
$a/\text{\AA}$	13.459(4)
$b/\text{\AA}$	13.517(6)
$c/\text{\AA}$	13.843(6)
α°	76.75(4)
β°	74.45(4)
γ°	78.76(4)
$V/\text{\AA}^3$	2337(2)
T/K	21.0
$\mu(\text{Mo-K}\alpha)/\text{cm}^{-1}$	4.935
No. of reflections measured	
Total	8595
Unique	8218
R_{int}	0.030
Residuals: R, R_w	0.053, 0.043

for **1a** are slightly shorter than those of **11**. The bond angles are similar, with deviations from the octahedral angles of 90 and 180° brought about by the small natural bite angle of L.

A number of η^3 -butenynyl complexes have been reported in the literature, including X-ray structures of complexes of tungsten,¹⁹ iron,^{5b,c} ruthenium^{2a,20} and osmium.²¹ The most relevant structures to compare with that of **1a** are the iron(II) η^3 -1,4-diphenylbutenynyl complex **12**^{5b,c} containing two dmpe ligands, and the ruthenium(II) η^3 -1,4-bis(trimethylsilyl)-butenynyl complex **13**^{2a} containing the tripodal tetradentate phosphine L¹ **8** (Table 3). The structures of **1a** and **12** are very similar. The Fe–C₁ bond is slightly longer in **12**, as is the C₂–C₃ bond. The C₁–C₂–C₃ bond angle is smaller in **1a** than in **12**.


The ruthenium butenynyl complex **13** exhibits the same regio- and stereo-chemistry as **1a**, with the alkyne *trans* to the central phosphorus and the metal *trans* to the non-hydrogen substituent on the double bond. The metal–butenynyl bonds are longer for **13**, as would be expected for the larger second-row metal. Bond lengths in the butenynyl fragment in **13** are similar to those of **1a** and **12**, however, **13** has a smaller R–C₁–C₂ bond angle and a larger C₁–C₂–C₃ bond angle.

Catalytic dimerisation of phenylacetylene

A number of transition metal complexes are catalysts for the head-to-head dimerisation of terminal alkynes to 1,4-disubstituted butenyne,^{1,2} including the ruthenium complex **13**.


Reaction of **1** with an excess of phenylacetylene results in the

Table 2 Comparison of selected bond lengths (Å) and angles (°) in the FeL fragments of $[\text{Fe}(\eta^3\text{-PhC}\equiv\text{C-C=CHPh})\text{L}]\text{BPh}_4$ **1a** and $[\text{FeCl}(\text{PPh}_3)\text{L}]\text{BPh}_4$ **11**^{9a}



	1a	11 ^{9a}
Fe–P _C	2.140(2)	2.202(2)
Fe–P _U	2.218(2)	2.234(2)
Fe–P _T (P1)	2.266(2)	2.312(2)
Fe–P _T (P4)	2.280(2)	2.312(2)
P _C –Fe–P _U	87.45(8)	83.85(9)
P _C –Fe–P _T (P1)	83.32(7)	83.66(9)
P _C –Fe–P _T (P4)	83.58(7)	83.24(9)
P _U –Fe–P _T (P1)	99.38(7)	98.71(9)
P _U –Fe–P _T (P4)	93.84(7)	93.40(9)
P _T –Fe–P _T	160.91(7)	161.07(9)

Table 3 Comparison of bond lengths (Å) and angles (°) in the metal–butenynyl fragments of $[\text{Fe}(\eta^3\text{-PhC}\equiv\text{C-C=CHPh})\text{L}]\text{BPh}_4$ **1a**, $[\text{Fe}(\eta^3\text{-PhC}\equiv\text{C-C=CHPh})(\text{dmpe})_2]\text{BPh}_4$ **12**^{5b,c} and $[\text{Ru}(\eta^3\text{-Me}_3\text{SiC}\equiv\text{C-C=CHSiMe}_3)\text{L}^1]\text{BPh}_4$ **13**^{2a}



	1a	12 ^{5b,c}	13 ^{2a}
M–C ₁	2.176(6)	2.309(8)	2.485(3)
M–C ₂	2.071(5)	2.093(7)	2.234(3)
M–C ₃	1.986(5)	1.985(7)	2.144(3)
C ₁ –C ₂	1.250(7)	1.249(9)	1.247(5)
C ₂ –C ₃	1.380(7)	1.396(10)	1.392(5)
C ₃ –C ₄	1.340(7)	1.338(9)	1.335(5)
R–C ₁ –C ₂	150.7(7)	152.9(7)	142.0(3)
C ₁ –C ₂ –C ₃	144.4(6)	149.1(7)	154.0(3)
C ₂ –C ₃ –C ₄	136.7(5)	132.5(7)	133.5(3)

slow formation of 1,4-diphenylbut-3-en-1-yne, which was identified by GC–MS and by ¹H NMR. Only the *Z* isomer^{2c} was formed in the reaction and the butenynyl complex **1a** was the only iron complex detected by ³¹P NMR during the course of the reaction. The reaction rate for this reaction at 80 °C was approximately one turnover per day (from the ¹H NMR spectrum). This turnover rate is slow compared with other systems, and the reaction was not further investigated.

Conclusions

The iron(II) butenynyl complexes $[\text{Fe}(\eta^3\text{-RC}\equiv\text{C-C=CHR})\text{L}]^+$ (R = Ph **1**; R = Bu^t **2**; R = *p*-HC≡CC₆H₄ **3**) have been prepared. Complex **1** exists as a pair of isomers (**1a** and **1b**) which differ in the stereochemistry of binding of the butenynyl ligand. Assignment of the relative stereochemistry of **1a** and **1b** was achieved by analysis of the 2D NOESY spectrum. The stereoisomers **1a** and **1b** are in equilibrium and exchange peaks in the NOESY spectrum provided information on the mechanism of exchange. A crystal structure of **1a** showed the solid state

structure to be that of the major solution state isomer. Complex **1** catalyses the stereospecific head-to-head dimerisation of phenylacetylene to *Z*-1,4-diphenylbut-1-en-3-yne.

Experimental

All synthetic manipulations involving air sensitive materials were carried out under an inert atmosphere of argon in an argon filled dry box or under a nitrogen atmosphere using standard Schlenk techniques. THF, benzene and hexane were dried over sodium before distillation from sodium and benzophenone under nitrogen. Ethanol and methanol were distilled from magnesium under nitrogen. The iron(II) dichloride complex FeCl_2L **5**,^{9a} the iron(II) chloro hydrido complex $\text{FeH}(\text{Cl})\text{L}$ **6**^{9a} and the iron(II) dihydrogen hydrido complex $[\text{FeH}(\text{H}_2)\text{L}]^+$ **6**^{9c} were prepared using previously reported methods. ¹H, ¹³C and ³¹P NMR spectra were recorded on Bruker AMX400 or AMX600 spectrometers at the temperatures quoted. ¹H and ¹³C chemical shifts were internally referenced to residual solvent resonances. ³¹P spectra were referenced to external neat trimethyl phosphite at δ 140.85. IR spectra were recorded on a Perkin-Elmer 1600 series FTIR. Mass spectra were recorded on a Finnigan MAT TSQ-46 (San Jose, CA, USA) spectrometer equipped with a desorption probe, with a source temperature of 140 °C and an electron energy of 100 eV. Chemical ionisation (CI) was used, with methane (>99.999%) as the ionisation gas. Elemental analyses were carried out at the Joint Elemental Analysis Facility, The University of Sydney. Melting points were recorded on a Gallenkamp heating stage and are uncorrected.

Crystal structure determination

The crystallographic data for **1a** (BPh₄ salt) are summarised in Table 1. A red–orange crystal of **1a** having approximate dimensions of 0.42 × 0.32 × 0.07 mm was mounted on an Enraf-Nonius CAD4 diffractometer employing graphite monochromated Mo-K α radiation. Triclinic cell constants were obtained from a least-squares refinement against the setting angles of 25 reflections in the range 16 < 2 θ < 27°. Diffraction data were collected at a temperature of 21 ± 1 °C using ω – θ scans to a maximum 2 θ value of 50°. The intensities of three representative reflections measured every hour did not change significantly during the course of the data collection. The data were corrected for Lorentz, polarisation and absorption (analytical) effects.

All calculations were performed using the teXsan¹⁰ crystallographic software package. The structure was solved by direct methods¹¹ and expanded using Fourier techniques.¹² Neutral atom scattering factors were taken from Cromer and Waber.¹³ Anomalous dispersion effects were included in the structure factor calculation,¹⁴ and the values for $\Delta f'$ and $\Delta f''$ were those of Creagh and McAuley.¹⁵ The values for the mass attenuation coefficients are those of Creagh and Hubbell.¹⁶ Non-hydrogen atoms were refined anisotropically and the hydrogen atoms were included in the full matrix least squares refinement at calculated positions with group temperature factors. An ORTEP¹⁷ representation of the complex is shown in Fig. 5.

CCDC reference number 186/1518.

See <http://www.rsc.org/suppdata/dt/1999/2557/> for crystallographic files in .cif format.

Preparations

[Fe(η^3 -PhC \equiv C–C=CHPh)L]⁺ **1.** Phenylacetylene (33 mg, 320 μmol) was added to a stirred solution of $\text{FeH}(\text{Cl})\text{L}$ **6** (34 mg, 87 μmol) in methanol (10 ml), resulting in a colour change from yellow to red. On addition of sodium tetraphenylborate (40 mg, 120 μmol) in methanol (5 ml), a red precipitate formed. The crude product was isolated by filtration, washed with methanol

(20 ml) and dried *in vacuo* to yield $[\text{Fe}(\eta^3\text{-PhC}\equiv\text{C}-\text{C}=\text{CHPh})\text{L}]^+$ **1** (BPh₄ salt) (72 mg, 94%), mp 242–244 °C.

The tetrafluoroborate salt of **1** was also prepared in an analogous way using sodium tetrafluoroborate in the place of sodium tetraphenylborate. MS (+ Cl, CH₄) *m/z* (>150): 558 (M + 1, 14), 557 (M, 36), 388 (10), 363 (30), 233 (14), 206 (17), 205 (95), 165 (100). IR ν_{max} (Nujol): 1578w, 1592w.

Major isomer 1a. ³¹P-{¹H} NMR (BF₄ salt, 162 MHz, acetone-*d*₆, 300 K): δ 171.9 (t, 1P, P_C, ²J_{P(C)-P(T)}} = 26.2), 49.5 (dd, 2P, P_T, ²J_{P(T)-P(U)}} = 34.8 Hz), 64.4 (t, 1P, P_U). ³¹P-{¹H} NMR (BF₄ salt, 162 MHz, acetone-*d*₆, 233 K): δ 171.7 (t, 1P, P_C, ²J_{P(C)-P(T)}} = 25.8), 50.5 (dd, 2P, P_T, ²J_{P(T)-P(U)}} = 35.3 Hz), 65.1 (t, 1P, P_U).

¹H-{³¹P} NMR spectrum (600 MHz, acetone-*d*₆, 303 K): δ 1.95, 2.56 (2 × m, 2 × 2H, –P_CCH₂CHHP_T–), 2.61, 2.86 (2 × m, 2 × 2H, –P_CCH₂CH₂P_T–), 2.27 (m, 2H, –P_CCH₂CH₂P_U–), 2.10 (m, 2H, –P_CCH₂CH₂P_U–), 0.92, 1.29 [2 × s, 2 × 6H, 2 × P_T(CH₃)], 1.85 [s, 6H, P_U(CH₃)₂], 7.51 (s, 1H, C=CH), 7.84 (d, 2H, CCPh_{ortho}), 7.70 (d, 2H, CCPh_{meta}), 7.62 (t, 1H, CCPh_{para}), 7.99 (d, 2H, C=CHPh_{ortho}), 7.57 (d, 2H, C=CHPh_{meta}), 7.39 (t, 1H, C=CHPh_{para}).

¹H-{³¹P} NMR (BF₄ salt, 600 MHz, acetone-*d*₆, 240 K): δ 1.91, 2.51 (2 × m, 2 × 2H, –P_CCH₂CHHP_T–), 2.57, 2.84 (2 × m, 2 × 2H, –P_CCH₂CH₂P_T–), 2.24 (m, 2H, –P_CCH₂CH₂P_U–), 2.05 (m, 2H, –P_CCH₂CH₂P_U–), 0.87, 1.26 [2 × s, 2 × 6H, 2 × P_T(CH₃)], 1.81 [s, 6H, P_U(CH₃)₂], 7.56 (s, 1H, C=CH), 7.84 (d, 2H, CCPh_{ortho}), 7.69 (d, 2H, CCPh_{meta}), 7.61 (t, 1H, CCPh_{para}), 7.99 (d, 2H, C=CHPh_{ortho}), 7.56 (d, 2H, C=CHPh_{meta}), 7.37 (t, 1H, C=CHPh_{para}).

¹³C-{¹H} NMR (chloride salt, 101 MHz, methanol-*d*₄, 300 K) δ 10.0 [t, –P_T(CH₃), ¹J_{P(T)-C}} = 11.4], 19.0 [t, –P_T(CH₃), ¹J_{P(T)-C}} = 7.6], 21.2 [d, –P_U(CH₃)₂, ¹J_{P(U)-C}} = 22.9], 26.3 (dt, –P_TCH₂CH₂P_C–, ¹J_{P(C)-C}} = 21.6, ²J_{P(T)-C}} = 7.6), 32.9 (dt, –P_TCH₂CH₂P_C–, ¹J_{P(T)-C}} = 16.5, ²J_{P(C)-C}} = 13.4), 29.1 (dd, –P_UCH₂CH₂P_C–, ¹J_{P(C)-C}} = 24.2, ²J_{P(U)-C}} = 15.9), 34.2 (dd, –P_UCH₂CH₂P_C–, ¹J_{P(U)-C}} = 29.2, ²J_{P(C)-C}} = 11.4), 42.6 (m, PhC \equiv C–, *J*_{P-C} < 2.5), 122.1 (dd, PhC \equiv C–, *J*_{P(C)-C}} = 10.1, *J*_{P(U)-C}} = 7.0), 165.4 (dd, Fe–C=CHPh, *J*_{P(C)-C}} = 10.2, *J*_{P(U)-C}} = 10.2, *J*_{P(T)-C}} = 16.5), 134.2 (d, Fe–C=CHPh, *J*_{P-C}} = 1.3), 131.3 (s, –C \equiv CPh_{ortho}), 130.5 (s, –C \equiv CPh_{meta}), 129.6 (s, –C \equiv CPh_{para}), 126.7 (s, –C=CHPh_{ortho}), 130.3 (s, –C=CHPh_{meta}), 128.0 (s, –C=CHPh_{para}), 133.0 (d, Ph_{ipso}, *J*_{P-C}} = 2.5), 139.1 (apparent q, Ph_{ipso}, *J*_{P-C}} = 1.9 Hz).

Minor isomer 1b. ³¹P-{¹H} NMR (BF₄ salt, 162 MHz, acetone-*d*₆, 300 K): δ 165.4 (br, 1P, P_C), 54.1 (dd, 2P, P_T, ²J_{P(C)-P(T)}} = 30.5, ²J_{P(T)-P(U)}} = 39.1 Hz), 78.9 (br, 1P, P_U).

³¹P-{¹H} NMR (BF₄ salt, 162 MHz, acetone-*d*₆, 233 K): δ 164.8 (t, 1P, P_C, ²J_{P(C)-P(T)}} = 30.5, ²J_{P(C)-P(U)}} = 12.9), 55.4 (dd, 2P, P_T, ²J_{P(T)-P(U)}} = 39.1 Hz), 79.3 (t, 1P, P_U).

¹H-{³¹P} NMR (BF₄ salt, 600 MHz, acetone-*d*₆, 303 K): δ 2.00, 2.31 (2 × m, 2 × 2H, –P_CCH₂CHHP_T–), 3.02, 3.12 (2 × m, 2 × 2H, –P_CCH₂CH₂P_T–), 2.44 (m, 2H, –P_CCH₂CH₂P_U–), 2.10 (m, 2H, –P_CCH₂CH₂P_U–), 0.68, 1.14 [2 × s, 2 × 6H, 2 × P_T(CH₃)], 2.01 [s, 6H, P_U(CH₃)₂], 7.84 (s, 1H, C=CH), 8.02 (d, 2H, CCPh_{ortho}), 7.68 (d, 2H, CCPh_{meta}), 7.52 (t, 1H, CCPh_{para}), 8.14 (d, 2H, C=CHPh_{ortho}), 7.63 (d, 2H, C=CHPh_{meta}), 7.42 (t, 1H, C=CHPh_{para}).

¹H-{³¹P} NMR (BF₄ salt, 600 MHz, acetone-*d*₆, 240 K): δ 1.97, 2.29 (2 × m, 2 × 2H, –P_CCH₂CHHP_T–), 3.00, 3.09 (2 × m, 2 × 2H, –P_CCH₂CH₂P_T–), 2.39 (m, 2H, –P_CCH₂CH₂P_U–), 2.05 (m, 2H, –P_CCH₂CH₂P_U–), 0.64, 1.12 [2 × s, 2 × 6H, 2 × P_T(CH₃)], 2.07 [s, 6H, P_U(CH₃)₂], 7.85 (s, 1H, C=CH), 8.02 (d, 2H, CCPh_{ortho}), 7.68 (d, 2H, CCPh_{meta}), 7.52 (t, 1H, CCPh_{para}), 8.14 (d, 2H, C=CHPh_{ortho}), 7.67 (d, 2H, C=CHPh_{meta}), 7.50 (t, 1H, C=CHPh_{para}).

[Fe(η^3 -Bu'C \equiv C–C=CHBu')L]⁺ **2. *tert*-Butylacetylene (10 mg, 120 μmol) was added to a stirred solution of $\text{FeH}(\text{Cl})\text{L}$ **6** (*ca.* 10 mg, 26 μmol) in methanol (5 ml), resulting in a change from yellow to dark orange. On addition of sodium tetraphenylborate (20 mg, 60 μmol) in methanol (5 ml), an**

orange precipitate formed. The crude product was isolated by filtration, washed with methanol (10 ml) and dried *in vacuo* to yield $[\text{Fe}(\eta^3\text{-Bu}^t\text{C}\equiv\text{C}-\text{C}=\text{CHBu}^t)\text{L}]^+$ **2** (BPh₄ salt).

³¹P-¹H} NMR (BPh₄ salt, 162 MHz, acetone-*d*₆, 300 K): δ 174.5 (t, 1P, P_C, ²J_{P(C)-P(T)}} = 25.3), 48.2 (dd, 2P, P_T, ²J_{P(T)-P(U)}} = 35.3), 60.40 (t, 1P, P_U). ³¹P-¹H} NMR spectrum (BPh₄ salt, 162 MHz, acetone-*d*₆, 233 K): δ 174.0 (t, 1P, P_C, ²J_{P(C)-P(T)}} = 24.8), 49.1 (dd, 2P, P_T, ²J_{P(T)-P(U)}} = 35.3 Hz), 61.0 (t, 1P, P_U).

$[\text{Fe}\{\eta^3\text{-HC}\equiv\text{CC}_6\text{H}_4\text{C}\equiv\text{C}-\text{C}=\text{CH}(\text{C}_6\text{H}_4)\text{C}\equiv\text{CH}\}\text{L}]^+$ **3**. 1,4-Diethynylbenzene (10 mg, 80 μmol) was added to a stirred solution of FeH(Cl)L **6** (ca. 10 mg, 26 μmol) in methanol (5 ml), resulting in a change from yellow to red. On addition of sodium tetraphenylborate (20 mg, 60 μmol) in methanol (5 ml) a red precipitate formed. The crude product was isolated by filtration, washed with methanol (10 ml) and dried *in vacuo* to yield $[\text{Fe}\{\eta^3\text{-HC}\equiv\text{CC}_6\text{H}_4\text{C}\equiv\text{C}-\text{C}=\text{CH}(\text{C}_6\text{H}_4)\text{C}\equiv\text{CH}\}\text{L}]^+$ **3** (BPh₄ salt).

Major isomer **3a**. ³¹P-¹H} NMR (BPh₄ salt, 162 MHz, acetone-*d*₆, 300 K): δ 171.0 (t, 1P, P_C, ²J_{P(C)-P(T)}} = 26.7, ²J_{P(C)-P(U)}} = 4.3), 48.9 (dd, 2P, P_T, ²J_{P(T)-P(U)}} = 35.8 Hz), 63.7 (t, 1P, P_U).

³¹P-¹H} NMR (BPh₄ salt, 162 MHz, acetone-*d*₆, 233 K): δ 170.5 (t, 1P, P_C, ²J_{P(C)-P(T)}} = 25.3, ²J_{P(C)-P(U)}} = 4.3), 49.9 (dd, 2P, P_T, ²J_{P(T)-P(U)}} = 36.2 Hz), 64.2 (t, 1P, P_U).

Minor isomer **3b**. ³¹P-¹H} NMR (BPh₄ salt, 162 MHz, acetone-*d*₆, 233 K): δ 163.7 (t, 1P, P_C, ²J_{P(C)-P(T)}} = 30.5, ²J_{P(C)-P(U)}} = 12.8), 55.5 (dd, 2P, P_T, ²J_{P(T)-P(U)}} = 36.2 Hz), 78.7 (t, 1P, P_U).

Acknowledgements

We gratefully acknowledge financial support from the Australian Research Council, the Australian Government for an Australian Postgraduate Award (R. J. S.) and the University of Sydney for an H. B. and F. M. Gritton Award (R. J. S.).

References

- See, for example: (a) A. Dobson, D. S. Moore, S. D. Robinson, M. B. Hursthouse and L. New, *Polyhedron*, 1985, **4**, 1119; (b) Y. Wakatsuki, H. Yamazaki, N. Kumegawa, T. Satoh and J. Y. Satoh, *J. Am. Chem. Soc.*, 1991, **113**, 9604; (c) M. Ishikawa, J. Ohshita, Y. Ito and A. Minato, *J. Organomet. Chem.*, 1988, **346**, C58; (d) M. Schafer, N. Mahr, J. Wolf and H. Werner, *Angew. Chem., Int. Ed. Engl.*, 1993, **32**, 1315; (e) A. M. Echavarren, J. Lopez, A. Santos and J. Montoya, *J. Organomet. Chem.*, 1991, **414**, 393; (f) I. P. Kovalev, K. V. Yevdakov, Y. A. Strelenko, M. G. Vinogradov and G. I. Nikishin, *J. Organomet. Chem.*, 1990, **386**, 139; (g) J. Ohshita, K. Furumori, A. Matsuguchi and M. Ishikawa, *J. Org. Chem.*, 1990, **55**, 3277; (h) C. Slugovc, K. Mereiter, E. Zobetz, R. Schmid and K. Kirchner, *Organometallics*, 1996, **15**, 5277.
- (a) C. Bianchini, M. Peruzzini, F. Zanobini, P. Frediani and A. Albinati, *J. Am. Chem. Soc.*, 1991, **113**, 5453; (b) C. Bianchini, C. Bohanna, M. A. Esteruelas, P. Frediani, A. Meli, L. A. Oro and M. Peruzzini, *Organometallics*, 1992, **11**, 3837; (c) P. Barbaro, C. Bianchini, M. Peruzzini, A. Polo, F. Zanobini and P. Frediani,

- Inorg. Chim. Acta*, 1994, **220**, 5; (d) C. Bianchini, P. Frediani, D. Masi, M. Peruzzini and F. Zanobini, *Organometallics*, 1994, **13**, 4616; (e) C. Bianchini, *Pure Appl. Chem.*, 1991, **63**, 829.
- L. Dahlenburg, K.-M. Frosin, S. Kerstan and D. Werner, *J. Organomet. Chem.*, 1991, **407**, 115.
- L. D. Field, A. V. George, E. Y. Malouf, I. H. M. Slip and T. W. Hambley, *Organometallics*, 1991, **10**, 3842.
- (a) L. D. Field, A. V. George, G. R. Purches and I. H. M. Slip, *Organometallics*, 1992, **11**, 3019; (b) A. Hills, D. L. Hughes, M. Jimenez-Tenorio, G. J. Leigh, C. A. McGeary, A. T. Rowley, M. Bravo, C. E. McKenna and M.-C. McKenna, *J. Chem. Soc., Chem. Commun.*, 1991, 522; (c) D. L. Hughes, M. Jimenez-Tenorio, G. J. Leigh and A. T. Rowley, *J. Chem. Soc., Dalton Trans.*, 1993, 3151.
- L. D. Field, A. V. George and T. W. Hambley, *Inorg. Chem.*, 1990, **29**, 4565.
- (a) C. Bianchini, A. Mealli, M. Peruzzini, F. Vizza and F. Zanobini, *Organometallics*, 1989, **8**, 2080; (b) C. Bianchini, A. Meli, M. Peruzzini, P. Frediani, C. Bohanna, M. A. Esteruelas and L. A. Oro, *Organometallics*, 1992, **11**, 138.
- N. Bampos, L. D. Field, B. A. Messerle and R. J. Smernik, *Inorg. Chem.*, 1993, **32**, 4084.
- (a) L. D. Field, B. A. Messerle, R. J. Smernik, T. W. Hambley and P. Turner, *Inorg. Chem.*, 1997, **36**, 2884; (b) L. D. Field, B. A. Messerle and R. J. Smernik, *Inorg. Chem.*, 1997, **36**, 5984; (c) R. J. Smernik, PhD Thesis, 1996, The University of Sydney.
- teXsan: Crystal Structure Analysis Package, Molecular Structure Corporation, The Woodlands, Texas, 1985 and 1992.
- SHELX86: G. M. Sheldrick, in *Crystallographic Computing 3*, eds. G. M. Sheldrick, C. Kruger and R. Goddard, Oxford University Press, 1985, pp. 175–188.
- DIRDIF94: P. T. Beurskens, G. Admiraal, G. Beurskens, W. P. Bosman, R. de Gelder, R. Israel and J. M. M. Smits, The DIRDIF-94 program system, Technical Report of the Crystallography Laboratory, University of Nijmegen, 1994.
- D. T. Cromer and J. T. Waber, *International Tables for X-ray Crystallography*, The Kynoch Press, Birmingham, 1974, vol. 4, Table 2.2A.
- J. A. Ibers and W. C. Hamilton, *Acta Crystallogr.*, 1964, **17**, 781.
- D. C. Creagh and W. J. McAuley, *International Tables for Crystallography*, ed. A. J. C. Wilson, Kluwer Academic Publishers, Boston, 1992, vol. C, Table 4.2.6.8, pp. 219–222.
- D. C. Creagh and J. H. Hubbell, *International Tables for Crystallography*, ed. A. J. C. Wilson, Kluwer Academic Publishers, Boston, 1992, vol. C, Table 4.2.4.3, pp. 200–206.
- C. K. Johnson, ORTEP, A Thermal Ellipsoid Plotting Program, Report ORNL-5138, Oak Ridge National Laboratories, Oak Ridge, TN, 1976.
- G. E. Herberich and W. Barlage, *Organometallics*, 1987, **6**, 1924.
- A. K. McMullen, J. P. Selegue and J.-G. Wang, *Organometallics*, 1991, **10**, 3421.
- N. W. Alcock, A. F. Hill, R. P. Melling and A. R. Thompsett, *Organometallics*, 1993, **12**, 641; G. Jia, J. C. Gallucci, A. L. Rheingold, B. S. Haggerty and D. W. Meek, *Organometallics*, 1991, **10**, 3459; G. Albertin, P. Amendola, S. Antoniutti, S. Ianelli, G. Pelizzi and E. Bordignon, *Organometallics*, 1991, **10**, 2876; C. Bianchini, P. Innocenti, M. Peruzzini, A. Romerosa and F. Zanobini, *Organometallics*, 1996, **15**, 272; C. Bianchini, D. Masi, M. Peruzzini, A. Romerosa and F. Zanobini, *Acta Crystallogr., Sect. C*, 1996, **52**, 1973.
- J. Gotzig, H. Otto and H. Werner, *J. Organomet. Chem.*, 1985, **287**, 247.

Paper 9/02784J