

Unusual pathways for the reaction between $[\text{MCl}_2(\text{Me}_2\text{SO})_4]$ ($\text{M} = \text{Os}, \text{Ru}$) and hydrazine dihydrochloride: deoxygenation of sulfoxides vs. coordination of hydrazinium

M. Fátima C. Guedes da Silva,^{a,b} Armando J. L. Pombeiro,^{a*} Silvano Geremia,^c Ennio Zangrando,^c Mario Calligaris,^{*c} Andrey V. Zinchenko^d and Vadim Yu. Kukushkin^{*e}

^a Centro de Química Estrutural, Complexo I, Instituto Superior Técnico, Av. Rovisco Pais, 1049-001 Lisbon, Portugal. E-mail: pombeiro@alfa.ist.utl.pt

^b Universidade Lusófona de Humanidades e Tecnologias, Av. Campo Grande 376, 1749-024, Lisbon, Portugal

^c Dipartimento di Scienze Chimiche, University of Trieste, via L. Giorgieri 1, 34127 Italy. E-mail: calligaris@univ.trieste.it

^d Inorganic Chemistry, St. Petersburg State Technological Institute, Zagorodny Pr. 49, 198013 St. Petersburg, Russian Federation

^e Department of Chemistry, St. Petersburg State University, 198904 Stary Petergof, Russian Federation. E-mail: kukushkin@VK2100.spb.edu

Received 12th October 1999, Accepted 3rd March 2000

Published on the Web 31st March 2000

Reaction of *trans*- $[\text{OsCl}_2(\text{Me}_2\text{SO})_4]$ **1** with excess $\text{N}_2\text{H}_4 \cdot 2\text{HCl}$ leads to the osmium(III) compound *mer*- $[\text{OsCl}_3(\text{NH}_3)_2(\text{Me}_2\text{S})]$ **2** in the course of concerted reactions, *i.e.* oxidation of the metal, deoxygenation of the sulfoxide, disproportionation of hydrazine and substitution by NH_3 . In contrast, interaction of *cis*- $[\text{RuCl}_2(\text{Me}_2\text{SO})_4]$ *cis*-**3** with hydrazine dihydrochloride brings about easy substitution (without redox conversion) leading to *mer*- $[\text{RuCl}_3(\text{N}_2\text{H}_5)(\text{Me}_2\text{SO})_2]$ **4** which is a rare example of a hydrazinium complex. X-Ray single-crystal diffraction analyses were performed on **1**, **2** and **4**. In **4**, one Ru–S bond is unusually short due to the enhanced π bonding contribution as a result of an intramolecular H-bond between the Me_2SO and the *cis* N_2H_5^+ group. The complexes show anodic oxidations and **4**, that in aqueous medium undergoes spontaneous dehydrochlorination, exhibits by controlled potential electrolysis a multi-electron oxidation process with anodically-induced H^+ loss, oxidation of the hydrazine ligand to N_2 and N-oxides, and of Me_2SO to SO_2 . The anodic waves of Me_2SO or H_2O solutions of *cis*-**3** were assigned to *trans*- $[\text{RuCl}_2(\text{Me}_2\text{SO}-\text{S})_4]$ *trans*-**3** and $[\text{RuCl}(\text{H}_2\text{O})_2(\text{Me}_2\text{SO}-\text{S})_3]^+$, respectively. The oxidation potential values were interpreted on the basis of redox potential–structure relationships and the Lever electrochemical parameter E_L was tentatively estimated for both S- and O-coordinated Me_2SO and for the hydrazinium ligand, showing that $\text{Me}_2\text{SO}-\text{S}$ in our complexes behaves as a significant π -electron acceptor and N_2H_5^+ as a rather weak net electron donor, and applied to predict the oxidation potential of some Ru and Os complexes.

Introduction


Dialkyl sulfoxides in general and dimethyl sulfoxide in particular are among the most studied solvents and reagents in organic chemistry.^{1–3} Despite the fact that their coordination chemistry started only in the early sixties⁴ and has a much poorer history, various reactions of sulfoxides with a variety of metal ions and complexes,⁵ including the versatile coordination modes of R_2SO ,^{6–8} the application of sulfoxide metal compounds as useful synthons in preparative coordination chemistry,⁹ their homogeneous catalytic properties¹⁰ and the potential of sulfoxide complexes in medicine,^{11,12} have been extensively investigated.

One of the most fundamental chemical properties of sulfoxides is their ability to undergo deoxygenation of the sulfinyl group to give the corresponding sulfides, a reaction that has been studied in depth in organic chemistry,^{1–3} while the deoxygenation of sulfoxide ligands has received much less attention and only one review on the subject has been written.¹³ Besides the data analysed in the article,¹³ new interesting examples of the deoxygenation have been reported in the last five years.^{14–23}

Inspection of the massive amount of structural data⁶ as well as theoretical calculations²⁴ indicate that the S=O bonds

are stronger in S-bound R_2SO ligands than those in free and O-coordinated sulfoxides and this indirectly implies that it is more difficult to abstract oxygen from S-bound sulfoxides than from both uncoordinated and O-bound ones. Indeed, data on the reduction of S-coordinated sulfoxides (in contrast to O-coordinated ligands) are rather scarce and restricted mainly to complexes of the platinum group metals.¹³ Most of the methods known so far for the reduction of the very strong S=O bond in S-coordinated R_2SO ligands are based upon an initial activation of the sulfinyl oxygen by a reagent having a highly reactive electrophilic centre. The activator, E^+ , binds to the oxygen atom, increases the partial positive charge on the sulfur atom and thus facilitates the reduction, Scheme 1.

The combination of an electrophilic activator and a reducing agent such as $\text{HX}-\text{Pt}^{\text{II}}$ ($\text{X} = \text{Cl}, \text{Br}$),²⁵ $\text{SOCl}_2-\text{Pt}^{\text{II}}$,²⁶ $\text{HCl}-\text{N}_2\text{H}_4$,¹⁹ $\text{PCl}_5-\text{Pr}^{\text{I}}\text{OH}$ ²⁷ proved to be useful in deoxygenations of S-coordinated sulfoxides. However, all these processes were carried out for Pt complexes only, while deoxygenation of other metal compounds that contain S-bound R_2SO ligands is still a very little known area. Thus, only reductive thermolysis studies²⁸ and deoxygenations involving HCl ,^{29,30} CO^{21} and PCl_3 ,^{22,23} were reported for both Os and Ru S-bound complexes.



Scheme 1

In view of our interest in the chemistry of metal sulfoxide complexes,^{6,7,9,11,13} we focused our efforts on deoxygenation of sulfoxide ligands in the Os^{II} and Ru^{II} compounds [MCl₂(Me₂SO)₄]. For this study we selected N₂H₄·2HCl as a deoxygenating reagent which was known to be efficient for the reduction of dimethyl sulfoxide ligands in [PtCl_n(Me₂SO)₂] (*n* = 2 or 4) to afford [PtCl₂(Me₂S)₂].¹⁹ In the case of the Os complex, we observed metal oxidation with concomitant reduction of Me₂SO to Me₂S and N₂H₄ to NH₃, while for the Ru^{II} compound the reaction proceeds in the unexpected direction giving a rare hydrazinium complex *mer*-[RuCl₃(N₂H₅)(Me₂SO-*S*)₂] without oxidation of the metal. In view of the different redox/substitution behaviours exhibited in these reactions by the Os^{II} and Ru^{II} sites, an electrochemical study was also performed, (i) indicating a higher lability of the latter towards ligand displacement but its higher stability towards oxidation, (ii) providing a distinction between the S- and O-coordination modes of Me₂SO and (iii) allowing us to tentatively estimate the corresponding values of the Lever *E_L* ligand parameter³¹ which was used for assignment of the binding mode of Me₂SO and the identification of new products. All these results along with structural data are reported below.

Experimental

Materials and instrumentation (chemical studies)

Hydrazine dihydrochloride (Reakhim) was used as received. *cis*-[RuCl₂(Me₂SO)₄] was prepared in accord with the published method.³² All other chemicals and solvents were obtained from commercial sources and were used as received. Decomposition points were determined on a Kofler table. For TLC, Silufol UV254 SiO₂-plates were used. Elemental analyses were carried out by the Microanalytical Service, St. Petersburg State Technological Institute. Positive-ion FAB mass spectra were obtained on a Trio 2000 instrument by bombarding 3-nitrobenzyl alcohol matrices of samples with 8 keV (*ca.* 1.28 × 10⁻¹⁵ J) Xe atoms. Mass calibration for data system acquisition was achieved using CsI. Infrared spectra (4000–220 cm⁻¹) were recorded on a Perkin-Elmer 983G spectrometer in KBr pellets. ¹H and ¹³C{¹H} NMR spectra (*δ* values relative to SiMe₄ as an internal reference) were measured on a Varian UNITY 300 spectrometer at ambient temperature.

Synthetic work and characterisation

***trans*-[OsCl₂(Me₂SO)₄] 1.** This was obtained by a slightly modified version of a literature method³³ using K₂[OsCl₆] instead of H₂[OsCl₆]. Found: C, 16.7; H, 4.0; Cl, 12.5; Os, 33.3. C₈H₂₄Cl₂O₄OsS₄ requires C, 16.8; H, 4.2; Cl, 12.4; Os, 33.2%. FAB⁺-MS, *m/z*: 574 [M]⁺. IR spectrum, cm⁻¹: 3041 m, 3014 m and 2929 w *ν*(C–H); 1081 vs and 1027 s *ν*(S=O). ¹H NMR spectrum in CDCl₃, *δ*: 3.47. ¹³C{¹H} NMR spectrum in CDCl₃, *δ*: 42.0. Crystals suitable for X-ray analysis were grown on slow evaporation of a chloroform solution.

***mer*-[OsCl₃(NH₃)₂(Me₂S)] 2.** A mixture of *trans*-[OsCl₂(Me₂SO)₄] **1** (0.20 g, 0.35 mmol) and N₂H₄·2HCl (0.40 g, 3.81

mmol) in water (2 ml) was heated at 90–95 °C for 5 h, cooled to room temperature and left to stand for 1 d. A crystalline precipitate was filtered off, washed on a filter with water (10 ml), ethanol (5 ml) and chloroform (5 ml) and dried in air at 20–25 °C. Yield of *mer*-[OsCl₃(NH₃)₂(Me₂S)] was 0.06 g, 43% based on Os. Found: C, 6.4; N, 6.8; Os, 48.7. C₂H₁₂N₂Cl₃OsS requires C, 6.1; N, 7.1; Os, 48.4%. FAB⁺-MS, *m/z*: 393 [M]⁺. This complex has no characteristic melting point, it decomposes above 225 °C. IR spectrum, cm⁻¹: 3291 m-s, 3233 m-s and 3180 m-s *ν*(N–H); 2998 w and 2929 w *ν*(C–H); 1618 m, br *δ*_{as}(HNH); 1333 vs and 1309 vs *δ*_s(HNH); 1028 m and 967 m *ρ*(SCH₃); 788 m, br *ρ*(NH₃). Solubility of the compound in the most common deuterated solvents is insufficient to measure NMR spectra even at high acquisition time. In dimethylformamide-*d*₇, where the complex exhibits a rather moderate solubility, signals due to the methyl groups overlap with solvent peaks and cannot be reliably attributed. Crystals suitable for X-ray analysis were grown on slow evaporation of DMF solution at room temperature.

***mer*-[RuCl₃(N₂H₅)(Me₂SO)₂] 4.** *cis*-[RuCl₂(Me₂SO)₄] *cis*-**3** (0.20 g, 0.41 mmol) and N₂H₄·2HCl (0.45 g, 4.29 mmol) in water (2 ml) were heated at 90–95 °C for 30 min, cooled to room temperature and left to stand for 1 h. A crystalline precipitate was filtered off, washed on a filter with water (1–2 ml), ethanol (5 ml) and chloroform (5 ml) and dried in air at 20–25 °C. Yield of *mer*-[RuCl₃(N₂H₅)(Me₂SO)₂] was 0.07 g, 50% based on Ru. Concentration of the filtrate on evaporation of the solvent at room temperature to *ca.* 1 ml led to the precipitation of a further crop (*ca.* 20–30%) of less pure complex. Found: C, 12.3; N, 7.2; Ru, 25.3. C₄H₁₇N₂Cl₃O₂RuS₂ requires C, 12.1; N, 7.1; Ru, 25.5%. FAB⁺-MS, *m/z*: 397 [M – H]⁺. This complex decomposes above 215 °C. IR spectrum, cm⁻¹: intense absorption in the range 3200–2900 cm⁻¹ due to C–H and N⁺–H stretches; 1084 and 1044 vs *ν*(S=O); the band at 1011 (vs), in accord with the data reported by Gajapathy *et al.*³⁴ can be attributed to the *ν*(N–N) stretch in coordinated hydrazinium ligand. ¹H NMR spectrum in D₂O/HCl, *δ*: 3.13 and 3.18 (Me). Crystals suitable for X-ray analysis were obtained directly from the reaction mixture.

Heating of *cis*-[RuCl₂(Me₂SO)₄] *cis*-**3** (0.20 g, 0.41 mmol) and N₂H₄·2HCl (0.45 g, 4.29 mmol) in water (2 ml) at 90–95 °C for 5 h (the reaction time for the osmium complex) resulted in the formation of [RuCl₂(NH₃)₄]Cl (35% yield) among other unidentified sulfur- and carbon-free products. Found: H, 5.0; N, 19.8; Ru, 36.4. H₁₂N₄Cl₃Ru requires H, 4.4; N, 20.3; Ru, 36.7%. FAB⁺-MS, *m/z*: 240 [M]⁺. IR spectrum, cm⁻¹: 3295 m-s, 3237 m-s and 3171 m-s *ν*(N–H); 1611 m, br *δ*_{as}(HNH); 1285 vs *δ*_s(HNH); 798 m, br *ρ*(NH₃).

Electrochemical studies

The electrochemical experiments were performed on an EG & G PARC 273 potentiostat/galvanostat connected to a PC computer through a GPIB interface (National Instruments PC-2A) or on an EG & G PAR 173 potentiostat/galvanostat and an EG & G PARC 175 Universal programmer. Cyclic voltammetry (CV), chronoamperometry (CA) and steady-state voltammetry (SSV) were undertaken in a two-compartment three-electrode cell, at platinum-disc working electrodes with a diameter of 0.5 mm (CV and CA) or 0.25 μm (SSV), probed by a Luggin capillary connected to a silver-wire pseudo-reference electrode; a platinum auxiliary electrode was employed. Controlled potential electrolyses (CPE) were carried out in a two-compartment three-electrode cell with platinum gauze working and counter electrodes in compartments separated by a glass frit; a Luggin capillary, probing the working electrode, was connected to a silver-wire pseudo-reference electrode. The electrochemical experiments were performed in a N₂ or argon atmosphere at room temperature. The potentials were measured in 0.2 mol dm⁻³ KCl (or KNO₃)/H₂O or Me₂SO (Ru complexes) or in

Table 1 Crystal data and structure refinement of compounds *mer*-[OsCl₃(NH₃)₂(SMe₂)] **2** and *mer*-[RuCl₃(N₂H₅)(Me₂SO-S)] **4**

	2	4
Formula	C ₂ H ₁₂ N ₂ Cl ₃ OsS	C ₄ H ₁₇ N ₂ Cl ₃ O ₂ RuS ₂
<i>M</i>	392.75	396.74
Crystal system	Monoclinic	Monoclinic
Space group	<i>P</i> ₂ / <i>c</i> (no. 14)	<i>C</i> 2/ <i>c</i> (no. 15)
<i>a</i> /Å	9.5765(10)	26.202(4)
<i>b</i> /Å	8.9186(8)	8.455(1)
<i>c</i> /Å	11.8001(9)	12.918(2)
Volume/Å ³	994.2(2)	2743.3(7)
<i>Z</i>	4	8
μ /mm ⁻¹	13.774	2.012
Reflections collected	3205	4407
Independent reflections	2896	3985
Reflections [<i>I</i> > 2σ(<i>I</i>)]	2594	3781
<i>R</i> (int)	0.0443	0.0349
<i>R</i> 1	0.0291	0.0247
<i>wR</i> 2	0.0771	0.0703

0.2 mol dm⁻³ [NBu₄][BF₄]/Me₂SO, DMF or CH₂Cl₂ (Os complexes) and are quoted relative to the saturated calomel electrode (SCE) by using as internal reference methylviologen (*E*^o = -0.68 V vs. SCE) for the aqueous medium, or [Fe(η⁵-C₅H₄COCl)(η⁵-C₅H₅)] (*E*^o = 0.67 or 0.53 V vs. SCE, in Me₂SO or CH₂Cl₂, respectively). The acid–base potentiometric titrations were carried out by using an aqueous solution of KOH. The results presented were corrected for background effects by performing also the titration of a blank aqueous solution of the electrolyte (0.2 mol dm⁻³ KCl) which had been electrolysed under identical conditions to those used for the corresponding complex solution.

The gaseous atmosphere of the electrochemically oxidised solution of the ruthenium complex and the solution itself were analysed by EI and FAB mass spectrometric measurements, respectively, performed on a Trio 2000 spectrometer. For the EI experiment, the sample was injected *via* an empty glass capillary column (temperatures of the injector, oven, interface and source were, respectively: 120, 50, 120 and 200 °C). Positive-ion FAB mass spectra were obtained by bombarding 3-nitrobenzyl alcohol matrices of the liquid electrolysed solution with 8 keV Xe atoms.

X-Ray crystallography of *trans*-[OsCl₂(Me₂SO-S)] **1**, *mer*-[OsCl₃(NH₃)₂(Me₂S)] **2** and *mer*-[RuCl₃(N₂H₅)(Me₂SO-S)] **4**

Diffraction measurements were carried out at room temperature on an Enraf-Nonius CAD4 diffractometer, equipped with graphite monochromator and Mo-Kα radiation (λ = 0.71069 Å). All the three structures were solved by conventional Patterson³⁵ and Fourier techniques³⁶ and refined by full-matrix anisotropic least-squares on *F*². In the final cycles the contribution of hydrogen atoms was held fixed at calculated positions. The structure of **1** presents the Me₂SO ligands disordered as already reported.³⁷ However, the set of diffraction data collected by us (39% larger) allowed a higher resolution and consequently a better refinement, avoiding the unreliable S–O bond distance of 1.77(1) Å previously found for the minor component (18%).³⁷ Crystal data and refinement parameters of the new structures **2** and **4** are reported in Table 1. Bond lengths and angles are given in Tables 2 and 3.

CCDC reference number 186/1886.

See <http://www.rsc.org/suppdata/dt/a9/a908170d/> for crystallographic files in .cif format.

Results and discussion

Deoxygenation of Me₂SO in the osmium complex

The starting material for the deoxygenation study, *trans*-[OsCl₂(Me₂SO-S)] **1**, was prepared by reacting K₂[OsCl₄] with

Table 2 Bond lengths (Å) and angles (°) for *mer*-[OsCl₃(NH₃)₂(SMe₂)] **2**

Os–Cl(1)	2.363(1)	Os–N(1)	2.131(4)
Os–Cl(2)	2.358(1)	Os–N(2)	2.099(4)
Os–Cl(3)	2.384(1)	S(1)–C(1)	1.802(7)
Os–S(1)	2.345(1)	S(1)–C(2)	1.791(6)
Cl(1)–Os–Cl(2)	174.26(5)	Cl(3)–Os–N(2)	174.1(2)
Cl(1)–Os–Cl(3)	91.66(5)	Cl(3)–Os–N(1)	88.1(1)
Cl(1)–Os–S(1)	86.63(5)	Cl(3)–Os–S(1)	89.65(5)
Cl(1)–Os–N(1)	90.5(1)	S(1)–Os–N(1)	176.3(1)
Cl(1)–Os–N(2)	88.5(2)	S(1)–Os–N(2)	96.3(2)
Cl(2)–Os–Cl(3)	94.08(5)	N(1)–Os–N(2)	86.0(2)
Cl(2)–Os–S(1)	93.54(4)	Os–S(1)–C(1)	110.8(3)
Cl(2)–Os–N(1)	89.5(1)	Os–S(1)–C(2)	110.5(2)
Cl(2)–Os–N(2)	85.8(1)	C(1)–S(1)–C(2)	99.7(4)

Table 3 Bond lengths (Å) and angles (°) for *mer*-[RuCl₃(N₂H₅)(Me₂SO-S)] **4**

Ru–Cl(1)	2.4377(5)	S(1)–C(1)	1.776(2)
Ru–Cl(2)	2.4139(6)	S(1)–C(2)	1.771(2)
Ru–Cl(3)	2.3970(6)	S(2)–O(2)	1.494(2)
Ru–S(1)	2.2420(5)	S(2)–C(3)	1.781(2)
Ru–S(2)	2.2187(5)	S(2)–C(4)	1.782(2)
Ru–N(1)	2.157(2)	N(1)–N(2)	1.443(2)
S(1)–O(1)	1.483(2)		
Cl(1)–Ru–Cl(2)	89.71(2)	S(2)–Ru–N(1)	92.02(5)
Cl(1)–Ru–Cl(3)	88.57(2)	Ru–S(1)–O(1)	119.27(7)
Cl(2)–Ru–Cl(3)	174.13(2)	Ru–S(1)–C(1)	112.02(8)
Cl(1)–Ru–S(1)	89.68(2)	Ru–S(1)–C(2)	112.33(9)
Cl(1)–Ru–S(2)	175.60(2)	Ru–S(2)–O(2)	115.02(7)
Cl(2)–Ru–S(1)	92.31(2)	Ru–S(2)–C(3)	115.12(8)
Cl(2)–Ru–S(2)	91.03(2)	Ru–S(2)–C(4)	114.94(8)
Cl(3)–Ru–S(1)	93.29(2)	Ru–N(1)–N(2)	119.4(1)
Cl(3)–Ru–S(2)	90.27(2)	O(1)–S(1)–C(1)	106.4(1)
Cl(1)–Ru–N(1)	83.71(5)	O(1)–S(1)–C(2)	106.0(1)
Cl(2)–Ru–N(1)	86.31(5)	C(1)–S(1)–C(2)	98.8(2)
Cl(3)–Ru–N(1)	87.93(5)	O(2)–S(2)–C(3)	104.7(1)
S(1)–Ru–S(2)	94.62(2)	O(2)–S(2)–C(4)	105.0(1)
S(1)–Ru–N(1)	173.25(5)	C(3)–S(2)–C(4)	100.5(1)

dimethyl sulfoxide in the presence of SnCl₂·2H₂O. On treatment of the dimethyl sulfoxide osmium(II) complex **1** with excess N₂H₄·2HCl in an aqueous solution at 90–95 °C for 5 h the osmium(III) complex *mer*-[OsCl₃(NH₃)₂(Me₂S)] **2** was formed and isolated as a solid in ca. 40% yield. IR monitoring of the residue (after isolation of **2** and evaporation of the solvent to dryness) shows no characteristic bands due to ν(N–N) stretch in the coordinated hydrazinium ligand.³⁴ The dimethyl sulfide complex **2** was characterised by elemental analyses, FAB-MS, IR and ¹H NMR spectroscopies (see Experimental section) and also by X-ray single-crystal diffraction analysis (see below).


Deoxygenations of osmium(II) sulfoxide complexes are known although scarce. Thus, it has been reported²⁸ that heating of the isomeric osmium(II) compounds [OsCl₂(Me₂SO-S)]₄ and [OsCl₂(Me₂SO-O)(Me₂SO-S)] at 150 °C *in vacuo* causes the disproportionation of dimethyl sulfoxide to dimethyl sulfone and the osmium(III) thioether complexes [Os₂Cl₄(μ-SMe)₂-(Me₂S)] and [Os₂Cl₄(Me₂SO-O)(μ-SMe)₂(Me₂S)], respectively. Chakravorty and colleagues^{16,17} have deoxygenated both dimethyl and dibenzyl sulfoxides on treatment with [OsBr₆]²⁻ in refluxing 2-methoxyethanol to give the osmium(III) complexes [OsBr₃(R₂S)] (R = Me, CH₂Ph) that contain coordinated thioethers. Eventually, one of us performed deoxygenation of *trans*-[OsCl₂(Me₂SO-S)] with HCl to yield [OsCl₄(Me₂S)]₂.²³

Currently, there is only one reported example indicating that hydrazine hydrochloride can be employed for reduction of both sulfoxide species and also the platinum(IV) centre in [PtCl₄-(Me₂SO)]₂ to give [PtCl₂(Me₂S)]₂,¹⁹ no ammonia containing products were detected in this reaction. Although disproportionation of free hydrazine to NH₃ and N₂ is well-known,³⁸ and

Table 4 M–S–C and C–S–C bond angles (°) in Ru and Os thioether complexes

Complex	M–S–C	C–S–C	Ref.
[RuBr ₃ (NO)(<i>n</i> -Pr ₂ S) ₂]	109.3(3), 109.8(3), 112.3(3), 106.9(3)	99.4(4), 100.8(5)	77
[RuBr ₃ (NO)(Et ₂ S) ₂ (Et ₂ SO)]	102.3(7)	99.3(9)	77
[Ru(OEP)(MeSDe) ₂] ^a	107.5(1), 111.2(1), 109.0(1), 108.1(1)	98.4(2), 101.0(2)	47
[Ru(OEP)(Ph ₂ S) ₂]	115.4(2), 110.9(2)	103.3(2)	47
[Ru(NH ₃) ₅ (MeSEt)] ³⁺	112.17(8), 107.86(8)	100.7(1)	48(a)
[Ru(NH ₃) ₅ (MeSEt)] ²⁺	113.4(2), 112.5(2)	99.5(3)	48(b)
[Ru(NH ₃) ₅ (Me ₂ S)] ³⁺	113.8(3) ^b	99.5(3)	48(a)
[OsCl ₃ (NH ₃) ₂ (Me ₂ S)]	110.5(2), 110.8(3)	99.7(4)	This work
[Ru(Cp)(chir)(MeSPh)] ^{+c}	110.2(4), 115.3(5)	99.2(5)	78
[Ru(Cp)(chir)(<i>i</i> -PrSMe)] ^{+c}	121.2(4), 106.1(3)	101.3(4)	78
	110.8(9) ^d	100.2(4) ^d	

^a De = *n*-decyl. ^b Possibly overestimated because of twinning. ^c chir = 2(*S*),3(*S*)-bis(diphenylphosphino)butane. ^d Unweighted average values with standard deviations in parentheses.

**Fig. 1** Perspective view of compound *mer*-[OsCl₃(NH₃)₂(Me₂S)] **2**.

documented even for osmium-mediated processes³⁹ (coordinated hydrazine is proved to be resistant towards reduction⁴⁰), formation of *mer*-[OsCl₃(NH₃)₂(Me₂S)] **2** from *trans*-[OsCl₂(Me₂SO-S)₄] **1** in concerted deoxygenation/oxidation/substitution processes is the first observation of this kind. We believe that deoxygenation proceeds *via* formation of electrophilically activated dimethyl sulfoxide^{3,13} (Scheme 1) and its reduction by hydrazine and/or by the osmium(II) centre. Taking into account that it is the high *trans*-effect of S-bound sulfoxides that labilises coordinated Me₂SO,⁴¹ it is perhaps reasonable to assume that the reaction might proceed dissociatively to give free Me₂SO species before deoxygenation. However, a possibility of intramolecular reaction (Scheme 1) should not be ruled out.

Molecular and crystal structures of *trans*-[OsCl₂(Me₂SO-S)₄] **1** and *mer*-[OsCl₃(NH₃)₂(Me₂S)] **2**

The structures of the osmium complexes appear of interest because of the paucity of available crystallographic data of both sulfoxide and sulfide derivatives which can provide some insight into the nature of the Os–S bonds and hence into the metal redox properties.

To the best of our knowledge, the only known crystal structures of S-bonded sulfoxide osmium complexes concern [(η⁶-*p*-cym)OsCl₂(Me₂SO-S)] (*p*-cym = *p*-cymene),⁴² *trans,cis*, *cis*-[Os(CN)₂(Me₂SO-S)₂L] (L = diphenylphenanthroline),⁴³ *trans*-[OsBr₂(Me₂SO-S)₄],⁴⁴ and *cis*- and *trans*-[OsCl₂(Me₂SO-S)₄].³⁷ The structure of the last complex, **1**, isostructural with the Br derivative,⁴⁴ and its ruthenium analogue,⁴⁵ have been already reported,³⁷ but because of static disorder, some bond

lengths and angles around the sulfur atoms were affected by large errors. More accurate and reliable data have been obtained now by us. The results are reported in the deposited .cif file. The average Os–S and S–O bond distances are 2.340 and 1.486 Å, respectively, close to the mean values of 2.34(1) and 1.48(1) Å found in the above Os(II)–Me₂SO-S complexes.⁶

The molecular structure of *mer,cis*-[OsCl₃(NH₃)₂(Me₂S)] **2** is shown in Fig. 1. Interestingly, the Os^{III}–S distance of 2.345(1) Å is shorter than those found in other Os^{III} thioether complexes, like *trans*-[Os(OEP)(PMS)₂][PF₆] (PMS = pentamethylene sulfide), 2.382(2) Å,⁴⁶ and *mer*-[OsBr₃{S(CH₂Ph)₂}₃], 2.387(5) Å (*trans* to Br) and 2.398(3) Å (*trans* to S).¹⁶ It is comparable, within experimental error, to the Os^{II}–S distance of 2.352(2) Å in *trans*-[Os(OEP)(PMS)₂].⁴⁶ The trend of the Os–S distances confirms the π accepting ability of the thioether ligands.^{46–48} However, thioethers display a lower π accepting ability with respect to sulfoxides, as evidenced by the comparison of the Ru–S distances in complexes containing both ligands, *e.g.* Ru–S(sulfide), 2.359(4) Å and Ru–S(sulfoxide), 2.298(4) Å, in *trans*-[RuCl₂L₂] (L = 1,5-dithiacyclooctane 1-oxide);⁴⁹ Ru–S(sulfide), 2.38(1) Å and Ru–S(sulfoxide), 2.26(3) Å, in *cis*-[RuBr₂(Me₂SO-S){EtS(CH₂)₃S(O)}₂].⁵⁰ This is consistent with the higher oxidation potential of **1** with respect to **2** (see Table 5).

As already observed, the coordinated thioether sulfur atom displays a pyramidal geometry with an average C–S–C angle of 100.2(4)° (Table 4). The orientation of the SC₂ group in (NH₃)₅Ru–sulfide complexes has been discussed in terms of bonding requirements of the thioether ligand and its steric repulsions with the *cis* amines which provoke a tilting of the SC₂ plane with respect to the RuN₃ plane.⁴⁸ This was evaluated by measuring the Ru–S–X angle, X being the midpoint between the two C atoms, which was found to be 130.1°, 132.0° in Ru^{II} complexes and 122.1°, 125.5° in Ru^{III} complexes.⁴⁸ In the present Os^{III} complex **2**, the Os–S–X angle is 123.1°, while the Os–S–C angles are 110.5(2)° and 110.8(3)°. These data suggest a tetrahedral hybridization of the sulfur atom, for which a M–S–X angle of 125.3° is expected. In **2**, the orientation of the Me₂S group with respect to the equatorial ligands is determined by van der Waals interactions between the CH₃ and NH₃ groups which lower the molecular potential energy, in spite of a slight repulsion between the two groups, as shown by the widening of the S(1)–Os–N(2) and S(1)–Os–Cl(2) angles with respect to S(1)–Os–Cl(1) and S(1)–Os–Cl(3). In fact, density functional calculations have shown that also in Ru–Me₂SO complexes steric interactions are prevalent over electronic factors.²⁴

Substitution of Me₂SO in the ruthenium complex and crystal structure of *mer*-[RuCl₃(N₂H₅)(Me₂SO-S)] **4**

In contrast to the reaction of *trans*-[OsCl₂(Me₂SO-S)₄] **1**, interaction of *cis*-[RuCl₂(Me₂SO-O)(Me₂SO-S)] *cis*-**3** with excess hydrazine dihydrochloride under the same reaction conditions




Fig. 2 Perspective view of compound *mer*-[RuCl₃(N₂H₅)(Me₂SO-S)₂] **4**, the dotted line indicating the intramolecular H-bond.

(but shorter reaction time) brings about fast substitution. The reaction is completed after 30 min and subsequent work up led to isolation of the hydrazinium Ru^{II} complex *mer*-[RuCl₃(N₂H₅)(Me₂SO-S)₂] **4** as a solid in good yield. Heating the Ru reaction mixture for 5 h gave the amine complex [RuCl₂(NH₃)₄]Cl along with other unidentified products in which we were unable to detect (by IR and NMR methods) either dimethyl sulfoxide or dimethyl sulfide products.

Complex **4** represents a rare example of a η^1 metal hydrazinium complex. In general, the chemistry of transition metal compounds containing hydrazine derivatives and other partially reduced dinitrogen ligands is of particular importance due to the fact that such species are relevant to intermediates in the nitrogen fixation process and that such complexes also display diverse and intriguing reactivity modes.⁵¹ The structure of the hydrazinium complex **4** was determined by X-ray diffraction.

As shown in Fig. 2, its molecular structure is characterised by the unusual η^1 metal coordination of the hydrazinium ion, *trans* to one Me₂SO-S ligand. This arrangement corresponds to the expected minimum molecular energy structure, in view of the presence of Cl and N donor ligands *trans* to the π accepting sulfoxide groups.⁶ The structure of this stereoisomer is further stabilised by a strong two-centre hydrogen bond between the terminal NH₃⁺ group and the oxygen atom of the *cis* Me₂SO-S ligand [N...O, 2.665(2) Å]. In fact, the Ru-S(1) bond distance of 2.2420(5) Å is within the range of Ru^{II}-S bond lengths (2.237(1)–2.2576(7) Å) found in *cis,cis,cis*-[RuCl₂L₂(Me₂SO-S)₂] complexes, with L = N-aromatic base, where a stabilising π back-donation is present.⁵² The Ru-S(2) distance of 2.2187(5) Å is significantly shorter, suggesting an enhanced π back-donation from Ru^{II} to the S(2)-sulfoxide empty orbitals. This could be due to the intramolecular N-H...O H-bonding which, reducing the bond order over the S(2)-O(2) bond, increases the positive charge on the sulfur atom enhancing the sulfoxide π accepting properties. However, a similar lengthening, due to H-bonding, has been observed in other Ru^{II}-Me₂SO-S complexes,⁵³ as well as in uncoordinated sulfoxides.⁶ Besides the N...O H-bond, all the other hydrogen atoms of the hydrazinium ion are involved in intermolecular H-bonds with the chloride ligands, with N...Cl distances ranging from 3.165 to 3.274 Å.

The N-N bond distance of 1.443(2) Å is comparable, within experimental error, to the value of 1.427(6) Å found in [Pd-(polyether)(N₂H₅)₂]²⁺, which represents the only other known crystal structure of a η^1 hydrazinium metal complex.⁵⁴ It is also close to the values found in the uncoordinated N₂H₅⁺ cation, e.g. 1.443(1),⁵⁵ 1.444(2),⁵⁶ and 1.450(3) Å,⁵⁷ and within




Fig. 3 Cyclic voltammograms of *mer*-[OsCl₃(NH₃)₂(Me₂S)] **2** (2.6 mmol dm⁻³ in Me₂SO with 0.2 mol dm⁻³ [NBu₄][BF₄]) at a platinum disc (*d* = 0.5 mm) working electrode: (a) anodic scan; (b) cathodic scan. Scan rate: 0.2 V s⁻¹.

the range of values observed for metal η^1 - and μ -coordinated hydrazine complexes, 1.439(10)–1.454(2) Å,^{58–60} and 1.43(1)–1.48(1) Å,^{61–64} respectively.

Electrochemical study

Osmium complexes. The cyclic voltammograms of *trans*-[OsCl₂(Me₂SO-S)₄] **1** and *mer*-[OsCl₃(NH₃)₂(Me₂S)] **2** (Fig. 3), in aprotic media, exhibit a single-electron (confirmed by CPE) reversible anodic wave (I) at $E_{1/2}^{\text{ox}} = 1.18$ or 0.48 V vs. SCE, respectively (Table 5). The lower oxidation potential of the latter complex agrees with the stronger net electron-donor character of its chloride, ammonia and dimethyl sulfide ligands compared with dimethyl sulfoxide in the former compound (see below).

Complex **2** also undergoes, by CV, a single-electron (confirmed by CPE) partially reversible cathodic wave (II, Fig. 3b) at $E_{1/2}^{\text{red}} = -1.20$ V. Upon scan reversal, following this wave, two anodic waves (III and IV), with a partial reversible character, are detected at $E_{1/2}^{\text{ox}} = -0.07$ and 0.06 V in Me₂SO (but not observed in DMF) due to the oxidation of products formed in the cathodic process in Me₂SO. CPE at the cathodic wave (II) of the starting Os^{III} complex in Me₂SO also generates the same species as shown by monitoring the electrolysis by CV (Fig. 4). Wave IV is due to an intermediate species conceivably the pentacoordinate [OsCl₂(NH₃)₂(Me₂S)],^{65–67} generated in the cathodic process, that converts into the final product oxidised at wave III (the latter wave is the only one detected upon standing the fully electrolysed solution, Fig. 4d). Attempts to isolate, in a pure form, this final product have failed, but we propose its formulation as [OsCl₂(NH₃)₂(Me₂S)(Me₂SO-S)] (formed by cathodically induced chloride loss with formation of the above intermediate, a type of reaction well documented,^{68,69} followed by addition of the Me₂SO molecule: overall Scheme 2), on the




Fig. 4 Cyclic voltammograms run along the cathodic controlled potential electrolysis (at -1.3 V) of a solution of *mer*-[OsCl₃(NH₃)₂(Me₂S)] **2** (2.6 mmol dm⁻³ in Me₂SO with 0.2 mol dm⁻³ [NBu₄][BF₄]) at a platinum disc ($d = 0.5$ mm) working electrode, after the consumption of (a) *ca.* 0.2 , (b) *ca.* 0.7 and (c) *ca.* 0.9 F mol⁻¹, and (d) upon standing the solution for 4 hours after completion of the electrolysis. Scan rate: 0.2 V s⁻¹.

Table 5 Cyclic voltammetric data^a for Os and Ru complexes

Complex	E_2^{ox} ($E_{p/2}^{\text{ox}}$)	E_2^{red} ($E_{p/2}^{\text{red}}$)
<i>trans</i> -[OsCl ₂ (Me ₂ SO-S) ₄] 1 ^b	1.18	
<i>mer</i> -[OsCl ₃ (NH ₃) ₂ (Me ₂ S)] 2 ^c	0.48 ^d	-1.20 ^e
<i>trans</i> -[RuCl ₂ (Me ₂ SO-S) ₄] <i>trans</i> - 3 ^{c,f}	(1.47) ^g	(-1.93)
<i>mer</i> -[RuCl ₃ (N ₂ H ₅)(Me ₂ SO-S) ₂] 4 ^h	(0.79) ⁱ	


^a Potentials (E_2 for reversible processes, or, in parentheses $E_{p/2}$ for irreversible ones) in V \pm 0.02 vs. SCE, measured at a Pt disc ($d = 0.5$ mm) electrode. ^b In 0.2 mol dm⁻³ [NBu₄][BF₄]/CH₂Cl₂. ^c In 0.2 mol dm⁻³ [NBu₄][BF₄]/Me₂SO. ^d Upon scan reversal following this anodic wave, cathodic waves are observed at $E_p^{\text{red}} = -0.54$ and -0.73 V. ^e Upon scan reversal following this cathodic wave, an anodic wave is observed at $E_3^{\text{ox}} = -0.07$ V which is followed by a less intense one at $E_2^{\text{ox}} = 0.06$ V, assigned (see text) to [OsCl₂(NH₃)₂(Me₂S)(Me₂SO-S)] and, tentatively, to [OsCl₂(NH₃)₂(Me₂S)], respectively. ^f Generated *in situ* from *cis*-**3** which, in 0.2 mol dm⁻³ KNO₃/H₂O, forms [RuCl(H₂O)₂(Me₂SO-S)₃]⁺ ($E_{p/2}^{\text{ox}} = 1.28$ V). ^g Upon scan reversal following this anodic wave, a cathodic wave is observed at $E_{p/2}^{\text{red}} = -0.39$ V. ^h In 0.2 mol dm⁻³ KCl/H₂O. ⁱ This wave is progressively replaced, with time, by a second one (which initially has a very weak current intensity) at $E_{1/2}^{\text{ox}} = 0.90$ V, assigned to [RuCl₂(N₂H₅)(Me₂SO-S)₂(H₂O)], due to the conversion of the starting complex into the latter (see text); in 0.2 mol dm⁻³ KNO₃/H₂O, the conversion is faster and in addition the formation of a third wave at $E_p^{\text{ox}} = 1.16$ V, involving the oxidation of liberated chloride ligand, is detected.

basis of the agreement (within the experimental error, ± 0.02 V) between the measured (-0.07 V vs. SCE) value of its oxidation potential and the predicted one (0.15 V vs. NHE = -0.09 V vs. SCE) by applying Lever's³¹ empirical expression (1) which

$$E = S_M (\sum E_L) + I_M \quad (1)$$

relates the redox potential of a complex (E , expressed in V vs. NHE) with the sum of the electrochemical E_L ligand parameter for all the ligands, and with the S_M and I_M metal centre parameters which depend upon the metal and redox couple, the spin state and stereochemistry.

For this purpose, we had to start with the estimate of E_L for S-coordinated Me₂SO, since for this ligand no reliable value was known.³¹ Hence, application of eqn. (1) to *trans*-[OsCl₂(Me₂SO-S)₄] **1**, considering our measured value of its oxidation poten-



Scheme 2


tial (1.43 V vs. NHE) and the known³¹ values of E_L (-0.24 V for the chloride ligand) and of S_M (1.01) and I_M (-0.40 V vs. NHE) for the Os^{III/II} redox couple, led to $E_L = 0.57$ V. This value is identical to those (0.55 – 0.58 V) we have estimated in a similar way for [RuCl₂(EtSCH₂CH₂SEt)(Me₂SO-S)₂],^{70a} [RuX₂(SMe₂)₃(Me₂SO-S)] ($X = \text{Cl}$ or Br)^{70a} and [Ru(NH₃)₅(Me₂SO-S)]²⁺,^{70b} which present clearly defined reversible anodic waves. Although this value is not unambiguously established,³¹ it is clearly high showing that Me₂SO-S behaves as a significant net electron π -acceptor/ σ -donor ligand in agreement with its expected (on the basis of X-ray data^{6,24}) significant π -electron acceptor ability. It is comparable to benzyl isocyanide ($E_L = 0.56$ V),³¹ being a weaker net electron donor than *e.g.* Me₂S ($E_L = 0.31$).³¹

[RuCl₂(Me₂SO)₄]. The cyclic voltammogram of a Me₂SO or an aqueous solution of *cis*-[RuCl₂(Me₂SO-O)(Me₂SO-S)] *cis*-**3** exhibits a single-electron irreversible anodic wave at $E_{p/2}^{\text{ox}} = 1.47$ or 1.28 V vs. SCE, respectively. These values are remarkably higher (by *ca.* 0.9 V) than those expected for this complex in these media (0.62 or 0.39 V, correspondingly) on the basis of eqn. (1) and the known³¹ values of S_M (0.97 or 1.14 , in organic or aqueous medium, respectively) and I_M (0.04 V or -0.35 V vs. NHE, in such media) for the Ru^{III/II} redox couple, and of E_L for the various ligands. Incidentally, the use of eqn. (1) required the previous estimate of E_L for O-coordinated Me₂SO by applying this expression to [Ru(NH₃)₅(Me₂SO-O)]²⁺ ($E^{\text{ox}} = 0.02$ V vs. NHE,⁷¹ $E_L(\text{NH}_3) = 0.07$ V³¹). Although one should be cautious about this estimated value ($E_L = -0.37$ V) since it is based on the oxidation potential of a single complex, it is drastically lower (by 0.94 V) than that of Me₂SO-S (see above), being intermediate between those of Cl⁻ (-0.24 V)³¹ and F⁻ (-0.42 V),³¹ showing that O-coordination results in a remarkable

enhancement of the net electron-donor ability of Me₂SO which then should not behave as an appreciable π-electron acceptor, in contrast to the S-bonded ligand.

However, the measured oxidation potentials of the anodic waves of the above solutions of *cis*-**3** are in close agreement with those estimated using eqn. (1), for the known⁴⁵ products of its spontaneous conversion in those media, *i.e.* *trans*-[RuCl₂(Me₂SO-S)₄] *trans*-**3** in Me₂SO and [RuCl(H₂O)₂(Me₂SO-S)₃]⁺ in H₂O (1.48 and 1.2 V vs. SCE, respectively). *trans*-**3** is oxidised at a more anodic potential (by *ca.* 0.3 V) than the analogous Os^{II} compound **1** (see above), thus accounting for the stronger resistance to oxidation of the Ru^{II} centre, in comparison with the Os^{II} site, in the above reactions with N₂H₄·2HCl.

[RuCl₃(N₂H₅)(Me₂SO)₂]. The cyclic voltammogram of an aqueous solution of *mer*-[RuCl₃(N₂H₅)(Me₂SO-S)₂] **4** also indicates its lability in solution, presenting one irreversible anodic wave (wave I) at ^IE_{p/2}^{ox} = 0.79 V vs. SCE that, with time, is progressively replaced by another one (wave II), at a slightly higher potential (^{II}E_{p/2}^{ox} = 0.90 V), with the occurrence of an isopotential point (at 0.87 V). Hence, in solution, the starting complex **4** (oxidised at wave I) converts into another species **5** (oxidised at wave II). This conversion conceivably involves dehydrochlorination (eqn. (2)) *via* [RuCl₃(N₂H₄)(Me₂SO-S)₂]⁻




to give [RuCl₂(N₂H₄)(Me₂SO-S)₂(H₂O)] **5**, on the basis of the following evidence: (i) acidic character, pH = 4.2, of a solution of **4** and titration (by a KOH solution) of 1H⁺/molecule; (ii) promotion of the conversion of **4** into **5** by addition of that base and on replacement of the KCl electrolyte by another one without Cl⁻, such as KNO₃, when a third anodic wave (III) then forms at ^{III}E_p^{ox} = 1.2 V, a value identical to that measured for the oxidation of Cl⁻ under a solution of [NEt₄]Cl under the same experimental conditions (0.2 mol dm⁻³ KNO₃/H₂O).

The anodic waves correspond to single-electron processes at relatively short time scales, as measured by chronoamperometry and steady-state voltammetry at an ultramicroelectrode⁷² (using methylviologen as the reference compound), but over the longer time scale of CPE (see below) a multi-electron process is involved. The oxidation potentials of our Ru^{II} complexes are comparable to (or even higher than) those reported, *ca.* 0.8 V, for [Ru(NH₃)₅(Me₂SO-S)]²⁺ and related complexes,^{70b,71,73} in which Me₂SO-S is behaving also as a significant π-electron acceptor. O-coordination would result in a dramatic cathodic shift (by *ca.* 0.6–1.0 V)^{70b,71,73} of the redox potential that would also occur if an S-bonded Me₂SO did not act as a π-electron acceptor as observed⁷³ for *mer*-[RuCl₃(Me₂SO-S)-(tmen)] (tmen = *N,N,N',N'*-tetramethylenediamine) with *E*_{1/2}(Ru^{III/II}) = -0.14 V vs. SCE.

The coordination electronic properties of the hydrazinium ligand can also be investigated in complex **4** by attempting to estimate its as yet unknown *E*_L value through the application of eqn. (1), considering our measured ^IE_{p/2}^{ox} = 1.04 V vs. NHE and our estimated *E*_L(Me₂SO-S) = 0.57 V. The obtained value, *E*_L(N₂H₅⁺) = 0.80 V, is one of the highest so far reported, suggesting that hydrazinium behaves as a rather weak overall electron-donor ligand, being comparable to *N*-methylpyrazinium (*E*_L = 0.79 V)³¹ and weaker than *e.g.* pyrimidinium (*E*_L = 0.43 V)³¹ or ammonia (0.07 V),³¹ but not so weak as the strong π-electron acceptors CO (0.99 V)³¹ or carbynes (*ca.* 1.2 V).⁷⁴ However, the estimated *E*_L value for N₂H₅⁺ should be taken cautiously since it is based on the oxidation potential of a single complex which, moreover, presents a ligand (Me₂SO) whose *E*_L value has not yet been fully established.

Exhaustive CPE at 0.90 V of a solution of **4** led to the consumption of *ca.* 5 F mol⁻¹, but the trend of the variation of current vs. charge indicates the initial involvement of a single electron process (see above). CV of the electrolysed solution showed the presence of a partially reversible cathodic wave at *E*_p^{red} = -0.44 V, due to H⁺ reduction, as corroborated (i) by its high cathodic shift upon replacement of the Pt by a vitreous C working electrode and (ii) by acid–base potentiometric titration, with a KOH solution, of the electrochemically oxidised solution of the complex (5H⁺/molecule of complex in a titration curve with only one very well defined end-point).

The acidity character of the hydrazinium or hydrazine ligand is enhanced upon oxidation of the complex, and the oxidised forms [RuCl₃(N₂H₅)(Me₂SO-S)₂]⁺ or [RuCl₂(N₂H₄)(Me₂SO-S)₂(H₂O)]⁺ (see above) generated at waves I or II, respectively, are expected to undergo ready H⁺ loss (Scheme 3 for the former



Scheme 3

case). The promotion of acidity upon further oxidation steps would account for the observed multi-deprotonation in the overall multi-electron anodic process. Related behaviour was reported by us⁷⁵ for aminocarbene ligands in some complexes of Pd^{II} or Pt^{II}, whereas the aminocarbyne CNH₂ ligating a Re phosphinic site is anodically converted into the corresponding isocyanide (CNH) and cyano complexes.⁷⁶

In order to try to identify the products formed during the CPE, this was also performed in a closed system, under argon, and the atmosphere in the electrochemical cell was then analysed by EI-MS that showed the presence of several volatile N- and S-containing species such as N₂, N₂O, N₂O₂, NO₂ and SO₂, therefore indicating an oxidative conversion of the hydrazine and the Me₂SO ligands into N₂ and N-oxides or SO₂, respectively. The oxidation of Me₂SO was only partial, and the analysis of the electrolysed solution, by FAB-MS, detected *e.g.* {RuCl₂O(Me₂SO)₂}⁺ (*m/z* 345), {Ru₂Cl₆(Me₂SO)}⁺ (*m/z* 498) and {Ru₂Cl₆O(Me₂SO)}⁺ (*m/z* 514).

Acknowledgements

This work has been partially supported by the Foundation for Science and Technology (FCT) and the PRAXIS XXI Programme (Portugal), the Ministero per l'Università e la Ricerca Scientifica e Tecnologica (Rome, Italy) and the Russian Fund for Basic Research. We thank Dr G. Wagner for NMR and Mr I. Marques for MS measurements.

References

- S. Oae, *Organic Sulphur Chemistry: Structure and Mechanism*, CRC Press, Boca Raton, FL, 1991, p. 312; M. Madesclaire, *Tetrahedron*, 1988, **44**, 6537; J. S. Grossert, in *Chemistry of Sulfoxes and Sulfoxides*, ed. S. Patai, Z. Rappoport and C. Stirling, Wiley, Chichester, 1988, p. 925.
- J. Drabowicz, H. Togo, M. Mikolajczyk and S. Oae, *Org. Prep. Proced. Int.*, 1984, **16**, 171 and references therein.
- A. J. Mancuso and D. Swern, *Synthesis*, 1981, 165.
- I. Lindquist and P. Einarsson, *Acta Chem. Scand.*, 1959, **13**, 420; F. A. Cotton and R. C. Elder, *J. Am. Chem. Soc.*, 1960, **82**, 2986; D. W. Meek, D. K. Straub and R. S. Drago, *Bull. Chem. Soc. Jpn.*, 1960, **33**, 861; F. A. Cotton and R. Francis, *J. Inorg. Nucl. Chem.*, 1961, **17**, 62.
- J. A. Davies, *Adv. Inorg. Chem. Radiochem.*, 1981, **24**, 115.

- 6 M. Calligaris and O. Carugo, *Coord. Chem. Rev.*, 1996, **153**, 83; M. Calligaris, *Croat. Chem. Acta*, 1999, **72**, 147.
- 7 S. Geremia, S. Mestroni, M. Calligaris and E. Alessio, *J. Chem. Soc., Dalton Trans.*, 1998, 2447.
- 8 T. Tanase, T. Aiko and Y. Yamamoto, *Chem. Commun.*, 1996, 2341.
- 9 E. Alessio, M. Bolle, B. Milani, G. Mestroni, P. Faleschini, S. Geremia and M. Calligaris, *Inorg. Chem.*, 1995, **34**, 4716; E. Alessio, B. Milani, M. Bolle, G. Mestroni, P. Faleschini, F. Todone, S. Geremia and M. Calligaris, *Inorg. Chem.*, 1995, **34**, 4722.
- 10 B. R. James, R. S. McMillan and K. J. Reimer, *J. Mol. Catal.*, 1976, **1**, 439; B. R. James and R. S. McMillan, *Can. J. Chem.*, 1977, **55**, 3927; D. P. Riley, *Inorg. Chem.*, 1983, **22**, 1965; D. P. Riley and R. E. Shumate, *J. Am. Chem. Soc.*, 1984, **106**, 3179; D. P. Riley and J. Lyon III, *J. Chem. Soc., Dalton Trans.*, 1991, 157; R. S. Srivastava, B. Milani, E. Alessio and G. Mestroni, *Inorg. Chim. Acta*, 1992, **191**, 15.
- 11 G. Mestroni, E. Alessio, M. Calligaris, W. M. Attia, F. Quadrioglio, S. Cauci, G. Sava, S. Zorzet, S. Pacor, C. Monti-Bragadin, M. Tamaro and L. Dolzani, *Prog. Clin. Biochem. Med.*, 1989, **10**, 71; G. Mestroni, E. Alessio, G. Sava, S. Pacor, M. Coluccia, A. Boccarelli, *Metal-Based Drugs*, 1994, **1**, 41; G. Sava, E. Alessio, A. Bergamo and G. Mestroni, *Top. Bioinorg. Chem.*, 2000, **2**, 143.
- 12 N. Farrell, in *Platinum, Gold, and Other Metal Chemotherapeutic Agents*, ed. S. Lippard, ACS Symposium Series 209, American Chemical Society, Washington, DC, 1983, p. 279.
- 13 V. Yu. Kukushkin, *Coord. Chem. Rev.*, 1995, **139**, 375.
- 14 J. B. Arterburn and M. C. Perry, *Tetrahedron Lett.*, 1996, **37**, 7941; J. B. Arterburn and S. L. Nelson, *J. Org. Chem.*, 1996, **61**, 2260; Y. M. Wang and M. Koreeda, *Synlett*, 1996, 885; X. T. Wang and L. K. Koo, *J. Org. Chem.*, 1998, **63**, 356; L. J. Laughlin and C. G. Young, *Inorg. Chem.*, 1996, **35**, 1050; M. A. Pietsch and M. B. Hall, *Inorg. Chem.*, 1996, **35**, 1273; D. D. DuMez and J. M. Mayer, *Inorg. Chem.*, 1995, **34**, 6396; R. H. Khan and R. C. Rastogi, *Indian J. Chem., Sect. B*, 1994, **33**, 293; H. Arzonmanian, R. Lopez and G. Agrifoglio, *Inorg. Chem.*, 1994, **33**, 3177.
- 15 F. Mohanazadeh, A. R. Momeni and Y. Ranjbar, *Tetrahedron Lett.*, 1994, **35**, 6127; M. Kobayashi, K. Kawakita, T. Yokota, K. Mannami, O. Yamamoto, H. Morikawa and H. Konishi, *Bull. Chem. Soc. Jpn.*, 1995, **68**, 1401; Y. Zhang, Y. Yu and W. Bao, *Synth. Commun.*, 1995, **25**, 1825; J. Q. Wang and Y. M. Zhang, *Synth. Commun.*, 1995, **25**, 3545; A. Leburn, E. Rantze, J.-L. Namy and H. B. Kagan, *New J. Chem.*, 1995, **19**, 699.
- 16 P. Ghosh, K. Pramanik and A. Chakravorty, *J. Chem. Soc., Chem. Commun.*, 1995, 477.
- 17 K. Pramanik, P. Ghosh and A. Chakravorty, *Inorg. Chem.*, 1998, **37**, 5678.
- 18 Yu. N. Kukushkin and Z. A. Khromenkova, *Zh. Obshch. Khim.*, 1996, **66**, 1752 and references therein.
- 19 Yu. N. Kukushkin, Z. A. Khromenkova and V. A. Esaulova, *Zh. Obshch. Khim.*, 1995, **65**, 1785.
- 20 A. D. Ryabov, G. M. Kazankov, I. M. Panyashkina, O. V. Grozovsky, O. G. Dyachenko, V. A. Polyakov and L. G. Kuzmina, *J. Chem. Soc., Dalton Trans.*, 1997, 4385.
- 21 S. Fabre, B. Findeis, D. J. M. Trösch, L. H. Gade, I. J. Scowen and M. McPartlin, *Chem. Commun.*, 1999, 577.
- 22 Yu. N. Kukushkin, M. V. Bavina and A. V. Zinchenko, *Zh. Obshch. Khim.*, 1995, **65**, 1788 and references therein.
- 23 Yu. N. Kukushkin, M. V. Bavina and A. V. Zinchenko, *Zh. Obshch. Khim.*, 1996, **66**, 337.
- 24 M. Stener and M. Calligaris, *J. Mol. Struct. (THEOCHEM)*, 2000, **497**, 91.
- 25 V. Yu. Kukushkin, V. K. Belsky, V. E. Kononov, E. A. Aleksandrova, E. Yu. Pankova and A. I. Moiseev, *Phosphorus Sulfur Silicon Relat. Elem.*, 1992, **69**, 103; Yu. N. Kukushkin and Z. A. Khromenkova, *Zh. Obshch. Khim.*, 1995, **65**, 1409.
- 26 V. Yu. Kukushkin and E. Yu. Pankova, *Koord. Khim.*, 1989, **15**, 531; *Russ. J. Coord. Chem. (Transl. of Khim.)*, 1989, **15**, 331.
- 27 V. Yu. Kukushkin, V. K. Belsky, E. A. Aleksandrova, E. Yu. Pankova, V. E. Kononov, V. N. Yakovlev and A. I. Moiseev, *Zh. Obshch. Khim.*, 1991, **61**, 318; *J. Gen. Chem. USSR (Engl. Transl.)*, 1991, **61**, 284; V. Yu. Kukushkin, E. Yu. Pankova, S. A. Simanova, S. S. Sotman, V. S. Fundamensky, I. G. Zenkevich and I. A. Poliakov, *Zh. Obshch. Khim.*, 1990, **60**, 587; *J. Gen. Chem. USSR (Engl. Transl.)*, 1990, **60**, 513.
- 28 P. G. Antonov and I. A. Amantova, *Zh. Obshch. Khim.*, 1988, **58**, 2523; *J. Gen. Chem. USSR (Engl. Transl.)*, 1988, **58**, 2245; Yu. N. Kukushkin, V. K. Krylov, Yu. E. Larionova and M. V. Bavina, *Zh. Obshch. Khim.*, 1995, **65**, 881.
- 29 J. S. Jaswal, S. J. Rettig and B. R. James, *Can. J. Chem.*, 1990, **68**, 1808; D. T. T. Yapp, J. Jaswal, S. J. Rettig, B. R. James and K. A. Skov, *Inorg. Chim. Acta*, 1990, **177**, 199.
- 30 O. V. Rudnitskaya and N. M. Sinitsyn, *Zh. Neorg. Khim.*, 1994, **39**, 925.
- 31 A. B. P. Lever, *Inorg. Chem.*, 1990, **29**, 1271.
- 32 I. P. Evans, A. Spencer and G. Wilkinson, *J. Chem. Soc., Dalton Trans.*, 1973, 204.
- 33 P. G. Antonov, Yu. N. Kukushkin, V. I. Konnov and Yu. P. Kostikov, *Koord. Khim.*, 1980, **6**, 1585.
- 34 D. Gajapathy, S. Govindarajan, K. C. Patil and H. Manohar, *Polyhedron*, 1983, **2**, 865.
- 35 G. M. Sheldrick, SHELXS-86, Program for structure solution, *Acta Crystallogr., Sect. A*, 1990, **46**, 467.
- 36 G. M. Sheldrick, SHELXL-97, Program for the Refinement of Crystal Structures, University of Göttingen, Germany, 1997.
- 37 A. M. McDonagh, M. G. Humphrey and D. C. R. Hockless, *Aust. J. Chem.*, 1998, **51**, 807.
- 38 D. M. Stanbury, *Prog. Inorg. Chem.*, 1998, **47**, 511 and references therein.
- 39 P. A. Lay and H. Taube, *Inorg. Chem.*, 1989, **28**, 3561.
- 40 T. Furuhashi, M. Kawano, Y. Koide, R. Somazawa and K. Matsumoto, *Inorg. Chem.*, 1999, **38**, 109.
- 41 L. I. Elding and Ö. Gröning, *Inorg. Chem.*, 1978, **17**, 1872; Y. Ducommun, L. Helm, A. E. Merbach, B. Hellquist and L. I. Elding, *Inorg. Chem.*, 1989, **28**, 377.
- 42 J. A. Cabeza, H. Adams and A. J. Smith, *Inorg. Chim. Acta*, 1986, **114**, L17.
- 43 J. Y. K. Cheng, K.-K. Cheung and C.-M. Che, *Chem. Commun.*, 1997, 623.
- 44 P. D. Robinson, C. C. Hinckley and A. Ikuo, *Acta Crystallogr., Sect. C*, 1989, **45**, 1079.
- 45 E. Alessio, G. Mestroni, G. Nardin, W. M. Attia, M. Calligaris, G. Sava and S. Zorzet, *Inorg. Chem.*, 1988, **27**, 4099.
- 46 W. R. Scheidt and H. Nasri, *Inorg. Chem.*, 1995, **34**, 2190.
- 47 B. R. James, A. Pacheco, S. J. Rettig and J. A. Ibers, *Inorg. Chem.*, 1988, **27**, 2414.
- 48 (a) K. Krogh-Jespersen, X. Zhang, J. D. Westbrook, R. Fikar, K. Nayak, W.-L. Kwik, J. A. Potenza and H. J. Schugar, *J. Am. Chem. Soc.*, 1989, **111**, 4082; (b) K. Krogh-Jespersen, X. Zhang, Y. Ding, J. D. Westbrook, J. A. Potenza and H. J. Schugar, *J. Am. Chem. Soc.*, 1992, **114**, 4345.
- 49 B. W. Arbuckle, P. K. Bharadwaj and W. K. Musker, *Inorg. Chem.*, 1991, **30**, 440.
- 50 D. P. Riley and J. D. Oliver, *Inorg. Chem.*, 1986, **25**, 1821.
- 51 G. Albertin, S. Antoniutti, A. Bacchi, M. Bergamo, E. Bordignon and G. Pelizzi, *Inorg. Chem.*, 1998, **37**, 479.
- 52 E. Alessio, M. Calligaris, M. Iwamoto and L. G. Marzilli, *Inorg. Chem.*, 1996, **35**, 2538; E. Alessio, E. Zangrando, R. Roppa and L. G. Marzilli, *Inorg. Chem.*, 1998, **37**, 2458.
- 53 S. Geremia, M. Calligaris, Yu. N. Kukushkin, A. V. Zinchenko and V. Yu. Kukushkin, *J. Mol. Struct.*, 2000, **516**, 49.
- 54 J. E. Kickham and S. J. Loeb, *Inorg. Chem.*, 1995, **34**, 5656.
- 55 H.-R. Bircher, P. Ochsenbein, J. Hauser and H.-B. Bürgi, *Acta Crystallogr., Sect. C*, 1996, **52**, 2002.
- 56 V. E. Shklover and Yu. T. Struchkov, *Acta Crystallogr., Sect. C*, 1985, **41**, 734.
- 57 E. Garcia, K.-Y. Lee and C. B. Storm, *Acta Crystallogr., Sect. C*, 1992, **48**, 1683.
- 58 D. Sellmann, W. Soglowek, F. Knoch, G. Ritter and J. Dengler, *Inorg. Chem.*, 1992, **31**, 3711.
- 59 M. T. Casey, P. Guinan, A. Canavan, M. McCann and C. Cardin, *Polyhedron*, 1991, **10**, 483.
- 60 E. N. Maslen, C. L. Raston, B. W. Skelton and A. H. White, *Aust. J. Chem.*, 1975, **28**, 739.
- 61 L. Blum, J. D. Williams and R. R. Schrock, *J. Am. Chem. Soc.*, 1984, **106**, 8316.
- 62 K. Matsumoto, H. Uemura and M. Kawano, *Chem. Lett.*, 1994, 1215.
- 63 K. Mashima, H. Kaneyoshi, S. Kaneko, K. Tani and A. Nakamura, *Chem. Lett.*, 1997, 569.
- 64 M. Kawano, C. Hoshino and K. Matsumoto, *Inorg. Chem.*, 1992, **31**, 5158.
- 65 The oxidation potential of this intermediate (0.06 V) does not differ substantially from that (-0.07 V vs. SCE, wave III) of its Me₂SO adduct [OsCl₂(NH₃)₂(Me₂S)(Me₂SO)], as expected in view of the similarities of (i) the oxidation potential values for the five-coordinate Re complex [ReCl(dppe)₂] (dppe = Ph₂PCH₂CH₂PPh₂, see ref. 66) ($E_p^{ox} = 0.09$ V vs. SCE, measured in this study, in 0.2 mol dm⁻³ [NBu₄][BF₄]/tetrahydrofuran) and for its alkyl isocyanide adducts [ReCl(CNR)(dppe)₂] (R = Bu^t or Me) ($E_{1/2}^{ox} = 0.06-0.08$ V vs. SCE, see ref. 67) and (see text) of (ii) the net electron-donor character of the Me₂SO and benzyl isocyanide ligands.

- 66 D. L. Hughes, A. J. L. Pombeiro, C. J. Pickett and R. L. Richards, *J. Organomet. Chem.*, 1983, **248**, C26.
- 67 A. J. L. Pombeiro, C. J. Pickett and R. L. Richards, *J. Organomet. Chem.*, 1982, **224**, 285.
- 68 T. A. Salih, M. T. Duarte, J. J. R. Fraústo da Silva, A. M. Galvão, M. F. C. Guedes da Silva, P. B. Hitchcock, D. L. Hughes, C. J. Pickett, A. J. L. Pombeiro and R. L. Richards, *J. Chem. Soc., Dalton Trans.*, 1993, 3015.
- 69 G. Costa, G. Balducci, E. Alessio, C. Tavagnacco and G. Mestroni, *J. Electroanal. Chem.*, 1990, **296**, 57.
- 70 (a) D. P. Riley and J. D. Olivier, *Inorg. Chem.*, 1986, **25**, 1814; (b) M. Sano and H. Taube, *Inorg. Chem.*, 1994, **33**, 705; M. Sano and A. Tomita, *Inorg. Chem.*, 1994, **33**, 5825.
- 71 A. Yeh, N. Scott and H. Taube, *Inorg. Chem.*, 1982, **21**, 2542.
- 72 C. Amatore, M. Azzabi, P. Calas, A. Jutand, C. Lefrou and Y. Rollin, *J. Electroanal. Chem.*, 1990, **288**, 45.
- 73 J. J. Rack and H. B. Gray, *Inorg. Chem.*, 1999, **38**, 2.
- 74 S. S. P. R. Almeida and A. J. L. Pombeiro, *Organometallics*, 1997, **16**, 4469.
- 75 R. Bertani, M. Mozzon, R. A. Michelin, F. Benetollo, G. Bombieri, T. J. Castilho and A. J. L. Pombeiro, *Inorg. Chim. Acta*, 1991, **189**, 175.
- 76 M. A. N. D. A. Lemos, M. F. C. Guedes da Silva and A. J. L. Pombeiro, *Inorg. Chim. Acta*, 1994, **226**, 9.
- 77 R. K. Coll, J. E. Fergusson, V. McKee, C. T. Page, W. T. Robinson and T. S. Keong, *Inorg. Chem.*, 1987, **26**, 106.
- 78 W. A. Schenk, J. Frisch, M. Dürr, N. Burzlaff, D. Stalke, R. Fleischer, W. Adam, F. Prechtel and A. K. Smertz, *Inorg. Chem.*, 1997, **36**, 2372.