

## Reductions by hyponitrite †

Ahmad M. Al-Ajlouni<sup>a</sup> and Edwin S. Gould<sup>b</sup><sup>a</sup> Department of Chemistry, Jordan University of Science and Technology, Irbid, Jordan<sup>b</sup> Department of Chemistry, Kent State University, Kent, Ohio 44242, USA

Received 24th November 1999, Accepted 4th February 2000

Published on the Web 29th March 2000

Hyponitrites, salts of the weak acid HO–N=N–OH, hyponitrous acid, react, in aqueous solution (pH 4–8), with an array of inorganic oxidants. With the high potential reagents Br<sub>2</sub> and [Fe(bipy)<sub>3</sub>]<sup>3+</sup> in excess, conversion is mainly to nitrate, whereas with I<sub>2</sub> and [IrCl<sub>6</sub>]<sup>2-</sup>, the product is nitrite. Kinetic acidity patterns indicate that nearly all oxidations proceed largely through the conjugate base, HN<sub>2</sub>O<sub>2</sub><sup>-</sup>. With I<sub>2</sub>, I<sub>3</sub><sup>-</sup>, and Br<sub>2</sub> there is also evidence also for a doubly deprotonated route, possibly involving a halogen–N<sub>2</sub>O<sub>2</sub><sup>2-</sup> complex. The 1e<sup>-</sup> metal oxidants [Fe(bipy)<sub>3</sub>]<sup>3+</sup> and [IrCl<sub>6</sub>]<sup>2-</sup> probably operate through a series of outer-sphere 1e<sup>-</sup> steps in which the initial transfer is rate-determining, whereas with I<sub>2</sub> and Br<sub>2</sub>, reaction *via* a halogen bridge, resulting in net transfer of X<sup>+</sup> (a 2e<sup>-</sup> change) is more likely. None of the reaction profiles exhibits irregularities attributable to the intervention of an intermediate formed or destroyed on a time scale commensurate with the initial reaction.

## Introduction

Although hyponitrites, salts of the weak acid acid HON=NOH, hyponitrous acid, have been known since 1871,<sup>2,3</sup> interest in this N(i) system has been greatly overshadowed by the vast body of attention bestowed on the higher oxidation states of nitrogen. Acidity constants (pK<sub>1</sub> = 7.18; pK<sub>2</sub> = 11.54) have been documented,<sup>4</sup> and a *trans*-configuration has been established, both for the dinegative anion,<sup>5</sup> and for the predominant form of the solid parent acid.<sup>5,6</sup>

In principle, a nitrogen(i) transient should be involved in the enzymatic oxidation of ammonia to nitrate (nitrification)<sup>7</sup> and in the biologically significant reductions of nitrite to ammonia<sup>8a</sup> or N<sub>2</sub>.<sup>8b</sup> However, attempts to demonstrate the intermediacy of hyponitrous acid or its anion in such biosystems<sup>7,9</sup> have yielded negative or questionable results, suggesting that such species, if they intervene at all, are unusually reactive.

We have extended our interest in the redox chemistry of the lower oxidation states of nitrogen<sup>10</sup> to the hyponitrite system, which, like hydrazine and hydroxylamine, is a powerful but selective reductant. Although detailed investigations into the decomposition of both hyponitrous acid and its salts have been carried out,<sup>11</sup> mechanistic studies of reactions with inorganic oxidants have been more fragmentary. Kinetic studies by Gupta and co-workers<sup>12</sup> embraced a variety of oxidants but generally dealt only with initial reaction rates and utilized quite limited variations in reaction media. The reaction between hyponitrous and nitrous acids (eqn. 1) has been examined at some length,<sup>13</sup>



and the N<sub>2</sub> formed has been shown<sup>14</sup> to arise from the N(i) acid. Hence this conversion may be considered to be a net reduction, rather than an oxidation, of hyponitrous acid.

## Experimental

## Materials

Iron(III) perchlorate [Fe(ClO<sub>4</sub>)<sub>3</sub>·6H<sub>2</sub>O], sodium *trans*-hyponitrite hydrate (Na<sub>2</sub>N<sub>2</sub>O<sub>2</sub>·xH<sub>2</sub>O), potassium hexachloro-

palladate(IV) (K<sub>4</sub>PdCl<sub>6</sub>), tris(2,2'-bipyridine)iron(III) hexafluorophosphate, thallium(III) perchlorate, sodium thiocyanate (NaSCN) (Aldrich products), bromine water (Fisher), sodium hexachloroiridate(IV) (Na<sub>2</sub>IrCl<sub>6</sub>·6H<sub>2</sub>O), and sodium hexachloroplatinate(IV) (Na<sub>2</sub>PtCl<sub>6</sub>) (Alfa products) were used as received.

Fresh stock solutions of sodium hyponitrite were prepared daily in aqueous base (1.0 M NaOH) and were standardized spectrophotometrically in 0.1 M NaOH (ε<sub>248</sub> 6.6 × 10<sup>3</sup> M<sup>-1</sup> cm<sup>-1</sup>).<sup>15</sup> Stock solutions of [Fe(bipy)<sub>3</sub>](PF<sub>6</sub>)<sub>3</sub> were prepared in dry acetonitrile and were filtered to remove insoluble impurities. Stock solutions of Na<sub>2</sub>PdCl<sub>6</sub> and Na<sub>2</sub>PtCl<sub>6</sub> were prepared in 0.1 M HCl.

## Stoichiometric studies

Stoichiometries of the oxidation of hyponitrite were generally determined with the oxidant in excess. The decrease in absorbance at λ<sub>max</sub> of the oxidant, after addition of a measured deficiency of hyponitrite and standing for 2–10 min, was used to estimate the loss of oxidant. These changes were compared with those seen when excess hyponitrite was added. Reasonable values could not be obtained for the oxidation by I<sub>3</sub><sup>-</sup> because of the competing reaction of the product, HNO<sub>2</sub>, with I<sup>-</sup> in the supporting medium, nor for oxidation by [PdCl<sub>6</sub>]<sup>2-</sup> because of the attendant decomposition of the oxidant to Cl<sub>2</sub>. In the latter case, however, the product was identified as [PdCl<sub>4</sub>]<sup>2-</sup> by its spectrum. Results for other systems are summarized in Table 1.

## Kinetic experiments

Reactions, under argon, were generally examined at or near λ<sub>max</sub> of the oxidant, using either a Shimadzu-1601 instrument or a Durrum-Gibson stopped-flow spectrophotometer interfaced with an OLIS computer system. Temperatures were kept at 24.0 ± 0.5 °C throughout the entire series of experiments. Ionic strength was regulated by addition of (NaClO<sub>4</sub> + HClO<sub>4</sub>) or (HCl + NaCl). Low pH values (0–2) were maintained with added HClO<sub>4</sub> or HCl, and higher pH's by phosphate, acetate or chloroacetate buffers. Solutions of hyponitrite, especially those held at pH 6–8, were used as quickly as possible after preparation to minimize decomposition of this reagent in solution. Reactions were run under pseudo-first order conditions with

† Electron Transfer, part 142. For part 141, see ref. 1.

**Table 1** Stoichiometry of oxidations of hyponitrite<sup>a</sup>

Oxidant (Ox)	pH	$\mu$ /M	$\lambda$ /nm	$\frac{\Delta[\text{Ox}]}{\Delta[\text{H}_2\text{N}_2\text{O}_2]}$
I <sub>2</sub>	4.0	0.20	456	1.85
Br <sub>2</sub>	0.7	0.20	395	3.8 <sup>b</sup>
[IrCl <sub>6</sub> ] <sup>2-</sup>	7.0	0.20	489	3.8
[Fe(bipy) <sub>3</sub> ] <sup>3+</sup>	6.0	1.0	525	7.4



<sup>a</sup> For reaction conditions, see Experimental section. <sup>b</sup> Value with Br<sub>2</sub> in excess. With hyponitrite in excess, the ratio  $[\text{HNO}_2]_{\text{formed}}/[\text{Br}_2]_{\text{consumed}}$  was  $1.2 \pm 0.1$ .

hyponitrite in greater than tenfold excess. Rate constants were obtained from the resulting profiles by nonlinear least squares fittings to the relationship describing first order decay. All reactions for which reproducible data could be obtained were first order both in hyponitrite and oxidant. Although many of the rates were strongly acid dependent, none of the reactions examined in buffered systems exhibited general acid or general base catalysis. In a few instances reactions were run without protection by argon; results were not significantly affected.



A number of oxidizing species reacted inconveniently slowly, or not at all, with hyponitrite under the conditions examined. Among these were [Fe(CN)<sub>6</sub>]<sup>3-</sup>, [Co(NH<sub>3</sub>)<sub>5</sub>Br]<sup>2+</sup>, [Fe(NCS)]<sup>2+</sup>, Fe(III)<sub>aq</sub> (pH 0.2), the Ni(IV) complex of 2,6-diacetylpyridine dioxime,<sup>16</sup> the Ag(III) complex of ethylenebis(biguanide),<sup>17</sup> and hydrogen peroxide (both in the presence and the absence of CH<sub>3</sub>ReO<sub>3</sub>).<sup>18</sup> In contrast, the reaction with Tl(ClO<sub>4</sub>)<sub>3</sub> (pH 2.8–4.7) was too rapid to follow, even using stopped-flow methods under second order conditions.

## Results and discussion

Stoichiometric results summarized in Table 1 indicate that under the conditions chosen, hyponitrous acid is oxidized principally to HNO<sub>2</sub> by I<sub>2</sub> ( $E^\circ$  0.54 V) and [IrCl<sub>6</sub>]<sup>2-</sup> (0.87 V),<sup>19</sup> even when these reagents are employed in excess (eqns. 2 and 3),



but is converted predominantly to NO<sub>3</sub><sup>-</sup> by the more powerful oxidants, Br<sub>2</sub> (1.06 V) and [Fe(bipy)<sub>3</sub>]<sup>3+</sup> (1.11 V)<sup>20</sup> (eqn. 4 and 5). However, under our kinetic conditions (reductant in excess),



oxidation by Br<sub>2</sub> yields mainly HNO<sub>2</sub>. Tripositive nitrogen doubtless intervenes in the H<sub>2</sub>N<sub>2</sub>O<sub>2</sub>–[Fe(bipy)<sub>3</sub>]<sup>3+</sup> reactions, but the oxidation of HNO<sub>2</sub> by this high potential chelate proceeds 40–60 times as rapidly as that of H<sub>2</sub>N<sub>2</sub>O<sub>2</sub> and hence is kinetically silent. None of the reaction profiles in the present study exhibits irregularities attributable to formation or destruction of a transient on a time scale comparable to that of the principal reaction.

### Oxidations by halogen species

Kinetic data for oxidations by I<sub>2</sub> and by I<sub>3</sub><sup>-</sup> are assembled in Table 2. Reactions are accelerated by increases in pH, and the trends in rate are decidedly sharper than that characterizing a

**Table 2** Kinetic data for the oxidation of hyponitrous acid with iodine and with triiodide<sup>a</sup>

A H <sub>2</sub> N <sub>2</sub> O <sub>2</sub> + I <sub>2</sub>		B H <sub>2</sub> N <sub>2</sub> O <sub>2</sub> + I <sub>3</sub> <sup>-b</sup>	
pH	$k/\text{M}^{-1} \text{s}^{-1c}$	pH	$k/\text{M}^{-1} \text{s}^{-1c}$
3.68	11 (15)	4.00	1.0 (0.81)
3.80	21 (24)	4.50	5.5 (4.5)
3.88	43 (32)	4.65	6.6 (8.1)
3.91	50 (36)	5.05	33 (42)
4.14	99 (87)	5.35	$1.2 \times 10^2$ ( $1.5 \times 10^2$ )
4.29	$1.8 \times 10^2$ ( $1.6 \times 10^2$ )	5.40	$1.6 \times 10^2$ ( $1.9 \times 10^2$ )
4.40	$2.5 \times 10^2$ ( $2.5 \times 10^2$ )	5.48	$2.3 \times 10^2$ ( $2.7 \times 10^2$ )
4.50	$3.8 \times 10^2$ ( $3.9 \times 10^2$ )	5.66	$5.5 \times 10^2$ ( $6.0 \times 10^2$ )
5.08	$4.5 \times 10^3$ ( $5.0 \times 10^3$ )	6.25	$8.0 \times 10^3$ ( $7.9 \times 10^3$ )
5.20	$8.8 \times 10^3$ ( $8.7 \times 10^3$ )	6.60	$3.4 \times 10^4$ ( $3.4 \times 10^4$ )
5.22	$1.00 \times 10^4$ ( $0.95 \times 10^4$ )	6.90	$1.1 \times 10^5$ ( $1.1 \times 10^5$ )
5.69	$7.3 \times 10^4$ ( $7.9 \times 10^4$ )		

<sup>a</sup> Reactions were run at  $24.5 \pm 0.5$  °C;  $\mu = 0.20$  M (NaClO<sub>4</sub>). Solutions were buffered using HOAc/OAc<sup>-</sup> or H<sub>2</sub>PO<sub>4</sub><sup>-</sup>/HPO<sub>4</sub><sup>2-</sup> (0.05 M).

<sup>b</sup>  $[\text{I}^-]_{\text{added}} = 0.05$  M. <sup>c</sup> Parenthetical values were calculated using eqn. 6 and parameters listed in Table 3.

simple inverse-[H<sup>+</sup>] relationship. Both oxidations conform to a binomial expression, eqn. 6, where K<sub>1</sub>, the first acidity constant

$$k = \frac{a[\text{H}^+] + bK_1}{[\text{H}^+](K_1 + [\text{H}^+])} \quad (6)$$

of the reductant, is taken as  $7 \times 10^{-8}$  M.<sup>4</sup> Refinement of data in terms of eqn. 6 yields the parameters listed in Table 3. Eqn. 6 implies a reaction proceeding through two acidity levels, formed by loss of successive protons from the principal reductant species, which itself is negligibly reactive. The relevant rate law (eqn. 7) in which the rate constants  $k_{\text{HA}}$  and  $k_{\text{A}}$  refer to the

$$k = \frac{1}{[\text{H}^+] + K_1} \left[ K_1 k_{\text{HA}} + \frac{k_{\text{A}} K_1 K_2}{[\text{H}^+]} \right] \quad (7)$$

contributing hyponitrite anions, is algebraically equivalent to eqn. 6.

The resulting bimolecular rate constant,  $k_{\text{HA}}$ , pertaining to uninegative hyponitrite, is  $2.3 \times 10^4$  M<sup>-1</sup> s<sup>-1</sup> for reduction of I<sub>2</sub> and  $7.2 \times 10^2$  M<sup>-1</sup> s<sup>-1</sup> for I<sub>3</sub><sup>-</sup>. As expected, the more electrophilic uncharged X<sub>2</sub> molecule is the more reactive.

Extension of this reasoning to the inverse-[H<sup>+</sup>] term in eqn. 7 to calculate rates for dinegative hyponitrite requires division by K<sub>1</sub>K<sub>2</sub> and leads to  $k_{\text{A}}$  values of  $1.3 \times 10^{10}$  M<sup>-1</sup> s<sup>-1</sup> for I<sub>3</sub><sup>-</sup> and  $2 \times 10^{12}$  M<sup>-1</sup> s<sup>-1</sup> for I<sub>2</sub>, both of these exceeding the diffusion controlled limit<sup>21</sup> for bimolecular reactions in water. We therefore conclude, with some hesitancy (for the data are suggestive rather than compelling) that this kinetic contribution stems instead either from deprotonation of a different species (possibly a halogen–HN<sub>2</sub>O<sub>2</sub><sup>-</sup> precursor) or from intervention of a reactive halogen–OH<sup>-</sup> adduct.

Rates for the much more rapid oxidation by Br<sub>2</sub> could be measured only at higher acidities (0.005–0.20 M H<sup>+</sup>). The three-term rate law, eqn. 8 in Table 4, tells us that an additional path ( $k = 9 \times 10^3$  M<sup>-1</sup> s<sup>-1</sup>) involving the parent acid, H<sub>2</sub>N<sub>2</sub>O<sub>2</sub>, has become operative. Moreover, the calculated kinetic contribution for HN<sub>2</sub>O<sub>2</sub><sup>-</sup> ( $1 \times 10^6$  M<sup>-1</sup> s<sup>-1</sup>) is over 40 times that for the corresponding component for I<sub>2</sub>. Here again, refinement indicates a doubly deprotonated route, the nature of which is subject to the same questions as those for oxidations by I<sub>2</sub> and I<sub>3</sub><sup>-</sup>.

### Reactions with metal-center oxidants

Data for oxidations by [Fe(bipy)<sub>3</sub>]<sup>3+</sup> and [IrCl<sub>6</sub>]<sup>2-</sup> appear in Table 5. The simpler rate expressions (Table 4, eqns. 9 and 10) correspond to partition of the reductant into parent acid and monoanion, and with the Fe(III) chelate, only the latter

**Table 3** Kinetic parameters for oxidation of hyponitrous acid<sup>a</sup>

Oxidant	$\mu/M$	Parameters <sup>b</sup>
I <sub>2</sub>	0.20	$K_1 = 7 \times 10^{-8} \text{ M}; k_{\text{HA}} = (2.3 \pm 0.1) \times 10^4 \text{ M}^{-1} \text{ s}^{-1}; b = (3.39 \pm 0.03) \times 10^{-7} \text{ s}^{-1}$
I <sub>3</sub> <sup>-</sup>	0.20	$k_{\text{HA}} = (7.2 \pm 0.5) \times 10^2 \text{ M}^{-1} \text{ s}^{-1}; b = (2.60 \pm 0.03) \times 10^{-9} \text{ s}^{-1}$
Br <sub>2</sub>	0.20	$k_{\text{HA}} = (9.0 \pm 0.3) \times 10^3 \text{ M}^{-1} \text{ s}^{-1}; k_{\text{HA}} = (1.03 \pm 0.02) \times 10^6 \text{ M}^{-1} \text{ s}^{-1}; b = (7.2 \pm 0.4) \text{ M s}^{-1}$
[Fe(bipy) <sub>3</sub> ] <sup>3+</sup>	1.0	$k_{\text{HA}} = (4.0 \pm 0.1) \times 10^4 \text{ M}^{-1} \text{ s}^{-1}$
[IrCl <sub>6</sub> ] <sup>2-</sup>	0.50	$k_{\text{HA}} = 0.83 \pm 0.02 \text{ M}^{-1} \text{ s}^{-1}; k_{\text{HA}} = 413 \pm 3 \text{ M}^{-1} \text{ s}^{-1}$
[PtCl <sub>6</sub> ] <sup>2-</sup>	0.50	$k_{\text{HA}} = 33 \pm 3 \text{ M}^{-1} \text{ s}^{-1}$
[PdCl <sub>6</sub> ] <sup>2-</sup>	1.0	$k_1 = 8 \times 10^4 \text{ M}^{-1} \text{ s}^{-1}; k_2 = 4 \times 10^3 \text{ M}^{-2} \text{ s}^{-1}$

<sup>a</sup> Reactions were run at  $24.5 \pm 0.5$  °C;  $\mu$  was generally regulated using NaClO<sub>4</sub>/HClO<sub>4</sub>. <sup>b</sup> Parameters are those in the rate laws listed in Table 4. <sup>c</sup>  $K_1$ , the first acidity constant for hyponitrous acid, is taken as  $7.0 \times 10^{-8} \text{ M}$  throughout (ref. 8).

**Table 4** Rate laws for oxidations of hyponitrous acid

Oxidant (Ox)	$\lambda/\text{nm}$	Rate law <sup>a</sup>	Eqn. no.
I <sub>2</sub>	456	Rate = [Ox][Red] ( $K_1 k_{\text{HA}} + b[\text{H}^+]^{-1}$ ) ( $[\text{H}^+] + K_1$ ) <sup>-1</sup>	7' <sup>b</sup>
I <sub>3</sub> <sup>-</sup>	352		7' <sup>b</sup>
Br <sub>2</sub>	395	Rate = [Ox][Red] ( $k_{\text{HA}} + K_1 k_{\text{HA}} [\text{H}^+]^{-1} + b[\text{H}^+]^{-2}$ )	8 <sup>c</sup>
[Fe(bipy) <sub>3</sub> ] <sup>3+</sup>	525	Rate = [Ox][Red] ( $K_1 k_{\text{HA}} ([\text{H}^+] + K_1)^{-1}$ )	9
[IrCl <sub>6</sub> ] <sup>2-</sup>	489	Rate = [Ox][Red] ( $k_{\text{HA}} [\text{H}^+] + K_1 k_{\text{HA}} ([\text{H}^+] + K_1)^{-1}$ )	10
[PtCl <sub>6</sub> ] <sup>2-</sup>	270	Rate = [Ox][Red] $K_1 k_{\text{HA}} [\text{H}^+]^{-1}$	11 <sup>c</sup>
[PdCl <sub>6</sub> ] <sup>2-</sup>	340	Rate = [Ox][Red] ( $k_1 + k_{\text{Cl}}[\text{Cl}]$ )	12 <sup>d</sup>


<sup>a</sup>  $K_1$  is the first acidity constant for hyponitrous acid. Parameters  $k_{\text{HA}}$  and  $k_{\text{HA}}$  refer to kinetic contributions associated with unionized hyponitrous acid and its mononegative anion, HN<sub>2</sub>O<sub>2</sub><sup>-</sup>. <sup>b</sup> Eqn. 7' is an adaption of eqn. 7 in the text with the kinetically inadmissible  $K_1 K_2 [\text{H}^+]^{-1}$  term (see text) replaced by parameter  $b$ . <sup>c</sup> Oxidations examined with  $[\text{H}^+] \gg K_1$ . <sup>d</sup> Reaction rate independent of  $[\text{H}^+]$  in the range 0.1–1.0 M.

**Table 5** Kinetic data for oxidation of hyponitrous acid with [Fe(bipy)<sub>3</sub>]<sup>3+</sup> and with [IrCl<sub>6</sub>]<sup>2-</sup><sup>a</sup>

A H <sub>2</sub> N <sub>2</sub> O <sub>2</sub> + [Fe(bipy) <sub>3</sub> ] <sup>3+</sup> <sup>b</sup>		B H <sub>2</sub> N <sub>2</sub> O <sub>2</sub> + [IrCl <sub>6</sub> ] <sup>2-</sup> <sup>c</sup>	
pH	$10^{-3}k/\text{M}^{-1} \text{ s}^{-1}$ <sup>d</sup>	pH	$k/\text{M}^{-1} \text{ s}^{-1}$
5.02	0.38 (0.38)	4.76	3.4 (4.0)
5.38	0.75 (0.85)	5.07	6.6 (6.0)
5.60	1.36 (1.39)	5.56	13.9 (14.2)
5.90	2.5 (2.7)	5.89	28 (27)
6.08	3.5 (3.9)	6.13	50 (44)
6.30	5.2 (6.1)	6.33	77 (64)
6.56	9.5 (9.9)	6.68	95 (117)
6.62	10.0 (11.0)	7.06	$2.0 \times 10^2$ ( $2.0 \times 10^2$ )
7.20	27 (24)	7.23	$2.4 \times 10^2$ ( $2.3 \times 10^2$ )
7.50	28 (30)		

<sup>a</sup> Reactions were run at  $24.5 \pm 0.5$  °C. Solutions were buffered using HOAc/OAc<sup>-</sup> or H<sub>2</sub>PO<sub>4</sub><sup>-</sup>/HPO<sub>4</sub><sup>2-</sup> (0.05 M). <sup>b</sup>  $\mu = 1.0 \text{ M}$  (NaClO<sub>4</sub>). At pH values below 5.0, the system was complicated by the reaction of HNO<sub>2</sub> with [Fe(bipy)<sub>3</sub>]<sup>3+</sup>. <sup>c</sup>  $\mu = 0.50 \text{ M}$  (NaClO<sub>4</sub>). Rates were not significantly affected by addition of 0.01–0.10 M Cl<sup>-</sup>. <sup>d</sup> Parenthetical values were calculated using the rate equations in Table 4 and the parameters in Table 3.

contributes significantly. The substitution-inert Fe(III) complex may be assumed to be an outer-sphere 1e<sup>-</sup> acceptor, and this is probably the case also for [IrCl<sub>6</sub>]<sup>2-</sup>, for the reducing agent is devoid of a positive center that might utilize a chloro bridge. Since there is no evidence for the intervention of a transient which is generated or consumed on a time scale similar to that of the primary conversion, the overall rates may be taken to be determined by the initial act of electron transfer (e.g. eqn. 13), with subsequent steps being much more rapid.



The greater net rate observed for Fe[(bipy)<sub>3</sub>]<sup>3+</sup> may be attributed<sup>22</sup> in part to its more positive potential ( $E^\circ_{\text{Fe(III)}} = 1.11 \text{ V}$ ;  $E^\circ_{\text{Ir(IV)}} = 0.87 \text{ V}$ ),<sup>19,20</sup> in part to its more rapid self-exchange ( $10^{8.5}$  vs.  $10^{5.4} \text{ M}^{-1} \text{ s}^{-1}$ ),<sup>19</sup> and, to a lesser extent, to partial anion-cation association in the Fe(III) system. The combination of these effects generates<sup>22</sup> a calculated kinetic advantage of  $10^4$




for [Fe(bipy)<sub>3</sub>]<sup>3+</sup>. Agreement with the observed ratio, ( $k_{\text{Fe(III)}}/k_{\text{Ir(IV)}} = 10^{2.0}$ , Table 5), although approximate, is comparable to agreements resulting from similar applications of the Marcus model<sup>23</sup> to other redox systems.<sup>24</sup>

A formal potential for the 1e<sup>-</sup> oxidation of HN<sub>2</sub>O<sub>2</sub><sup>-</sup> has not yet been reported but probably lies near 0.70 V, slightly below that of Fe<sup>III,II</sup>(aq). This estimate, in conjunction with the “ $k_{\text{HA}}$ ” values in Table 3, leads to calculated self-exchange rates in the range  $10^{-4.8 \pm 1.5} \text{ M}^{-1} \text{ s}^{-1}$  for (HN<sub>2</sub>O<sub>2</sub>)<sup>-1,0</sup>.

Rate constants for oxidations by the 2e<sup>-</sup> reagents [PtCl<sub>6</sub>]<sup>2-</sup> ( $E^\circ_{\text{Pt(IV)}} = 0.73 \text{ V}$ ) and [PdCl<sub>6</sub>]<sup>2-</sup> ( $E^\circ_{\text{Pd(IV)}} = 1.47 \text{ V}$ )<sup>25</sup> are seen to be similar to those of single electron oxidants of similar 1e<sup>-</sup> potential. Outer-sphere 2e<sup>-</sup> transactions, if they occur at all, should be considerably slower, reflecting a more severe Franck–Condon barrier. For these oxidants, however, reaction through a chloro bridge is reasonable, but a sequence of 1e<sup>-</sup> transfers (with the initial step rate determining), cannot be excluded. Evidence for an odd-electron Pt(III) transient has been cited for other reactions.<sup>26,27</sup>

Does the oxidation by I<sub>2</sub> involve successive 1e<sup>-</sup> transactions? We think not. The high rate constant for the principal path,  $2 \times 10^4 \text{ M}^{-1} \text{ s}^{-1}$ , considered in conjunction with the weakly oxidizing potential for I<sub>2</sub>/I<sub>2</sub><sup>-</sup> (0.21 V),<sup>28</sup> points instead to utilization of an iodine bridge (I<sup>+</sup> transfer) (eqn. 14), analogous to that demonstrated<sup>29</sup> for the HOCl–NO<sub>2</sub><sup>-</sup> reaction.

Subsequent hydrolysis of the N–I species (eqn. 15) and further attack on the trioxodinitrate intermediate (N<sub>2</sub>O<sub>3</sub><sup>2-</sup>) (eqn. 16)



may be taken to be rapid<sup>10c</sup> in relation to the proposed initial step. Analogous halogen-bridged sequences may be assigned to the slower oxidation by I<sub>3</sub><sup>-</sup> and, at least in part, to the more rapid reaction of Br<sub>2</sub>. As with oxidations of trioxodinitrate (hyponitrate),<sup>10c</sup> cleavage of the N=N double bond to yield the predominant product, nitrous acid, probably reflects the diminution of bonding action by the attached strongly electro-negative oxygen atoms.

In summary, hyponitrite, like trioxodinitrate,<sup>10c</sup> exhibits considerable versatility of action in its redox behavior. Reaction sequences may be initiated by transfer of a single electron to an outer-sphere type oxidant when the latter features a sufficiently positive potential. Oxidants having a somewhat lower potential may operate *via* 2e<sup>-</sup> paths provided that the redox step also entails transfer of a bridging atom to the reductant. With Br<sub>2</sub>, both paths may compete.

## Acknowledgements

We are grateful to the National Science Foundation for support of this work (grant CHE-9714981) and to Mrs Arla McPherson for technical assistance.

## References

- 1 S. Swavey and E. S. Gould, *Inorg. Chem.*, in the press.
- 2 E. Divers, *Proc. R. Soc. London, A*, 1871, **29**, 425.
- 3 (a) For a critical summary of the older literature, see: D. M. Yost and H. Russell, *Systematic Inorganic Chemistry of the Fifth and Sixth Group Nonmetallic Elements*, Prentice-Hall, New York, 1944, pp. 52–57; (b) For a more recent review, see: M. N. Hughes, *Q. Rev., Chem. Soc.*, 1968, **22**, 1.
- 4 M. N. Hughes and G. Stedman, *J. Chem. Soc.*, 1963, 1239.
- 5 D. J. Millen, C. N. Polydoropoulos and D. Watson, *J. Chem. Soc.*, 1960, 687.
- 6 M. N. Hughes, *J. Inorg. Nucl. Chem.*, 1967, **29**, 1376; G. E. McGraw, D. L. Bernitt and I. C. Hisatsune, *Spectrochim. Acta, Part A*, 1967, **23**, 25.
- 7 See, for example: T. Hofmann and H. Lees, *Biochem. J.*, 1954, **54**, 579.
- 8 (a) Y. Osajima and K. Yamafuji, *Enzymol.*, 1962, **24**, 57; (b) R. T. St. John and T. C. Hollocher, *J. Biol. Chem.*, 1977, **252**, 212.
- 9 See, for example: D. S. Frear and R. C. Burrell, *Plant Physiol.*, 1957, **33**, 105; D. P. Huckelsby and E. J. Hewitt, *Biochem. J.*, 1970, **119**, 615; T. C. Hollocher, E. Garber, A. J. Cooper and R. E. Reiman, *J. Biol. Chem.*, 1980, **255**, 5027.
- 10 (a) V. S. Srinivasan and E. S. Gould, *Inorg. Chem.*, 1981, **20**, 3176; (b) N. Rajasekar, R. Subramaniam and E. S. Gould, *Inorg. Chem.*, 1983, **22**, 1100; (c) A. M. Al-Ajlouni and E. S. Gould, *Inorg. Chem.*, 1999, **38**, 1592.
- 11 Work before 1968 has been reviewed by Hughes.<sup>3b</sup> For more recent studies, see: M. J. Akhtar; J. A. Balschi and F. T. Bonner, *Inorg. Chem.*, 1982, **21**, 2216; J. D. Abata, M. P. Dziobak, M. Nachbor and G. D. Mendenhall, *J. Phys. Chem.*, 1989, **93**, 3368; N. Arulsamy, D. S. Bohle, J. A. Imonigie and E. S. Sagan, *Inorg. Chem.*, 1999, **38**, 2716; E. L. Loechler, A. M. Schneider, D. B. Schwartz and T. C. Hollocher, *J. Am. Chem. Soc.*, 1987, **109**, 3076.
- 12 M. R. Goyal, R. K. Mittal, Y. K. Gupta, *Indian J. Chem., Sect. A*, 1988, **27**, 487 and 584; M. R. Goyal, P. Bhatnagar, R. K. Mittal and Y. K. Gupta, *Indian J. Chem., Sect. A*, 1989, **28**, 382; P. Bhatnagar, R. K. Mittal and Y. K. Gupta, *J. Chem. Soc., Dalton Trans.*, 1990, 3669; P. Bhatnagar, R. K. Mittal and Y. K. Gupta, *Indian J. Chem., Sect. A*, 1991, **30**, 187; 1993, **32**, 195.
- 13 M. N. Hughes and G. Stedman, *J. Chem. Soc.*, 1963, 4230; J. R. Buchholtz and R. E. Powell, *J. Am. Chem. Soc.*, 1965, **87**, 2350; M. N. Hughes, P. E. Wimbledon and G. Stedman, *J. Chem. Soc., Dalton Trans.*, 1989, 533.
- 14 F. T. Bonner, C. E. Donald and M. N. Hughes, *J. Chem. Soc., Dalton Trans.*, 1989, 527.
- 15 C. N. Polydoropoulos and S. D. Voliotis, *Anal. Chim. Acta*, 1968, **40**, 170; C. N. Conner, C. E. Donald, M. N. Hughes and C. Sami, *Polyhedron*, 1989, **8**, 2621.
- 16 S. Mandal and E. S. Gould, *Inorg. Chem.*, 1995, **34**, 3993.
- 17 S. P. Ghosh, M. C. Ghosh and E. S. Gould, *Inorg. Chim. Acta*, 1994, **225**, 83.
- 18 See, for example: A. Al-Ajlouni and J. H. Espenson, *J. Org. Chem.*, 1996, **61**, 3969.
- 19 M. C. Ghosh and E. S. Gould, *J. Am. Chem. Soc.*, 1993, **115**, 3167.
- 20 P. George, I. H. Hanania and D. H. Irvine, *J. Chem. Soc.*, 1957, 3048.
- 21 See, for example: J. H. Espenson, *Chemical Kinetics and Reaction Mechanisms*, McGraw-Hill, New York, 2nd edn., 1995, p. 201.
- 22 See, for example: G. Lappin, *Redox Mechanisms in Inorganic Chemistry*, Ellis Norwood, New York, 1994, ch. 2.
- 23 R. A. Marcus, *Annu. Rev. Phys. Chem.*, 1964, **15**, 155.
- 24 See, for example: D. H. Macartney, *Inorg. Chem.*, 1986, **25**, 2222; K. R. Howes, A. Bakac and J. H. Espenson, *Inorg. Chem.*, 1988, **27**, 791.
- 25 F. Colon, in *Standard Potentials in Aqueous Solution*, ed. A. J. Bard, R. Parsons and J. Jordon, Marcel Dekker, New York, 1985, ch. 12.
- 26 C. S. Glennon, T. D. Hand and A. G. Sykes, *J. Chem. Soc., Dalton Trans.*, 1980, 19.
- 27 A. J. Peloso, *J. Chem. Soc., Dalton Trans.*, 1987, 1473.
- 28 D. M. Stanbury, *Adv. Inorg. Chem.*, 1989, **33**, 59.
- 29 D. W. Johnson and D. W. Margerum, *Inorg. Chem.*, 1991, **30**, 4845.