

Synthesis, characterisation and reactions of ruthenium(II) complexes based upon $[\text{RuL}^3]^{2+}$ ($\text{L}^3 =$ tripodal triseleno- or tritelluro-ether) fragments. Structures of $[\text{RuCl}_2(\text{PPh}_3)\{\text{MeC}(\text{CH}_2\text{SeMe})_3\}]$ and $[\text{RuCl}_2(\text{dmsO})\{\text{MeC}(\text{CH}_2\text{SeMe})_3\}]$

William Levason, Simon D. Orchard and Gillian Reid

Department of Chemistry, University of Southampton, Southampton, UK SO17 1BJ

Received 15th September 2000, Accepted 27th October 2000

First published as an Advance Article on the web 1st December 2000

The reaction of $[\text{RuCl}_2(\text{PPh}_3)_3]$ with tripodal Group 16 donor ligands $\text{L}^3 \{\text{MeC}(\text{CH}_2\text{E})_3$ ($\text{E} = \text{Se}$ or Te) and $\text{MeC}(\text{CH}_2\text{TePh})_3\}$ gave $[\text{RuCl}_2(\text{PPh}_3)\text{L}^3]$ complexes which have been characterised by elemental analysis, IR and NMR spectroscopy and ES^+ mass spectrometry. The structure of $[\text{RuCl}_2(\text{PPh}_3)\{\text{MeC}(\text{CH}_2\text{SeMe})_3\}]$ reveals a distorted octahedral geometry with a facially co-ordinated triselenoether. The reaction of $[\text{RuCl}_2(\text{dmsO})_4]$ with L^3 gave $[\text{RuCl}_2(\text{dmsO})\text{L}^3]$ which have similarly been characterised, including a crystal structure of $[\text{RuCl}_2(\text{dmsO})\{\text{MeC}(\text{CH}_2\text{SeMe})_3\}]$, which is *fac*-octahedral with S-bonded dmsO. The $[\text{RuCl}_2(\text{dmsO})\text{L}^3]$ species react with $\text{Ag}[\text{CF}_3\text{SO}_3]$ in MeCN to produce $[\text{Ru}(\text{MeCN})_3\text{L}^3]^{2+}$ ($\text{L}^3 = \text{MeC}(\text{CH}_2\text{SeMe})_3$ or $\text{MeC}(\text{CH}_2\text{TePh})_3$). The MeCN is labile and readily replaced by a second tridentate ligand to give mixed tripod ligand complexes including $[\text{Ru}\{\text{MeC}(\text{CH}_2\text{SeMe})_3\}\{\text{MeC}(\text{CH}_2\text{SeMe})_3\}][\text{CF}_3\text{SO}_3]_2$ and $[\text{Ru}\{\text{MeC}(\text{CH}_2\text{SeMe})_3\}\{\text{MeC}(\text{CH}_2\text{TePh})_3\}][\text{CF}_3\text{SO}_3]_2$. Attempts to generate hydride species by reaction of $[\text{Ru}(\text{MeCN})_3\text{L}^3]^{2+}$ with NaBH_4 in ethanol bring about decomposition.

Introduction

The development of ruthenium based catalysts is a very active area. Good examples are provided by ruthenium(II) complexes of the tripodal triphosphine, $\text{MeC}(\text{CH}_2\text{PPh}_2)_3$ (triphos),¹ notably the work of Bianchini *et al.* who have developed ruthenium (and Rh and Ir) systems which mimic metal catalysed hydrodesulfurisation processes.²⁻⁴ The chemistry of Ru^{II} with thioether ligands, in particular the macrocyclic [9]aneS₃, has received considerable attention,⁵⁻⁸ with complexes such as $[\text{Ru}(\text{[9]aneS}_3)_2]^{2+}$ and $[\text{RuCl}(\text{[9]aneS}_3)(\text{Me}_2\text{SO})_2]^+$ being reported. The species $[\text{RuCl}_2(\text{PPh}_3)(\text{[9]aneS}_3)]$ and $[\text{RuX}(\text{CS})(\text{PPh}_3)(\text{[9]aneS}_3)][\text{PF}_6]$ ($\text{X} = \text{H}, \text{Cl}, \text{SCN}$ or $\text{C}_6\text{H}_4\text{Me-4}$) have been prepared as part of an investigation into organometallic macrocycle chemistry.⁶ These studies have also reported the σ -vinyl and σ -aryl complexes $[\text{Ru}(\text{CH}=\text{CH}_2)(\text{CO})(\text{PPh}_3)(\text{[9]aneS}_3)]^+$ and $[\text{Ru}(\text{C}_6\text{H}_4\text{Me-4})(\text{CO})(\text{PPh}_3)(\text{[9]aneS}_3)]^+$.

The chemistry of ruthenium(II) with the heavier seleno- and telluro-ether ligands has generally been limited to the preparation and characterisation of bidentate analogues including $[\text{RuCl}_2(\text{L-L})_2]$ ($\text{L-L} =$ diseleno- or ditelluro-ether) and $[\text{RuCl}(\text{PPh}_3)(\text{L-L})][\text{PF}_6]$ ($\text{L-L} =$ ditelluroether).^{9,10} The crystal structures of the macrocyclic *cis*- $[\text{RuCl}_2(\text{[16]aneSe}_4)]$ and *trans*- $[\text{RuCl}(\text{PPh}_3)(\text{[16]aneSe}_4)][\text{PF}_6]$ have also been reported.¹¹ The recent successful preparation of low and medium oxidation state rhodium and iridium organometallic complexes with the Group 16 tripodal ligands $\text{L}^3 \{\text{L}^3 = \text{MeC}(\text{CH}_2\text{E})_3$ ($\text{E} = \text{Se}$ or Te) or $\text{MeC}(\text{CH}_2\text{TePh})_3\}$ ¹² led us to investigate their reaction chemistry on ruthenium(II) centres. Our investigation into homoleptic platinum metal complexes with L^3 reported the synthesis of the complexes $[\text{Ru}(\text{L}^3)_2]^{2+}$ ($\text{L}^3 = \text{MeC}(\text{CH}_2\text{E})_3$ ($\text{E} = \text{S}, \text{Se}$ or Te) or $\text{MeC}(\text{CH}_2\text{TePh})_3$).¹³ Here we report on the preparation and reactions of species containing the $[\text{RuL}^3]^{2+}$ fragment.

Experimental

The complexes $[\text{RuCl}_2(\text{PPh}_3)_3]^{14}$ and $[\text{RuCl}_2(\text{dmsO})_4]^{15}$ were

prepared by literature procedures, as were the ligands $\text{MeC}(\text{CH}_2\text{SeMe})_3$,¹⁶ $\text{MeC}(\text{CH}_2\text{TeMe})_3$,¹⁷ $\text{MeC}(\text{CH}_2\text{TePh})_3$,¹⁸ and $\text{MeC}(\text{CH}_2\text{TePh})_3$.¹⁹ Physical measurements were made as described previously.^{10,12}

Preparations

$[\text{RuCl}_2(\text{PPh}_3)\{\text{MeC}(\text{CH}_2\text{SeMe})_3\}]$. $[\text{RuCl}_2(\text{PPh}_3)_3]$ (208 mg, 2.2×10^{-4} mol) was added to $\text{MeC}(\text{CH}_2\text{SeMe})_3$ (77 mg, 2.2×10^{-4} mol) in dry CH_2Cl_2 (40 cm³) and stirred at room temperature for 18 h to give an orange solution. This was reduced to ca. 2 cm³ *in vacuo* and diethyl ether (10 cm³) added to precipitate an orange solid. Yield 113 mg, 66% (Found: C, 39.9; H, 4.4. Calc. for $\text{C}_{26}\text{H}_{33}\text{Cl}_2\text{PRuSe}_3$: C, 39.8; H, 4.2%). ¹H NMR (CDCl_3 , 300 K): δ 1.37 (s, 1H, CCH₃), 1.6–2.0 (m, 3H, SeCH₃), 2.3–2.6 (m, 2H, SeCH₂) and 7.2–8.2 (m, 5H, Ph). ⁷⁷Se-¹H NMR ($\text{CH}_2\text{Cl}_2\text{-CH}_3\text{OH-CDCl}_3$, 300 K): δ 165, 168, 171, 245, 247, 272 and 275. ³¹P-¹H NMR ($\text{CH}_2\text{Cl}_2\text{-CH}_3\text{OH-CDCl}_3$, 300 K): δ 35.2 and 34.4. ES^+ (MeCN): $m/z = 792$ and 751; calc. for $[\text{RuCl}_2(\text{PPh}_3)\{\text{MeC}(\text{CH}_2\text{SeMe})_3\}(\text{NCMe})]^+$ 794 and $[\text{RuCl}_2(\text{PPh}_3)\{\text{MeC}(\text{CH}_2\text{SeMe})_3\}]^+$ 753. IR: 3050w, 2962w, 2940w, 1481m, 1433m, 1358s, 1090s, 989m, 907w, 834m, 746m, 697s, 614w, 523s, 499m, 459m, 422m, 290m and 216m cm⁻¹.

$[\text{RuCl}_2(\text{PPh}_3)\{\text{MeC}(\text{CH}_2\text{TeMe})_3\}]$. This was prepared similarly as a light brown solid (61%) (Found: C, 31.9; H, 3.5. Calc. for $\text{C}_{26}\text{H}_{33}\text{Cl}_2\text{PRuTe}_3 \cdot \text{CH}_2\text{Cl}_2$: C, 31.9; H, 3.3%). ES^+ (MeCN): $m/z = 938$ and 897; calc. for $[\text{RuCl}_2(\text{PPh}_3)\{\text{MeC}(\text{CH}_2\text{TeMe})_3\}(\text{NCMe})]^+$ 944 and $[\text{RuCl}_2(\text{PPh}_3)\{\text{MeC}(\text{CH}_2\text{TeMe})_3\}]^+$ 903. IR: 3051w, 2922w, 1481m, 1432s, 1360s, 1267w, 1217w, 1190w, 1090s, 998m, 835s, 744s, 697s, 614w, 526s, 459w, 309m and 223m cm⁻¹.

$[\text{RuCl}_2(\text{PPh}_3)\{\text{MeC}(\text{CH}_2\text{TePh})_3\}]$. This was prepared similarly as an orange solid (72%) (Found: C, 43.7; H, 3.1. Calc. for $\text{C}_{41}\text{H}_{39}\text{Cl}_2\text{PRuTe}_3$: C, 44.1; H, 3.5%). ¹²⁵Te-¹H NMR ($\text{CH}_2\text{Cl}_2\text{-CH}_3\text{OH-CDCl}_3$, 300 K): δ 566, 570, 741, 742 and

770. ^{31}P - $\{^1\text{H}\}$ NMR (CH_2Cl_2 - CH_3OH - CDCl_3 , 300 K): δ 25.6. ES^+ (MeCN): $m/z = 1122$ and 1081; calc. for $[\text{Ru}^{102}\text{Ru}^{35}\text{Cl}(\text{PPh}_3)_3\{\text{MeC}(\text{CH}_2^{130}\text{TePh})_3\}(\text{NCMe})]^+$ 1130 and $[\text{Ru}^{102}\text{Ru}^{35}\text{Cl}(\text{PPh}_3)_3\{\text{MeC}(\text{CH}_2^{130}\text{TePh})_3\}]^+$ 1089. IR: 3052w, 1571m, 1476m, 1432s, 1358s, 1263w, 1187w, 1090s, 1017m, 998m, 834w, 797w, 735s, 694s, 524s, 456m and 250 cm^{-1} .

[RuCl₂(dmsO){MeC(CH₂SeMe)₃]. $[\text{RuCl}_2(\text{dmsO})_4]$ (40 mg, 8.3×10^{-5} mol) was added to dry toluene (40 cm^3) and heated to 100 °C for 10 min. The resulting suspension was allowed to cool, $\text{MeC}(\text{CH}_2\text{SeMe})_3$ (29 mg, 8.3×10^{-5} mol) in toluene (5 cm^3) added and the mixture heated to 100 °C for 24 h. The resulting precipitate was filtered off and washed with diethyl ether (10 cm^3) to give an orange solid (30 mg, 60%) (Found: C, 20.3; H, 3.8. Calc. for $\text{C}_{10}\text{H}_{24}\text{Cl}_2\text{ORuSSe}_3$: C, 20.0; H, 4.0%). ^1H NMR (CDCl_3 , 300 K): δ 1.34 (s, 1H, CCH_3), 2.1–2.6 (m, 3H, SeCH_3), 2.61 (s, 2H, CH_3S) and 3.35–3.51 (m, 2H, SeCH_2). ^{77}Se - $\{^1\text{H}\}$ NMR (CH_2Cl_2 - CDCl_3 , 300 K): δ 168, 170, 218, 219, 229 and 244. FAB MS (3-nitrobenzyl alcohol): $m/z = 601$, 567 and 523; calc. for $[\text{Ru}^{102}\text{Ru}^{35}\text{Cl}_2(\text{dmsO})\{\text{MeC}(\text{CH}_2^{80}\text{SeMe})_3\}]^+$ 604, $[\text{Ru}^{102}\text{Ru}^{35}\text{Cl}(\text{dmsO})\{\text{MeC}(\text{CH}_2^{80}\text{SeMe})_3\}]^+$ 569 and $[\text{Ru}^{102}\text{Ru}^{35}\text{Cl}_2\{\text{MeC}(\text{CH}_2^{80}\text{SeMe})_3\}]^+$ 526. IR: 2950w, 1413m, 1358s, 1262m, 1076s, 1017m, 924w, 834w, 802w, 713w, 678w, 614w, 540w, 427m and 238 cm^{-1} .

[RuCl₂(dmsO){MeC(CH₂TeMe)₃]. This was prepared similarly as a brown solid (61%) (Found: C, 16.5; H, 3.5. Calc. for $\text{C}_{10}\text{H}_{24}\text{Cl}_2\text{ORuSTe}_3$: C, 16.1; H, 3.2%). ^1H NMR (CDCl_3 , 300 K): δ 1.26 (s, 1H, CCH_3), 2.1–2.4 (m, 3H, TeCH_3), 2.63 (s, 2H, CH_3S) and 3.40–3.55 (m, 2H, TeCH_2). ^{130}Te - $\{^1\text{H}\}$ NMR (CH_2Cl_2 - CDCl_3 , 300 K): δ 222 see text. FAB MS (3-nitrobenzyl alcohol): $m/z = 748$; calc. for $[\text{Ru}^{102}\text{Ru}^{35}\text{Cl}_2(\text{dmsO})\{\text{MeC}(\text{CH}_2^{130}\text{TeMe})_3\}]^+$ 754. IR: 2925w, 1359s, 1095s, 1018m, 996m, 835m, 682w, 613w, 536w, 425w and 236 cm^{-1} .

[RuCl₂(dmsO){MeC(CH₂TePh)₃]. This was prepared similarly except an orange solution was produced upon heating for 24 h. The solvent volume was reduced *in vacuo* to 5 cm^3 and diethyl ether added to give an orange solid (69%) (Found: C, 31.8; H, 3.3. Calc. for $\text{C}_{25}\text{H}_{30}\text{Cl}_2\text{ORuSTe}_3$: C, 32.2; H, 3.2%). ^1H NMR (CDCl_3 , 300 K): δ 1.26 (s, 1H, CCH_3), 2.57 (s, 2H, CH_3S), 3.10–3.50 (m, 2H, TeCH_2) and 6.8–8.2 (m, 5H, TePh). ^{130}Te - $\{^1\text{H}\}$ NMR (CH_2Cl_2 - CDCl_3 , 300 K): δ 570, 677 and 737. FAB MS (3-nitrobenzyl alcohol): $m/z = 821$; calc. for $[\text{Ru}^{102}\text{Ru}^{35}\text{Cl}\{\text{MeC}(\text{CH}_2^{130}\text{TePh})_3\}]^+$ 827. IR: 3050w, 2951w, 1570w, 1475m, 1432m, 1359s, 1262m, 1089s, 1017s, 998s, 802m, 740m, 693m, 612w, 541w, 455w, 421w and 253 cm^{-1} .

[Ru(NCMe)₃{MeC(CH₂SeMe)₃][CF₃SO₃]₂. $[\text{RuCl}_2\{\text{dmsO}\}\{\text{MeC}(\text{CH}_2\text{SeMe})_3\}]$ (34 mg, 5.7×10^{-5} mol) was added to AgCF_3SO_3 (29 mg, 1.1×10^{-4} mol) in MeCN (40 cm^3). The mixture was refluxed for 2 h, cooled and filtered to remove the precipitated AgCl. The solvent volume was reduced *in vacuo* to 2 cm^3 and diethyl ether added to give a light yellow solid (40 mg, 80%) (Found: C, 21.9; H, 3.2; N, 4.8. Calc. for $\text{C}_{16}\text{H}_{27}\text{F}_6\text{N}_3\text{O}_6\text{RuS}_2\text{Se}_3$: C, 22.0; H, 3.1; N, 4.8%). ^1H NMR ($(\text{CD}_3)_2\text{CO}$, 300 K): δ 1.47 (s, 1H, CCH_3), 2.42 (s, 3H, NCCH_3), 2.51 (s, 3H, SeCH_3) and 2.85 (m, 2H, SeCH_2). ^{77}Se - $\{^1\text{H}\}$ NMR (MeCN- CDCl_3 , 300 K): δ 159. ES^+ (MeCN): $m/z = 288$ and 267; calc. for $[\text{Ru}^{102}\text{Ru}(\text{NCMe})_3\{\text{MeC}(\text{CH}_2^{80}\text{SeMe})_3\}]^{2+}$ 290 and $[\text{Ru}^{102}\text{Ru}(\text{NCMe})_2\{\text{MeC}(\text{CH}_2^{80}\text{SeMe})_3\}]^{2+}$ 269. IR: 2312w, 1360s, 1263s, 1225m, 1150m, 1098m, 1032m, 991w, 836w, 638s and 518 cm^{-1} .

[Ru(NCMe)₃{MeC(CH₂TePh)₃][CF₃SO₃]₂. This was prepared similarly as an orange solid (53%) (Found: C, 30.5; H, 2.4; N, 3.3. Calc. for $\text{C}_{31}\text{H}_{33}\text{F}_6\text{N}_3\text{O}_6\text{RuS}_2\text{Te}_3$: C, 30.9; H, 2.7; N, 3.5%). ^1H NMR ($(\text{CD}_3)_2\text{CO}$, 300 K): δ 1.91 (s, 1H, CCH_3), 2.29 (s, 3H, NCCH_3), 2.90 (s, 2H, TeCH_2) and 7.5–7.8 (m, 5H, TePh). ^{125}Te - $\{^1\text{H}\}$ NMR (MeCN- CDCl_3 , 300 K): δ 531.

ES^+ (MeCN): $m/z = 453$, 432 and 414; calc. for $[\text{Ru}^{102}\text{Ru}(\text{NCMe})_3\{\text{MeC}(\text{CH}_2^{130}\text{TePh})_3\}]^{2+}$ 458, $[\text{Ru}^{102}\text{Ru}(\text{NCMe})_2\{\text{MeC}(\text{CH}_2^{130}\text{TePh})_3\}]^{2+}$ 437 and $[\text{Ru}^{102}\text{Ru}(\text{NCMe})\{\text{MeC}(\text{CH}_2^{130}\text{TePh})_3\}]^{2+}$ 417. IR: 2315w, 1478w, 1435w, 1358m, 1276s, 1154s, 1093m, 1032s, 998m, 834w, 745m, 693m, 638s, 574w, 518m and 458 cm^{-1} .

[Ru{MeC(CH₂SMe)₃}{MeC(CH₂SeMe)₃][CF₃SO₃]₂. $\text{MeC}(\text{CH}_2\text{SMe})_3$ (17 mg, 7.9×10^{-5} mol) was added to $[\text{Ru}(\text{NCMe})_3\{\text{MeC}(\text{CH}_2\text{SeMe})_3\}][\text{CF}_3\text{SO}_3]_2$ (69 mg, 7.9×10^{-5} mol) in CH_3OH (30 cm^3) and the reaction mixture refluxed for 18 h. After cooling the solvent volume was reduced *in vacuo* to 5 cm^3 and diethyl ether added to precipitate a light yellow solid (60 mg, 79%) (Found: C, 22.5; H, 3.5. Calc. for $\text{C}_{18}\text{H}_{36}\text{F}_6\text{O}_6\text{RuS}_5\text{Se}_3$: C, 22.5; H, 3.8%). ^1H NMR ($(\text{CD}_3)_2\text{CO}$, 300 K): δ 1.26 (s, 1H, $\text{CH}_3\text{C}(\text{CH}_2\text{SCH}_3)_3$), 1.38 (s, 1H, $\text{CH}_3\text{C}(\text{CH}_2\text{SeCH}_3)_3$), 2.34 (s, 3H, SeCH_3), 2.52 (s, 3H, SCH_3) and 2.7–2.9 (m, 4H, SeCH_2 , SCH_2). ^{77}Se - $\{^1\text{H}\}$ NMR (MeNO₂- CDCl_3 , 300 K): δ 123. ES^+ (MeCN): $m/z = 811$ and 331; calc. for $[\text{Ru}^{102}\text{Ru}\{\text{MeC}(\text{CH}_2\text{SMe})_3\}\{\text{MeC}(\text{CH}_2^{80}\text{SeMe})_3\}][\text{CF}_3\text{SO}_3]_2$ 815 and $[\text{Ru}^{102}\text{Ru}\{\text{MeC}(\text{CH}_2\text{SMe})_3\}\{\text{MeC}(\text{CH}_2^{80}\text{SeMe})_3\}]^{2+}$ 333. IR: 2940w, 1461w, 1420m, 1358m, 1262s, 1227m, 1166m, 1096m, 1032s, 976w, 639s and 518 cm^{-1} .

[Ru{MeC(CH₂TePh)₃}{MeC(CH₂SeMe)₃][CF₃SO₃]₂. This was similarly prepared *via* the reaction of $[\text{Ru}(\text{NCMe})_3\{\text{MeC}(\text{CH}_2\text{TePh})_3\}][\text{CF}_3\text{SO}_3]_2$ with $\text{MeC}(\text{CH}_2\text{SeMe})_3$ (73%) (Found: C, 27.4; H, 2.8. Calc. for $\text{C}_{33}\text{H}_{42}\text{F}_6\text{O}_6\text{RuS}_2\text{Se}_3\text{Te}_3$: C, 27.6; H, 2.9%). ^1H NMR ($(\text{CD}_3)_2\text{CO}$, 300 K): δ 1.16 (s, 2H, CH_3), 2.06 (s, 3H, SeCH_3), 2.4–2.9 (m, 4H, SeCH_2 , TeCH_2) and 7.5–8.0 (m, 5H, TePh). ^{77}Se - $\{^1\text{H}\}$ NMR (MeNO₂- CDCl_3 , 300 K): δ 128. ^{125}Te - $\{^1\text{H}\}$ NMR (MeNO₂- CDCl_3 , 300 K): δ 485. ES^+ (MeCN): $m/z = 569$; calc. for $[\text{Ru}^{102}\text{Ru}\{\text{MeC}(\text{CH}_2\text{TePh})_3\}\{\text{MeC}(\text{CH}_2^{80}\text{SeMe})_3\}]^{2+}$ 573. IR: 2929w, 1572w, 1476w, 1433w, 1358s, 1262s, 1224m, 1156m, 1096m, 1030s, 997m, 910w, 834w, 738m, 693m, 638s, 573w, 518m and 456 cm^{-1} .

X-Ray crystallographic studies

Details of the crystallographic data collection and refinement parameters for $[\text{RuCl}_2(\text{PPh}_3)_3\{\text{MeC}(\text{CH}_2\text{SeMe})_3\}]$ and $[\text{RuCl}_2(\text{dmsO})\{\text{MeC}(\text{CH}_2\text{SeMe})_3\}]$ are given in Table 1. The crystals were grown *via* vapour diffusion of diethyl ether into a solution of $[\text{RuCl}_2(\text{PPh}_3)_3\{\text{MeC}(\text{CH}_2\text{SeMe})_3\}]$ in CH_2Cl_2 -MeOH and by slow evaporation of CH_2Cl_2 from a solution of $[\text{RuCl}_2(\text{dmsO})\{\text{MeC}(\text{CH}_2\text{SeMe})_3\}]$ in CH_2Cl_2 -MeOH. Data collection used a Rigaku AFC7S four circle diffractometer operating at 150 K, with graphite-monochromated Mo- $\text{K}\alpha$ X-radiation ($\lambda = 0.71073$ Å). Structure solution and refinement were routine.^{20,21} Crystal data are given in Table 1, and selected bond lengths and angles in Tables 2 and 3.

CCDC reference number 186/2252.

See <http://www.rsc.org/suppdata/dt/b0/b007487j/> for crystallographic files in .cif format.

Results and discussion

[RuCl₂(PPh₃)L³]

The reaction of $[\text{RuCl}_2(\text{PPh}_3)_3]$ with ditelluroether ligands forms $[\text{RuCl}(\text{PPh}_3)(\text{L}-\text{L})][\text{PF}_6]$ ($\text{L}-\text{L} = \text{RTe}(\text{CH}_2)_2\text{TeR}$, $\text{R} = \text{Me}$ or Ph , and $\text{C}_6\text{H}_4(\text{TeMe})_2$ -*o*).¹⁰ We were interested in the reaction of Group 16 tripodal ligands with $[\text{RuCl}_2(\text{PPh}_3)_3]$ and their chemistry, since such species would allow the study of complexes containing both phosphine and Group 16 donors, and provide complexes containing the $[\text{RuL}^3]^{2+}$ ($\text{L}^3 = \text{MeC}(\text{CH}_2\text{EMe})_3$ ($\text{E} = \text{Se}$ or Te) or $\text{MeC}(\text{CH}_2\text{TePh})_3$) fragment, upon which further chemistry may be undertaken. Reaction of $[\text{RuCl}_2(\text{PPh}_3)_3]$ with 1 mol equivalent of L^3 in CH_2Cl_2 at room temperature gave an orange (selenoether) or brown (telluroether) solution. After reduction of the solvent volume and addition of diethyl ether, the complexes $[\text{RuCl}_2(\text{PPh}_3)\text{L}^3]$

Table 1 Crystallographic data

	[RuCl ₂ (PPh ₃){MeC(CH ₂ SeMe) ₃ }	[RuCl ₂ (dmsO){MeC(CH ₂ SeMe) ₃ }
Formula	C ₂₆ H ₃₃ Cl ₂ PRuSe ₃	C ₁₀ H ₂₄ Cl ₂ ORuSSe ₃
Formula weight	785.38	601.22
Crystal system	Orthorhombic	Monoclinic
Space group	<i>Pbca</i>	<i>Cc</i>
<i>a</i> /Å	16.001(7)	10.13(1)
<i>b</i> /Å	22.237(7)	13.486(6)
<i>c</i> /Å	15.686(5)	13.396(7)
β /°		101.49(5)
<i>V</i> /Å ³	5581(3)	1794(2)
<i>Z</i>	8	4
Observed reflections	5533	1733
Observed reflections [<i>I</i> _o > 2σ(<i>I</i> _o)]	2695	1417
<i>R</i>	0.036	0.040
<i>R</i> _w	0.037	0.054

were obtained in good yield. The ES⁺ mass spectra (MeCN solution) showed clusters of peaks with the correct *m/z* and isotope patterns for [RuCl(NCMe)(PPh₃)L³⁺]. A further cluster of peaks corresponding to [RuCl(PPh₃)L³⁺] was also observed. Elemental analysis confirmed the identity of the complexes. Although stable in the solid state, these complexes were found to be unstable in solution, even when thoroughly degassed with N₂, rapidly giving green solutions, assigned to ruthenium(III) species. Since such species are paramagnetic this led to complications when recording NMR spectra. To inhibit this process methanol (*ca.* 10%) was added to solutions of the complexes in CH₂Cl₂ for the multinuclear NMR studies where long accumulations were necessary. Even with these precautions the telluroether complexes showed NMR spectra consistent only with decomposition products.

The ¹H NMR spectra were recorded from freshly prepared solutions under N₂ and are, as expected, complicated due to the different environments for the tripod donor arms, and the potential presence of both *syn* and *anti* invertomers, since inversion at an Ru^{II}-Se/TeR₂ centre is expected to be slow.^{9,10} Sharp resonances that may be assigned to PPh₃ and the tripod ligand were apparent for the selenoether complex, however only broad resonances, possibly associated with a paramagnetic species, were observed for the telluroether complexes. The ³¹P-{¹H} NMR spectra of the telluroether complexes only showed resonances corresponding to oxidised phosphine (Ph₃PO δ 26). This behaviour is common and has been observed for other ruthenium complexes,²² although the reaction appears to be extremely rapid for these species. The selenoether complex exhibited two resonances in the ³¹P-{¹H} NMR spectrum of approximately equal intensity at δ 34.4 and 35.2, shifts consistent with co-ordinated PPh₃ and probably indicating the presence of two invertomers. The ⁷⁷Se-{¹H} NMR spectrum of [RuCl₂(PPh₃){MeC(CH₂SeMe)₃} was also recorded and showed seven resonances over the range of 100 ppm, although ²*J*_{Se-P} were poorly resolved. This is consistent with the inequivalence of the tripod arms with both Se-*trans*-Cl and Se-*trans*-P environments, together with the presence of both the *syn* and *anti* invertomers.

Interestingly the complexes [RuCl₂(PPh₃)([9]aneS₃)], [RuCl(PPh₃)([14]aneS₄)]⁺⁷ and [RuCl(PPh₃)([16]aneSe₄)]⁺⁶ have been observed to be stable in solution and therefore similar in behaviour to [RuCl₂(PPh₃){MeC(CH₂SeMe)₃}].

Despite their obvious vulnerability to oxidation and dissociation of PPh₃, it was hoped that by replacement of the phosphine and chloride co-ligands with labile solvent molecules such as a [RuL³⁺]²⁺ based system might be acquired. Unfortunately the reaction of [RuCl₂(PPh₃)L³] with 2 mol equivalents of Ag[CF₃SO₃] in refluxing MeCN led to formation of dark grey materials (which decomposed rapidly to black oils). These showed no selenium or tellurium isotope pattern in the electro-spray mass spectra, hence indicating that the target complexes




Fig. 1 Structure of [RuCl₂(PPh₃){MeC(CH₂SeMe)₃} with the numbering scheme adopted. Ellipsoids are drawn at 40% probability and H atoms omitted for clarity.

[Ru(NCMe)₃L³⁺]²⁺ or [Ru(NCMe)₂(PPh₃)L³⁺]²⁺ had not been obtained.

Crystal structure of [RuCl₂(PPh₃){MeC(CH₂SeMe)₃}

Crystals of the complex were grown *via* slow diffusion of diethyl ether into a solution of the complex in MeOH-CH₂Cl₂ under N₂. The structure (Fig. 1, Table 2) shows Ru^{II} co-ordinated to all three arms of the facially bound selenoether ligand, with the Cl and PPh₃ auxiliary ligands completing the distorted octahedral geometry, *d*(Ru-Se) = 2.429(1), 2.423(1) and 2.492(1) Å with the longer bond *trans* to PPh₃, consistent with the higher *trans* influence of PPh₃ compared to Cl, *d*(Ru-Cl) = 2.453(2) and 2.454(2) Å, *d*(Ru-P) = 2.336(2) Å. The majority of the angles around Ru^{II} are close to the 90 or 180° expected for a regular octahedron, although Se(1)-Ru(1)-P(1) 99.91(6)° is noticeably larger. The Ru-Se bond lengths may be compared with those in *trans*-[RuCl₂{PhSe(CH₂)₂SePh}₂] (2.433(1)-2.460(1) Å)⁹ and *trans*-[RuCl(PPh₃)([16]aneSe₄)]⁺ (2.465(3)-2.497(3) Å),¹¹ with the Ru-P and two Ru-Cl bond lengths also consistent with those found in *trans*-[RuCl(PPh₃)([16]aneSe₄)]⁺ (*d*(Ru-P) = 2.307(6); *d*(Ru-Cl) = 2.499(5) Å). The methyl substituents on the selenoether adopt the *syn* arrangement.

[RuCl₂(dmsO)L³]

The sensitivity of the chloro-phosphine complexes was thought to be due to the presence of the phosphine ligand, and by using

Table 2 Selected bond lengths (Å) and angles (°) for [RuCl₂(PPh₃)₃]{MeC(CH₂SeMe)₃}

Ru(1)–Se(1)	2.429(1)	Ru(1)–Se(2)	2.423(1)
Ru(1)–Se(3)	2.492(1)	Ru(1)–Cl(1)	2.453(2)
Ru(1)–Cl(2)	2.454(2)	Ru(1)–P(1)	2.336(2)
Se(1)–Ru(1)–Se(2)	89.34(4)	Se(1)–Ru(1)–Se(3)	87.47(3)
Se(1)–Ru(1)–Cl(2)	86.83(6)	Se(1)–Ru(1)–P(1)	99.91(6)
Se(2)–Ru(1)–Se(3)	93.53(4)	Se(2)–Ru(1)–Cl(1)	89.13(6)
Se(2)–Ru(1)–P(1)	92.86(6)	Se(3)–Ru(1)–Cl(1)	83.85(5)
Se(3)–Ru(1)–Cl(2)	84.15(6)	Cl(1)–Ru(1)–Cl(2)	94.33(7)
Cl(1)–Ru(1)–P(1)	88.95(7)	Cl(2)–Ru(1)–P(1)	89.94(7)

an alternative ruthenium(II) precursor these difficulties should be avoided. Similar work on complexes with MeC(CH₂PPh₂)₃ has shown that [RuCl₂(dmsO)₄] provides a convenient route into such chemistry, avoiding the use of phosphine co-ligands.²³ Treatment of [RuCl₂(dmsO)₄] with 1 mol equivalent of L³ in toluene at 100 °C for 24 h afforded the complexes [RuCl₂(dmsO)₃L³]. For the ligands MeC(CH₂EMe)₃ (E = Se or Te) the complexes were precipitated as orange or brown powders respectively. For L³ = MeC(CH₂TePh)₃, an orange solution was obtained from which the complex was isolated on concentration.

FAB mass spectrometry showed clusters of peaks with the correct *m/z* and isotope patterns for [RuCl₂(dmsO){MeC(CH₂EMe)₃}]⁺ (E = Se or Te). For [RuCl₂(dmsO){MeC(CH₂TePh)₃}] the molecular ion was not observed, however clusters of peaks were observed corresponding to [RuCl{MeC(CH₂TePh)₃}]⁺. The IR spectra showed dmsO ligands ($\nu(\text{SO})$ 1080–1090 cm⁻¹) indicative of S-bound dmsO.²² The ¹H NMR spectra were again complex but resonances associated with the tripod and dmsO ligands were apparent and, in contrast to the previous dichloro-triphenylphosphine species, these complexes were found to be stable in solution. Interestingly, from the reactions of [RuCl₂(dmsO)₄] with MeC(CH₂EPh₂)₃ (E = P or As) the chloro-bridge dimer [Ru₂(μ-Cl)₃{MeC(CH₂PPh₂)₃}₂]⁺ is obtained with the phosphine, although with the arsine [RuCl₂(dmsO){MeC(CH₂AsPPh₂)₃}] is isolated.²³


The ⁷⁷Se-¹H or ¹²⁵Te-¹H NMR spectra were also recorded. For the selenoether complex six resonances were observed (δ 168, 170, 218, 219, 229 and 244), with similar shifts to those observed for the dichloro-phosphine complex, showing inequivalence of the tripod donors (*trans*-dmsO and *trans*-Cl) and the presence of both *syn* and *anti* invertomers. However since seven resonances are predicted for the presence of the three possible isomers this indicates coincidence of two of the Se-*trans*-dmsO signals. The MeC(CH₂TeMe)₃ complex was highly insoluble in non-co-ordinating solvents and hence the spectrum obtained was too weak to provide useful information. For the MeC(CH₂TePh)₃ complex three resonances were observed of similar intensity which may be assigned to the presence of one major invertomer.

Structure of [RuCl₂(dmsO){MeC(CH₂SeMe)₃}

Crystals were grown by slow evaporation of a solution of the complex in CH₃OH–CH₂Cl₂. The structure (Fig. 2, Table 3) shows the ruthenium co-ordinated to all three arms of the selenoether, with the methyl groups adopting the *syn* arrangement. The octahedral co-ordination sphere is completed by two chlorines, and one dmsO molecule co-ordinated *via* the sulfur atom. Spectroscopic data for [RuCl₂(dmsO){MeC(CH₂AsPh₂)₃}] also indicated S-bonded dmsO, although the crystal structure was not reported.²³ The *d*(Ru–Se) = 2.455(2), 2.417(2) and 2.466(2) Å and *d*(Ru–Cl) = 2.441(4) and 2.448(4) Å are comparable to those in [RuCl₂(PPh₃)₃]{MeC(CH₂SeMe)₃} (above) and [Ru{MeC(CH₂SeMe)₃}]₂²⁺.¹³

Table 3 Selected bond lengths (Å) and angles (°) for [RuCl₂(dmsO){MeC(CH₂SeMe)₃}

Ru(1)–Se(1)	2.455(2)	Ru(1)–Se(2)	2.417(2)
Ru(1)–Se(3)	2.466(2)	Ru(1)–Cl(1)	2.441(4)
Ru(1)–Cl(2)	2.448(4)	Ru(1)–S(1)	2.258(4)
Se(1)–Ru(1)–Se(2)	91.29(6)	Se(1)–Ru(1)–Se(3)	91.75(8)
Se(1)–Ru(1)–Cl(1)	85.7(1)	Se(1)–Ru(1)–Cl(2)	89.4(1)
Se(2)–Ru(1)–Se(3)	87.49(6)	Se(2)–Ru(1)–Cl(1)	92.77(10)
Se(2)–Ru(1)–S(1)	92.6(1)	Se(3)–Ru(1)–Cl(2)	89.7(1)
Se(3)–Ru(1)–S(1)	94.0(1)	Cl(1)–Ru(1)–Cl(2)	90.1(1)
Cl(1)–Ru(1)–S(1)	88.5(1)	Cl(2)–Ru(1)–S(1)	87.0(1)

**Fig. 2** Structure of [RuCl₂(dmsO){MeC(CH₂SeMe)₃}. Details as in Fig. 1.

[Ru(NCMe)₃L³]²⁺ Complexes

Reaction of [RuCl₂(dmsO){MeC(CH₂ER)₃}] (E = Se, R = Me; E = Te, R = Ph) with 2 mol equivalents of Ag[CF₃SO₃] in refluxing MeCN for 2 h gave a light yellow solution and white precipitate. After removal of the AgCl through filtration, reduction of the solvent volume *in vacuo* and addition of diethyl ether, the complexes [Ru(NCMe)₃{MeC(CH₂ER)₃]-[CF₃SO₃]₂ were obtained in good yield as yellow (selenoether) or orange (telluroether) solids. Unfortunately the MeC(CH₂TeMe)₃ complex could not be isolated despite numerous attempts, including the use of TlPF₆ instead of Ag[CF₃SO₃]. The reasons for this are unclear. The characterisation of these complexes was straightforward due to the higher symmetry compared to the previous species. The ES⁺ mass spectra showed clusters of peaks with the correct isotopic distribution for doubly charged species [Ru(NCMe)₃{MeC(CH₂ER)₃]²⁺ and [Ru(NCMe)₂{MeC(CH₂ER)₃]²⁺. IR spectroscopy displayed peaks associated with the tripod ligand and CF₃SO₃⁻ anion, along with weak bands assigned to the co-ordinated MeCN ($\nu(\text{CN})$ 2310 cm⁻¹).

¹H NMR spectra showed signals assigned to the tripod ligand adopting the *syn* arrangement with a further resonance at δ 2.29 (telluroether) or 2.42 (selenoether) assigned to the co-ordinated MeCN molecules. These are comparable with that for the MeC(CH₂PPh₂)₃ complex where $\delta(\text{CH}_3\text{CN})$ 2.34.²³ The ⁷⁷Se-¹H and ¹²⁵Te-¹H NMR spectra showed just one resonance probably indicating the presence of the *syn* inver-

omer, since fast inversion is unlikely with a weak *trans* donor MeCN. Both signals are to low frequency of the corresponding chloro-dmsO species, consistent with substitution of the electronegative chloride ligands with acetonitrile; they are however to high frequency of those of the homoleptic ruthenium(II) seleno- and telluro-ether complexes.¹²

The reaction of $[\text{RuCl}_2(\text{dmsO})\{\text{MeC}(\text{CH}_2\text{SeMe})_3\}]$ with two molar equivalents of $[\text{Ag}(\text{CF}_3\text{SO}_3)]$ in acetone was also studied, with the aim of preparing the tris(acetone) derivative $[\text{Ru}(\text{Me}_2\text{CO})_3\{\text{MeC}(\text{CH}_2\text{SeMe})_3\}]^{2+}$. The product obtained was found to be extremely unstable upon isolation, although the mass spectrum was recorded confirming its identity, with rapid oxidation to ruthenium(III) species occurring. However, this intermediate may be of use since it is stable in solution under N_2 .

One aim of this research was to obtain a reactive $[\text{RuL}^3]^{2+}$ fragment upon which further chemistry could be conducted. Therefore we wished to confirm that the acetonitrile ligands could be substituted easily by other ligands, obviously a prerequisite if these complexes were to be able to carry out reaction chemistry. Addition of one mol equivalent of $\text{MeC}(\text{CH}_2\text{SMe})_3$ to $[\text{Ru}(\text{NCMe})_3\{\text{MeC}(\text{CH}_2\text{SeMe})_3\}][\text{CF}_3\text{SO}_3]_2$ in methanol and reflux for 18 h led to isolation of the light yellow complex $[\text{Ru}\{\text{MeC}(\text{CH}_2\text{SMe})_3\}\{\text{MeC}(\text{CH}_2\text{SeMe})_3\}][\text{CF}_3\text{SO}_3]_2$. Crystals of this complex were obtained† and confirmed the expected cation, but due to disorder of the tripod ligands across the crystallographic inversion centre the structure is not described. The complex $[\text{Ru}\{\text{MeC}(\text{CH}_2\text{TePh})_3\}\{\text{MeC}(\text{CH}_2\text{SeMe})_3\}][\text{CF}_3\text{SO}_3]_2$ was obtained similarly via the reaction of $[\text{Ru}(\text{NCMe})_3\{\text{MeC}(\text{CH}_2\text{TePh})_3\}][\text{CF}_3\text{SO}_3]_2$ with 1 mol equivalent of $\text{MeC}(\text{CH}_2\text{SeMe})_3$. IR spectra of these products displayed peaks associated with the co-ordinated tripodal ligands and CF_3SO_3^- anion, with the ES^+ mass spectra showing clusters of peaks corresponding to the doubly charged cations. The ^1H NMR spectra were complex due to the number of overlapping signals, however resonances associated with both ligands in each complex could be identified. The $^{77}\text{Se}-\{^1\text{H}\}$ and $^{125}\text{Te}-\{^1\text{H}\}$ NMR spectra showed one resonance for each nucleus corresponding to the presence of the *syn* invertomers. For the $[\text{Ru}\{\text{MeC}(\text{CH}_2\text{SMe})_3\}\{\text{MeC}(\text{CH}_2\text{SeMe})_3\}]^{2+}$ complex $\delta(^{77}\text{Se}-\{^1\text{H}\})$ 123, a similar shift to that of the homoleptic complex $[\text{Ru}\{\text{MeC}(\text{CH}_2\text{SeMe})_3\}]^{2+}$ (120).¹³ The $[\text{Ru}\{\text{MeC}(\text{CH}_2\text{TePh})_3\}\{\text{MeC}(\text{CH}_2\text{SeMe})_3\}]^{2+}$ complex shows $\delta(^{77}\text{Se}-\{^1\text{H}\})$ 128 and $\delta(^{125}\text{Te}-\{^1\text{H}\})$ 485; both shifts are similar to those observed for the respective homoleptic Se_6 or Te_6 donor species reported previously.¹³

Having established the lability of the acetonitrile ligands and hence the availability of the $[\text{RuL}^3]^{2+}$ fragment, we were interested to study the reaction of these species with NaBH_4 in the expectation of generating hydride species, so important for hydrogenation and hydrodesulfurisation catalysis. Initially, an excess of solid NaBH_4 was added slowly to a solution of $[\text{Ru}(\text{NCMe})_3\{\text{MeC}(\text{CH}_2\text{ER})_3\}][\text{CF}_3\text{SO}_3]_2$ ($\text{E} = \text{Se}, \text{R} = \text{Me}; \text{E} = \text{Te}, \text{R} = \text{Ph}$) in dry ethanol at room temperature. A gas was evolved immediately along with precipitation of a black solid. Attempts to identify this product were unsuccessful, with the mass and ^1H NMR spectra showing no peaks that could be assigned to a tripod-containing product. It is likely that this product is largely ruthenium metal, obviously in contrast to the chemistry observed with $\text{MeC}(\text{CH}_2\text{PPh}_2)_3$, and is probably as a result of the poorer σ -donor/ π -acceptor ligand properties of the

Group 16 tripods. In an attempt to avoid the decomposition, the reaction was repeated by adding NaBH_4 to a slurry of $[\text{Ru}(\text{NCMe})_3\{\text{MeC}(\text{CH}_2\text{ER})_3\}][\text{CF}_3\text{SO}_3]_2$ in ethanol at -78°C . No reaction was observed until the mixture was allowed to warm slowly, when a black precipitate was again formed indicating decomposition. This rather disappointing result does not necessarily exclude these complexes as potential catalysts, since co-ordination of the substrate may well stabilise the ruthenium centre. The complex $[\text{Ru}(\text{NCMe})_3\{\text{MeC}(\text{CH}_2\text{SeMe})_3\}]^{2+}$ did not react with carbon monoxide in CH_2Cl_2 solution at ambient temperatures, however reaction of the cation with PMe_3 in acetone resulted in replacement of MeCN by the phosphine. Details of these and related reactions will be reported in due course.

Acknowledgements

We thank EPSRC for support and Johnson Matthey for loan of the ruthenium salts.

References

- H. A. Meyer and W. C. Kaska, *Chem. Rev.*, 1994, **94**, 1239.
- C. Bianchini, A. Meli, M. Peruzzini, F. Vizza and F. Zanobini, *Coord. Chem. Rev.*, 1992, **120**, 193.
- C. Bianchini, D. Masi, A. Meli, M. Peruzzini, F. Vizza and F. Zanobini, *Organometallics*, 1998, **17**, 2495; C. Bianchini, A. Meli, S. Moneti and F. Vizza, *Organometallics*, 1998, **17**, 2636.
- C. Bianchini, A. Meli, S. Moneti, W. Oberhauser, F. Vizza, V. Herrera, A. Fuentes and R. Sanchez-Delgado, *J. Am. Chem. Soc.*, 1999, **121**, 7071.
- M. N. Bell, A. J. Blake, H.-J. Kuppers, M. Schröder and K. Wieghardt, *Angew. Chem., Int. Ed. Engl.*, 1987, **26**, 250.
- N. W. Alcock, J. C. Cannadine, G. R. Clark and A. F. Hill, *J. Chem. Soc., Dalton Trans.*, 1993, 1131.
- A. L. Hector and A. F. Hill, *Inorg. Chem.*, 1995, **34**, 3797.
- J. C. Cannadine, A. F. Hill, A. J. P. White, D. J. Williams and J. D. E. T. Wilton-Ely, *Organometallics*, 1996, **15**, 5409.
- N. R. Champness, W. Levason, S. R. Preece and M. Webster, *Polyhedron*, 1994, **13**, 881.
- W. Levason, S. D. Orchard, G. Reid and V.-A. Tolhurst, *J. Chem. Soc., Dalton Trans.*, 1999, 2071.
- W. Levason, J. J. Quirk, G. Reid and S. M. Smith, *J. Chem. Soc., Dalton Trans.*, 1997, 3719.
- W. Levason, S. D. Orchard, G. Reid and J. M. Street, *J. Chem. Soc., Dalton Trans.*, 2000, 2537.
- W. Levason, S. D. Orchard and G. Reid, *Chem. Commun.*, 1999, 1071; W. Levason, S. D. Orchard and G. Reid, *Inorg. Chem.*, 2000, **39**, 3853.
- T. A. Stephenson and G. Wilkinson, *J. Inorg. Nucl. Chem.*, 1966, **28**, 945.
- I. P. Evans, A. Spencer and G. Wilkinson, *J. Chem. Soc., Dalton Trans.*, 1973, 204.
- R. Ali, S. J. Higgins and W. Levason, *Inorg. Chim. Acta*, 1984, **84**, 65.
- D. J. Gulliver, E. G. Hope, W. Levason, G. L. Marshall, S. G. Murray and D. M. Potter, *J. Chem. Soc., Perkin Trans. 2*, 1984, 429.
- E. G. Hope, T. Kemmitt and W. Levason, *Organometallics*, 1988, **7**, 78.
- J. Connolly, A. R. J. Genge, W. Levason, S. D. Orchard, S. J. A. Pope and G. Reid, *J. Chem. Soc., Dalton Trans.*, 1999, 2343.
- PATY, The DIRDIF Program System, P. T. Beurskens, G. Admiraal, G. Beurskens, W. P. Bosman, S. Garcia-Granda, R. O. Gould, J. M. M. Smits and C. Smykalla, Technical Report of the Crystallography Laboratory, University of Nijmegen, 1992.
- TEXSAN, Crystal Structure Analysis Package, Molecular Structure Corporation, Houston, TX, 1995.
- B. A. Moyer, B. K. Sipe and T. J. Meyer, *Inorg. Chem.*, 1981, **20**, 1475.
- L. F. Rhodes, C. Sorato, L. M. Venanzi and F. Bachechi, *Inorg. Chem.*, 1988, **27**, 604.

† Triclinic, space group $P\bar{1}$, $a = 8.791(2)$, $b = 11.406(8)$, $c = 8.555(2)$ Å, $\alpha = 107.53(3)$, $\beta = 91.38(2)$, $\gamma = 106.72(3)^\circ$, $V = 777.6(6)$ Å³.