

This article was downloaded by:

On: 22 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Journal of Asian Natural Products Research

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713454007>

Constituents of *Clerodendrum bungei*

Hui Yang^a; Ai-Jun Hou^a; Shuang-Xi Mei^a; Han-Dong Sun^a; Chun-Tao Che^b

^a Laboratory of Phytochemistry, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming, Yunnan, China ^b School of Chinese Medicine, The Chinese University of Hong Kong, Shatin, N.T., Hong Kong, China

Online publication date: 09 September 2010

To cite this Article Yang, Hui, Hou, Ai-Jun, Mei, Shuang-Xi, Sun, Han-Dong and Che, Chun-Tao(2002) 'Constituents of *Clerodendrum bungei*', *Journal of Asian Natural Products Research*, 4: 3, 165 – 169

To link to this Article: DOI: 10.1080/1028602021000000053

URL: <http://dx.doi.org/10.1080/1028602021000000053>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

CONSTITUENTS OF *CLERODENDRUM BUNGEI*

HUI YANG^{a,b}, AI-JUN HOU^a, SHUANG-XI MEI^a, HAN-DONG SUN^{a,*} and
CHUN-TAO CHE^c

^aLaboratory of Phytochemistry, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming 650204, Yunnan, China; ^bDepartment of Applied Chemistry, College of Life Science and Chemistry, Yunnan University, Kunming 650091, Yunnan, China; ^cSchool of Chinese Medicine, The Chinese University of Hong Kong, Shatin, N.T., Hong Kong, China

(Received 10 September 2001; Revised 19 September 2001; In final form 28 November 2001)

Two new compounds, 5-*O*-ethylcleroindicin D (**1**) and bungein A (**2**), together with 12 known compounds (**3–14**), were isolated from the aerial parts of the medicinal plant *Clerodendrum bungei*. The structures of **1** and **2** were elucidated as a perhydrobenzofuran derivative and a peroxide dimer by spectral and chemical evidence. Compounds **3–14** have been obtained from this species for the first time.

Keywords: Verbenaceae; *Clerodendrum bungei*; 5-*O*-Ethylcleroindicin D; Bungein A

INTRODUCTION

In a previous paper [1], we reported the isolation and structural elucidation of several perhydrobenzofuran derivatives from *Clerodendrum indicum* (Verbenaceae). Another species of the same genus, *Clerodendrum bungei* Steud., which is distributed widely in China, has long been used in folk medicine to treat headache, dizziness, furuncle, and hysteroptosis [2,3]. A few papers concerning the chemical constituents of this plant have been reported [3–5]. In the present study, a methanol extract of *C. bungei* was separated to afford two new compounds, 5-*O*-ethylcleroindicin D (**1**) and bungein A (**2**) (Scheme 1), together with 12 known compounds, including acteoside (**3**) [6], betulinic acid (**4**) [7], cleroindicin A (**5**), cleroindicin C (**6**), cleroindicin E (**7**), cleroindicin F (**8**) [1], clerosterol (**9**) [8], clerosterol 3 β -*O*- β -D-glucopyranoside (**10**) [8], martinoid (**11**) [9], octadecnoic acid (**12**), *n*-pentacosane (**13**) [10], and 5,7,4'-trihydroxyflavone (**14**) [11]. We report here the isolation and structural elucidation of these compounds.

RESULTS AND DISCUSSION

Compound **1** was obtained as a colorless oil and assigned a molecular formula of C₁₀H₁₆O₄ on the basis of its negative HRFABMS ([M – 1][–] at *m/z* 199.0949). Its IR spectrum

*Corresponding author. Tel.: +86-871-5223251. Fax: +86-871-5216343. E-mail: hdsun@mail.kib.ac.cn

TABLE I NMR data for 5-*O*-ethylcleroindicin D (**1**)*

Position	^{13}C	$^1\text{H}^\dagger$	COSY	HMBC ‡
2	66.6 t	3.96, m	3	9
3	39.2 t	2.18, m	2	5, 9
4	79.1 s			2, 6 _{ax} , 6 _{eq} , 8 _{ax} , 8 _{eq}
5	79.3 d	3.77, dd (6.7, 3.4)	6 _{ax} , 6 _{eq}	3, 9, α_a , α_b
6 _{ax}	40.6 t	2.86, dd (16.6, 3.4)	5, 6 _{eq}	8 _{ax} , 8 _{eq}
6 _{eq}		2.77, dd (16.6, 6.7)	5, 6 _{ax}	
7	207.6 s			5, 9
8 _{ax}	43.0 t	2.72, dd (16.8, 4.7)	8 _{eq} , 9	6 _{ax} , 6 _{eq}
8 _{eq}		3.05, dd (16.8, 4.7)	8 _{ax} , 9	
9	83.8 d	4.22, t (4.7)	8 _{ax} , 8 _{eq}	2, 3, 5
α_a	65.9 t	3.58, m	α_b , β	5
α_b		3.46, m	α_a , β	
β	15.7 q	1.07, t (8.2)	α_a , α_b	


* In $\text{C}_5\text{D}_5\text{N}$, ^1H at 500 MHz, ^{13}C at 125 MHz.

† Couplings in Hz; chemical shifts of overlapped signals determined from 2D NMR data.

‡ Long-range correlation of ^1H signals to ^{13}C , in $\text{C}_5\text{D}_5\text{N}$, ^1H at 500 MHz, ^{13}C at 125 MHz.

displayed absorption bands for hydroxyl (3383 cm^{-1}) and carbonyl (1698 cm^{-1}) groups. A comparison of the MS and NMR spectral data of **1** with those of cleroindicin D [1] indicated that the two compounds were structurally related, except for the presence of an oxymethylene (δ 65.9, C- α) and a methyl group (δ 15.7, C- β) in **1**. Analysis of the 2-dimensional NMR spectra of **1** (Table I) clearly suggested that **1** possessed the same carbon skeleton as cleroindicin D (perhydrobenzofuran), and an ethoxyl group was assignable to the C-5 (δ 79.3, d) position based on the observation of long-range carbon–proton correlation (HMBC) between C- α and H-5. Consequently, the structure of **1** was determined as 5-*O*-ethylcleroindicin D.

Bungein A (**2**) was obtained as an optically inactive colorless wax. Its molecular formula ($\text{C}_{16}\text{H}_{18}\text{O}_4$) was deduced from the negative HRFABMS ($[\text{M} - 1]^-$ at m/z 273.1218). The IR absorption bands revealed the presence of hydroxyl ($3147\text{--}3391\text{ cm}^{-1}$) and aromatic (1599 , 1512 and 819 cm^{-1}) functional groups. The UV spectrum of **2** displayed absorption maxima at 224 (4.16), 278.5 (4.09) and 281.5 (4.37) nm, suggesting the presence of an aromatic structure. Interestingly, in its ^{13}C NMR spectrum only six carbon signals, namely, two quaternary carbons (δ 156.5 and 130.8), two methines (δ 130.7 and 115.8), and two methylenes (δ 64.2 and 39.3) were observed. Such a simplified spectrum led to the assumption that **2** has a symmetrical structure. Indeed, the ^1H NMR spectrum exhibited four groups of proton signals, i.e. aromatic protons at δ 7.03 (4H, dd, $J = 8.4, 2.9\text{ Hz}$, H-3, 5, 3', 5') and 6.73 (4H, dd, $J = 8.4, 2.9\text{ Hz}$, H-2, 6, 2', 6'), oxymethylenes at δ 3.69 (4H, t, $J = 7.2\text{ Hz}$, H- β , β'), and methylenes at δ 2.70 (4H, t, $J = 7.2\text{ Hz}$, H- α , α'). Analysis of the proton coupling patterns of **2** led to the conclusion that the molecule was composed of two units of 2-phenylethanol derivative. Additionally, Compound **2** produced a positive reaction with the Farbentwickler 3 reagent (Merck), whereas with the above reagent after treatment with triphenyl phosphine it showed a negative reaction. Such experimental results indicated that **2** was a peroxide [12]. This was further confirmed by a downfield shift of the aromatic carbon signals at δ 156.5 (s, C-1 and C-1') due to the effect of peroxide bridge. From the above discussion, two possible Structures **2** and **A** (Scheme 1) could be suggested for the compound. The EIMS data $\{m/z$ 274 $[\text{M}]^+$ (27), 256 $[\text{M} - \text{H}_2\text{O}]^+$ (4.5), 243 $[\text{M} - \text{CH}_2\text{OH}]^+$ (45), and 225 $[m/z$ 256 $-\text{CH}_2\text{OH}$ or m/z 243 $-\text{H}_2\text{O}]^+$ (23) $\}$ excluded the possibility of structure A, since it would not produce the fragmentation patterns as those listed above. Hence, all available evidence suggested the structure of bungein A as depicted in **2**.



SCHEME 1

In addition to the two new constituents, Compounds **3–14** were purified from the plant extract. These constituents are reported from this species for the first time. Their structures were identified by direct comparison with reference samples, or by comparing their physical and spectral properties with those reported in the literature.

EXPERIMENTAL SECTION

General Experimental Procedures

Optical rotations were measured on a JASCO-20C digital polarimeter. UV spectra were determined on a UV 210A spectrometer and IR spectra on Bio-Rad FTS-135 spectrometer. 1D- and 2D-NMR spectra were taken on a DRX-500 instrument with TMS as internal reference. MS were recorded on a VG Auto Spec-3000 mass spectrometer.

Plant Material

The plant material was collected in Meishan County, Sichuan Province, China, in August 1997, and identified as *Clerodendrum bungei* Steud. by Prof. Ri-Zheng Fang (Kunming Institute of Botany). A voucher specimen (KIB 97-8-16 Fang) was deposited in the Herbarium of the Department of Taxonomy, Kunming Institute of Botany, Chinese Academy of Sciences.

Extraction and Isolation

The air-dried and powdered aerial parts of *C. bungei* (6.0 kg) were extracted with MeOH (3 × 20 l) under reflux to afford a crude extract (600.5 g). The extract was dissolved in 90% MeOH and defatted with petroleum-ether (60–90°C). After removal of excessive MeOH, the residue was partitioned with EtOAc (6 × 1.5 l) and *n*-BuOH (4 × 1.5 l), successively. The EtOAc extract (50.0 g) was subjected to chromatography over silica gel (1.3 kg, 200–300 mesh), eluting with CHCl₃–Me₂CO (1:0, 9:1, 8:2, 7:3, 6:4, 3:7, and 0:1) and MeOH, to afford nine fractions (A–I).

From fraction A, *n*-pentacosane (109 mg) was obtained by column chromatography on silica gel eluting with petroleum ether–CHCl₃ (7:3). Fraction C was subjected to medium-pressure column chromatography on silica gel, developing with CHCl₃–MeOH (90:1 and 30:1) and petroleum ether–Et₂O (5:5, 6:4, 8:2 and 2:8) to yield 5-*O*-ethylcleroindicin D (1,134 mg), cleroindicin C (356 mg), cleroindicin F (116 mg), bungein A (2, 12 mg), 5,7,4'-trihydroxyflavone (17 mg), betulinic acid (7 mg), and octadecenoic acid (16 mg). Fraction F was purified by silica gel column chromatography eluted with CHCl₃–MeOH (9:1) and petroleum ether–*i*-PrOH (8:2) to afford compounds cleroindicin E (42 mg) and cleroesterol (31 mg). From fractions G and H, cleroindicin A (40 mg), cleroesterol 3β-*O*-β-D-glucopyranoside (8 mg), martinosiol (565 mg), and acteoside (5.4 g) were obtained following repeated chromatography on silica gel, reversed-phase silica gel (RP-18) and MCI-gel CHP-20P [eluents: CHCl₃–MeOH (9:1), CHCl₃–MeOH–H₂O (9:2:0.1) and MeOH–H₂O (3:7 and 4:6)].

5-*O*-ethylcleroindicin D (I). C₁₀H₁₆O₄, colorless oil, [α]_D¹³ –0.27 (c 0.93, CHCl₃). IR (film) ν_{max} 3383, 2975, 2907, 2869, 1698, 1411, 1352, 1286, 1189, 1127, 1110, 1060, 996 cm⁻¹; ¹H- and ¹³C-NMR, see Table I. EIMS (70 eV) *m/z* 200 [M]⁺ (26), 182 [M – H₂O]⁺ (10), 154 [M – EtOH]⁺ (21), 138 (18), 127 (29), 115 (95), 107 (23), 99 (45), 86 (100), 73 (37), 55 (56); negative HRFABMS *m/z* 199.0949 [M – 1]⁻ (Calcd for C₁₀H₁₅O₄: 199.0970).

Bungein A (2). C₁₆H₁₈O₄, colorless wax, UV (MeOH) λ_{max} (log ε) 281.5 (4.37), 278.5 (4.09), 224 (4.16), 202 (4.15) nm; IR (film) ν_{max} 3391, 3147, 3024, 2955, 2928, 2881, 2702, 2679, 1884, 1599, 1512, 1453, 1364, 1233, 1054, 819, 555 cm⁻¹; ¹H-NMR (CD₃COCD₃, 400 MHz) δ 7.03 (4H, dd, *J* = 8.4, 2.9 Hz, H-3, 5, 3', 5'), 6.73 (4H, dd, *J* = 8.4, 2.9 Hz, H-2, 6, 2', 6'), 3.69 (4H, t, *J* = 7.2 Hz, H-β, β'), 2.70 (4H, t, *J* = 7.2 Hz, H-α, α'); ¹³C-NMR (CD₃COCD₃, 100 MHz) δ 156.5 (s, C-1, 1'), 130.8 (s, C-4, 4'), 130.6 (d, C-3, 5, 3', 5'), 115.8 (d, C-2, 6, 2', 6'), 64.2 (t, C-β, β'), 39.3 (t, C-α, α'); EIMS (70 eV) *m/z* 274 [M]⁺ (27), 256 [M – H₂O]⁺ (4.5), 243 [M – CH₂OH]⁺ (45), 225 [*m/z* 256 – CH₂OH or *m/z* 243 – H₂O]⁺ (23), 168 (65), 153 (41), 138 (19), 121 (29), 107 (100), 97 (44), 77 (34); negative HRFABMS *m/z* 273.1218 [M – 1]⁻ (Calcd for C₁₆H₁₇O₄: 273.1204).

Acknowledgement

The work was supported by a grant from the National Natural Science Foundation of the People's Republic of China (Grant No. 29772039).

References

- [1] Tian, J., Zhao, Q.S., Zhang, H.J., Lin, Z.W. and Sun, H.S. (1997), *J. Nat. Prod.* **60**, 766–769.
- [2] Kunming Institute of Botany, Chinese Academy of Sciences (1977) *Flora Yunnanica* (Science Press, Beijing) Vol. 1, pp. 468–469.
- [3] Zhou, P.C., Pang, Z.H., Hao, H.F. and Pu, Q.L. (1982), *Acta Bot. Sin.* **24**, 564–567.
- [4] He, L., Chen, Y.Z., Ding, L.S. and Li, P.G. (1997), *Zhong Caoyao* **28**, 395–396.

- [5] Fan, T.P., Min, Z.H. and Iinuma, M. (1999), *Chem. Pharm. Bull.* **47**, 1797–1798.
- [6] Atsushi, S. and Takahiko, K. (1983), *Bull. Chem. Soc. Jpn* **56**, 1573–1574.
- [7] Kitajima, J., Shindo, M. and Tanaka, Y. (1990), *Chem. Pharm. Bull.* **38**, 714–716.
- [8] Goswami, P., Kotoky, J., Chen, Z.N. and Lu, Y. (1996), *Phytochemistry* **41**, 279–281.
- [9] Tian, J. and Sun, H.D. (1995), *Acta Bot. Yunnan.* **17**, 103–108.
- [10] Wu, S.J. (1980), *Zhong Caoyao* **11**, 99–101.
- [11] Ternai, B. and Markham, K.R. (1976), *Tetrahedron* **32**, 565–569.
- [12] Lou, S.D. and Zhang, Q.F. (1997), *Nat. Prod. Res. Dev.* **9**, 7–9.