

This article was downloaded by:

On: 22 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Journal of Asian Natural Products Research

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713454007>

Triterpenes from *Tripterygium wilfordii* Hook

J. -H. Yang^a; S. -D. Luo^b; Y. -S. Wang^b; J. -F. Zhao^a; H. -B. Zhang^a; L. Li^a

^a School of Pharmacy, Center for Advanced Studies of Medicinal and Organic Chemistry, Yunnan University, Kunming, Yunnan, China ^b State Key Laboratory of Phytochemistry and Plant Resources in West China, Kunming Institute of Botany, Graduate School of the Chinese Academy of Sciences, Chinese Academy of Sciences, Kunming, Yunnan, China

To cite this Article Yang, J. -H. , Luo, S. -D. , Wang, Y. -S. , Zhao, J. -F. , Zhang, H. -B. and Li, L.(2006) 'Triterpenes from *Tripterygium wilfordii* Hook', Journal of Asian Natural Products Research, 8: 5, 425 — 429

To link to this Article: DOI: 10.1080/10286020500172665

URL: <http://dx.doi.org/10.1080/10286020500172665>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

Triterpenes from *Tripterygium wilfordii* Hook

J.-H. YANG^{†‡}, S.-D. LUO[†], Y.-S. WANG[†], J.-F. ZHAO[‡], H.-B. ZHANG[‡] and L. LI^{‡*}

[†]State Key Laboratory of Phytochemistry and Plant Resources in West China, Kunming Institute of Botany, Graduate School of the Chinese Academy of Sciences, Chinese Academy of Sciences, Kunming, Yunnan 650204, China

[‡]School of Pharmacy, Center for Advanced Studies of Medicinal and Organic Chemistry, Yunnan University, Kunming, Yunnan 650091, China

(Received 1 November 2004; revised 29 December 2004; in final form 8 January 2005)

Two new friedelane-type triterpenes, tripterfrianons A (**1**) and B (**2**), along with six known triterpenoids, friedelin (**3**), canophyllal (**4**), canophyllalic acid (**5**), 3-oxo-29-hydroxyfriedelane (**6**), wilforlide A (**7**), wilforlide B (**8**), have been isolated from the EtOH extract of the roots of *Tripterygium wilfordii* Hook.f. Compounds **4**, **5**, **6** were isolated for the first time from this plant. The new triterpenes **1** and **2** exhibited mild cytotoxic activity against human Hela cell lines *in vitro*. The assay showed the IC₅₀ of **1** and **2** were 8.5 and 25 µg/mL, respectively.

Keywords: *Tripterygium wilfordii* Hook.f.; Celastraceae; Triterpenes; Tripterfrianons A–B

1. Introduction

Tripterygium wilfordii Hook.f., known as ‘Lei-Gong-Teng’, is widely distributed in southern China as a Chinese folk medicine. Since the 1960s, it has been found to possess antitumor, anti-inflammatory, anti-HIV and immunosuppressive activities. Lei-Gong-Teng and its preparation have been used for the treatment of various diseases, including dermatitis, rheumatoid arthritis, systemic acne roscoe and nephritis [1,2]. Many pentacyclic triterpenes were previously isolated from this plant. In the course of our continuing search for active constituents, two new friedelane-type triterpenes, tripterfrianons A (**1**) and B (**2**), along with six known triterpenoids, friedelin (**3**), canophyllal (**4**), canophyllalic acid (**5**), 3-oxo-29-hydroxyfriedelane (**6**) [3], wilforlide A (**7**), wilforlide B (**8**) [4], were isolated from the EtOH extract of the roots of *Tripterygium wilfordii* Hook.f. Compounds **4**, **5**, **6** were isolated for the first time from this plant. Compound **1** is the first report of a friedelane-type triterpenoid with a C-30 formyl function from natural source. The new triterpenes **1** and **2** exhibited mild cytotoxic activity against human Hela cell lines *in vitro*. The assay showed the IC₅₀ of **1** and **2** were 8.5 and 25 µg/mL, respectively.


*Corresponding author. E-mail: liliang5758@sina.com; luoshide@hotmail.com

2. Results and discussion

Compound **1** was obtained as white amorphous powder and has a molecular formula of $C_{30}H_{48}O_3$ based on HR-ESI-MS at m/z 479.3528 $[M + Na]^+$. The IR (KBr) spectrum showed absorption bands at 3442 ($-OH$), 2933, 2867, 1716 ($-C=O$) cm^{-1} . The ^{13}C and DEPT NMR spectra showed 30 carbon signals including six methyls, twelve methylenes, four methines, six quaternary carbons and two carbonyls (δ_C 213.4, 208.9). The 1H NMR spectrum showed signals for an aldehyde (δ_H 9.45, s, 1H) and a hydroxymethylene group at δ_H 3.26 (brs, 2H). The HMQC and HMBC analyses revealed that **1** was of the friedelane-type pentacyclic triterpenes [5]. The NMR spectra of **1** were very similar to those of known compound canophyllal (**4**) (see table 1). So the aldehyde group was assigned to C-28, which is coincident with HMBC. In the HMBC experiment, the aldehyde proton signal at δ_H 9.45 (1H, s) was correlated to C-17 (δ_C 48.6s), C-18 (δ_C 35.9d), C-16 (δ_C 28.6t). The difference was the absence of one methyl group and the presence of a hydroxymethylene group (δ_C 74.7t, δ_H 3.26, brs, 2H) in compound **1**. The IR spectrum of **1** showed hydroxyl absorption at 3442 cm^{-1} . A characteristic peak at m/z 273 arising from the cleavage of C13–C18 and C14–C15 bonds and a base peak at m/z 427 resulting from loss of an aldehyde group (C17–CHO) as seen in a typical friedelan fragmentation pattern led to the conclusion that the CH_2OH group was located to D/E rings. Because of the substitution of hydroxyl group, the carbon signal of C-20 at δ_C 33.7s was shifted downfield from δ_C 28.7s in comparisons to **4**.


Table 1. The ^{13}C NMR data for compounds **1**, **2**, **4**, **6** ($CDCl_3$, δ in ppm).

Carbon	1	2	4	6
1	22.7t	22.0t	22.6t	22.7t
2	41.4t	41.2t	41.9t	41.6d
3	213.4s	212.8s	213.4s	213.7s
4	58.6d	57.9d	58.6d	58.6d
5	42.3s	41.9s	42.3s	42.5s
6	41.9t	41.0t	41.4t	41.9t
7	18.4t	18.0t	18.4t	18.6t
8	53.2d	53.1d	53.2d	53.8d
9	38.0s	37.2s	37.5s	37.8s
10	59.6d	59.2d	59.7d	59.8d
11	35.7t	35.3t	35.8t	36.0t
12	31.0t	29.8t	31.0t	30.9t
13	37.6s	38.0s	38.1s	38.6s
14	39.3s	39.6s	39.1s	40.3s
15	28.6t	32.3t	28.4t	33.5t
16	32.8t	35.5t	32.7t	36.2t
17	48.6s	30.1s	48.1s	31.0s
18	35.9d	41.6d	36.8d	42.2d
19	29.8T	30.3t	35.3t	30.1t
20	33.7s	31.4s	33.8t	33.1s
21	28.0t	27.7t	28.7s	28.2t
22	33.7t	38.8t	32.8t	39.9t
23	7.2q	6.6q	7.2q	7.2q
24	15.0q	14.4q	15.0q	15.0q
25	17.5q	17.6q	17.6q	18.3q
26	20.5q	18.2q	19.2q	18.8q
27	19.2q	20.7q	20.4q	21.1q
28	208.9d	31.8q	209.4d	32.5q
29	24.4q	25.9q	34.9q	75.1t
30	74.7t	74.3t	29.8q	26.2q
HCO-		161.2d		

Figure 1. Structures of compounds **1**–**6**.

This suggested that the CH₂OH group was at E ring. The position of hydroxyl group at C-30 was unambiguously elucidated by the ¹H NMR analysis. As reported in the literature, there is a minor difference in the coupling pattern of the hydroxymethylene signal at C-29 or C-30 [6]. The supplementary two-proton singlet at δ_H 3.26 (2H, brs) indicated that the hydroxyl group was attached at C-30, which correlated with a hydroxymethyl at C-30 (δ_C 74.7) in the HMQC experiment. Compound **6** with a 29-hydroxy group exhibited two AB doublet signal at δ_H 3.44, 3.39 (each 1H, d, $J = 10.4$ Hz). This was further supported by the HMBC spectrum, which had long-range correlations between H-30 with C-29, C-20, C-19 and C-21 (figure 2). Thus, the structure of **1** was elucidated as 3-oxo-30-hydroxyfriedelan-28-al (figure 1).

Compound **2** was isolated as amorphous powder and showed a molecular formula of C₃₁H₅₀O₃ based on HR-ESI-MS at m/z 493.3682 [M + Na]⁺. The IR (KBr) spectrum showed absorption bands at 2929, 2869, 1750, 1731 (C=O), 1388, 1162 cm⁻¹. The ¹H NMR spectrum showed signals for six methyl singlets (δ_H 0.72, 0.87, 1.03 × 2, 1.07, 1.21), one methyl doublet (δ_H 0.89, d, $J = 6.0$ Hz) and a formate proton singlet at δ_H 8.14 [7]. The ¹³C and DEPT NMR spectra displayed the presence of seven methyls, twelve methylenes, one of which was oxygenated, five methines, six quaternary carbons, one carbonyl (δ 212.8) and one formyl carbon (δ 161.2) [7]. The EI-MS showed a characteristic peak at m/z 273 as seen in a typical friedelane fragmentation pattern. All these evidences suggested that **2** was also a friedelane-type triterpenes. Similar to **1**, the hydroxyl is substituted at C-30. The HMBC experiment supported that the methyl group at C-30 has been oxygenated into a hydroxymethylene unit, indicated by a correlation between H-30 and C-29, C-20 and the formyl carbon. Meanwhile, the correlation from the formyl proton to C-30 supported that the formyl group was linked to C-30 hydroxymethylene (figure 2). Moreover, the correlation between the formyl proton and H-30 confirmed the placement of the —OCHO at C-30 in the

Figure 2. Selective HMBC correlations of **1** and **2**.

^1H - ^1H COSY spectrum. Therefore, the structure of **2** was deduced to be 3-oxo-30-hydroxyfriedelane formate (figure 1). So far, friedelane-type triterpene with aldehyde group has been reported [3,5], but this is the first report of a friedelane-type triterpene with a C-30 formyl function from natural sources.

Six known compounds from *Tripterygium wilfordii* Hook.f. were identified by spectroscopic evidence or direct comparison with authentic samples as friedelin (**3**), canophyllal (**4**), canophyllalic acid (**5**), 3-oxo-29-hydroxyfriedelane (**6**) [3], wilforlide A (**7**), wilforlide B (**8**) [4]. Among them, compounds **4**, **5**, **6** were isolated for the first time from this plant.

The new triterpenes **1** and **2** exhibited mild cytotoxic activity against human Hela cell lines *in vitro*. The assay showed the IC_{50} of **1** and **2** were 8.5 and 25 $\mu\text{g/mL}$, respectively.

3. Experimental

3.1 General experimental procedures

IR spectra were taken on a Nicolet AVATAR-360 spectrophotometer, ν_{max} in cm^{-1} . ^1H and ^{13}C NMR spectra were measured with a Bruker AV 300 (300 MHz for ^1H and 75 MHz for ^{13}C) and AV-500 spectrometer. Chemical shifts (δ) are in ppm relative to TMS as internal standard and coupling constant (J) are in Hz. The EI-MS was obtained on VG-Autospec 3000 mass spectrometers. Optical rotation was measured on a Jasco-20 MC Polarimeter. Commercial Silica gel plates (Qing Dao Marine Chemical Group Co.) were used for TLC. The chromatograms were sprayed with 10% H_2SO_4 and heated at 80°C to detect the spots.

3.2 Plant material

The plant sample was collected from Lincang county of Yunnan province in October 2001 and identified by Professor Zhi-Hao Hu of the department of botany, Yunnan University. A voucher specimen (No. 200110) is deposited in the School of Pharmacy, Yunnan University.

3.3 Extraction and isolation

The powdered roots of *Tripterygium wilfordii* Hook.f. (10.0 kg) were repeatedly extracted with EtOH at room temperature. The extract was then concentrated under reduced pressure to give brown syrup, which was partitioned with solvents into petroleum ether-soluble (50 g), EtOAc-soluble (120 g) and *n*-BuOH-soluble (80 g) fractions. The PE-soluble portion was subjected to silica gel column chromatography eluting with PE-EtOAc (10:1–1:1), by which nine fractions (I–IX) were obtained. Fractions II and III were resubmitted to silica gel column chromatography to yield compounds **1** (20 mg), **3** (25 mg), **4** (18 mg), **5** (15 mg) and **6** (21 mg), respectively. The EtOAc-soluble portion was subjected to silica gel column chromatography washing with CHCl_3 -MeOH (99:1–1:1), by which 15 fractions (I–XV) were obtained. Fraction II (4.0 g) was resubmitted to silica gel column chromatography (petroleum ether-EtOAc (5:1, 3:1, 1:1) to yield **2** (7 mg). Fraction III (8.0 g) was subjected to chromatography over silica gel eluting with CHCl_3 -MeOH (30:1, 20:1, 10:1) to yield compounds **7** (50 mg) and **8** (20 mg), respectively.

3.4 Cytotoxic activity

Hela (human carcinoma of the cervix) cell lines were grown as a monolayer in Dulbecco's modified eagle's medium, DMEM (Gibco), supplemented with 10% new-born calf serum

(Gibco) and 1% of penicillin-streptomycin mixture (10,000 UI/ml). The cells were maintained at 37°C in 5% CO₂ and 90% humidity. The cytotoxic activity was assessed using colorimetric MTT reduction assay [8]. The percentage viability was plotted against the compound concentrations, and the 50% cell viability (IC₅₀) was calculated from the curve. All the experiments were repeated three times. The assay showed the IC₅₀ of **1** and **2** were 8.5 and 25 µg/mL, respectively.

3.5 Tripterfrielanon A (1)

White amorphous powder. $[\alpha]_D^{24} - 10.12$ (c 0.35, CHCl₃). IR $\nu_{\max}^{\text{KBr}} \text{ cm}^{-1}$: 3442 (—OH), 2933, 2867, 1716 (—C=O). ¹H NMR (δ : ppm, CDCl₃): 0.68 (3H, s, 26 β -CH₃); 0.71 (3H, s, 24 β -CH₃); 0.84 (3H, s, 25 β -CH₃); 0.87 (3H, s, $J = 6.7$ Hz, 23 β -CH₃); 0.98 (3H, s, 29 β -CH₃); 1.08 (3H, s, 27 α -CH₃); 3.26 (2H, s, 30-CH₂); 9.45 (1H, s, 28-CHO). ¹³C NMR data (see table 1). EI MS m/z : 456 (7), 438 (10), 427 (100), 273 (25), 109 (60). HR-ESI-MS m/z : 479.3528 [M + Na]⁺ (calcd for C₃₀H₄₈O₃, 479.3501).

3.6 Tripterfrielanon B (2)

White amorphous powder. $[\alpha]_D^{25} - 22.8^\circ$ (c 0.20, CHCl₃). IR $\nu_{\max}^{\text{KBr}} \text{ cm}^{-1}$: 2929, 2869, 1750, 1731 (C=O), 1388, 1162. ¹H NMR (δ : ppm, CDCl₃): 0.89 (3H, d, $J = 6.0$ Hz, 23 β -CH₃); 0.72 (3H, s, 24 β -CH₃); 0.87 (3H, s, 25 β -CH₃); 1.03 (3H, s, 26 β -CH₃); 0.99 (3H, s, 27 α -CH₃); 1.21 (3H, s, 28-CH₃); 1.07 (3H, s, 29 β -CH₃); 3.84 (2H, s, 30-CH₂); 8.14 (1H, s, HCO-). ¹³C NMR data (see table 1). EI MS m/z : 470 (18), 455 (12), 273 (28), 123 (100), 135 (46), 109 (83), 81 (92), 67. HR-ESI-MS m/z : 493.3682 [M + Na]⁺ (calcd for C₃₁H₅₀O₃, 493.3681).

Acknowledgements

The authors thank Mr. Ji-Hua YU for collecting plant material and Mr. Shao-You LI for cytotoxicity assay.

References

- [1] Y. Kashiwada, M. Nishizawa, T. Yamagishi, T. Tanaka, G.L. Nanaka, L. Cosentino, J.V. Snider, K.H. Lee. *J. Nat. Prod.*, **58**, 392 (1995).
- [2] W.Z. Qin, C.H. Liu, S.M. Yang, G.D. Zhu, F.Y. Han, L. Fang, S.F. Feng. *Chinese Med J.*, **94**, 827 (1981).
- [3] X.Q. Liu, D.G. Wu. *Chin. Med.*, **24**, 395 (1993).
- [4] G.W. Qin, X.M. Yang, W.H. Gu, B.D. Wang, Z.X. Chen, R.X. Guo, K.W. Shao. *Acta Chimica Sinica*, **40**, 637 (1982).
- [5] C.W. Chang, T.S. Wu, Y.S. Hsieh, S.C. Kuo, P.D. Chao. *J. Nat. Prod.*, **62**, 327 (1999).
- [6] H. Nozaki, H. Suzuki, T. Hirayama, R. Kasai, R.Y. Wu, K.H. Lee. *Phytochemistry*, **25**, 479 (1986).
- [7] S. Vasanth, A.B. Kundu. *J. Nat. Prod.*, **55**, 1149 (1992).
- [8] T. Mosmann. *Immunol. Methods*, **65**, 55 (1983).