

This article was downloaded by:

On: 23 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Journal of Carbohydrate Chemistry

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713617200>

PRACTICAL β -STEREOSELECTIVE O-GLYCOSYLATION OF PHENOLS WITH PENTA-O-ACETYL- β -D-GLUCOPYRANOSE

Yeon Soo Lee^a; Eun Suk Rho^a; Yong Ki Min^a; Bum Tae Kim^a; Ki Ho Kim^b

^a Korea Research Institute of Chemical Technology, Taejeon, Korea ^b Bioland Ltd., Chonan-city, Chungcheongnam-do, Korea

Online publication date: 31 July 2001

To cite this Article Lee, Yeon Soo , Rho, Eun Suk , Min, Yong Ki , Kim, Bum Tae and Kim, Ki Ho(2001) 'PRACTICAL β -STEREOSELECTIVE O-GLYCOSYLATION OF PHENOLS WITH PENTA-O-ACETYL- β -D-GLUCOPYRANOSE', *Journal of Carbohydrate Chemistry*, 20: 6, 503 – 506

To link to this Article: DOI: 10.1081/CAR-100106933

URL: <http://dx.doi.org/10.1081/CAR-100106933>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

COMMUNICATION

**PRACTICAL β -STEREOSELECTIVE
O-GLYCOSYLATION OF PHENOLS WITH
PENTA-O-ACETYL- β -D-GLUCOPYRANOSE**

Yeon Soo Lee,^{1,*} Eun Suk Rho,¹ Yong Ki Min,¹ Bum Tae Kim,¹
and Ki Ho Kim²

¹Korea Research Institute of Chemical Technology, Yusong-ku,
Taejon, 305-343, Korea


²Bioland Ltd. 39-4 Songjeong-ri, Byongchon-myun, Chonan-city,
Chungcheongnam-do, 330-860, Korea

O-Glycosylation of phenol derivatives is an important synthetic method for natural products containing carbohydrate molecules.¹ Although various *O*-glycosylation methods have been studied in order to improve stereoselectivity and yield by changing the leaving groups, catalysts and solvents, previous methods are not always satisfactory in terms of ease and practicality.^{1–7} Recently Oyama *et al.* reported the glycosylation of phenols with 2,3,4,6-tetra-*O*-acetyl- α -D-glucopyranosyl fluoride in the presence of $\text{BF}_3 \cdot \text{OEt}_2$ and hindered base in order to achieve high β -stereoselectivity.^{7,8} However, it is not easy to obtain the starting glycosylating reagent, 2,3,4,6-tetra-*O*-acetyl- α -D-glucopyranosyl fluoride.⁸

Compared to other leaving groups, an acetoxy group can be a very convenient anomeric leaving group in glycosylation reactions. For example penta-*O*-acetyl- β -D-glucopyranose can be prepared more easily than other glucosyl donors. We have therefore tried to develop a new β -stereoselective *O*-glycosylation of phenols by using penta-*O*-acetyl- β -D-glucopyranose. The results of our glycosylation experiments of phenol are shown in **Table 1**.

It is well known that the glycosylation of phenols with penta-*O*-acetyl- β -D-glucopyranose gives phenyl β -glucosides as major products, due to the neighbouring group effect of the 2-*O*-acetyl group. A serious drawback of the method is conversion to some extent of the kinetically stable phenyl β -glucoside **3b** into the thermodynamically stable α -anomer **3a** during the glycosylation reaction (entry 1).⁹ However, glycosylation of phenol with penta-*O*-acetyl- β -D-glucose in the presence of $\text{BF}_3 \cdot \text{Et}_2\text{O}$ and organic base results in both high yield and high β -

Table 1. Glycosylation of Phenol with Penta-*O*-acetyl- β -D-glucopyranose



	BF ₃ ·OEt ₂ (eq)	base (eq)	rxn time (h)	yield ^a (%)	ratio ^b (3a/3b)
1	2.5	0	5.5	72	18/82
2	2.5	TEA (0.1)	3	89	7/93
3	2.5	TEA (0.5)	3	92	2/98
4	2.5	TMG ^c (0.5)	5	89	4/96
5	2.5	2,6-lutidine (0.5)	5.5	91	3/97
6	2.5	DTBMP ^d (0.5)	7	88	3/97
7	2.5	TEA (1)	5	87	3/97
8	1.5	TEA (1)	11	72	1/99
9	1.5	TEA (0.5)	5.5	87	2/98
10	4	TEA (2)	2	80	2/98
11	10	TEA (1)	0.5	92	9/91

a: isolated yield. b: isolated ratio. c: 1,1,3,3-tetramethylguanidine. d: 2,6-di-*tert*-butyl-4-methylpyridine.

stereoselectivity compared to glycosylation without organic base as shown in **Table 1**.


Glycosylation of phenol in the presence of 2.5 equivalents of BF₃·Et₂O and 0.5 equivalent of triethylamine to penta-*O*-acetyl- β -D-glucopyranose gave the highest yield and excellent β -stereoselectivity (entry 3). The addition of more than 0.5 equivalent of triethylamine decreased the reaction rate and the yield, but still gave high β -stereoselectivity (entry 7). It is reported that base can abstract the proton quickly from the adduct intermediate cation⁷ of phenol in the glycosylation of phenols with 2,3,4,6-tetra-*O*-acetyl- α -D-glucopyranosyl fluoride, and prevent isomerization of the β -isomer to the α -anomer.

We have also examined the glycosylation of several substituted phenols with penta-*O*-acetyl- β -D-glucopyranose (**Table 2**). As shown in **Table 2**, the glycosylation of phenols substituted with an electron withdrawing group retarded the reaction rate (**2h-2j**). The reaction rate was also affected by the location of substituent, the order of reaction rate being; ortho < meta < para, irrespective of substituent structure.

By taking advantage of this method, we have synthesized arbutin,¹⁰ a widely used natural whitening agent. Deacetylation¹¹ of phenyl- β -glucoside **4b** with sodium methoxide in methanol affords arbutin as shown in the scheme below.




Table 2. Glycosylation of Several Phenols with Penta-*O*-acetyl- β -D-glucopyranose



2	R	rxn time (h)	yield (%) ^a	product ratio (α/β) ^b
a	H	3	92	3a/3b =2/98
b	4-CH ₃ COO	24	92	4a/4b =2/98
c	4-Cl	7	91	5a/5b =1/99
d	4-MeO	8	90	6a/6b =2/98
e	2-Br	27	86	7a/7b =1/99
f	3-Br	10	88	8a/8b =2/98
g	4-Br	6	91	9a/9b =2/98
h	2-NO ₂ ^c	120	75	10a/10b =2/98
i	3-NO ₂	92	86	11a/11b =1/99
j	4-NO ₂	41	91	12a/12b =2/98
k	3-C ₆ H ₅	3	83	13a/13b =2/98

a: The yields were isolated b: The ratios of α/β were determined by gas chromatography.
c: 5 equivalents of BF₃·OEt₂ and 1 equivalent of TEA were used.



Scheme.

EXPERIMENTAL

A typical procedure for the glucosylation is as follows: to a mixture of penta-*O*-acetyl- β -D-glucopyranose (10 mmol) and phenol derivative (15 mmol) under a nitrogen atmosphere, was added triethylamine (5 mmol) in methylene chloride (20 mL) and then BF₃·OEt₂ (25 mmol) in methylene chloride (5 mL) over 30 min. The reaction mixture was then kept stirring for the time shown in **Table 2**. Saturated aqueous sodium bicarbonate (20 mL) was added to the reaction mixture, and the reaction mixture was then extracted with ethyl acetate (30 mL \times 2). The organic layer was dried over magnesium sulfate and the filtrate was concentrated in a vacuum evaporator. The residue was chromatographed on silica gel to give the phenyl glucoside product.



ACKNOWLEDGMENT

The authors are grateful to the Ministry of Commerce, Industry and Energy of Korea for financial support of this work.

REFERENCES

1. Toshima, K.; Tatsuta, K. Recent Progress in *O*-Glycosylation Methods and Its Application to Natural Products Synthesis. *Chem. Rev.* **1993**, *93*, 1503–1531.
2. Matsumoto, T.; Katsuki, M.; Suzuki, K. Rapid *O*-Glycosidation of Phenols with Glycosidyl Fluoride by using the Combinational Activator, Cp₂HfCl₂-AgClO₄. *Chem. Lett.* **1989**, 437–440.
3. Smits, E.; Engberts, Jan B. F. N.; Kellogg, Richard M.; van Doren, Henk A. Reliable Method for the Synthesis of Aryl β-D-glucopyranosides, Using Boron Trifluoride-Diethyl Ether as Catalyst. *J. Chem. Soc., Perkin Trans.1* **1996**, 2873–2877.
4. Li, Z-J.; Cai, L-N.; Cai, M-S. Studies on Glycosides X. An Alternate Method for Highly Stereoselective Synthesis of Alkyl-β-d-Glucopyranosides. *Synth. Commun.*, **1992**, *22*, 2121–2124.
5. Clerici, F.; Gelmi, M. L.; Mottadelli, S. Glycosides. Part 1. New Synthesis of 1,2-*trans*-*O*-acetyl Glycosides, *via* Tributyltin Phenoxides. *J. Chem. Soc., Perkin Trans.1* **1994**, 985–988.
6. Oyama, K-I.; Kondo, T., Highly Efficient β-Glucosylation of the Acidic Hydroxyl Groups, Phenol and Carboxylic Acid with a Peracetylated Glucosyl Fluoride Using a Combination of BF₃·Et₂O and DTBMP as a Promoter. *Synlett.* **1999**, 1627–1629.
7. Yamaguchi, M.; Horiguchi, A.; Fukuda, A.; Minami, T. Novel Synthesis of Aryl 2,3,4,6-Tetra-*O*-Acetyl-d-glucopyranosides. *J. Chem. Soc., Perkin Trans.1* **1990** 1079–1082.
8. Shimizu, M.; Togo, H.; Yokoyama, M. Chemistry of Glucosyl Fluoride. *Synthesis*, **1998**, 799–822.
9. It was also proved in our experiments that phenyl-β-glucoside (**3a**) was not isomerized to phenyl-α-glucoside (**3b**) in the presence of BF₃·Et₂O (2.5 eq) and triethylamine (0.5 eq) in methylene chloride, but 50% of phenyl-β-glucoside was isomerized to phenyl-α-glucoside in the presence of only BF₃·Et₂O (2.5 eq) within 10 h at rt.
10. Maeda, K.; Fukuda, M. Arbutin: Mechanism of Its Depigmenting Action in Melanocyte Culture. *J. Pharmacol. Exp. Ther.* **1996**, *276*, 765–769.
11. Hirokai, T.; Shigeru, I. A Process for the Preparation of Arbutin and Analogs as Drugs. *Kokai Tokyo Koho JP.*, 87,226,974, Oct 05, 1987.; *Chem. Abstr.* **1988**, *109*, 6894k.

Received March 30, 2001

Accepted May 31, 2001



Request Permission or Order Reprints Instantly!

Interested in copying and sharing this article? In most cases, U.S. Copyright Law requires that you get permission from the article's rightsholder before using copyrighted content.

All information and materials found in this article, including but not limited to text, trademarks, patents, logos, graphics and images (the "Materials"), are the copyrighted works and other forms of intellectual property of Marcel Dekker, Inc., or its licensors. All rights not expressly granted are reserved.

Get permission to lawfully reproduce and distribute the Materials or order reprints quickly and painlessly. Simply click on the "Request Permission/Reprints Here" link below and follow the instructions. Visit the [U.S. Copyright Office](#) for information on Fair Use limitations of U.S. copyright law. Please refer to The Association of American Publishers' (AAP) website for guidelines on [Fair Use in the Classroom](#).

The Materials are for your personal use only and cannot be reformatted, reposted, resold or distributed by electronic means or otherwise without permission from Marcel Dekker, Inc. Marcel Dekker, Inc. grants you the limited right to display the Materials only on your personal computer or personal wireless device, and to copy and download single copies of such Materials provided that any copyright, trademark or other notice appearing on such Materials is also retained by, displayed, copied or downloaded as part of the Materials and is not removed or obscured, and provided you do not edit, modify, alter or enhance the Materials. Please refer to our [Website User Agreement](#) for more details.

[Order now!](#)

Reprints of this article can also be ordered at

<http://www.dekker.com/servlet/product/DOI/101081CAR100106933>