

This article was downloaded by:

On: 23 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Journal of Carbohydrate Chemistry

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713617200>

EFFICIENT SYNTHESIS OF POLYLACTOSAMINE STRUCTURES THROUGH REGIOSELECTIVE GLYCOSYLATIONS[1]

Therese Buskas^a; Peter Konradsson^b; Stefan Oscarson^a

^a Department of Organic Chemistry, Arrhenius Laboratory, Stockholm University, Stockholm, Sweden

^b Department of Chemistry, University of Linköping, Linköping, Sweden

Online publication date: 30 November 2001

To cite this Article Buskas, Therese , Konradsson, Peter and Oscarson, Stefan(2001) 'EFFICIENT SYNTHESIS OF POLYLACTOSAMINE STRUCTURES THROUGH REGIOSELECTIVE GLYCOSYLATIONS[1]', *Journal of Carbohydrate Chemistry*, 20: 7, 569 – 583

To link to this Article: DOI: 10.1081/CAR-100108275

URL: <http://dx.doi.org/10.1081/CAR-100108275>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

EFFICIENT SYNTHESIS OF POLYLACTOSAMINE STRUCTURES THROUGH REGIOSELECTIVE GLYCOSYLATIONS¹

Therese Buskas,¹ Peter Konradsson,² and Stefan Oscarson¹

¹Department of Organic Chemistry, Arrhenius Laboratory, Stockholm University, S-106 91 Stockholm, Sweden

²Department of Chemistry, University of Linköping, S-581 83 Linköping, Sweden

ABSTRACT

Di-, tri- and tetramers of β -(1 \rightarrow 3)-linked *N*-acetylglucosamine residues have been synthesised as their methyl glycosides, to be used in ITC binding studies to various galectins. The synthetic strategy involves two types of regioselective glycosylations: couplings of a galactosyl donor to 3,4-diol *N*-tetrachlorophthalimido glucose acceptors to give the lactosamine monomer building blocks, and subsequent formation of the oligomers through consecutive couplings of lactosamine donors to 2',3',4'-lactosamine acceptors, with high selectivity for the desired products.

INTRODUCTION

Polyglucosamines consist of β -(1 \rightarrow 3)-linked *N*-acetylglucosamine chains attached to sphingolipids and proteins. They are the backbones of keratan sulfates and are present at the surface of mammalian cells with or without distal decorations (sulfate groups, single monosaccharides, and numerous oligosaccharide determinants).² A few chemical syntheses of polyglucosamine structures have already been published, the tetramer by Alais and Veyrieres,^{3–5} Srivastava and Hindsgaul⁶ and Shimizu et al.,⁷ the latter performed on solid phase, and the trimer by Nilsson and Norberg⁸ as an intermediate in the synthesis of a trimeric Lewis x structure. Also enzymatic syntheses have been described.⁹




Figure 1.

Galectins, named after their preferential binding to β -galactose motifs, are known to have a number of important biological functions,^{10,11} and to further investigate these, the study of their binding to poly-lactosamine structures was important. Lactosamine oligomers with varying length were essential for these studies. Consequently, the di- and trimer of β -(1 \rightarrow 3)-linked *N*-acetyllactosamine residues (**17** and **20**, Figure 1) were synthesised. These will primarily be used in isothermal titration calorimetry (ITC) studies of their binding to galectins, to render possible the determination of the size of the binding site and the thermodynamics of the binding.


RESULTS AND DISCUSSION

Although a number of syntheses of 2-amino-2-deoxylactose from a disaccharide precursor have been published,^{3,12–14} the alternative to start from monosaccharides is still a relevant option. The “drawback” of having to perform an extra glycosylation reaction is balanced by the advantage of the easy access to monosaccharide precursors, protected in a suitable way for subsequent reactions. Especially with phthalimido-protected derivatives the latter approach is feasible, since it has been found that the coupling between galactosyl donors and, easily obtainable, 3,4-diol *N*-phthalimido glucose acceptors is high-yielding and very regioselective for the 4-position due to the bulkiness of the phthalimido group.^{15–17} This regioselectivity is even more pronounced for tetrachlorophthalimido compounds.¹⁸

Regioselective benzylation of the tin-activated pentenyl glycoside **1**¹⁹ gave directly the 3,4-diol **2** in 69% yield (Scheme 1). This procedure was found to be more convenient than the two-step one, involving 4,6-*O*-benzylidene formation and reductive opening used earlier.²⁰ The silver triflate-promoted coupling between **2** and galactosyl donor **3** (obtained from the known corresponding thioethyl glycoside²¹ by bromine treatment) proceeded in high yield (87%) and with regioselectivity. The 1 \rightarrow 4-linkage in the product **4** was proven by the downfield shift of the H-3 proton in the benzoylated derivative **5**. To allow for more flexibility in the synthetic strategy, thioglycoside donors were also synthesised (Scheme 2). Hence, derivatives **10** and **12** were synthesised in a parallel fashion from the same galactosyl donor (**3**) and the 3,4-diol thioglycoside acceptors **7** and **8**,¹⁷ respectively. Once more the glycosylations gave high regioselectivity and yield of products.

Since the primary use of the target compounds was to perform ITC-studies of






Scheme 1. *i:* $(\text{Bu}_3\text{Sn})_2\text{O}$, MeOH; *ii:* BnBr, QBr, 90 °C ; *iii:* Br_2 , CH_2Cl_2 ; *iv:* AgOTf, CH_2Cl_2 , 3Å MS, -40°C; *v:* BzCl, pyridine.


their binding to various animal lectins, and accordingly no conjugate formation was necessary, it was decided to synthesise them as their methyl β -glycosides. Therefore, the pentenyl glycoside **5** was transformed into the methyl glycoside **13** (93%) by treatment with 5 equiv of MeOH in the presence of NIS/TESOTf (triethylsilyl triflate) (Scheme 3). If a larger excess of MeOH was used, the major reaction was addition of MeOH to the double bond.

Another known type of regioselective glycosylation, preferentially used in sialylation of galactose derivatives, was utilised to synthesise oligomers. In 2,3,4-triol galactose acceptors a high selectivity for the 3-hydroxyl group is usually observed in glycosylations, sialylation on 2,4-protected derivatives being often impossible.²² Already anticipating this route when choosing galactosyl donor **3**, the acetyl groups of compound **13** were now removed chemoselectively by treatment with HCl in MeOH²³ to give the 2',3',4'-triol **14** (82%). In the coupling reaction to give the dimer it was most important to match the reactivity of the reacting species, acceptor, donor and promoter, to get a high yield of product (Table 1, Scheme 4). However, all the glycosylations were regioselective for the



Scheme 2. *i:* $(\text{Bu}_3\text{SnO})_2$, MeOH; *ii:* BnBr (5 eq), TEABr, toluene, 90 °C; *iii:* AgOTf, CH_2Cl_2 , 3Å MS, -40°C; *iv:* BzCl, pyridine.













Scheme 3. *i*: MeOH, NIS, TESOTf, CH₂Cl₂, 3 Å MS; *ii*: 5% HCl/MeOH.


3-position. The use of an effective promoter combined with especially tetra-chlorophthalimido donors gave mainly decomposition of the donor. If the promoter was changed into a less active one or if a phthalimido donor was employed, good yields of the desired β -(1 \rightarrow 3)-linked products **15** and **16** were obtained.

Choosing the most high-yielding conditions (donor **10**, promoter MeOTf) from this dimer glycosylation study, syntheses of a trimer and a tetramer were attempted (Scheme 5). Chemoselective removal of the three acetates from derivative **15**, again proceeded smoothly to give an 88% yield of compound **18**. Reasoning

Table 1. Glycosylation of Acceptor **14**. General Conditions: 1.7 Equiv of the Donor, 3 Å MS, CH₂Cl₂, Ar. Bromide Donors were Prepared by Br₂ Treatment of the Corresponding Thioglycoside Just Prior to Coupling. ^aThe Donor Decompose

Donor	Promoter	temp	yield (%) / product
	NIS/TESOTf	rt	17%/ 15
	IDCT	rt	44%/ 15
	NIS/TMSOTf	rt	a
	MeOTf	rt	69%/ 15
	DMTST	rt	61%/ 15
	AgOTf	-40 °C	a
	NIS/TESOTf	rt	50%/ 16
	AgOTf	-40 °C	62%/ 16






Scheme 4.


that glycosylation at the 3'-position should even more decrease the already low nucleophilicity of the 2'- and 4'-hydroxyl groups, pentaol **18** was tried as an acceptor in a glycosylation employing the conditions used above, and was found to give the trimer **19** in a most satisfactory yield (61%). Continued use of this methodology then effectively gave first the heptaol acceptor **21** (81%) and then the tetramer **22** (60%, 69% calculated on consumed acceptor).

Since the 3-position of the glucosamine residue, as discussed and shown, is quite unreactive also, perhaps its protection was not necessary, i.e., the regioselective glycosylation could be performed on a 3,2',3',4'-tetraol acceptor using a 3-OH donor. This approach would have the advantage of fewer reaction steps (no benzylation) and more easily performed deacetylation steps (nonchemoselective). To test this hypothesis the pentenyl glycoside **4** was transformed into the methyl glycoside **23** using the same conditions as for the benzyolated analogue **5** above, and in comparable yield (88%) (Scheme 6). Removal of the acetyl protecting groups then gave the tetraol **24** in almost quantitative yield (97%). Coupling between acceptor **24** and the non-benzyolated donor **9** using methyl triflate as promoter pro-



Scheme 5. *i*: HCl/MeOH-CH₂Cl₂ 1:1; *ii*: MeOTf, CH₂Cl₂, 3Å MS; *iii*: H₂NNH₂-H₂O, MeCN-THF-EtOH 2:1:1, 70 °C; *iv*: Ac₂O, pyridine; *v*: H₂, Pd/C; *vi*: 1M NaOMe, MeOH.





Scheme 6. *i*: MeOH, NIS, TESOTf, CH₂Cl₂, 3 Å MS; *ii*: 5% HCl/MeOH; *iii*: 9, MeOTf, CH₂Cl₂, 3 Å MS.

ceeded to give one major product, which by NMR was shown to be the expected β -(1 \rightarrow 3)-linked tetrasaccharide **25** (46%), proving this to be an attractive alternative pathway to lactosamine oligomers.

Deprotection of the oligomers **15** and **19** was accomplished using standard conditions, hydrazine hydrate at elevated temperature followed by acetylation, hydrogenolysis and deacetylation with sodium methoxide (Scheme 5). Although rather straightforward, the yield in the deprotection is acceptable but not high, as is also the case with deprotections of similar compounds found in the literature.^{3–8} The yields are normally in the range of 40–50% corresponding to an 80–84% yield in each step in a four-step deprotection scheme.

EXPERIMENTAL

General Methods. Organic solutions were dried over MgSO₄ before concentrations, which were performed under reduced pressure at $\leq 40^\circ\text{C}$ (water bath). TLC was performed on silica gel 60 F₂₅₄ (Merck) with detection by UV light and/or charring with 8% sulfuric acid, ninhydrin or AMC (ammonium molybdate 10g, cerium(IV) sulfate 2 g, dissolved in aq 10% H₂SO₄ 2 L). Silica gel (0.0404–0.063 mm, Amicon) was used for column chromatography. NMR spectra were recorded in CDCl₃ at 25 °C (internal standard Me₄Si, $\delta = 0.00$) unless otherwise stated, using a Varian Mercury 300 at 300 MHz (¹H) or 75 MHz (¹³C) or a Varian Inova 400 at 400 MHz (¹H) or 100 MHz (¹³C).

Pent-4-enyl 6-O-Benzyl-2-deoxy-2-tetrachlorophthalimido- β -D-glucopyranoside (2). Bis(tributyltin) oxide (14.5 mL, 28.54 mmol) was added to a solution of **1**¹⁹ (10.5 g, 20.39 mmol) in dry MeOH (120 mL) stirred under argon. After stirring at 75°C for 2 h the reaction mixture was concentrated and dried *in vacuo*. The oily tin ether complex was dissolved in benzyl bromide (40 mL) and tetrabutylammonium bromide (7.22 g, 22.4 mmol) was added. After stirring at 90°C for 20 h benzyl bromide was distilled from the reaction vessel. Flash chromatography (two columns: toluene-EtOAc 20:1 \rightarrow 5:1 and toluene-EtOAc 5:1) of



the oily residue furnished **2** (8.51 g, 14.07 mmol, 69%) having NMR data in agreement with those reported elsewhere.²⁰

Pent-4-enyl (2,3,4-Tri-*O*-acetyl-6-*O*-benzyl- β -D-galactopyranosyl)-(1 \rightarrow 4)-3-*O*-benzoyl-6-*O*-benzyl-2-deoxy-2-tetrachlorophthalimido- β -D-glucopyranoside (5**).** A solution of ethyl 2,3,4-tri-*O*-acetyl-6-*O*-benzyl-1-thio- β -D-galactopyranoside²¹ (6.59 g, 14.96 mmol) and bromine (2.3 mL, 44.89 mmol) in CH₂Cl₂ (100 mL) was stirred under argon for 30 min at room temperature. The mixture was then concentrated and co-concentrated twice with toluene (Na-dried). A solution of the residue and **2** (6.87 g, 11.26 mmol) in CH₂Cl₂ (150 mL) was stirred with 3Å MS for 30 min at -60°C before AgOTf (7.30 g, 28.40 mmol) was added. The reaction was quenched with triethylamine (6 mL) after 1 h and the mixture was stirred for another 15 min, filtered through Celite and concentrated. Flash chromatography of the residue (toluene-EtOAc 6:1) gave pent-4-enyl (2,3,4-tri-*O*-acetyl-6-*O*-benzyl- β -D-galactopyranosyl)-(1 \rightarrow 4)-6-*O*-benzyl-2-deoxy-2-tetrachlorophthalimido- β -D-glucopyranoside (**4**) (9.51 g, 9.79 mmol, 87%). NMR (CDCl₃): ¹³C, δ 21.1 (2C), 21.2 (CH₃ acetyl), 29.1, 30.4 (CH₂ pentenyl), 56.7, 67.4, 68.0, 68.2, 69.0, 69.1, 69.6, 71.0, 72.5, 73.6, 73.8, 74.4, 81.74 (C-2-6, 2'-6', OCH₂ pentenyl, CH₂ benzyl), 98.0 (C-1), 101.3 (C-1'), 115.0 (CH₂ = pentenyl), 125.6-140.1 (aromatic C, CH = pentenyl), 162.4, 162.5 (C=O TCP), 169.4, 170.1, 170.3 (C=O acetyl); ¹H, δ 1.53-1.64 (m, 2H, CH₂ pentenyl), 1.89-1.94 (m, 2H, CH₂ pentenyl), 1.97, 1.98, 2.06 (s, 9H, CH₃ acetyl), 3.41-3.87 (m, 8H, OCH₂ pentenyl, H-4, 5, 5', 6, and 6'), 4.17 (dd, 1H, H-2), 4.25-4.55 (5H, H-1', 3, CH₂ benzyl), 4.71 (d, 1H, CH₂ benzyl), 4.82-4.87 (m, 2H, CH₂ = pentenyl), 4.94 (dd, 1H, H-3'), 5.09 (d, 1H, *J*_{1,2} 8.42 Hz, H-1), 5.17 (dd, 1H, H-2'), 5.37 (d, 1H, H-4'), 5.58-5.74 (m, 1H, CH = pentenyl), 7.15-7.36 (m, 10H, aromatic H). Benzoyl chloride (1.63 mL, 14.0 mmol) and pyridine (2.3 mL) were added to a stirred solution of **4** (1.38 g, 1.40 mmol) in dry CH₂Cl₂ (20 mL) and the reaction mixture was stirred at ambient temperature overnight and then concentrated. The residue was dissolved in CH₂Cl₂ and the organic layer was washed with aq 1M HCl, satd aq NaHCO₃, and H₂O, dried and concentrated. Flash chromatography (toluene-EtOAc 8:1) of the residue gave **5** (1.35 g, 1.24 mmol, 88%). NMR (CDCl₃): ¹³C, δ 21.1 (2C), 21.3 (CH₃ acetyl), 29.1, 30.4 (CH₂ pentenyl), 56.1, 66.3, 67.2, 67.8, 69.5, 70.0, 71.6, 72.3, 73.5, 74.2, 75.1, 75.9 (C-2-6, 2'-6', OCH₂ pentenyl, CH₂ benzyl), 98.2 (C-1), 100.7 (C-1'), 115.2 (CH₂ = pentenyl), 127.8-140.4 (aromatic C, CH = pentenyl), 162.6, 162.8 (C=O TCP) 165.6 (C=O benzoyl), 169.2, 170.2 (2C) (C=O acetyl); ¹H, δ 1.56-1.64 (m, 2H, CH₂ pentenyl), 1.85, 1.92, 1.94 (s, 9H, CH₃ acetyl), 1.89-1.98 (m, 2H, CH₂ pentenyl), 3.47-3.51 (m, 1H, OCH₂ pentenyl), 2.70 (dd, 1H), 2.91 (dd, 1H) 3.65-3.69 (m, 1H), 3.74-3.87 (m, 3H, OCH₂ pentenyl, H-6), 4.07 (d, 1H, CH₂ benzyl), 4.16 (dd, 1H, H-4), 4.25 (d, 1H, CH₂ benzyl), 4.38 (dd, 1H, H-2), 4.45 (d, 1H, *J*_{1,2} 8.06 Hz, H-1'), 4.54 (d, 1H, CH₂ benzyl), 4.74-4.89 (m, 4H, H-3', CH₂ benzyl, CH₂ = pentenyl), 4.94 (dd, 1H, H-2'), 5.22 (d, 1H, H-4'), 5.35 (d, 1H, *J*_{1,2} 8.42 Hz, H-1), 5.64-4.74 (m, 1H, CH = pentenyl), 5.88 (dd, 1H, H-3) 7.14-8.17 (m, 15H, aromatic H). MS (MALDI-TOF): (M+Na)⁺ 1108.1.



Ethyl (2,3,4-Tri-*O*-acetyl-6-*O*-benzyl- β -D-galactopyranosyl)-(1 \rightarrow 4)-3-*O*-benzoyl-6-*O*-benzyl-2-deoxy-2-tetrachlorophthalimido-1-thio- β -D-glucopyranoside (10). Bis(tributyltin) oxide (3.53 mL, 6.93 mmol) was added to a solution of **6**²⁴ (3.09 g, 6.3 mmol) in dry MeOH (60 mL) stirred under argon. The reaction was stirred at 75°C for 2 h, concentrated and dried in vacuo. The yellow oil was dissolved in toluene (Na-dried) (70 mL) and benzyl bromide (3.75 mL, 31.5 mmol) and tetrabutylammonium bromide (2.13 g, 6.62 mmol) were added. The reaction was kept at 90°C for 21 h and then concentrated. Purification by flash chromatography (toluene-EtOAc 4:1) furnished ethyl 6-*O*-benzyl-2-deoxy-2-tetrachlorophthalimido-1-thio- β -D-glucopyranoside (**7**) (2.45 g, 4.22 mmol, 67%). NMR (CDCl₃): ¹³C, δ 15.3 (CH₃CH₂S), 24.5 (CH₂S), 56.2, 70.6, 72.4, 73.7, 74.0, 78.3 (C-2–6, CH₂ benzyl), 81.1 (C-1), 125.5–140.4 (aromatic C). Ethyl 2,3,4-tri-*O*-acetyl-6-*O*-benzyl-1-thio- β -D-galactopyranoside²¹ (1.9 g, 4.34 mmol) was converted into the corresponding bromosugar **3**, by treatment with bromine (0.45 mL, 8.68 mmol). A mixture of **3**, acceptor **7** (1.8 g, 3.1 mmol) and 3 Å MS in CH₂Cl₂ (40 mL) was stirred at –60°C for 30 min, whereafter AgOTf (2.23 g, 8.68 mmol) was added. After 45 min the reaction was quenched by addition of triethylamine (1 mL) and the mixture was diluted with CH₂Cl₂ and filtered through Celite and concentrated. Flash chromatography (toluene-EtOAc 6:1) gave ethyl (2,3,4-tri-*O*-acetyl-6-*O*-benzyl- β -D-galactopyranosyl)-(1 \rightarrow 4)-6-*O*-benzyl-2-deoxy-2-tetrachlorophthalimido-1-thio- β -D-glucopyranoside (**9**) (2.61 g, 2.73 mmol, 88%). NMR (CDCl₃): ¹³C, δ 15.3 (CH₃CH₂S), 20.9, 21.0, 21.1 (CH₃ acetyl), 24.3 (CH₂S), 56.0, 67.6, 68.2, 68.5, 69.3, 70.6, 71.2, 72.7, 73.8, 74.0, 78.6, 81.0 (C-2–6, 2'-6', CH₂ benzyl), 81.6 (C-1), 101.5 (C-1'), 125.9–140.3 (aromatic C), 162.9, 163.4 (C=O TCP), 169.3, 170.0, 170.2 (C=O acetyl). Benzoyl chloride (1.40 mL, 12.0 mmol) was added to a stirred solution of **9** (2.3 g, 2.40 mmol) in pyridine (35 mL). After stirring at room temperature overnight the reaction mixture was concentrated and worked up as described for **5**. Flash chromatography (toluene-EtOAc 4:1) of the crude product afforded **10** (2.17 g, 2.04 mmol, 85%). [α]_D +69°(c 1.1, CHCl₃) NMR (CDCl₃): ¹³C, α 15.0 (CH₃CH₂S), 20.4, 20.5, 20.7 (CH₃ acetyl), 24.2 (CH₂S), 54.5, 65.6, 66.6, 67.4, 69.4, 71.0 (2C), 72.3, 72.9, 73.6, 74.9, 78.7 (C-2–6, 2'-6', CH₂ benzyl), 80.5 (C-1), 100.0 (C-1'), 125.0–140.1 (aromatic C), 162.1, 162.6 (C=O TCP), 165.0 (C=O benzoyl), 168.6, 169.6 (C=O acetyl); ¹H, δ 1.24 (t, 3H, CH₃CH₂S), 1.83, 1.91, 1.94 (s, 9H, CH₃ acetyl), 2.64–2.75 (m, 3H, CH₂S), 2.89–2.92 (m, 1H), 3.30–3.34 (m, 1H), 3.70–3.79 (3H, H-5, H-6), 4.09 (d, 1H, CH₂ benzyl), 4.19 (dd, 1H, H-4), 4.26 (d, 1H, CH₂ benzyl), 4.46–4.55 (m, 3H, H-2, H-1', CH₂ benzyl), 4.76–4.79 (m, 2H, H-3', CH₂ benzyl), 4.95 (dd, 1H, H-4'), 5.47 (d, 1H, *J*_{1,2} 10.62 Hz, H-1), 5.95 (dd, 1H, H-3), 7.12–8.10 (m, 15H, aromatic H).

Anal. Calcd for C₄₉H₄₇O₁₅NCl₄: C, 55.33; H, 4.45%. Found: C, 55.19; H, 4.48%.

Ethyl (2,3,4-Tri-*O*-acetyl-6-*O*-benzyl- β -D-galactopyranosyl)-(1 \rightarrow 4)-3-*O*-benzoyl-6-*O*-benzyl-2-deoxy-2-phthalimido-1-thio- β -D-glucopyranoside (12). A solution of ethyl 2,3,4-tri-*O*-acetyl-6-*O*-benzyl-1-thio- β -D-galactopyranoside²¹ (3.45 g, 7.83 mmol) and bromine (0.8 mL, 15.65 mmol) in CH₂Cl₂ (60 mL) was



stirred for 30 min at room temperature and under argon. The mixture was then concentrated and co-evaporated twice from toluene (Na-dried). A solution of the residue and ethyl 6-*O*-benzyl-2-deoxy-2-phthalimido-1-thio- β -D-glucopyranoside¹⁷ (**8**, 2.89 g, 6.53 mmol) in CH₂Cl₂ (75 mL) was stirred with 3Å MS for 30 min at -60°C before AgOTf (4.0 g, 15.67 mmol) was added. After 1 h, triethylamine (3 mL) was added and the mixture was allowed to attain room temperature, whereafter it was filtered through Celite and concentrated. Flash chromatography of the residue (toluene-EtOAc 6:1) gave ethyl (2,3,4-tri-*O*-acetyl-6-*O*-benzyl- β -D-galactopyranosyl)-(1→4)-6-*O*-benzyl-2-deoxy-2-phthalimido-1-thio- β -D-glucopyranoside (**11**) (3.70 g, 4.51 mmol, 69%). NMR (CDCl₃): ¹³C, δ 15.4 (CH₃CH₂S), 20.9 (2C), 21.1 (CH₃ acetyl), 24.2 (CH₂S), 55.5, 67.6, 67.7, 68.5, 69.3, 71.0, 71.2, 72.6, 73.8, 73.9, 78.6, 81.3 (C-2-6, 2'-6', CH₂ benzyl), 81.7 (C-1), 101.5 (C-1'), 123.4-138.3 (aromatic C), 167.8, 168.2 (C=O NPhth), 169.4, 170.0, 170.2 (C=O acetyl). Benzoyl chloride (2.09 mL, 18.04 mmol) was added to a stirred solution of **11** (3.70 g, 4.51 mmol) in pyridine (100 mL). After stirring in room temperature for 15 h the reaction was concentrated and then worked up as described for compound **5**. Flash chromatography (toluene-EtOAc 6:1) gave **12** (3.63 g, 3.92 mmol, 87%). NMR (CDCl₃): ¹³C, δ 15.6 (CH₃CH₂S), 20.9, 20.9, 21.1 (CH₃ acetyl), 24.5 (CH₂S), 54.2, 66.1, 67.1, 68.0, 69.9, 71.5, 72.8 (2C), 73.3, 74.0, 75.8, 79.2 (C-2-6, 2'-6', CH₂ benzyl), 81.3 (C-1), 100.57 (C-1'), 125.5-138.0 (aromatic C), 165.2 (C=O benzoyl), 167.5, 167.8 (C=O NPhth), 169.1, 170.1 (2C) (C=O acetyl); ¹H, δ 1.23 (t, 3H, CH₃CH₂S), 1.84, 1.91, 1.94 (s, 9H, CH₃ acetyl), 2.61-2.72 (m, 3H, CH₂S), 2.90 (dd, 1H), 3.30 (dd, 1H), 3.74-3.80 (m, 3H, H-5, 6), 4.07 (d, 1H, CH₂ benzyl), 4.18 (dd, 1H, H-4), 4.25 (d, 1H, CH₂ benzyl), 4.47-4.55 (m, 3H, H-1 (*J*_{1,2} 10.2 Hz), H-2, CH₂ benzyl), 4.75-4.79 (m, 2H, H-3', CH₂ benzyl), 4.96 (dd, 1H, H-2'), 5.22 (d, 1H, H-4'), 5.50 (d, 1H, *J*_{1,2} 10.6 Hz, H-1'), 6.03 (dd, 1H, H-3), 7.14-8.12 (m, 19H, aromatic H). MS (MALDI-TOF): (M+Na)⁺ 948.3; (M+K)⁺ 964.3.

Methyl (6-*O*-Benzyl- β -D-galactopyranosyl)-(1→4)-3-*O*-benzoyl-6-*O*-benzyl-2-deoxy-2-tetrachlorophthalimido- β -D-glucopyranoside (14**).** A solution of **5** (2.23 g, 2.05 mmol) and dry MeOH (0.412 mL, 10.25 mmol) in CH₂Cl₂ (100 mL) was stirred under argon for 30 minutes, whereafter NIS (0.922 g, 4.10 mmol) and TESOTf (0.927 mL, 4.10 mmol) were added. After 15 min TLC indicated complete reaction, and the mixture was diluted with CH₂Cl₂, filtered through Celite, washed with satd aq Na₂S₂O₃ and satd aq NaHCO₃, dried and concentrated. Flash chromatography (toluene-EtOAc 5:1) yielded methyl (2,3,4-tri-*O*-acetyl-6-*O*-benzyl- β -D-galactopyranosyl)-(1→4)-3-*O*-benzoyl-6-*O*-benzyl-2-deoxy-2-tetrachlorophthalimido- β -D-glucopyranoside (**13**) (1.97 g, 1.91 mmol, 93%). NMR (CDCl₃): ¹³C, δ 20.8 (2C), 21.0 (CH₃ acetyl), 55.7, 57.1, 67.9, 66.9, 67.4, 69.7, 71.2, 71.3, 71.9, 73.2, 73.9, 74.8, 75.5 (C-2-6, 2'-6', CH₂ benzyl, CH₃O), 98.8 (C-1), 100.3 (C-1'), 125.3-140.1 (aromatic C), 162.3, 162.4 (C=O TCP), 165.3 (C=O benzoyl), 168.9, 169.9 (2C) (C=O acetyl); ¹H, δ 1.85, 1.91, 1.94 (s, 9H, CH₃ acetyl), 2.66 (dd, 1H), 2.89 (dd, 1H), 3.26-3.41 (m, 1H), 3.47 (s, 3H, CH₃O), 3.68 (m, 1H, H-5), 3.80 (m, 2H, H-6), 4.08 (d, 1H, CH₂ benzyl), 4.17 (dd, 1H, H-2), 4.25 (d, 1H, CH₂ benzyl), 4.38 (dd, 1H, H-4), 4.44 (d, 1H, *J*_{1,2} 7.87 Hz,



H-1'), 4.55 (dd, 1H, H-3'), 4.80 (d, 1H, CH₂ benzyl), 4.94 (dd, 1H, H-2'), 5.22 (d, 1H, H-4'), 5.28 (d, 1H, J_{1,2} 8.42 Hz, H-1), 5.89 (dd, 1H, H-3). Disaccharide **13** (1.74 g, 1.68 mmol) was dissolved in 5% HCl/MeOH (70 mL) and the mixture was stirred for 15 h and then concentrated. Flash chromatography (toluene-EtOAc 1:1–1:3) of the residue afforded **14** (1.25 g, 1.38 mmol, 82%). [α]_D +35° (c 1.0, CHCl₃) NMR: ¹³C (CDCl₃), δ 55.7, 57.3, 67.6, 68.3, 68.5, 72.3, 72.4, 72.9, 73.3 (2C), 73.7, 74.8, 76.9 (C-2–6, 2'-6', CH₂ benzyl, CH₃O), 98.9 (C-1), 103.7 (C-1'), 127.3–140.0 (aromatic C), 162.4, 162.5 (C=O TCP), 166.3 (C=O benzoyl); ¹H (CD₃OD + 2 drops of CDCl₃), δ 2.90 (dd, 1H), 3.09 (dd, 1H), 3.22–3.34 (m, 5H, H-3', CH₃O), 3.39 (dd, 1H, H-2'), 3.65 (d, 1H, H-4'), 3.82 (m, 1H, H-5), 3.95 (dd, 1H, H-6), 4.12 (dd, 1H, H-6), 4.17–4.34 (m, 4H, CH₂ benzyl, H-1', 2), 4.65 (dd, 2H, CH₂ benzyl), 5.30 (d, 1H, H-1), 5.85 (dd, 1H, H-3).

Anal. Calcd for C₄₂H₃₈O₁₂NCl₄: C, 56.64; H, 4.30%. Found: C, 56.44; H, 4.50%.

Methyl (2,3,4-Tri-O-acetyl-6-O-benzyl-β-D-galactopyranosyl)-(1→4)-(3-O-benzoyl-6-O-benzyl-2-deoxy-2-tetrachlorophthalimido-β-D-glucopyranosyl)-(1→3)-(6-O-benzyl-β-D-galactopyranosyl)-(1→4)-(3-O-benzoyl-6-O-benzyl-2-deoxy-2-tetrachlorophthalimido-β-D-glucopyranoside) (15). Methyl triflate (0.28 mL, 1.5 mmol) was added to a stirred mixture of **14** (0.417 g, 0.46 mmol), **10** (0.685 g, 0.644 mmol) and 3 Å MS in CH₂Cl₂. The reaction mixture was stirred under argon for 18 h, quenched by addition of triethylamine (1 mL), filtered through Celite and concentrated. Flash chromatography of the residue gave **15** (0.615 g, 0.322 mmol, 69%). NMR (CDCl₃): ¹³C, δ 20.9 (2C), 21.1 (CH₃ acetyl), 55.8, 57.2, 66.1, 67.0, 67.5, 67.6, 68.3, 69.9, 71.1, 71.4, 71.5, 71.8, 72.4, 72.8, 73.4, 73.4, 73.8, 74.0, 74.4, 74.9, 75.6, 76.7, 77.5 (C-2–6, 2'-6', 2''-6'', 2—6'', CH₃O, CH₂ benzyl), 99.1, 99.2, 100.5, 103.3 (C-1–1'''), 125.4–140.2 (aromatic C), 162.3, 163.7 (C=O TCP), 165.3, 165.7 (C=O benzoyl), 169.0, 170.0 (2C) (C=O acetyl). An aliquot of **15** was acetylated (pyridine-acetic anhydride 3:2) to give material having NMR (CDCl₃): ¹H, δ 1.86, 1.90, 1.91, 1.97 (CH₃ acetyl), 2.54–2.68 (m, 2H), 2.86–2.92 (m, 2H), 2.23–3.30 (m, 2H), 3.44 (s, 3H, CH₃O), 3.55–3.61 (m, 2H, H-3'), 3.68–3.77 (m, 5H), 4.05–4.24 (m, 8H, H-4, 4'', 2'', 1', CH₂ benzyl), 4.35 (dd, 1H, H-2), 4.46–4.62 (m, 4H, H-1''', 2', CH₂ benzyl), 4.75–4.82 (m, 3H, H-3''', CH₂ benzyl), 4.97 (dd, 1H, H-2'''), 5.13 (d, 1H, J_{1,2} 8.42 Hz, H-1''), 5.20–5.22 (m, 2H, H-4''', 1) 5.34 (d, 1H, H-4'), 5.74 (dd, 1H, H-3), 5.94 (dd, 1H, H-3''), 7.12–7.83 (m, 30H, aromatic H). The signals for H-2' (4.61 ppm) and H-4' (5.34 ppm) in the acetylated tetrasaccharide was shifted downfield, proving selective coupling at 3'-OH of the acceptor.

Methyl (2,3,4-Tri-O-acetyl-6-O-benzyl-β-D-galactopyranosyl)-(1→4)-(3-O-benzoyl-6-O-benzyl-2-deoxy-2-phthalimido-β-D-glucopyranosyl)-(1→3)-(6-O-benzyl-β-D-galactopyranosyl)-(1→4)-(3-O-benzoyl-6-O-benzyl-2-deoxy-2-tetrachlorophthalimido-β-D-glucopyranoside) (16). A solution of **12** (0.90 g, 0.97 mmol) and bromine (0.1 mL, 1.95 mmol) in CH₂Cl₂ (10 mL) was stirred under argon for 30 min. The mixture was then concentrated and co-concentrated



twice with toluene (Na-dried). A solution of the obtained glycosyl bromide and **14** (0.50 g, 0.55 mmol) in CH_2Cl_2 -MeCN 3:1 (20 mL) was stirred with 3Å MS for 30 min at -40°C before AgOTf (0.51 g, 1.98 mmol) was added. After stirring at -40°C for 1 h the reaction was allowed to warm to -10°C , and after an additional 4 h Et_3N (1 mL) was added. The mixture was stirred for another 15 min then filtered through Celite and concentrated. Flash chromatography (toluene-EtOAc 3:1) of the residue yielded 594 mg of **16** (0.34 mmol, 62%). NMR (CDCl_3): ^{13}C , δ 20.8, 20.9, 21.1 (CH_3 acetyl), 55.1, 55.8, 57.2, 66.1, 67.0, 67.5, 68.2, 69.9, 71.0, 71.4, 71.5, 71.8, 72.5 (2C), 72.7 (2C), 73.3, 73.4, 73.6, 74.0, 74.6, 75.0, 75.9, 76.6, 82.3 (C-2-6, 2'-6', 2''-6'', 2-6''', CH_3O), 99.0, 99.2, 100.6, 103.4 (C-1-1'''), 123.7-138.2 (aromatic C), 162.4, 162.6 (C=O TCP), 165.1, 165.7 (C=O benzoyl), 167.6, 167.8 (C=O NPhth) 169.1, 170.1 (2C) (C=O acetyl). An aliquot of **16** was treated with benzoyl chloride (10 equiv) in pyridine and then acetylated (pyridine-Ac₂O 3:2) to give the corresponding 2'-O-acetylated-4'-O-benzoylated tetrasaccharide with NMR (CDCl_3): ^1H , δ 1.70, 1.84, 1.91, 1.95 (s, 12H, CH_3 acetyl), 2.52-2.62 (m, 2H), 2.79-2.90 (m, 2H), 3.20-3.25, 3.35-3.43 (m, 5H, H-5', 5'', CH_3O), 3.57 (m, 1H, H-5), 3.65-3.78 (m, 7H, H-3'), 3.99-4.25 (m, 8H, H-4, 4'', 2, 1'), 4.37 (dd, 1H, H-2''), 4.46-4.52 (m, 4H, H-1''', CH_2 benzyl), 4.72-4.86 (m, 4H, H-3''', 2', CH_2 benzyl), 4.96 (dd, 1H, H-2'''), 5.20-5.24 (m, 2H, H-4''', 1'' ($J_{1,2}$ 8.24 Hz)), 4.36 (d, 1H, $J_{1,2}$ 8.24 Hz), 5.61 (d, 1H, H-4'), 5.79 (dd, 1H, H-3''), 5.92 (dd, 1H, H-3), 6.86-8.14 (m, 34H, aromatic H). The downfield shift for H-2' (4.72-4.85 ppm) and H-4' (5.61 ppm) in the acylated tetrasaccharide confirmed the (1→3)-linkage.

Methyl (β -D-Galactopyranosyl)-(1→4)-(2-acetamido-2-deoxy- β -D-glucopyranosyl)-(1→3)-(β -D-galactopyranosyl)-(1→4)-2-acetamido-2-deoxy- β -D-glucopyranoside (17**).** Hydrazine hydrate (0.145 mL, 3 mmol) was added to a solution of tetrasaccharide **15** (0.130 g, 0.068 mmol) in MeCN-THF-EtOH (2:1:1, 4 mL). After stirring at 70°C for 18 h, the mixture was concentrated and co-concentrated with EtOH. The residue was purified on a short silica gel column (EtOAc-MeOH-H₂O 7:2:1). The obtained material was dissolved in pyridine (3 mL) and acetic anhydride (1 mL), and the mixture was stirred overnight and concentrated. Flash chromatography (CHCl_3 -MeOH 30:1→10:1) followed by further purification by size exclusion chromatography (LH-20 Sephadex, eluted with MeOH) of the residue gave the fully acetylated tetrasaccharide (63 mg, 0.044 mmol, 65%), which was hydrogenolysed (120 psi) over Pd/C (10%) in a solution of EtOAc-MeOH-H₂O (7:2:1, 3 mL). After 12 h the mixture was filtered through Celite and concentrated. The residue was dissolved in MeOH and a catalytic amount of 1 M NaOMe in MeOH was added. The mixture was stirred for 5 h, then neutralised with Dowex H⁺ ion exchange resin, filtered and concentrated. Size exclusion chromatography (Biogel P2 column, eluted with H₂O containing 1% *n*-BuOH) gave **17** (25 mg, 0.033 mmol, 49% from **15**). $[\alpha]_D -12^\circ$ (*c* 1.1, H₂O), [Lit.³ -10° (*c* 1.0, H₂O)]. NMR (D₂O): ^1H (selected data) δ 2.04, 2.05 (s, 6H, CH_3 acetyl), 3.51 (s, 3H, CH_3O), 4.48 (2d, 3H), 4.71 (2d, 1H). NMR data were in agreement with those reported elsewhere.^{3,6,8}



Methyl (β -D-galactopyranosyl)-(1 \rightarrow 4)-(2-acetamido-2-deoxy- β -D-glucopyranosyl)-(1 \rightarrow 3)-(β -D-galactopyranosyl)-(1 \rightarrow 4)-(2-acetamido-2-deoxy- β -D-glucopyranosyl)-(1 \rightarrow 3)-(β -D-galactopyranosyl)-(1 \rightarrow 4)-2-acetamido-2-deoxy- β -D-glucopyranoside (20**).** Tetrasaccharide **15** (71 mg, 0.037 mmol) was dissolved in 5% HCl/MeOH-CH₂Cl₂ (1:1, 4 mL), and the solution stirred under argon for 37 h, then concentrated and purified by flash chromatography (toluene-EtOAc 2:3) to afford methyl (6-*O*-benzyl- β -D-galactopyranosyl)-(1 \rightarrow 4)-(3-*O*-benzoyl-6-*O*-benzyl-2-deoxy-2-tetrachlorophthalimido- β -D-glucopyranosyl)-(1 \rightarrow 3)-(6-*O*-benzyl- β -D-galactopyranosyl)-(1 \rightarrow 4)-(3-*O*-benzoyl-6-*O*-benzyl-2-deoxy-2-tetrachlorophthalimido- β -D-glucopyranoside) (**18**) (58 mg, 0.033 mmol, 88%). NMR (CDCl₃): ¹³C, δ 55.7 (2C), 57.3, 67.5, 67.7, 67.8, 68.2, 68.3, 68.5, 70.9, 72.2, 72.5, 72.8, 72.9, 73.4, 73.4, 73.7, 73.8, 74.4, 74.6, 76.6, 77.5, 77.7, 82.4 (C-2-6, 2'-6', 2''-6'', 2-6''', CH₃O), 99.0, 99.2, 103.3, 103.8 (C-1-1'''), 127.2-140.3 (aromatic C), 162.7, 163.4 (C=O TCP), 165.7, 165.7 (C=O benzoyl). A mixture of **18** (52 mg, 0.029 mmol), **10** (53 mg, 0.05 mmol), and 3 Å MS in CH₂Cl₂ (2 mL) was treated with methyl triflate (13 μ L, 0.116 mmol) under argon. After stirring for 48 h, Et₃N (0.250 mL) was added and stirring was continued for 15 min. The mixture was then diluted with CH₂Cl₂, filtered through Celite, and the solvent removed under reduced pressure. Flash chromatography (toluene-EtOAc 3:1) of the residue gave methyl (2,3,4-tri-*O*-acetyl-6-*O*-benzyl- β -D-galactopyranosyl)-(1 \rightarrow 4)-(3-*O*-benzoyl-6-*O*-benzyl-2-deoxy-2-tetrachlorophthalimido- β -D-glucopyranosyl)-(1 \rightarrow 3)-(6-*O*-benzyl- β -D-galactopyranosyl)-(1 \rightarrow 4)-(3-*O*-benzoyl-6-*O*-benzyl-2-deoxy-2-tetrachlorophthalimido- β -D-glucopyranosyl)-(1 \rightarrow 3)-(6-*O*-benzyl- β -D-galactopyranosyl)-(1 \rightarrow 4)-(3-*O*-benzoyl-6-*O*-benzyl-2-deoxy-2-tetrachlorophthalimido- β -D-glucopyranoside) (**19**) (49 mg, 0.018 mmol, 61%). NMR (CDCl₃): ¹³C, 20.8, 21.1, 21.0 (CH₃ acetyl), 55.7 (3C), 57.3, 66.0, 67.0, 67.4, 67.6, 68.1, 68.2, 69.8, 71.0, 71.4, 71.5, 71.8, 72.1, 72.3, 72.7, 73.3, 73.4, 73.6, 73.7, 73.9, 74.0, 74.3, 74.4, 74.9, 75.6, 76.3, 77.6, 82.4 (C-2-6, 2'-6', 2''-6'', 2-6''', 2''''-6''', 2''''-6''''', CH₃O, CH₂ benzyl), 99.0, 99.1, 99.3, 100.5, 103.3 (2C) (C-1-1'''''), 125.5-140.3 (aromatic C), 162.7, 163.3 (C=O TCP), 165.3, 165.6, 165.7 (C=O benzoyl), 170.0, 170.1 (C=O acetyl). An aliquot of **19** was acetylated (pyridine-Ac₂O 3:2). ¹H and COSY NMR experiments showed downfield shifts for H-2', 2''' (4.50-4.66 ppm), H-4', 4''' (5.29-5.34 ppm), whereas H-3', 3''' (3.5-3.7 ppm) which proved the (1 \rightarrow 3)-linkage. Hexasaccharide **19** (100 mg, 0.036 mmol) was deprotected as described for compound **15** to give 16 mg (0.013 mmol, 39%) of **20**. [α]_D -8° (c 1.0, H₂O), [Lit.³ -6° (c 1.0, H₂O)]. NMR (D₂O): ¹H, (selected data) δ 2.03 (3s, 9H, CH₃ acetyl), 3.50 (s, 3H, CH₃O), 3.93 (d, 1H), 4.16 (d, 2H), 4.48 (2d, 4H), 4.71 (2d, 2H). NMR data were in agreement with those reported elsewhere.^{3,6,8}

Methyl (2,3,4-Tri-*O*-acetyl-6-*O*-benzyl- β -D-galactopyranosyl)-(1 \rightarrow 4)-(3-*O*-benzoyl-6-*O*-benzyl-2-deoxy-2-tetrachlorophthalimido- β -D-glucopyranosyl)-(1 \rightarrow 3)-(6-*O*-benzyl- β -D-galactopyranosyl)-(1 \rightarrow 4)-(3-*O*-benzoyl-6-*O*-benzyl-2-deoxy-2-tetrachlorophthalimido- β -D-glucopyranosyl)-(1 \rightarrow 3)-(6-*O*-benzyl- β -D-galactopyranosyl)-(1 \rightarrow 4)-(3-*O*-benzoyl-6-*O*-benzyl-2-deoxy-2-tetrachlorophthalimido- β -D-glucopyranosyl)-(1 \rightarrow 3)-*O*-(6-*O*-benzyl- β -D-



galactopyranosyl)-(1→4)-*O*-(3-*O*-benzoyl-6-*O*-benzyl-2-deoxy-2-tetrachlorophthalimido-β-D-glucopyranoside) (**22**). 5% HCl/MeOH (1 mL) was added to a stirred solution of **19** (50 mg, 0.018 mmol) in CH₂Cl₂ (1 mL). After 42 h the mixture was concentrated and the residue subjected to flash chromatography (toluene-EtOAc 1:1) to give **21** (39 mg, 0.015 mmol, 81%). NMR (selected data) (CDCl₃): ¹H δ 3.41 (s, 3H, CH₃O), 5.21, 5.48, 5.58 (d, 3H, H-1, 1'', 1'''), 5.86, 5.99 (dd, 3H, H-3, 3'', 3'''). Compound **21** (39 mg) and **10** (27 mg, 0.026 mmol) was dissolved in CH₂Cl₂ (1 mL) and stirred with 3Å MS for 30 min when methyl triflate (9 μL, 0.075 mmol) was added. After 24 h the reaction was quenched by addition of Et₃N (0.1 mL), and the mixture diluted with CH₂Cl₂, filtered through Celite and concentrated. Flash chromatography (toluene-EtOAc 3:1→2:1) of the crude product gave **22** (31 mg, 0.009 mmol, 60%). Further elution (toluene-EtOAc 1:1) rendered 7 mg of unreacted **19** (0.002 mmol, 13%). **22**: NMR (CDCl₃): ¹³C, 20.9, 21.0, 21.1 (CH₃ acetyl), 55.8 (4C), 57.2, 66.1, 67.0, 67.1, 67.5, 67.6, 67.7, 68.1, 68.3, 69.8, 71.0, 71.3, 71.5, 71.8, 72.0, 72.4, 72.8, 73.4 (2C), 73.6, 73.7, 74.0, 74.4, 74.9, 75.6, 76.3, 76.6, 76.7, 77.5, 82.4 (C-2-6, 2'-6', 2''-6'', 2'''-6''', 2''''-6''''', 2'''''-6'''''), CH₂ benzyl, CH₃O, 99.0 (2C), 99.2 (2C), 100.5, 103.3 (3C) (C-1-1') 128.3-140.1 (aromatic C), 162.7, 163.4 (C=O TCP), 165.3, 165.6 (2C), 165.7 (C=O benzoyl), 169.0, 170.0, 170.1 (C=O acetyl).

An aliquot of **22** was acetylated and used for NMR analysis to prove the selectivity in the coupling. ¹H and COSY NMR experiments showed the characteristic downfield shifts for H-2', 2'', 2''', 4', 4'' and 4''' compared to H-3', 3'', and 3''', and it could be concluded that the coupling was selective for 3''''-OH.

Methyl (6-*O*-benzyl-β-D-galactopyranosyl)-(1→4)-6-*O*-benzyl-2-deoxy-2-tetrachlorophthalimido-β-D-glucopyranoside (24). 120 mg (0.122 mmol) of **4** was treated with dry MeOH (25 μL, 0.61 mmol), NIS (55 mg, 0.244 mmol) and TESOTf (55 μL, 0.244 mmol) in CH₂Cl₂ and then worked up as described for **13**. Flash chromatography (toluene-EtOAc 5:1) gave methyl (2,3,4-tri-*O*-acetyl-6-*O*-benzyl-β-D-galactopyranosyl)-(1→4)-6-*O*-benzyl-2-deoxy-2-tetrachlorophthalimido-β-D-glucopyranoside (**23**) (99 mg, 0.107 mmol, 88%). NMR (CDCl₃): ¹³C δ 20.9, 21.0, 21.1 (CH₃ acetyl), 56.8, 57.0, 67.6, 68.2, 68.3, 69.3, 69.7, 71.2, 72.6, 73.7, 73.9, 74.6, 81.9 (C-2-6, C-2'-6', CH₂ benzyl, CH₃O), 99.1, 101.5 (C-1, C-1'), 125.5-138.2 (aromatic C), 169.3, 170.0, 170.1 (C=O acetyl). Disaccharide **23** (95 mg, 0.102 mmol) was dissolved in 5% HCl/MeOH and stirred at room temperature for 36 h. The mixture was then concentrated and the residue subjected to flash chromatography (toluene-EtOAc 1:2) to give tetraol **24** (79 mg, 0.099 mmol, 97%). NMR (CDCl₃): ¹³C δ 56.8, 56.9, 69.0, 69.2, 69.5, 69.6, 71.5, 73.7, 73.8, 73.9 (2C), 74.5, 82.9 (C-2-6, C-2'-6', CH₃O, CH₂ benzyl), 99.0, 103.9 (C-1, C-1'), 127.5-140.1 (aromatic C), 163.1 (C=O TCP).

Methyl (2,3,4-Tri-*O*-acetyl-6-*O*-benzyl-β-D-galactopyranosyl)-(1→4)-(6-*O*-benzyl-2-deoxy-2-tetrachlorophthalimido-β-D-glucopyranosyl)-(1→3)-(6-*O*-benzyl-β-D-galactopyranosyl)-(1→4)-(6-*O*-benzyl-2-deoxy-2-tetrachlorophthalimido-β-D-glucopyranoside) (25). Methyl triflate (40 μL, 0.35 mmol), was added to a stirred solution of thioglucoside **9** (117 mg, 0.122 mmol),



24 (70 mg, 0.087 mmol) and 3 Å MS in CH₂Cl₂ (3 mL). The reaction mixture was stirred under argon for 24 h, quenched by addition of triethylamine (0.25 mL), filtered through Celite and concentrated. Flash chromatography (toluene-EtOAc 4:1–3:1) gave **25** (68 mg, 40 mmol, 46%). NMR (CDCl₃): ¹³C δ 20.9, 21.0, 21.1, 56.6, 56.7, 57.0, 67.6, 68.2, 68.3, 68.5, 68.8, 69.3, 69.6, 69.7, 69.9, 70.3, 71.2, 72.4, 73.5, 73.6, 73.8, 73.9, 74.0, 74.4, 77.5, 81.9, 83.1, 84.7 (C-2–6, 2'-6', 2''-6'', 2—C-6''', CH₃O, CH₂ benzyl), 99.0, 99.3, 101.4, 104.0 (C-1–1'''), 127.4–140.2 (aromatic C), 163.5, 163.9 (C=O TCP), 169.2, 170.0, 170.1 (C=O acetyl). An aliquot of **25** was acetylated (pyridine:Ac₂O 3:2) to give the corresponding 3,2',4',3''-O-acetylated tetrasaccharide with NMR (CDCl₃): ¹H δ 1.77, 1.82, 1.90, 1.96, 1.99, 2.01, 2.04 (CH₃ acetyl), 3.25 (m, 1H), 3.33–3.58 (m, 15H), 3.62–3.68 (m, 3H, H-2 GlcNTCP, H-4×2 GlcNTCP), 4.16 (dd, 1H, H-2 GlcNTCP), 4.20 (d, 1H, H-1'), 4.34–4.66 (m, 6H, H-1''', 2'), 4.77 (m, 2H), 4.91 (dd, 1H, H-3'''), 5.02 (dd, 1H, H-2'''), 5.11 (d, 1H, H-1 GlcNTCP), 5.15 (d, 1H, H-1 GlcNTCP), 5.41–5.47 (m, 3H, H-3 GlcNTCP, H-4', 4'''), 5.66 (dd, 1H, H-3 GlcNTCP). The downfield shift for H-3, 2', 4' and 3'' confirmed the regioselective coupling.

ACKNOWLEDGMENTS

This is a collaboration with Professor Fred Brewer at the Albert Einstein College of Medicine, New York. Financial support from the Swedish Natural Science Research Council is gratefully acknowledged.

REFERENCES

1. The manuscript is dedicated to Professor Joachim Thiem on the occasion of his 60th birthday.
2. Renkonen, O. Enzymatic glycosylations with glycosyltransferases. In *Carbohydrates in Chemistry and Biology. Part I: Chemistry of Saccharides*; Ernst, B; Hart, G. W.; Sinay, P. (Eds); Wiley-WCH, Weinheim, **2000**; Vol. 2, pp 647–661.
3. Alais, J.; Veyrières, A. Block synthesis of a hexasaccharide hapten of I-blood group antigen. *Tetrahedron Lett.* **1983**, *24*, 5223–5226.
4. Alais, J.; Veyrières, A. Synthesis of an octasaccharide fragment of the poly-lactosamine series by a blockwise approach. *Tetrahedron Lett.* **1987**, *28*, 3345–3348.
5. Alais, J.; Veyrières, A. Synthesis of linear tetrasaccharide, hexasaccharide, and octasaccharide fragments of the I-blood group active poly-(N-acetyl-lactosamine) series-blockwise methods for the synthesis of repetitive oligosaccharide sequences. *Carbohydr. Res.* **1990**, *207*, 11–31.
6. Srivastava, G.; Hindsgaul, O. Synthesis of poly-lactosamine oligomers by disaccharide polymerization. *J. Carbohydr. Chem.* **1991**, *10*, 927–933.
7. Shimizu, H.; Ito, Y.; Kanie, O.; Ogawa, T. Solid phase synthesis of a poly-lactosamine oligosaccharide. *Bioorg. Med. Chem. Lett.* **1996**, *6*, 2841–2846.
8. Nilsson, M.; Norberg, T. Synthesis of carbohydrate derivatives corresponding to a tumor-associated glycolipid: a trimeric Lewis x nonasaccharide and a trimeric N-acetyl lactosamine hexasaccharide. *J. Carbohydr. Chem.* **1989**, *8*, 613–627.



9. Salminen, H.; Ahokas, K.; Niemela, R.; Penttila, L.; Maaheimo, H.; Helin, J.; Costello, C. E.; Renkonen, O. Improved enzymic synthesis of a highly potent oligosaccharide antagonist of L-selectin. *FEBS Lett.* **1997**, *419*, 220–226.
10. Barrondes, S. H.; Cooper, D. N. W.; Gitt, M. A.; Leffler, H. Galectins-Structure and function of a large family of animal lectins. *J. Biol. Chem.* **1994**, *269*, 20807–20810.
11. Perillo, N. L.; Marcus, M. E.; Baum, L.G. Galectins: Versatile modulators of cell adhesion, cell proliferation, and cell death. *J. Mol. Med.* **1998**, *76*, 402–412.
12. Arnarp, J.; Lönngren, J. Synthesis of a tri-saccharide, a penta-saccharide, and a hepta-saccharide containing terminal *N*-acetyl- β -D-lactosaminyl residues, part of the complex-type carbohydrate moiety of glycoproteins. *J. Chem. Soc. Perkin Trans 1* **1981**, 2070–2074.
13. Kaji, E.; Lichtenthaler, F. W. Expedient conversion of lactose into versatile derivatives of lactosamine and β -D-galactosyl-(1 \rightarrow 4)-D-mannosamine. *J. Carbohydr. Chem.* **1995**, *4*, 791–803.
14. Wrodnigg, T. M.; Stütz, A. E. The Heyns rearrangement revisited: An exceptionally simple two-step chemical synthesis of D-lactosamine from lactulose. *Angew. Chem. Int. Ed.* **1999**, *38*, 827–828.
15. Reddy, G. V.; Jain, R. K.; Locke and R. D.; Matta, K. L. Synthesis of precursors for the dimeric 3-*O*-SO₃Na Lewis X and Lewis A structures. *Carbohydr. Res.* **1996**, *280*, 261–276.
16. Numomura, S.; Iida, M.; Numata, M.; Sugimoto, M.; Ogawa, T. Total synthesis of sulfated Le(X) pentasoyl ceramide. *Carbohydr. Res.* **1994**, *263*, C1–C6.
17. Zhang, Y.-M.; Brodsky, A.; Sinaÿ, P.; Saint-Marcoux, G.; Perly, B. Synthesis of a nonasaccharide with two Lewis X trisaccharides anchored onto a branched trimannoside. *Tetrahedron: Asymmetry* **1995**, *6*, 1195–1216.
18. Ellervik, U.; Magnusson, G. A high yielding chemical synthesis of a sialyl Lewis X tetrasaccharide and Lewis X trisaccharide; Example of regio- and stereodifferentiated glycosylations. *J. Org. Chem.* **1998**, *63*, 9314–9322.
19. Debenham, J. S.; Debenham, S. D.; Fraser-Reid, B. *N*-Tetrachlorophthaloyl (TCP) for ready protection/deprotection of amino sugar glycosides. *Bioorg. Med. Chem.* **1996**, *4*, 1909–1918.
20. Busk, T.; Konradsson, P. Synthesis of oligosaccharides designed to form micelles, corresponding to structures found in ovarian cyst fluid. *J. Carbohydr. Chem.* **2000**, *19*, 25–51.
21. Sato, S.; Ito, Y.; Ogawa, T. Synthetic studies on cell-surface glucans 48. Stereocontrolled and regio-controlled, total synthesis of the LeB antigen, III⁴FucIV²FucLcOse₄Cer. *Carbohydr. Res.* **1986**, *155*, C1–C5.
22. Okamoto, K.; Goto, T. Glycosidation of sialic acid. *Tetrahedron* **1990**, *46*, 5835–5837.
23. Byramova, N. E.; Ovchinnikov, M. V.; Backinovskiy, L. V.; Kochetkov, N. K. Selective removal of *O*-acetyl groups in the presence of *O*-benzoyl groups by acid-catalyzed methanolysis. *Carbohydr. Res.* **1983**, *124*, C8–C11.
24. Olsson, L.; Kelberlay, S.; Jia, Z. J.; Fraser-Reid, B. Access to tetrachlorophthalimide-protected ethyl 2-amino-2-deoxy-1-thio- β -D-glucopyranosides. *Carbohydr. Res.* **1998**, *314*, 273–276.

Received March 23, 2001

Accepted July 25, 2001



Request Permission or Order Reprints Instantly!

Interested in copying and sharing this article? In most cases, U.S. Copyright Law requires that you get permission from the article's rightsholder before using copyrighted content.

All information and materials found in this article, including but not limited to text, trademarks, patents, logos, graphics and images (the "Materials"), are the copyrighted works and other forms of intellectual property of Marcel Dekker, Inc., or its licensors. All rights not expressly granted are reserved.

Get permission to lawfully reproduce and distribute the Materials or order reprints quickly and painlessly. Simply click on the "Request Permission/Reprints Here" link below and follow the instructions. Visit the [U.S. Copyright Office](#) for information on Fair Use limitations of U.S. copyright law. Please refer to The Association of American Publishers' (AAP) website for guidelines on [Fair Use in the Classroom](#).

The Materials are for your personal use only and cannot be reformatted, reposted, resold or distributed by electronic means or otherwise without permission from Marcel Dekker, Inc. Marcel Dekker, Inc. grants you the limited right to display the Materials only on your personal computer or personal wireless device, and to copy and download single copies of such Materials provided that any copyright, trademark or other notice appearing on such Materials is also retained by, displayed, copied or downloaded as part of the Materials and is not removed or obscured, and provided you do not edit, modify, alter or enhance the Materials. Please refer to our [Website User Agreement](#) for more details.

[Order now!](#)

Reprints of this article can also be ordered at

<http://www.dekker.com/servlet/product/DOI/101081CAR100108275>