

This article was downloaded by:

On: 23 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Journal of Carbohydrate Chemistry

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713617200>

Subject Index to Volume 22

Online publication date: 12 March 2003

To cite this Article (2003) 'Subject Index to Volume 22', Journal of Carbohydrate Chemistry, 22: 9, 989 – 996

To link to this Article: DOI: 10.1081/CAR-120028479

URL: <http://dx.doi.org/10.1081/CAR-120028479>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

Subject Index to Volume 22

A

- Acetalization, non-classical, epimerization by, 801–825
- N*-Acetyl- α -D-glucosamine 1-phosphate, synthesis of *C*-glycosyl phosphate and phosphonate analogues of, 645–656
- Acyl group migrations, in transesterification of sucrose in organic medium, 657–669
- Alkylation reactions, of 1,2-*O*-stannylene acetals of D-arabinofuranose, regioselectivity in, 963–981
- Allolactose, synthesis from amygdalin, 267–274
- Allyl glycoside, synthesis of the blocked pentasaccharide derivative related to the repeating unit of the O-antigen from *Shigella dysenteriae* type 3 in the form of, 285–296
- Aminolysis, of D-glucaric acid esters, mechanisms for the formation of diamides and polyamides by, 903–918
- Aminopyranosides, unsaturated, synthesis as transition state mimics for glycosidases, 705–717
- Amygdalin, synthesis of allolactose from, 267–274
- Angucyclines, total synthesis of, 579–592
- α -1,2-Anhydrosugars, to *C*-glycosides, influence of Lewis acids and nucleophiles on stereochemistry of, 549–564
- Antibiotic 100-1, synthesis of, 579–592
- Anti-Gal antibodies, frontal affinity chromatography coupled to mass spectrometry for K_d determination and screening of α -Gal derivatives binding to, 347–376
- D-Arabinofuranose, regioselectivity in alkylation reactions of 1,2-*O*-stannylene acetals of, 963–981
- D-Arabinofuranosyl side-chains, synthesis of laminarin oligosaccharide derivatives with, 275–283
- 4-Aryl-1-thioxo[1,2,4]triazolo[4,3-*a*]quinazolin-5(4H)-one, glycosylation of, 79–94
- Asymmetric dihydroxylation, sharpless, facial selectivity for styrene derivatives, 25–34
- Asymmetric total synthesis, of Brousso-netine C, 719–732

B

- Benzothiazole sulfonyl ethyl *C*-glycosides, radical mediated diastereoselective synthesis of, 423–431

989

DOI: 10.1081/CAR-120028479
Copyright © 2003 by Marcel Dekker, Inc.

0732-8303 (Print); 1532-2327 (Online)
www.dekker.com



- 1,2-*O*-Benzylidene hexopyranoses, synthesis of, 143–147
- Branched cyclomalto-oligosaccharides, syntheses using the trichloroacetimidate method, 297–308
- Broussonetine C, asymmetric total synthesis of, 719–732
- γ -Butyrolactones, [3,3,0]-fused, stereoselective synthesis of, 501–511

C

- Capsular polysaccharide
of *Cryptococcus neoformans*, synthesis of, 565–577
linear, solution conformation from a soybean-nodulating bacterium, 765–779
- Carbocyclization, of C-4 branched deoxy-sugars for synthesis of chiral cyclopentenes, 377–383
- 3-*C*-Carboxymethylgalactosyl lipid, synthesis and selectin-blocking activity of, 513–520
- Cell wall rhamnogalactomannan, from *Cubonia bulbifera*, structure of, 603–611
- Cellulose, crosslinking by olefin metathesis, 47–55
- Chiral cyclopentenes, synthesis via carbocyclization of C-4 branched deoxysugars, 377–383
- β -1-*N*-Chloroacetamido derivatives, of D-glucose and D-galactose, crystal structures of, 891–901
- Combinatorial chemistry
on solid phase, D-glucose as a regioselectively addressable scaffold for, 57–71
synthesis of D-glucosamine derived 3,4-epoxides as potential scaffolds for, 179–184
- Conformational analysis, of C-trehaloses using molecular mechanics calculations, 407–421
- Crosslinking, of cellulose by olefin metathesis, 47–55

- Cryptococcus neoformans* capsular polysaccharide structures, synthesis of, 565–577
- Crystal structures
of β -1-*N*-chloroacetamido derivatives of D-glucose and D-galactose, 891–901
of methyl α -D-mannofuranoside, 593–601
- Cubonia bulbifera*, structure of a cell wall rhamnogalactomannan from, 603–611
- Cyclomalto-oligosaccharides, branched, syntheses using the trichloroacetimidate method, 297–308

D

- Diamides, mechanisms for formation by aminolysis of D-glucaric acid esters, 903–918
- Diamido-di-D-glucaric acid monomers, symmetrical, alternating stereoregular *head, tail–tail, head*-poly(alkylene D-glucarimides) derived from a homologous series of, 123–142
- (*E*)-1,2-Dideoxy-1-dimethylamino-4,5:6,7-di-*O*-isopropylidene-D-arabino-hept-1-en-3-ulose, for synthesis of acyclic C-nucleoside analogues, 537–548
- Difructose dianhydride, structure of, 1–8
- N,N*-Dimethylthioformamide, synthesis of tetra-*O*-acetyl-1-thio- α -D-glucopyranose by reaction with tetra-*O*-acetyl- α -D-glucopyranosyl bromide, 73–78
- Diphenylphosphenium cation, reactions with glycals for regioselective synthesis of phosphorylated sugars, 443–457
- 1,3-Dipolar cycloaddition, of nitrones to 2,3-unsaturated sugar 1,5-lactones, double asymmetric induction in, 613–629



Disaccharides, β -(1 \rightarrow 3)-linked, MM3 force-field calculation of the 2D- and 3D- potential energy surfaces of, 217–239

Double asymmetric induction, in 1,3-dipolar cycloaddition of nitrones to 2,3-unsaturated sugar 1,5-lactones, 613–629

E

Epi-2-deoxystreptamine and analogs, protected, synthesis of, 733–742

Epimerization, by non-classical acetalization, 801–825

3,4-Epoxides, D-glucosamine derived, synthesis as potential scaffolds for combinatorial chemistry, 179–184

D-Erythritols, enantiomerically pure alkylated, synthesis from D-xylose, 35–46

Escherichia coli type O26:H, synthesis of the trisaccharide repeating unit of the O-antigen related to, 95–107

F

Facial selectivity, of the sharpless asymmetric dihydroxylation of styrene derivatives, 25–34

Frontal affinity chromatography, coupled to mass spectrometry, 347–376

Fucoidan fragments, synthesis, NMR, and conformational studies of, 109–122

α -L-Fucosidase, from *Alcaligenes* sp. for glycosidase-catalyzed synthesis of fucosyl di- and trisaccharide derivatives, 309–316

Fucosyl di- and trisaccharide derivatives, glycosidase-catalyzed synthesis using α -L-fucosidase from *Alcaligenes* sp., 309–316

(1 \rightarrow 3)-Fucotrioxide fragments, linear 4,4',4''-tri-O-sulfated and parent non-sulfated, synthesis, NMR,

and conformational studies of, 109–122

[3,3,0]-Fused γ -butyrolactones, stereoselective synthesis of, 501–511

G

D-Galactose, conversion to D-threo-hexos-2,3-diulose by fungal pyranose oxidase, 207–216

D-Galactose, crystal structures of β -1-N-chloroacetamido derivatives of, 891–901

Ganglioside analogs, lacto- and neolactoseries, containing N-glycolylneuraminic acid, 919–937

Ganglioside GM3, neoglycolipid analogues of the oligosaccharide portion of, 395–406

D-Glucaric acid esters, mechanisms for the formation of diamides and polyamides by aminolysis of, 903–918

D-Glucosamine derivatives

- 3,4-epoxides, synthesis as potential scaffolds for combinatorial chemistry, 179–184
- O-glycosylated, synthesis by Heyns rearrangement of the corresponding O-glycosyl fructoses, 253–265

D-Glucose

- crystal structures of β -1-N-chloroacetamido derivatives of, 891–901
- micellar effects on oxidation by chromic acid in perchloric acid medium, 843–866
- as a regioselectively addressable scaffold for combinatorial chemistry on solid phase, 57–71

β -D-Glucose oligosaccharides, 3,6-branched, synthesis of, 881–890

Glucuronate glycosyl donors, comparison of, 939–962

Glycals, reactions with diphenylphosphonium cation for regioselective synthesis of phosphorylated sugars, 443–457



- N*-Glycolylneuraminic acid, synthesis of lacto- and neolacto-series ganglioside analogs containing, 919–937
- Glycosidase-catalyzed synthesis, of fucosyl di- and trisaccharide derivatives using α -L-fucosidase from *Alcaligenes* sp., 309–316
- Glycosidases, synthesis of unsaturated aminopyranosides as transition state mimics for, 705–717
- N*-Glycoside analogs, synthesis via thionolactones, 743–751
- C-Glycosides
- benzothiazole sulfonyl ethyl, radical mediated diastereoselective synthesis of, 423–431
 - influence of Lewis acids and nucleophiles on stereochemistry of, 549–564
 - of methyl α -D-altropyranosyl-(1→4)- α -D-glucopyranoside, 671–683
- Glycosyl donors
- glucuronate, comparison of, 939–962
 - silver triflate as a mild alternative catalyst for glycosylation using trichloroacetimidates as, 385–393
- O*-Glycosyl fructoses, Heyns rearrangement for synthesis of *O*-glycosylated D-glucosamine derivatives, 253–265
- C-Glycosyl phosphate and phosphonate, synthesis of, 645–656
- Glycosylations
- of 4-aryl-1-thioxo[1,2,4]triazolo[4,3-a]quinazolin-5(4H)-one, 79–94
 - multigram scale, of scented alcohols employing phase-transfer reactions, 9–23
 - using trichloroacetimidates as glycosyl donors, silver triflate as a mild alternative catalyst for, 385–393
- H
- Heterobifunctional candidate ligands, of P-selectin containing both sulfated

- Lewis X trisaccharide and various sulfated peptides, 481–500
- Hexa-*O*-acetyl- α -D-fructofuranose- β -D-fructofuranose 1,2':2,6'-dianhydride, structure of, 1–8
- D*-threo-Hexos-2,3-diulose, fungal pyranose oxidase conversion of D-galactose to, 207–216
- Heyns rearrangement, of the corresponding *O*-glycosyl fructoses for synthesis of *O*-glycosylated D-glucosamine derivatives, 253–265
- Highly deoxygenated sugars, synthesis of chiral cyclopentenones via carbocyclization of C-4 branched deoxy-sugars, 377–383

I

- Influenza A viruses, synthesis of lacto- and neolacto-series ganglioside analogs containing *N*-glycolylneuraminic acid as probes for investigation of specific receptor structures recognized by, 919–937
- Iso-*C*-nucleoside analogues, synthesis of (methyl 3-*O*-benzyl-4,6-*O*-benzylidene-2-deoxy- α -D-altropyranosid-2-yl)thiophene derivatives as precursors of, 171–178

L

- Laminarin oligosaccharide derivatives, with D-arabinofuranosyl side-chains, 275–283
- Lewis acids, influence on stereochemistry of α -1,2-anhydrosugars to C-glycosides, 549–564
- Linkers, commercially available, comparison for mannose-BSA conjugates in reactivity and bioactivity, 317–329
- Lipase mutants, site directed, application on regioselective acylation of monosaccharides, 631–644



- Lipoarabinomannan, mannosylated and appropriately functionalized for neoglycoconjugate preparation, synthesis of oligosaccharide fragments of, 149–170, 459–480
- Liquid crystals, enantiopure trioxadecalin, influence of the nature of the phenyl substituent on the mesogenic properties of, 685–703

M

- Macrocyclic derivatives, containing a sucrose unit, 753–763
- D-Mannitol, selective silylation assisted by phenylboronic acid, 867–879
- Mannose-BSA conjugates, comparison between commercially available linkers in reactivity and bioactivity, 317–329
- Mass spectrometry, coupled to frontal affinity chromatography, 347–376
- Metathesis, of olefin, crosslinking of cellulose by, 47–55
- 1,2-*O*-*p*-Methoxybenzylidene hexopyranoses, synthesis of, 143–147
- Methyl α -D-altropyranosyl-(1 \rightarrow 4)- α -D-glucopyranoside, synthesis of the C-glycoside of, 671–683
- (Methyl 3-*O*-benzyl-4,6-*O*-benzylidene-2-deoxy- α -D-altropyranosid-2-yl)thiophene derivatives, synthesis as precursors of iso-*C*-nucleoside analogues, 171–178
- Methyl α - and β -D-glucopyranosides, synthesis of nitrogen heterocycles from, 433–442
- (4-*O*-Methyl-D-glucurono)-D-xylan, isolation and characterization from the skin of *Opuntia ficus-indica* prickly pear fruits, 331–337
- Methyl α -D-mannofuranoside, crystal structure and solid-state ^{13}C NMR of, 593–601
- Methyl (*R*)-pyruvate ester, synthesis of a blocked tetrasaccharide related to the repeating unit of the antigen from *Shigella dysenteriae* type 9 in the form of, 521–535
- Micellar effects, on the oxidation of D-glucose by chromic acid in perchloric acid medium, 843–866
- MM3 force-field, for calculation of the 2D- and 3D-potential energy surfaces of β -(1 \rightarrow 3)-linked disaccharides, 217–239
- Molecular mechanics calculations, for conformational analysis of *C*-trehaloses, 407–421
- Monosaccharides, application of site-directed lipase mutants on regioselective acylation of, 631–644
- Multigram scale glycosylations, of scented alcohols employing phase-transfer reactions, 9–23

N

- Neoglycoconjugate, synthesis of oligosaccharide fragments of mannosylated lipoarabinomannan appropriately functionalized for preparation of, 149–170, 459–480
- Neoglycolipid analogues, of the oligosaccharide portion of ganglioside GM3, 395–406
- Nitrogen heterocycles, synthesis from methyl α - and β -D-glucopyranosides, 433–442
- Nitrones, double asymmetric induction in 1,3-dipolar cycloaddition to 2,3-unsaturated sugar 1,5-lactones, 613–629
- Nucleophiles
influence on stereochemistry of α -1,2-anhydrosugars to C-glycosides, 549–564
reactions with reactive intermediates in the 3,4,6-tri-*O*-benzyl-D-glucal-TfOH-*n*-Bu₄NI reaction system, 827–841



C-Nucleoside analogues, acyclic, synthesis using (*E*)-1,2-dideoxy-1-dimethylamino-4,5:6,7-di-*O*-isopropylidene-*D*-arabino-hept-1-en-3-ulose, 537–548

O

O-antigen

related to the enterohemorrhagic *Escherichia coli* type O26:H, synthesis of the trisaccharide repeating unit of, 95–107

from *Shigella dysenteriae* type 3, synthesis of the blocked pentasaccharide derivative related to the repeating unit in the form of its allyl glycoside, 285–296

Olefin metathesis, crosslinking of cellulose by, 47–55

Oligosaccharides

3,6-branched β -*D*-glucose, synthesis of, 881–890

fragments, of mannosylated lipoarabinomannan appropriately functionalized for neoglycoconjugate preparation, 149–170, 459–480

of ganglioside GM3, neoglycolipid analogues of, 395–406

identification of individual sugar components with 2D selective-TOCSY-DQFCOSY, 339–345

laminarin derivatives, with *D*-arabino-furanosyl side-chains, 275–283

Opuntia ficus-indica prickly pear fruits, isolation and characterization of (4-*O*-methyl-*D*-glucurono)-*D*-xy-lan from the skin of, 331–337

Oxidation, of *D*-glucose by chromic acid in perchloric acid medium, micellar effects on, 843–866

P

Pentasaccharide derivative, blocked, related to the repeating unit of the

O-antigen from *Shigella dysenteriae* type 3 in the form of its allyl glycoside, 285–296

Phase-transfer glycosylations, of scented alcohols, 9–23

Phenyl substituent, influence of the nature on the mesogenic properties of enantiopure trioxadecalin derived liquid crystals, 685–703

Phenylboronic acid, selective silylation of *D*-mannitol assisted by, 867–879

Phosphonylated sugars, regioselective synthesis from reactions of glycols with diphenylphosphonium cation, 443–457

Piperidines, highly functionalized, synthesis via a tandem retro-Michael-[2 + 3]-cycloaddition, 241–252

Poly(alkylene *D*-glucarimides), alternating stereoregular *head, tail-tail, head*, derived from a homologous series of symmetrical diamido-*D*-glucaric acid monomers, 123–142

Polyamides, mechanisms for formation by aminolysis of *D*-glucaric acid esters, 903–918

Prickly pear fruits, isolation and characterization of (4-*O*-methyl-*D*-glucurono)-*D*-xy-lan from the skin of, 331–337

Pyranose oxidase, fungal, for conversion of *D*-galactose to *D*-*threo*-hexos-2,3-diulose, 207–216

R

Radical mediated diastereoselective synthesis, of benzothiazole sulfonyl ethyl C-glycosides, 423–431

Reactive intermediates, in the 3,4,6-tri-*O*-benzyl-*D*-glucal-TfOH-*n*-Bu₄NI reaction system, reactions of nucleophiles with, 827–841

Regioselective acylation, of monosaccharides, application of site-directed lipase mutants on, 631–644

Rhamnogalactomannan, from *Cubonia bulbifera*, structure of, 603–611

S

Scaffolds, for combinatorial chemistry
on solid phase, D-glucose as, 57–71
synthesis of D-glucosamine derived
3,4-epoxides as, 179–184

Scented alcohols, phase-transfer glycosylations of, 9–23

P-Selectin, containing both sulfated Lewis X trisaccharide and various sulfated peptides, syntheses of heterobifunctional candidate ligands of, 481–500

Selectin-blocking activity, of an analog of sulfate, 513–520

Selective silylation, of D-mannitol assisted by phenylboronic acid, 867–879

Selective-TOCSY-DQF COSY, 2D, for identification of individual sugar components in oligosaccharides, 339–345

Shigella dysenteriae

type 3, synthesis of the blocked pentasaccharide derivative related to the repeating unit of the O-antigen in the form of allyl glycoside, 285–296

type 9, synthesis of a blocked tetrasaccharide related to the repeating unit of antigen from in the form of its methyl (*R*)-pyruvate ester and 2-(trimethylsilyl)ethyl glycoside, 521–535

Silver triflate, as a mild alternative catalyst for glycosylation conditions using trichloroacetimidates as glycosyl donors, 385–393

Site-directed lipase mutants, application on regioselective acylation of monosaccharides, 631–644

Solid-phase synthesis, of 1-thio-β-D-GlcNAc carbohydrate mimetic library, 781–800

Solid-state ¹³C NMR, of methyl α-D-mannofuranoside, 593–601

Solution conformation, of linear capsular polysaccharide from a soybean-nodulating bacterium, 765–779

Soybean-nodulating bacterium, solution conformation of linear capsular polysaccharide from, 765–779

1,2-*O*-Stannylene acetals, of D-arabinofuranose, regioselectivity in alkylation reactions of, 963–981

Stereoregular *head, tail–tail, head*-poly(alkylene D-glucarimides), derived from a homologous series of symmetrical diamido-di-D-glucaric acid monomers, 123–142

Stereoselective synthesis, of [3,3,0]-fused γ-butyrolactones, 501–511

Styrene derivatives, facial selectivity of the sharpless asymmetric dihydroxylation of, 25–34

Sucrose, transesterification in organic medium, 657–669

Sucrose unit, synthesis of macrocyclic derivatives containing, 753–763

Sugar 1,5-lactones, 2,3-unsaturated, double asymmetric induction in 1,3-dipolar cycloaddition of nitrones to, 613–629

Sulfated Lewis X trisaccharide, syntheses of heterobifunctional candidate ligands of P-selectin containing, 481–500

Sulfated peptides, syntheses of heterobifunctional candidate ligands of P-selectin containing, 481–500

T

Tandem retro-Michael-[2 + 3]-cycloaddition, for synthesis of highly functionalized piperidines, 241–252

Taste properties, of 4,1',4',6'-tetrahalodeoxysucrose analogues, 185–206

Tetra-*O*-acetyl-α-D-glucopyranosyl bromide, synthesis of tetra-*O*-acetyl-



- 1-thio- α -D-glucopyranose by reaction with *N,N*-dimethylthioformamide, 73–78
- Tetra-*O*-acetyl-1-thio- α -D-glucopyranose, synthesis by reaction of tetra-*O*-acetyl- α -D-glucopyranosyl bromide with *N,N*-dimethylthioformamide, 73–78
- 4,1',4',6'-Tetrahalodeoxysucrose analogues, synthesis and taste properties of, 185–206
- Tetrasaccharide, blocked, related to the repeating unit of the antigen from *Shigella dysenteriae* type 9 in the form of its methyl (*R*)-pyruvate ester and 2-(trimethylsilyl)ethyl glycoside, 521–535
- 1-Thio- β -D-GlcNAc carbohydrate mimetic library, solid-phase synthesis of, 781–800
- Thionolactones, synthesis of *N*-glycoside analogs via, 743–751
- D-Threitols, enantiomerically pure alkylated, synthesis from D-xylose, 35–46
- Transesterification, of sucrose in organic medium, 657–669
- C-Trehaloses, conformational analysis using molecular mechanics calculations, 407–421
- 3,4,6-Tri-*O*-benzyl-D-glucal-TfOH-*n*-Bu₄NI reaction system, reactions of nucleophiles with reactive intermediates in, 827–841
- Trichloroacetimidate method, for syntheses of branched cyclomalto-oligosaccharides, 297–308
- Trichloroacetimidates, as glycosyl donors, silver triflate as a mild alternative catalyst for glycosylation using, 385–393
- 2-(Trimethylsilyl)ethyl glycoside, synthesis of a blocked tetrasaccharide related to the repeating unit of the antigen from *Shigella dysenteriae* type 9 in the form of, 521–535
- Trioxadecalin derived liquid crystals, enantiopure, influence of the nature of the phenyl substituent on the mesogenic properties of, 685–703
- Trisaccharide repeating unit synthesis, of the O-antigen related to the enterohemorrhagic *Escherichia coli* type O26:H, 95–107

U

- Unsaturated aminopyranosides, synthesis as transition state mimics for glycosidases, 705–717

X

- D-Xylose, synthesis of enantiomerically pure alkylated D-erythritols and D-threitols from, 35–46

